

19 Continuity

20 Continuity II

21 Continuity III

Continuous Functions



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I


Instructor: David Earn

Lecture 18
Continuity
Monday 25 February 2019

Announcements

- A preliminary version of [Assignment 4](#) has been posted on the course web site. More problems will be added soon.
Due Friday 8 March 2019 at 1:25pm via [crowdmark](#).
BUT you should do **questions 1 and 2** before Test #1.
- **Math 3A03 Test #1**
Monday 4 March 2019 at 7:00pm in MDCL 1110
(room is booked for 90 minutes; you should not feel rushed)

Limits of functions



Limits of functions

Definition (Limit of a function on an interval (a, b))

Let $a < x_0 < b$ and $f : (a, b) \rightarrow \mathbb{R}$. Then f is said to **approach the limit L as x approaches x_0** , often written “ $f(x) \rightarrow L$ as $x \rightarrow x_0$ ” or

$$\lim_{x \rightarrow x_0} f(x) = L,$$

iff for all $\varepsilon > 0$ there exists $\delta > 0$ such that if $0 < |x - x_0| < \delta$ then $|f(x) - L| < \varepsilon$.

Shorthand version:

$$\forall \varepsilon > 0 \exists \delta > 0 \text{ } 0 < |x - x_0| < \delta \implies |f(x) - L| < \varepsilon.$$

Limits of functions

The function f need not be defined on an entire interval. It is enough for f to be defined on a set with at least one accumulation point.

Definition (Limit of a function with domain $E \subseteq \mathbb{R}$)

Let $E \subseteq \mathbb{R}$ and $f : E \rightarrow \mathbb{R}$. Suppose x_0 is a point of accumulation of E . Then f is said to **approach the limit L as x approaches x_0** , i.e.,

$$\lim_{x \rightarrow x_0} f(x) = L,$$

iff for all $\varepsilon > 0$ there exists $\delta > 0$ such that if $x \in E$, $x \neq x_0$, and $|x - x_0| < \delta$ then $|f(x) - L| < \varepsilon$.

Shorthand version:

$$\forall \varepsilon > 0 \exists \delta > 0 \text{ } \vdash \left(x \in E \wedge 0 < |x - x_0| < \delta \right) \implies |f(x) - L| < \varepsilon.$$

Limits of functions

Example

Prove directly from the [definition of a limit](#) that

$$\lim_{x \rightarrow 3} (2x + 1) = 7.$$

(solution on board)

Proof that $2x + 1 \rightarrow 7$ as $x \rightarrow 3$.

We must show that $\forall \varepsilon > 0 \exists \delta > 0$ such that $0 < |x - 3| < \delta \implies |(2x + 1) - 7| < \varepsilon$. Given ε , to determine how to choose δ , note that

$$|(2x + 1) - 7| < \varepsilon \iff |2x - 6| < \varepsilon \iff 2|x - 3| < \varepsilon \iff |x - 3| < \frac{\varepsilon}{2}$$

Therefore, given $\varepsilon > 0$, let $\delta = \frac{\varepsilon}{2}$. Then $|x - 3| < \delta \implies |(2x + 1) - 7| = |2x - 6| = 2|x - 3| < 2\frac{\varepsilon}{2} = \varepsilon$, as required. □

Limits of functions

Example

Prove directly from the [definition of a limit](#) that

$$\lim_{x \rightarrow 2} x^2 = 4.$$

(solution on board)

(and on next slide)

Limits of functions

Proof that $x^2 \rightarrow 4$ as $x \rightarrow 2$.

We must show that $\forall \varepsilon > 0 \exists \delta > 0$ such that $0 < |x - 2| < \delta \implies |x^2 - 4| < \varepsilon$. Given ε , to determine how to choose δ , note that

$$|x^2 - 4| < \varepsilon \iff |(x - 2)(x + 2)| < \varepsilon \iff |x - 2| |x + 2| < \varepsilon.$$

We can make $|x - 2|$ as small as we like by choosing δ sufficiently small. Moreover, if x is close to 2 then $x + 2$ will be close to 4, so we should be able to ensure that $|x + 2| < 5$. To see how, note that

$$\begin{aligned} |x + 2| < 5 &\iff -5 < x + 2 < 5 \iff -9 < x - 2 < 1 \\ &\iff -1 < x - 2 < 1 \iff |x - 2| < 1. \end{aligned}$$

Therefore, given $\varepsilon > 0$, let $\delta = \min(1, \frac{\varepsilon}{5})$. Then

$$|x^2 - 4| = |(x - 2)(x + 2)| = |x - 2| |x + 2| < \frac{\varepsilon}{5} 5 = \varepsilon. \quad \square$$



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 19
Continuity II
Wednesday 27 February 2019

Announcements

- A preliminary version of [Assignment 4](#) has been posted on the course web site. More problems will be added soon.
Due Friday 8 March 2019 at 1:25pm via [crowdmark](#).
BUT you should do **questions 1 and 2** before Test #1.
- **Math 3A03 Test #1**
Monday 4 March 2019 at 7:00pm in MDCL 1110
(room is booked for 90 minutes; you should not feel rushed)
 - Test will cover everything up to the end of the topology section.
- Niky Hristov will hold extra office hours this Friday 1 March 2019, 11:30am–12:30pm and immediately before class on the day of the test, *i.e.*, Monday 4 March 2019, 10:30–11:30am.
- Solutions to $\lim_{x \rightarrow 3} (2x + 1) = 7$ and $\lim_{x \rightarrow 2} x^2 = 4$ are now in the slides for the previous lecture.

Limits of functions

Rather than the ε - δ definition, we can exploit our experience with sequences to define “ $f(x) \rightarrow L$ as $x \rightarrow x_0$ ”.

Definition (Limit of a function via sequences)

Let $E \subseteq \mathbb{R}$ and $f : E \rightarrow \mathbb{R}$. Suppose x_0 is a point of accumulation of E . Then

$$\lim_{x \rightarrow x_0} f(x) = L$$

iff for every sequence $\{e_n\}$ of points in $E \setminus \{x_0\}$,

$$\lim_{n \rightarrow \infty} e_n = x_0 \implies \lim_{n \rightarrow \infty} f(e_n) = L.$$

Limits of functions

Lemma (Equivalence of limit definitions)

The ε - δ definition of limits and the sequence definition of limits are equivalent.

(solution on board)

Note: The definition of a limit via sequences is sometimes easier to use than the ε - δ definition.

Proof of Equivalence of ε - δ definition and sequence definition of limit.

Proof (ε - $\delta \implies$ seq).

Suppose the ε - δ definition holds and $\{e_n\}$ is a sequence in $E \setminus \{x_0\}$ that converges to x_0 . Given $\varepsilon > 0$, there exists $\delta > 0$ such that if $0 < |x - x_0| < \delta$ then $|f(x) - L| < \varepsilon$. But since $e_n \rightarrow x_0$, given $\delta > 0$, there exists $N \in \mathbb{N}$ such that, for all $n \geq N$, $|e_n - x_0| < \delta$. This means that if $n \geq N$ then $x = e_n$ satisfies $0 < |x - x_0| < \delta$, implying that we can put $x = e_n$ in the statement $|f(x) - L| < \varepsilon$. Hence, for all $n \geq N$, $|f(e_n) - L| < \varepsilon$. Thus,

$$e_n \rightarrow x_0 \implies f(e_n) \rightarrow L,$$

as required. □


Proof of Equivalence of ε - δ definition and sequence definition of limit.

Proof ($\text{seq} \implies \varepsilon$ - δ) via contrapositive.

Suppose that as $x \rightarrow x_0$, $f(x) \not\rightarrow L$ according to the ε - δ definition. We must show that $f(x) \not\rightarrow L$ according to the **sequence definition**.

Since the ε - δ **criterion** does not hold, $\exists \varepsilon > 0$ such that $\forall \delta > 0$ there is some $x_\delta \in E$ for which $0 < |x_\delta - x_0| < \delta$ and yet $|f(x_\delta) - L| \geq \varepsilon$. This is true, in particular, for $\delta = 1/n$, where n is any natural number. Thus, $\exists \varepsilon > 0$ such that: $\forall n \in \mathbb{N}$, there exists $x_n \in E$ such that $0 < |x_n - x_0| < 1/n$ and yet $|f(x_n) - L| \geq \varepsilon$. This demonstrates that there is a sequence $\{x_n\}$ in $E \setminus \{x_0\}$ for which $x_n \rightarrow x_0$ and yet $f(x_n) \not\rightarrow L$. Hence, $f(x) \not\rightarrow L$ as $x \rightarrow x_0$ according to the **sequence criterion**, as required. \square

One-sided limits



One-sided limits

Definition (Right-Hand Limit)

Let $f : E \rightarrow \mathbb{R}$ be a function with domain E and suppose that x_0 is a point of accumulation of $E \cap (x_0, \infty)$. Then we write

$$\lim_{x \rightarrow x_0^+} f(x) = L$$

if for every $\varepsilon > 0$ there is a $\delta > 0$ so that

$$|f(x) - L| < \varepsilon$$

whenever $x_0 < x < x_0 + \delta$ and $x \in E$.

One-sided limits

One-sided limits can also be expressed in terms of sequence convergence.

Definition (Right-Hand Limit – sequence version)

Let $f : E \rightarrow \mathbb{R}$ be a function with domain E and suppose that x_0 is a point of accumulation of $E \cap (x_0, \infty)$. Then we write

$$\lim_{x \rightarrow x_0^+} f(x) = L$$

if for every decreasing sequence $\{e_n\}$ of points of E with $e_n > x_0$ and $e_n \rightarrow x_0$ as $n \rightarrow \infty$,

$$\lim_{n \rightarrow \infty} f(e_n) = L.$$

Infinite limits

Definition (Right-Hand Infinite Limit)

Let $f : E \rightarrow \mathbb{R}$ be a function with domain E and suppose that x_0 is a point of accumulation of $E \cap (x_0, \infty)$. Then we write

$$\lim_{x \rightarrow x_0^+} f(x) = \infty$$

if for every $M > 0$ there is a $\delta > 0$ such that $f(x) \geq M$ whenever $x_0 < x < x_0 + \delta$ and $x \in E$.

Properties of limits

There are theorems for limits of functions of a real variable that correspond (and have similar proofs) to the various results we proved for limits of sequences:

- Uniqueness of limits
- Algebra of limits
- Order properties of limits
- Limits of absolute values
- Limits of Max/Min

See Chapter 5 of textbook for details.



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 20
Continuity III
Friday 1 March 2019

Announcements

- All of [Assignment 4](#) has now been posted on the course web site. Due Friday 8 March 2019 at 1:25pm via [crowdmark](#). BUT you should do **questions 1 and 2** before Test #1.
- **Math 3A03 Test #1**
Monday 4 March 2019 at 7:00pm in MDCL 1110
(room is booked for 90 minutes; you should not feel rushed)
 - Test will cover everything up to the end of the topology section.
- Niky Hristov will hold an extra office hour immediately before class on Monday, the day of the test, *i.e.*, Monday 4 March 2019, **10:30–11:30am**.
- I will also hold my usual office hour on Monday, **1:30–2:30pm**.

SCIENCE



Faculty of Science

Graduate Studies Open House

Get a head start on grad school & learn about everything from **#Gradlife** to crafting the perfect application

Tuesday, March 12th 2019
CIBC Hall
5:00PM - 6:30PM

Contact: trepanr@mcmaster.ca

Last time...

- Equivalence of ε - δ definition and sequence definition of limit.
- One-sided limit from the right.

Limits of compositions of functions

When is $\lim_{x \rightarrow x_0} g(f(x)) = g\left(\lim_{x \rightarrow x_0} f(x)\right)$?

Theorem (Limit of composition)

Suppose

$$\lim_{x \rightarrow x_0} f(x) = L.$$

If g is a function defined in a neighborhood of the point L and

$$\lim_{z \rightarrow L} g(z) = g(L)$$

then

$$\lim_{x \rightarrow x_0} g(f(x)) = g\left(\lim_{x \rightarrow x_0} f(x)\right) = g(L).$$

(Textbook (TBB) §5.2.5)

Limits of compositions of functions – more generally

Note: It is a little more complicated to generalize the statement of this theorem so as to minimize the set on which g must be defined but the proof is no more difficult.

Theorem (Limit of composition)

Let $A, B \subseteq \mathbb{R}$, $f : A \rightarrow \mathbb{R}$, $f(A) \subseteq B$, and $g : B \rightarrow \mathbb{R}$. Suppose x_0 is an accumulation point of A and

$$\lim_{x \rightarrow x_0} f(x) = L.$$

Suppose further that g is defined at L . If L is an accumulation point of B and


$$\lim_{z \rightarrow L} g(z) = g(L),$$

or $\exists \delta > 0$ such that $f(x) = L$ for all $x \in (x_0 - \delta, x_0 + \delta) \cap A$, then

$$\lim_{x \rightarrow x_0} g(f(x)) = g\left(\lim_{x \rightarrow x_0} f(x)\right) = g(L).$$


Continuity

Intuitively, a function f is **continuous** if you can draw its graph without lifting your pencil from the paper...



Continuity

and **discontinuous** otherwise. . .



Continuity

In order to develop a rigorous foundation for the theory of functions, we need to be more precise about what we mean by “continuous”.

The main challenge is to define “continuity” in a way that works consistently on sets other than intervals (and generalizes to spaces that are more abstract than \mathbb{R}).

We will define:

- continuity at a single point;
- continuity on an open interval;
- continuity on a closed interval;
- continuity on more general sets.

Pointwise continuity

Definition (Continuous at an interior point of the domain of f)

If the function f is defined in a neighbourhood of the point x_0 then we say f is **continuous at x_0** iff

$$\lim_{x \rightarrow x_0} f(x) = f(x_0).$$

This definition works more generally provided x_0 is a point of accumulation of the domain of f (notation: $\text{dom}(f)$).

We will also consider a function to be continuous at any isolated point in its domain.

Pointwise continuity

Definition (Continuous at any $x_0 \in \text{dom}(f)$ – limit version)

If $x_0 \in \text{dom}(f)$ then f is **continuous at x_0** iff x_0 is either an isolated point of $\text{dom}(f)$ or x_0 is an accumulation point of $\text{dom}(f)$ and $\lim_{x \rightarrow x_0} f(x) = f(x_0)$.

Definition (Continuous at any $x_0 \in \text{dom}(f)$ – sequence version)

If $x_0 \in \text{dom}(f)$ then f is **continuous at x_0** iff for any sequence $\{x_n\}$ in $\text{dom}(f)$, if $x_n \rightarrow x_0$ then $f(x_n) \rightarrow f(x_0)$.

Definition (Continuous at any $x_0 \in \text{dom}(f)$ – ε - δ version)

If $x_0 \in \text{dom}(f)$ then f is **continuous at x_0** iff for any $\varepsilon > 0$ there exists $\delta > 0$ such that if $x \in \text{dom}(f)$ and $|x - x_0| < \delta$ then $|f(x) - f(x_0)| < \varepsilon$.

Pointwise continuity

Example

Suppose $f : A \rightarrow \mathbb{R}$. In which cases is f continuous on A ?

- $A = (0, 1) \cup \{2\}$, $f(x) = x$;
- $A = \{0\} \cup \{\frac{1}{n} : n \in \mathbb{N}\} \cup \{2\}$, $f(x) = x$;
- $A = \{\frac{1}{n} : n \in \mathbb{N}\} \cup \{2\}$, $f(x) = \text{whatever you like}$.

Example

Is it possible for a function f to be discontinuous at every point of \mathbb{R} and yet for its restriction to the rational numbers ($f|_{\mathbb{Q}}$) to be **continuous** at every point in \mathbb{Q} ?

Extra Challenge Problem:

Prove or disprove: There is a function $f : \mathbb{R} \rightarrow \mathbb{R}$ that is continuous at every irrational number and discontinuous at every rational number.

Continuity on an interval

Definition (Continuous on an open interval)

The function f is said to be **continuous on** (a, b) iff

$$\lim_{x \rightarrow x_0} f(x) = f(x_0) \quad \text{for all } x_0 \in (a, b).$$

Definition (Continuous on a closed interval)

The function f is said to be **continuous on** $[a, b]$ iff it is continuous on the open interval (a, b) , and

$$\lim_{x \rightarrow a^+} f(x) = f(a) \quad \text{and} \quad \lim_{x \rightarrow b^-} f(x) = f(b).$$

Continuity on an arbitrary set $E \subseteq \mathbb{R}$

Definition (Continuous on a set E)

The function f is said to be **continuous on E** iff f is continuous at each point $x \in E$.

Example

- Every polynomial is continuous on \mathbb{R} .
- Every rational function is continuous on its domain (i.e., avoiding points where the denominator is zero).

These facts are painful to prove directly from the definition. But they follow easily if we know that the composition of continuous functions is continuous (which is true under natural conditions) and we have the theorem on the algebra of limits.

Continuity of compositions of functions

Theorem (Continuity of $f \circ g$ at a point)

If g is continuous at x_0 and f is continuous at $g(x_0)$ then $f \circ g$ is continuous at x_0 .

Consequently, if g is continuous at x_0 and f is continuous at $g(x_0)$ then

$$\lim_{x \rightarrow x_0} f(g(x)) = f\left(\lim_{x \rightarrow x_0} g(x)\right).$$

Theorem (Continuity of $f \circ g$ on a set)

If g is continuous on $A \subseteq \mathbb{R}$ and f is continuous on $g(A)$ then $f \circ g$ is continuous on A .

Continuity of compositions of functions

Example

Use the theorem on continuity of $f \circ g$, and the theorem on the algebra of limits, to prove that

- 1 the polynomial $x^8 + x^3 + 2$ is continuous on \mathbb{R} ;
- 2 the rational function $\frac{x^2 + 2}{x^2 - 2}$ is continuous on $\mathbb{R} \setminus \{-\sqrt{2}, \sqrt{2}\}$.
- 3 the function $\sqrt{\frac{x^2 + 2}{x^2 - 2}}$ is continuous on its domain.