

2 Differentiation

3 Differentiation II

4 Differentiation III

Differentiation



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 2
Differentiation
Wednesday 8 January 2025

Survey to do right now

- Please go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Survey 2**
- .

Announcements

Announcements

- Results of Survey 1

Announcements

- Results of Survey 1
- Results of Survey 2

Background / reminder

Definition (Cauchy sequence)

Background / reminder

Definition (Cauchy sequence)

A sequence $\{s_n\}$

Background / reminder

Definition (Cauchy sequence)

A sequence $\{s_n\}$ is said to be

Background / reminder

Definition (Cauchy sequence)

A sequence $\{s_n\}$ is said to be a *Cauchy sequence*

Background / reminder

Definition (Cauchy sequence)

A sequence $\{s_n\}$ is said to be a *Cauchy sequence* iff

Background / reminder

Definition (Cauchy sequence)

A sequence $\{s_n\}$ is said to be a *Cauchy sequence* iff for all $\varepsilon > 0$

Background / reminder

Definition (Cauchy sequence)

A sequence $\{s_n\}$ is said to be a *Cauchy sequence* iff for all $\varepsilon > 0$ there exists $N \in \mathbb{N}$

Background / reminder

Definition (Cauchy sequence)

A sequence $\{s_n\}$ is said to be a *Cauchy sequence* iff for all $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that

Background / reminder

Definition (Cauchy sequence)

A sequence $\{s_n\}$ is said to be a *Cauchy sequence* iff for all $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that if $m \geq N$

Background / reminder

Definition (Cauchy sequence)

A sequence $\{s_n\}$ is said to be a *Cauchy sequence* iff for all $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that if $m \geq N$ and $n \geq N$

Background / reminder

Definition (Cauchy sequence)

A sequence $\{s_n\}$ is said to be a *Cauchy sequence* iff for all $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that if $m \geq N$ and $n \geq N$ then

Background / reminder

Definition (Cauchy sequence)


A sequence $\{s_n\}$ is said to be a *Cauchy sequence* iff for all $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that if $m \geq N$ and $n \geq N$ then $|s_n - s_m| < \varepsilon$.

Poll: another background check

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Background: Cauchy sequences**
- .

The Derivative

The Derivative



The Derivative

The Derivative

Definition (Derivative)

The Derivative

Definition (Derivative)

Let f be defined on an interval I

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$.

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0 , denoted by $f'(x_0)$

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0 , denoted by $f'(x_0)$, is defined as

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) =$$

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0}$$

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$$

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite.

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The *derivative* of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite we say that f is

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The **derivative** of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite we say that f is **differentiable** at x_0 .

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The **derivative** of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite we say that f is **differentiable** at x_0 . If f is differentiable at every point of a set $E \subseteq I$

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The **derivative** of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite we say that f is **differentiable** at x_0 . If f is differentiable at every point of a set $E \subseteq I$, we say that f is

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The **derivative** of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite we say that f is **differentiable** at x_0 . If f is differentiable at every point of a set $E \subseteq I$, we say that f is differentiable on E .

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The **derivative** of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite we say that f is **differentiable** at x_0 . If f is differentiable at every point of a set $E \subseteq I$, we say that f is differentiable on E . If E is all of I

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The **derivative** of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite we say that f is **differentiable** at x_0 . If f is differentiable at every point of a set $E \subseteq I$, we say that f is differentiable on E . If E is all of I , we simply say that f is

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The **derivative** of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite we say that f is **differentiable** at x_0 . If f is differentiable at every point of a set $E \subseteq I$, we say that f is differentiable on E . If E is all of I , we simply say that f is a **differentiable function**.

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The **derivative** of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite we say that f is **differentiable** at x_0 . If f is differentiable at every point of a set $E \subseteq I$, we say that f is differentiable on E . If E is all of I , we simply say that f is a **differentiable function**.

Note: “Differentiable” and “a derivative exists” always mean that

The Derivative

Definition (Derivative)

Let f be defined on an interval I and let $x_0 \in I$. The **derivative** of f at x_0 , denoted by $f'(x_0)$, is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this limit exists or is infinite. If $f'(x_0)$ is finite we say that f is **differentiable** at x_0 . If f is differentiable at every point of a set $E \subseteq I$, we say that f is differentiable on E . If E is all of I , we simply say that f is a **differentiable function**.

Note: “Differentiable” and “a derivative exists” always mean that the derivative is finite.

The Derivative

Example

The Derivative

Example

$$f(x) = x^2.$$

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) =$$

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2}$$

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2}$$

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2}$$

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x + 2)(x - 2)}{x - 2}$$

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x + 2)(x - 2)}{x - 2} = \lim_{x \rightarrow 2} x + 2$$

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x + 2)(x - 2)}{x - 2} = \lim_{x \rightarrow 2} x + 2 = 4$$

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x + 2)(x - 2)}{x - 2} = \lim_{x \rightarrow 2} x + 2 = 4$$

Note:

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x + 2)(x - 2)}{x - 2} = \lim_{x \rightarrow 2} x + 2 = 4$$

Note:

- In the first two limits, we must have $x \neq 2$.

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x + 2)(x - 2)}{x - 2} = \lim_{x \rightarrow 2} x + 2 = 4$$

Note:

- In the first two limits, we must have $x \neq 2$.
- But in the third limit, we just plug in $x = 2$.

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x + 2)(x - 2)}{x - 2} = \lim_{x \rightarrow 2} x + 2 = 4$$

Note:

- In the first two limits, we must have $x \neq 2$.
- But in the third limit, we just plug in $x = 2$.
- Two things are equal, but in one $x \neq 2$ and in the other $x = 2$.

The Derivative

Example

$f(x) = x^2$. Find $f'(2)$.

$$f'(2) = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x + 2)(x - 2)}{x - 2} = \lim_{x \rightarrow 2} x + 2 = 4$$

Note:

- In the first two limits, we must have $x \neq 2$.
- But in the third limit, we just plug in $x = 2$.
- Two things are equal, but in one $x \neq 2$ and in the other $x = 2$.
- Good illustration of why it is important to define the meaning of limits rigorously.

Poll

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Derivatives: Differentiable at 0**
- .

The Derivative

The Derivative

Example

The Derivative

Example

Let f be defined in a neighbourhood I of 0

The Derivative

Example

Let f be defined in a neighbourhood I of 0, and suppose $|f(x)| \leq x^2$ for all $x \in I$.

The Derivative

Example

Let f be defined in a neighbourhood I of 0, and suppose $|f(x)| \leq x^2$ for all $x \in I$. Is f necessarily differentiable at 0?

The Derivative


Example

Let f be defined in a neighbourhood I of 0, and suppose $|f(x)| \leq x^2$ for all $x \in I$. Is f necessarily differentiable at 0? e.g.,

The Derivative

Example

Let f be defined in a neighbourhood I of 0, and suppose $|f(x)| \leq x^2$ for all $x \in I$. Is f necessarily differentiable at 0? e.g.,



The Derivative

The Derivative

Example (Trapping principle)

The Derivative

Example (Trapping principle)

$$\text{Suppose } f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$\forall x \neq 0 :$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| =$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| =$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| =$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| = \left| x \sin \frac{1}{x^2} \right| \leq$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| = \left| x \sin \frac{1}{x^2} \right| \leq$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| = \left| x \sin \frac{1}{x^2} \right| \leq |x|$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| = \left| x \sin \frac{1}{x^2} \right| \leq |x|$$

Therefore:

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| = \left| x \sin \frac{1}{x^2} \right| \leq |x|$$

Therefore:

$$|f'(0)| =$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| = \left| x \sin \frac{1}{x^2} \right| \leq |x|$$

Therefore:

$$|f'(0)| = \left| \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} \right| =$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| = \left| x \sin \frac{1}{x^2} \right| \leq |x|$$

Therefore:

$$|f'(0)| = \left| \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} \right| = \lim_{x \rightarrow 0} \left| \frac{f(x) - f(0)}{x - 0} \right| \leq$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| = \left| x \sin \frac{1}{x^2} \right| \leq |x|$$

Therefore:

$$|f'(0)| = \left| \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} \right| = \lim_{x \rightarrow 0} \left| \frac{f(x) - f(0)}{x - 0} \right| \leq \lim_{x \rightarrow 0} |x| =$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| = \left| x \sin \frac{1}{x^2} \right| \leq |x|$$

Therefore:

$$|f'(0)| = \left| \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} \right| = \lim_{x \rightarrow 0} \left| \frac{f(x) - f(0)}{x - 0} \right| \leq \lim_{x \rightarrow 0} |x| = 0.$$

The Derivative

Example (Trapping principle)

Suppose $f(x) = \begin{cases} x^2 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$ Then:

$$\forall x \neq 0 : \left| \frac{f(x) - f(0)}{x - 0} \right| = \left| \frac{f(x)}{x} \right| = \left| \frac{x^2 \sin \frac{1}{x^2}}{x} \right| = \left| x \sin \frac{1}{x^2} \right| \leq |x|$$

Therefore:

$$|f'(0)| = \left| \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} \right| = \lim_{x \rightarrow 0} \left| \frac{f(x) - f(0)}{x - 0} \right| \leq \lim_{x \rightarrow 0} |x| = 0.$$

$\therefore f$ is differentiable at 0 and $f'(0) = 0$. □

The Derivative

The Derivative

Definition (One-sided derivatives)

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$.

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The *right-hand derivative* of f at x_0

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The *right-hand derivative* of f at x_0 , denoted by $f'_+(x_0)$

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The *right-hand derivative* of f at x_0 , denoted by $f'_+(x_0)$, is the limit

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The **right-hand derivative** of f at x_0 , denoted by $f'_+(x_0)$, is the limit

$$f'_+(x_0) = \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0},$$

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The **right-hand derivative** of f at x_0 , denoted by $f'_+(x_0)$, is the limit

$$f'_+(x_0) = \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The **right-hand derivative** of f at x_0 , denoted by $f'_+(x_0)$, is the limit

$$f'_+(x_0) = \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this one-sided limit exists or

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The **right-hand derivative** of f at x_0 , denoted by $f'_+(x_0)$, is the limit

$$f'_+(x_0) = \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this one-sided limit exists or is infinite.

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The **right-hand derivative** of f at x_0 , denoted by $f'_+(x_0)$, is the limit

$$f'_+(x_0) = \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this one-sided limit exists or is infinite. Similarly, the **left-hand derivative** of f at x_0

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The **right-hand derivative** of f at x_0 , denoted by $f'_+(x_0)$, is the limit

$$f'_+(x_0) = \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this one-sided limit exists or is infinite.

Similarly, the **left-hand derivative** of f at x_0 , denoted by $f'_-(x_0)$

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The **right-hand derivative** of f at x_0 , denoted by $f'_+(x_0)$, is the limit

$$f'_+(x_0) = \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this one-sided limit exists or is infinite.

Similarly, the **left-hand derivative** of f at x_0 , denoted by $f'_-(x_0)$, is the limit

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The **right-hand derivative** of f at x_0 , denoted by $f'_+(x_0)$, is the limit

$$f'_+(x_0) = \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this one-sided limit exists or is infinite.

Similarly, the **left-hand derivative** of f at x_0 , denoted by $f'_-(x_0)$, is the limit

$$f'_-(x_0) = \lim_{x \rightarrow x_0^-} \frac{f(x) - f(x_0)}{x - x_0}.$$

The Derivative

Definition (One-sided derivatives)

Let f be defined on an interval I and let $x_0 \in I$. The **right-hand derivative** of f at x_0 , denoted by $f'_+(x_0)$, is the limit

$$f'_+(x_0) = \lim_{x \rightarrow x_0^+} \frac{f(x) - f(x_0)}{x - x_0},$$

provided either that this one-sided limit exists or is infinite.

Similarly, the **left-hand derivative** of f at x_0 , denoted by $f'_-(x_0)$, is the limit


$$f'_-(x_0) = \lim_{x \rightarrow x_0^-} \frac{f(x) - f(x_0)}{x - x_0}.$$

Note: If x_0 is not an endpoint of the interval I then f is differentiable at x_0 iff $f'_+(x_0) = f'_-(x_0) \neq \pm\infty$.

The Derivative

Example


$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$



The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$




- Same slope from left and right.

The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$




- Same slope from left and right. Why isn't f differentiable???

The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$




- Same slope from left and right. Why isn't f differentiable???
- $\lim_{x \rightarrow 0^-} f'(x) =$

The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$




- Same slope from left and right. Why isn't f differentiable???
- $\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^+} f'(x) =$

The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$




- Same slope from left and right. Why isn't f differentiable???
- $\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0} f'(x) =$

The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$




- Same slope from left and right. Why isn't f differentiable???
- $\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0} f'(x) = 1.$

The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$




- Same slope from left and right. Why isn't f differentiable???
- $\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0} f'(x) = 1.$
- $f'_-(0) =$

The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$




- Same slope from left and right. Why isn't f differentiable???
- $\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0} f'(x) = 1.$
- $f'_-(0) = f'_+(0) =$

The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$




- Same slope from left and right. Why isn't f differentiable???
- $\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0} f'(x) = 1.$
- $f'_-(0) = f'_+(0) = f'(0) =$

The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$




- Same slope from left and right. Why isn't f differentiable???
- $\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0} f'(x) = 1.$
- $f'_-(0) = f'_+(0) = f'(0) = \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} =$

The Derivative

Example

$$f(x) = \begin{cases} x - 1, & x < 0 \\ 0, & x = 0, \\ x + 1, & x > 0 \end{cases}$$



- Same slope from left and right. Why isn't f differentiable???
- $\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0} f'(x) = 1.$
- $f'_-(0) = f'_+(0) = f'(0) = \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \infty.$

The Derivative

The Derivative

- Higher derivatives:

The Derivative

- Higher derivatives: we write

The Derivative

- Higher derivatives: we write
 - $f'' = (f')'$ if f' is differentiable;

The Derivative

- Higher derivatives: we write
 - $f'' = (f')'$ if f' is differentiable;
 - $f^{(n+1)} = (f^{(n)})'$ if $f^{(n)}$ is differentiable.

The Derivative

- Higher derivatives: we write
 - $f'' = (f')'$ if f' is differentiable;
 - $f^{(n+1)} = (f^{(n)})'$ if $f^{(n)}$ is differentiable.
- Other standard notation for derivatives:

The Derivative

- Higher derivatives: we write
 - $f'' = (f')'$ if f' is differentiable;
 - $f^{(n+1)} = (f^{(n)})'$ if $f^{(n)}$ is differentiable.
- Other standard notation for derivatives:

$$\frac{df}{dx} = f'(x)$$

The Derivative

- Higher derivatives: we write
 - $f'' = (f')'$ if f' is differentiable;
 - $f^{(n+1)} = (f^{(n)})'$ if $f^{(n)}$ is differentiable.
- Other standard notation for derivatives:

$$\frac{df}{dx} = f'(x)$$

$$D = \frac{d}{dx}$$

The Derivative

- Higher derivatives: we write
 - $f'' = (f')'$ if f' is differentiable;
 - $f^{(n+1)} = (f^{(n)})'$ if $f^{(n)}$ is differentiable.
- Other standard notation for derivatives:

$$\frac{df}{dx} = f'(x)$$

$$D = \frac{d}{dx}$$

$$D^n f(x) = \frac{d^n f}{dx^n} = f^{(n)}(x)$$

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of sequences)

Suppose $\{s_n\}$ and $\{t_n\}$ are *convergent sequences* and $C \in \mathbb{R}$.

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of sequences)

Suppose $\{s_n\}$ and $\{t_n\}$ are *convergent sequences* and $C \in \mathbb{R}$.

$$\mathbf{1} \quad \lim_{n \rightarrow \infty} C s_n = C \left(\lim_{n \rightarrow \infty} s_n \right) ;$$

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of sequences)

Suppose $\{s_n\}$ and $\{t_n\}$ are *convergent sequences* and $C \in \mathbb{R}$.

1 $\lim_{n \rightarrow \infty} C s_n = C \left(\lim_{n \rightarrow \infty} s_n \right) ;$

2 $\lim_{n \rightarrow \infty} (s_n + t_n) = \left(\lim_{n \rightarrow \infty} s_n \right) + \left(\lim_{n \rightarrow \infty} t_n \right) ;$

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of sequences)

Suppose $\{s_n\}$ and $\{t_n\}$ are *convergent sequences* and $C \in \mathbb{R}$.

$$\mathbf{1} \quad \lim_{n \rightarrow \infty} C s_n = C \left(\lim_{n \rightarrow \infty} s_n \right) ;$$

$$\mathbf{2} \quad \lim_{n \rightarrow \infty} (s_n + t_n) = \left(\lim_{n \rightarrow \infty} s_n \right) + \left(\lim_{n \rightarrow \infty} t_n \right) ;$$

$$\mathbf{3} \quad \lim_{n \rightarrow \infty} (s_n - t_n) = \left(\lim_{n \rightarrow \infty} s_n \right) - \left(\lim_{n \rightarrow \infty} t_n \right) ;$$

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of sequences)

Suppose $\{s_n\}$ and $\{t_n\}$ are *convergent sequences* and $C \in \mathbb{R}$.

$$1 \quad \lim_{n \rightarrow \infty} C s_n = C \left(\lim_{n \rightarrow \infty} s_n \right) ;$$

$$2 \quad \lim_{n \rightarrow \infty} (s_n + t_n) = \left(\lim_{n \rightarrow \infty} s_n \right) + \left(\lim_{n \rightarrow \infty} t_n \right) ;$$

$$3 \quad \lim_{n \rightarrow \infty} (s_n - t_n) = \left(\lim_{n \rightarrow \infty} s_n \right) - \left(\lim_{n \rightarrow \infty} t_n \right) ;$$

$$4 \quad \lim_{n \rightarrow \infty} (s_n t_n) = \left(\lim_{n \rightarrow \infty} s_n \right) \left(\lim_{n \rightarrow \infty} t_n \right) ;$$

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of sequences)

Suppose $\{s_n\}$ and $\{t_n\}$ are *convergent sequences* and $C \in \mathbb{R}$.

1 $\lim_{n \rightarrow \infty} C s_n = C \left(\lim_{n \rightarrow \infty} s_n \right) ;$

2 $\lim_{n \rightarrow \infty} (s_n + t_n) = \left(\lim_{n \rightarrow \infty} s_n \right) + \left(\lim_{n \rightarrow \infty} t_n \right) ;$

3 $\lim_{n \rightarrow \infty} (s_n - t_n) = \left(\lim_{n \rightarrow \infty} s_n \right) - \left(\lim_{n \rightarrow \infty} t_n \right) ;$

4 $\lim_{n \rightarrow \infty} (s_n t_n) = \left(\lim_{n \rightarrow \infty} s_n \right) \left(\lim_{n \rightarrow \infty} t_n \right) ;$

5 if $t_n \neq 0$ for all n and $\lim_{n \rightarrow \infty} t_n \neq 0$ then

$$\lim_{n \rightarrow \infty} \left(\frac{s_n}{t_n} \right) = \frac{\lim_{n \rightarrow \infty} s_n}{\lim_{n \rightarrow \infty} t_n} .$$

(TBB §2.7, and problem 2.7.4)

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of functions)

Suppose $f, g : \mathbb{R} \rightarrow \mathbb{R}$, $x_0 \in \mathbb{R}$, the limits as $x \rightarrow x_0$ of $f(x)$ and $g(x)$ both exist, and $C \in \mathbb{R}$.

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of functions)

Suppose $f, g : \mathbb{R} \rightarrow \mathbb{R}$, $x_0 \in \mathbb{R}$, the limits as $x \rightarrow x_0$ of $f(x)$ and $g(x)$ both exist, and $C \in \mathbb{R}$.

$$\mathbf{1} \quad \lim_{x \rightarrow x_0} C f(x) = C \left(\lim_{x \rightarrow x_0} f(x) \right) ;$$

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of functions)

Suppose $f, g : \mathbb{R} \rightarrow \mathbb{R}$, $x_0 \in \mathbb{R}$, the limits as $x \rightarrow x_0$ of $f(x)$ and $g(x)$ both exist, and $C \in \mathbb{R}$.

$$\mathbf{1} \quad \lim_{x \rightarrow x_0} C f(x) = C \left(\lim_{x \rightarrow x_0} f(x) \right) ;$$

$$\mathbf{2} \quad \lim_{x \rightarrow x_0} (f(x) + g(x)) = \left(\lim_{x \rightarrow x_0} f(x) \right) + \left(\lim_{x \rightarrow x_0} g(x) \right) ;$$

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of functions)

Suppose $f, g : \mathbb{R} \rightarrow \mathbb{R}$, $x_0 \in \mathbb{R}$, the limits as $x \rightarrow x_0$ of $f(x)$ and $g(x)$ both exist, and $C \in \mathbb{R}$.

$$\mathbf{1} \quad \lim_{x \rightarrow x_0} C f(x) = C \left(\lim_{x \rightarrow x_0} f(x) \right) ;$$

$$\mathbf{2} \quad \lim_{x \rightarrow x_0} (f(x) + g(x)) = \left(\lim_{x \rightarrow x_0} f(x) \right) + \left(\lim_{x \rightarrow x_0} g(x) \right) ;$$

$$\mathbf{3} \quad \lim_{x \rightarrow x_0} (f(x) - g(x)) = \left(\lim_{x \rightarrow x_0} f(x) \right) - \left(\lim_{x \rightarrow x_0} g(x) \right) ;$$

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of functions)

Suppose $f, g : \mathbb{R} \rightarrow \mathbb{R}$, $x_0 \in \mathbb{R}$, the limits as $x \rightarrow x_0$ of $f(x)$ and $g(x)$ both exist, and $C \in \mathbb{R}$.

$$1 \quad \lim_{x \rightarrow x_0} C f(x) = C \left(\lim_{x \rightarrow x_0} f(x) \right) ;$$

$$2 \quad \lim_{x \rightarrow x_0} (f(x) + g(x)) = \left(\lim_{x \rightarrow x_0} f(x) \right) + \left(\lim_{x \rightarrow x_0} g(x) \right) ;$$

$$3 \quad \lim_{x \rightarrow x_0} (f(x) - g(x)) = \left(\lim_{x \rightarrow x_0} f(x) \right) - \left(\lim_{x \rightarrow x_0} g(x) \right) ;$$

$$4 \quad \lim_{x \rightarrow x_0} (f(x)g(x)) = \left(\lim_{x \rightarrow x_0} f(x) \right) \left(\lim_{x \rightarrow x_0} g(x) \right) ;$$

REMINDER: Algebra of limits

Theorem (Algebraic operations on limits of functions)

Suppose $f, g : \mathbb{R} \rightarrow \mathbb{R}$, $x_0 \in \mathbb{R}$, the limits as $x \rightarrow x_0$ of $f(x)$ and $g(x)$ both exist, and $C \in \mathbb{R}$.

$$1 \quad \lim_{x \rightarrow x_0} C f(x) = C \left(\lim_{x \rightarrow x_0} f(x) \right) ;$$

$$2 \quad \lim_{x \rightarrow x_0} (f(x) + g(x)) = \left(\lim_{x \rightarrow x_0} f(x) \right) + \left(\lim_{x \rightarrow x_0} g(x) \right) ;$$

$$3 \quad \lim_{x \rightarrow x_0} (f(x) - g(x)) = \left(\lim_{x \rightarrow x_0} f(x) \right) - \left(\lim_{x \rightarrow x_0} g(x) \right) ;$$

$$4 \quad \lim_{x \rightarrow x_0} (f(x)g(x)) = \left(\lim_{x \rightarrow x_0} f(x) \right) \left(\lim_{x \rightarrow x_0} g(x) \right) ;$$

$$5 \quad \text{if } g(x) \neq 0 \text{ for } x \in (x_0 - \delta, x_0 + \delta) \text{ for some } \delta > 0, \text{ and} \\ \lim_{x \rightarrow x_0} g(x) \neq 0 \text{ then } \lim_{x \rightarrow x_0} \left(\frac{f(x)}{g(x)} \right) = \frac{\lim_{x \rightarrow x_0} f(x)}{\lim_{x \rightarrow x_0} g(x)} .$$

The Derivative

The Derivative

Theorem (Differentiable $\implies$ continuous)

The Derivative

Theorem (Differentiable $\implies$ continuous)

If f is defined in a neighbourhood I of x_0 and f is differentiable at x_0 then f is continuous at x_0 .

The Derivative

Theorem (Differentiable $\implies$ continuous)

If f is defined in a neighbourhood I of x_0 and f is differentiable at x_0 then f is continuous at x_0 .

Proof.

Must show $\lim_{x \rightarrow x_0} f(x) = f(x_0)$

The Derivative

Theorem (Differentiable $\implies$ continuous)

If f is defined in a neighbourhood I of x_0 and f is differentiable at x_0 then f is continuous at x_0 .

Proof.

Must show $\lim_{x \rightarrow x_0} f(x) = f(x_0)$, i.e., $\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = 0$

The Derivative

Theorem (Differentiable $\implies$ continuous)

If f is defined in a neighbourhood I of x_0 and f is differentiable at x_0 then f is continuous at x_0 .

Proof.

Must show $\lim_{x \rightarrow x_0} f(x) = f(x_0)$, i.e., $\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = 0$.

$$\lim_{x \rightarrow x_0} (f(x) - f(x_0)) =$$

The Derivative

Theorem (Differentiable $\implies$ continuous)

If f is defined in a neighbourhood I of x_0 and f is differentiable at x_0 then f is continuous at x_0 .

Proof.

Must show $\lim_{x \rightarrow x_0} f(x) = f(x_0)$, i.e., $\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = 0$.

$$\begin{aligned}\lim_{x \rightarrow x_0} (f(x) - f(x_0)) &= \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \times (x - x_0) \right) \\ &= \end{aligned}$$

The Derivative

Theorem (Differentiable $\implies$ continuous)

If f is defined in a neighbourhood I of x_0 and f is differentiable at x_0 then f is continuous at x_0 .

Proof.

Must show $\lim_{x \rightarrow x_0} f(x) = f(x_0)$, i.e., $\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = 0$.

$$\begin{aligned}\lim_{x \rightarrow x_0} (f(x) - f(x_0)) &= \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \times (x - x_0) \right) \\ &= \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \right) \times \lim_{x \rightarrow x_0} (x - x_0)\end{aligned}$$

The Derivative

Theorem (Differentiable $\implies$ continuous)

If f is defined in a neighbourhood I of x_0 and f is differentiable at x_0 then f is continuous at x_0 .

Proof.

Must show $\lim_{x \rightarrow x_0} f(x) = f(x_0)$, i.e., $\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = 0$.

$$\begin{aligned}\lim_{x \rightarrow x_0} (f(x) - f(x_0)) &= \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \times (x - x_0) \right) \\ &= \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \right) \times \lim_{x \rightarrow x_0} (x - x_0) \\ &= f'(x_0) \times 0 =\end{aligned}$$

The Derivative

Theorem (Differentiable $\implies$ continuous)

If f is defined in a neighbourhood I of x_0 and f is differentiable at x_0 then f is continuous at x_0 .

Proof.

Must show $\lim_{x \rightarrow x_0} f(x) = f(x_0)$, i.e., $\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = 0$.

$$\begin{aligned}\lim_{x \rightarrow x_0} (f(x) - f(x_0)) &= \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \times (x - x_0) \right) \\ &= \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \right) \times \lim_{x \rightarrow x_0} (x - x_0) \\ &= f'(x_0) \times 0 = 0,\end{aligned}$$

The Derivative

Theorem (Differentiable $\implies$ continuous)

If f is defined in a neighbourhood I of x_0 and f is differentiable at x_0 then f is continuous at x_0 .

Proof.

Must show $\lim_{x \rightarrow x_0} f(x) = f(x_0)$, i.e., $\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = 0$.

$$\begin{aligned}\lim_{x \rightarrow x_0} (f(x) - f(x_0)) &= \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \times (x - x_0) \right) \\ &= \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \right) \times \lim_{x \rightarrow x_0} (x - x_0) \\ &= f'(x_0) \times 0 = 0,\end{aligned}$$

where we have used the theorem on the algebra of limits. □



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 3
Differentiation II
Friday 10 January 2025

Announcements

Announcements

- Lectures are being live streamed and recorded as of today.

Announcements

- Lectures are being live streamed and recorded as of today.
- Recordings will be available 24 hours after lectures.

Announcements

- Lectures are being live streamed and recorded as of today.
- Recordings will be available 24 hours after lectures.
 - Go to <https://echo360.ca>.

Announcements

- Lectures are being live streamed and recorded as of today.
- Recordings will be available 24 hours after lectures.
 - Go to <https://echo360.ca>.
 - Sign in with your `macID@mcmaster.ca` e-mail address.

Announcements

- Lectures are being live streamed and recorded as of today.
- Recordings will be available 24 hours after lectures.
 - Go to <https://echo360.ca>.
 - Sign in with your `macID@mcmaster.ca` e-mail address.
 - Click on the Courses tab.

Announcements

- Lectures are being live streamed and recorded as of today.
- Recordings will be available 24 hours after lectures.
 - Go to <https://echo360.ca>.
 - Sign in with your `macID@mcmaster.ca` e-mail address.
 - Click on the Courses tab.
 - Select "MATH 3A03 - WINTER 2025"

Announcements

- Lectures are being live streamed and recorded as of today.
- Recordings will be available 24 hours after lectures.
 - Go to <https://echo360.ca>.
 - Sign in with your `macID@mcmaster.ca` e-mail address.
 - Click on the Courses tab.
 - Select "MATH 3A03 - WINTER 2025"
- Course evaluation scheme has changed

Announcements

- Lectures are being live streamed and recorded as of today.
- Recordings will be available 24 hours after lectures.
 - Go to <https://echo360.ca>.
 - Sign in with your `macID@mcmaster.ca` e-mail address.
 - Click on the Courses tab.
 - Select "MATH 3A03 - WINTER 2025"
- Course evaluation scheme has changed (next two slides).

Announcements

- Lectures are being live streamed and recorded as of today.
- Recordings will be available 24 hours after lectures.
 - Go to <https://echo360.ca>.
 - Sign in with your `macID@mcmaster.ca` e-mail address.
 - Click on the Courses tab.
 - Select "MATH 3A03 - WINTER 2025"
- Course evaluation scheme has changed (next two slides).
 - The [course web site](#) and [online syllabus](#) have not yet been updated to reflect these changes, but that will happen soon (hopefully over the weekend).

Course evaluation will be revised as follows:

Course evaluation will be revised as follows:

- We will not have quizzes

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025
- 50% for final exam in April

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025
- 50% for final exam in April
- Note: If your final exam mark is better than your midterm test mark

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025
- 50% for final exam in April
- Note: If your final exam mark is better than your midterm test mark then the final exam mark will replace the midterm test mark.

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025
- 50% for final exam in April
- Note: If your final exam mark is better than your midterm test mark then the final exam mark will replace the midterm test mark.
- Important: Do NOT skip the midterm.

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025
- 50% for final exam in April
- Note: If your final exam mark is better than your midterm test mark then the final exam mark will replace the midterm test mark.
- Important: Do NOT skip the midterm. Even if you don't feel well prepared,

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025
- 50% for final exam in April
- Note: If your final exam mark is better than your midterm test mark then the final exam mark will replace the midterm test mark.
- Important: Do NOT skip the midterm. Even if you don't feel well prepared, write it for practice

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025
- 50% for final exam in April
- Note: If your final exam mark is better than your midterm test mark then the final exam mark will replace the midterm test mark.
- Important: Do NOT skip the midterm. Even if you don't feel well prepared, write it for practice so you are better prepared for writing the final exam.

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025
- 50% for final exam in April
- Note: If your final exam mark is better than your midterm test mark then the final exam mark will replace the midterm test mark.
- Important: Do NOT skip the midterm. Even if you don't feel well prepared, write it for practice so you are better prepared for writing the final exam.
- If you must miss the midterm

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025
- 50% for final exam in April
- Note: If your final exam mark is better than your midterm test mark then the final exam mark will replace the midterm test mark.
- Important: Do NOT skip the midterm. Even if you don't feel well prepared, write it for practice so you are better prepared for writing the final exam.
- If you must miss the midterm (e.g., illness

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025
- 50% for final exam in April
- Note: If your final exam mark is better than your midterm test mark then the final exam mark will replace the midterm test mark.
- Important: Do NOT skip the midterm. Even if you don't feel well prepared, write it for practice so you are better prepared for writing the final exam.
- If you must miss the midterm (e.g., illness or accepting a Nobel prize),

Course evaluation will be revised as follows:

- We will not have quizzes
- 5% for participating in at least 80% of in-class polls
- 15% for participating in assignments, based on multiple choice (MC) questions:

$$\text{assignment mark} = \frac{\text{number MC questions answered}}{\text{total number MC questions assigned}}$$

- 30% for midterm test on Thurs 27 Feb 2025
- 50% for final exam in April
- Note: If your final exam mark is better than your midterm test mark then the final exam mark will replace the midterm test mark.
- Important: Do NOT skip the midterm. Even if you don't feel well prepared, write it for practice so you are better prepared for writing the final exam.
- If you must miss the midterm (e.g., illness or accepting a Nobel prize), your final exam mark will replace it.

Tentative plan for assignments

Tentative plan for assignments

- There will be regular assignments.

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)).

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven't had time to think about this yet”.

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven't had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#).

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven't had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks.

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven't had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven't had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you're not sure if your proof is complete,

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven't had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you're not sure if your proof is complete, or you got stuck and don't know how to complete it,

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven't had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you're not sure if your proof is complete, or you got stuck and don't know how to complete it, make that clear in the document that you submit on [crowdmark](#),

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven’t had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you’re not sure if your proof is complete, or you got stuck and don’t know how to complete it, make that clear in the document that you submit on [crowdmark](#), so the TA can focus on the help you need.

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven’t had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you’re not sure if your proof is complete, or you got stuck and don’t know how to complete it, make that clear in the document that you submit on [crowdmark](#), so the TA can focus on the help you need.
- Always try your best to solve problems on your own first.

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven't had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you're not sure if your proof is complete, or you got stuck and don't know how to complete it, make that clear in the document that you submit on [crowdmark](#), so the TA can focus on the help you need.
- Always try your best to solve problems on your own first. But if you used stackexchange

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven’t had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you’re not sure if your proof is complete, or you got stuck and don’t know how to complete it, make that clear in the document that you submit on [crowdmark](#), so the TA can focus on the help you need.
- Always try your best to solve problems on your own first. But if you used stackexchange or ChatGPT

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven't had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you're not sure if your proof is complete, or you got stuck and don't know how to complete it, make that clear in the document that you submit on [crowdmark](#), so the TA can focus on the help you need.
- Always try your best to solve problems on your own first. But if you used stackexchange or ChatGPT or whatever

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven’t had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you’re not sure if your proof is complete, or you got stuck and don’t know how to complete it, make that clear in the document that you submit on [crowdmark](#), so the TA can focus on the help you need.
- Always try your best to solve problems on your own first. But if you used stackexchange or ChatGPT or whatever for help,

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven’t had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you’re not sure if your proof is complete, or you got stuck and don’t know how to complete it, make that clear in the document that you submit on [crowdmark](#), so the TA can focus on the help you need.
- Always try your best to solve problems on your own first. But if you used stackexchange or ChatGPT or whatever for help, provide a URL to your source if possible,

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven’t had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you’re not sure if your proof is complete, or you got stuck and don’t know how to complete it, make that clear in the document that you submit on [crowdmark](#), so the TA can focus on the help you need.
- Always try your best to solve problems on your own first. But if you used stackexchange or ChatGPT or whatever for help, provide a URL to your source if possible, so it is easier for the TA to provide the help you need.

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven’t had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you’re not sure if your proof is complete, or you got stuck and don’t know how to complete it, make that clear in the document that you submit on [crowdmark](#), so the TA can focus on the help you need.
- Always try your best to solve problems on your own first. But if you used stackexchange or ChatGPT or whatever for help, provide a URL to your source if possible, so it is easier for the TA to provide the help you need.
- Make the best possible use of the TA’s time

Tentative plan for assignments

- There will be regular assignments.
- Each question will have a multiple choice component (probably on [childsmath](#)). Only participation counts for marks; you will get the same credit for correct and incorrect answers, or for selecting “I haven’t had time to think about this yet”.
- Optionally, full solutions/proofs can be written up and submitted on [crowdmark](#). Feedback will be given, but no marks. The purpose is to help you prepare better for the test and exam.
- If you’re not sure if your proof is complete, or you got stuck and don’t know how to complete it, make that clear in the document that you submit on [crowdmark](#), so the TA can focus on the help you need.
- Always try your best to solve problems on your own first. But if you used stackexchange or ChatGPT or whatever for help, provide a URL to your source if possible, so it is easier for the TA to provide the help you need.
- Make the best possible use of the TA’s time: say what you think you do or don’t understand.

Last time...

Last time...

- Definition of the derivative.

Last time...

- Definition of the derivative.
 - Example: Trapping Principle

Last time...

- Definition of the derivative.
 - Example: Trapping Principle
- Defined one-sided derivatives

Last time...

- Definition of the derivative.
 - Example: Trapping Principle
- Defined one-sided derivatives
 - Example

Last time...

- Definition of the derivative.
 - Example: Trapping Principle
- Defined one-sided derivatives
 - Example
- Proved differentiable $\implies$ continuous.

More on the derivative

More on the derivative

Theorem (Algebra of derivatives)

More on the derivative

Theorem (Algebra of derivatives)

Suppose f and g are defined on an interval I and $x_0 \in I$.

More on the derivative

Theorem (Algebra of derivatives)

Suppose f and g are defined on an interval I and $x_0 \in I$. If f and g are differentiable at x_0 then

More on the derivative

Theorem (Algebra of derivatives)

Suppose f and g are defined on an interval I and $x_0 \in I$. If f and g are differentiable at x_0 then $f + g$ and fg are differentiable at x_0 .

More on the derivative

Theorem (Algebra of derivatives)

Suppose f and g are defined on an interval I and $x_0 \in I$. If f and g are differentiable at x_0 then $f + g$ and fg are differentiable at x_0 . If, in addition, $g(x_0) \neq 0$ then

More on the derivative

Theorem (Algebra of derivatives)

Suppose f and g are defined on an interval I and $x_0 \in I$. If f and g are differentiable at x_0 then $f + g$ and fg are differentiable at x_0 . If, in addition, $g(x_0) \neq 0$ then f/g is differentiable at x_0 .

More on the derivative

Theorem (Algebra of derivatives)

Suppose f and g are defined on an interval I and $x_0 \in I$. If f and g are differentiable at x_0 then $f + g$ and fg are differentiable at x_0 . If, in addition, $g(x_0) \neq 0$ then f/g is differentiable at x_0 . Under these conditions:

More on the derivative

Theorem (Algebra of derivatives)

Suppose f and g are defined on an interval I and $x_0 \in I$. If f and g are differentiable at x_0 then $f + g$ and fg are differentiable at x_0 . If, in addition, $g(x_0) \neq 0$ then f/g is differentiable at x_0 . Under these conditions:

1 $(cf)'(x_0) = cf'(x_0)$ for all $c \in \mathbb{R}$;

More on the derivative

Theorem (Algebra of derivatives)

Suppose f and g are defined on an interval I and $x_0 \in I$. If f and g are differentiable at x_0 then $f + g$ and fg are differentiable at x_0 . If, in addition, $g(x_0) \neq 0$ then f/g is differentiable at x_0 . Under these conditions:

- 1** $(cf)'(x_0) = cf'(x_0)$ for all $c \in \mathbb{R}$;
- 2** $(f + g)'(x_0) = (f' + g')(x_0)$;

More on the derivative

Theorem (Algebra of derivatives)

Suppose f and g are defined on an interval I and $x_0 \in I$. If f and g are differentiable at x_0 then $f + g$ and fg are differentiable at x_0 . If, in addition, $g(x_0) \neq 0$ then f/g is differentiable at x_0 . Under these conditions:

- 1** $(cf)'(x_0) = cf'(x_0)$ for all $c \in \mathbb{R}$;
- 2** $(f + g)'(x_0) = (f' + g')(x_0)$;
- 3** $(fg)'(x_0) = (f'g + fg')(x_0)$;

More on the derivative

Theorem (Algebra of derivatives)

Suppose f and g are defined on an interval I and $x_0 \in I$. If f and g are differentiable at x_0 then $f + g$ and fg are differentiable at x_0 . If, in addition, $g(x_0) \neq 0$ then f/g is differentiable at x_0 . Under these conditions:

- 1** $(cf)'(x_0) = cf'(x_0)$ for all $c \in \mathbb{R}$;
- 2** $(f + g)'(x_0) = (f' + g')(x_0)$;
- 3** $(fg)'(x_0) = (f'g + fg')(x_0)$;
- 4** $\left(\frac{f}{g}\right)'(x_0) = \left(\frac{gf' - fg'}{g^2}\right)(x_0) \quad (g(x_0) \neq 0).$

(TBB Theorem 7.7, p. 408)

The Derivative

The Derivative

Theorem (Chain rule)

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$.

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$ then the composite function $h = g \circ f$ is

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$ then the composite function $h = g \circ f$ is differentiable at x_0

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$ then the composite function $h = g \circ f$ is differentiable at x_0 and

$$h'(x_0) =$$

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$ then the composite function $h = g \circ f$ is differentiable at x_0 and

$$h'(x_0) = (g \circ f)'(x_0) =$$

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$ then the composite function $h = g \circ f$ is differentiable at x_0 and

$$h'(x_0) = (g \circ f)'(x_0) = g'(f(x_0))f'(x_0).$$

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$ then the composite function $h = g \circ f$ is differentiable at x_0 and

$$h'(x_0) = (g \circ f)'(x_0) = g'(f(x_0))f'(x_0).$$

Informally, if $y = f(x)$ and $z = g(y)$

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$ then the composite function $h = g \circ f$ is differentiable at x_0 and

$$h'(x_0) = (g \circ f)'(x_0) = g'(f(x_0))f'(x_0).$$

Informally, if $y = f(x)$ and $z = g(y)$ then

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$ then the composite function $h = g \circ f$ is differentiable at x_0 and

$$h'(x_0) = (g \circ f)'(x_0) = g'(f(x_0))f'(x_0).$$

Informally, if $y = f(x)$ and $z = g(y)$ then $\frac{dz}{dx} = \frac{dz}{dy} \frac{dy}{dx}$.

The Derivative

Theorem (Chain rule)

Suppose f is defined in a neighbourhood U of x_0 and g is defined in a neighbourhood V of $f(x_0)$ such that $f(U) \subseteq V$. If f is differentiable at x_0 and g is differentiable at $f(x_0)$ then the composite function $h = g \circ f$ is differentiable at x_0 and

$$h'(x_0) = (g \circ f)'(x_0) = g'(f(x_0))f'(x_0).$$

Informally, if $y = f(x)$ and $z = g(y)$ then $\frac{dz}{dx} = \frac{dz}{dy} \frac{dy}{dx}$.

(TBB §7.3.2, p. 411)

Why the chain rule is plausible

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0}$$

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0}$$

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} =$$

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{x - x_0}$$

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)}.$$

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0}$$

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0}$$

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{\textcolor{red}{f(x)} - \textcolor{red}{f(x_0)}} \cdot \frac{\textcolor{red}{f(x)} - \textcolor{red}{f(x_0)}}{x - x_0} \quad (\spadesuit)$$

Recall:

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{\color{red}f(x) - f(x_0)} \cdot \frac{\color{red}f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists $\implies$

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{\color{red}f(x) - f(x_0)} \cdot \frac{\color{red}f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists $\implies f$ continuous at x_0

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists $\implies f$ continuous at x_0
 $\implies f(x) \rightarrow f(x_0)$ as $x \rightarrow x_0$.

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists $\implies f$ continuous at x_0
 $\implies f(x) \rightarrow f(x_0)$ as $x \rightarrow x_0$.

Can we take the limit

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists $\implies f$ continuous at x_0
 $\implies f(x) \rightarrow f(x_0)$ as $x \rightarrow x_0$.

Can we take the limit as $x \rightarrow x_0$

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists $\implies f$ continuous at x_0
 $\implies f(x) \rightarrow f(x_0)$ as $x \rightarrow x_0$.

Can we take the limit as $x \rightarrow x_0$ and conclude that

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists $\implies f$ continuous at x_0
 $\implies f(x) \rightarrow f(x_0)$ as $x \rightarrow x_0$.

Can we take the limit as $x \rightarrow x_0$ and conclude that $(g \circ f)'(x_0) = g'(f(x_0)) \cdot f'(x_0)$?

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists $\implies f$ continuous at x_0
 $\implies f(x) \rightarrow f(x_0)$ as $x \rightarrow x_0$.

Can we take the limit as $x \rightarrow x_0$ and conclude that $(g \circ f)'(x_0) = g'(f(x_0)) \cdot f'(x_0)$?

■ What if $f(x) = 0$ for all x ?

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists $\implies f$ continuous at x_0
 $\implies f(x) \rightarrow f(x_0)$ as $x \rightarrow x_0$.

Can we take the limit as $x \rightarrow x_0$ and conclude that $(g \circ f)'(x_0) = g'(f(x_0)) \cdot f'(x_0)$?

- What if $f(x) = 0$ for all x ?
- What if f is a constant function?

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists $\implies f$ continuous at x_0
 $\implies f(x) \rightarrow f(x_0)$ as $x \rightarrow x_0$.

Can we take the limit as $x \rightarrow x_0$ and conclude that $(g \circ f)'(x_0) = g'(f(x_0)) \cdot f'(x_0)$?

- What if $f(x) = 0$ for all x ?
- What if f is a constant function?
- What if $f(x) = f(x_0)$ for some $x \neq x_0$?

Why the chain rule is plausible

The derivative of $g \circ f$ at x_0 is the limit as $x \rightarrow x_0$ of the difference quotient

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = \frac{g(f(x)) - g(f(x_0))}{f(x) - f(x_0)} \cdot \frac{f(x) - f(x_0)}{x - x_0} \quad (\spadesuit)$$

Recall: $f'(x_0)$ exists $\implies f$ continuous at x_0
 $\implies f(x) \rightarrow f(x_0)$ as $x \rightarrow x_0$.

Can we take the limit as $x \rightarrow x_0$ and conclude that $(g \circ f)'(x_0) = g'(f(x_0)) \cdot f'(x_0)$?

- What if $f(x) = 0$ for all x ?
- What if f is a constant function?
- What if $f(x) = f(x_0)$ for some $x \neq x_0$?
- Can we use $(\spadesuit)$ to prove the chain rule?

Poll

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Derivatives: Chain Rule**
- .

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$,

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$,

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$.

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

1 for all $\varepsilon > 0$

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

1 *for all $\varepsilon > 0$ there exists*

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

- 1** *for all $\varepsilon > 0$ there exists $\delta > 0$*

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

1 *for all $\varepsilon > 0$ there exists $\delta > 0$ such that*

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

1 *for all $\varepsilon > 0$ there exists $\delta > 0$ such that if*

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

- 1** *for all $\varepsilon > 0$ there exists $\delta > 0$ such that if $0 < |x - x_0| < \delta$*

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

1 *for all $\varepsilon > 0$ there exists $\delta > 0$ such that if $0 < |x - x_0| < \delta$ then*

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

- 1** *for all $\varepsilon > 0$ there exists $\delta > 0$ such that if $0 < |x - x_0| < \delta$ then $|f(x) - L| < \varepsilon$.*

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

- 1** *for all $\varepsilon > 0$ there exists $\delta > 0$ such that if $0 < |x - x_0| < \delta$ then $|f(x) - L| < \varepsilon$.*
- 2** *for every sequence $\{x_n\}$ of points in $I \setminus \{x_0\}$,*

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

- 1 for all $\varepsilon > 0$ there exists $\delta > 0$ such that if $0 < |x - x_0| < \delta$ then $|f(x) - L| < \varepsilon$.
- 2 for every sequence $\{x_n\}$ of points in $I \setminus \{x_0\}$,

$$\lim_{n \rightarrow \infty} x_n = x_0 \implies \lim_{n \rightarrow \infty} f(x_n) = L.$$

REMINDER: limits of functions

Theorem (Equivalence of ε - δ and sequence definitions of limits)

Let $a < x_0 < b$, $I = (a, b)$, and $f : I \setminus \{x_0\} \rightarrow \mathbb{R}$. Then the following two definitions of

$$\lim_{x \rightarrow x_0} f(x) = L$$

are equivalent:

- 1 for all $\varepsilon > 0$ there exists $\delta > 0$ such that if $0 < |x - x_0| < \delta$ then $|f(x) - L| < \varepsilon$.
- 2 for every sequence $\{x_n\}$ of points in $I \setminus \{x_0\}$,

$$\lim_{n \rightarrow \infty} x_n = x_0 \implies \lim_{n \rightarrow \infty} f(x_n) = L.$$

Note: The deleted neighbourhood $(I \setminus \{x_0\})$ can be replaced by any set on which f is defined and x_0 is an accumulation point.

Proof of the chain rule.

Proof of the chain rule.

- 1 Suppose there is an open interval I

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$.

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in ()

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get the **chain rule**.

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get the **chain rule**.
- 2 Next suppose that

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get the **chain rule**.
- 2 Next suppose that no open interval

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in () and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists.

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$.

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore,

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in () and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in $(\spadesuit)$ and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0,

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in () and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in () and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in () and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in () and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in () and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$.

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in () and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore,

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in () and we get the **chain rule**.
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists,

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in () and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#),

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) =$$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) = \lim_{n \rightarrow \infty} \frac{f(x_n) - f(x_0)}{x_n - x_0}$$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) = \lim_{n \rightarrow \infty} \frac{f(x_n) - f(x_0)}{x_n - x_0} =$$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) = \lim_{n \rightarrow \infty} \frac{f(x_n) - f(x_0)}{x_n - x_0} = \lim_{n \rightarrow \infty} \frac{0}{x_n - x_0}$$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) = \lim_{n \rightarrow \infty} \frac{f(x_n) - f(x_0)}{x_n - x_0} = \lim_{n \rightarrow \infty} \frac{0}{x_n - x_0} = 0.$$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) = \lim_{n \rightarrow \infty} \frac{f(x_n) - f(x_0)}{x_n - x_0} = \lim_{n \rightarrow \infty} \frac{0}{x_n - x_0} = 0.$$

We can also show that

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) = \lim_{n \rightarrow \infty} \frac{f(x_n) - f(x_0)}{x_n - x_0} = \lim_{n \rightarrow \infty} \frac{0}{x_n - x_0} = 0.$$

We can also show that $(g \circ f)'(x_0) = 0$

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) = \lim_{n \rightarrow \infty} \frac{f(x_n) - f(x_0)}{x_n - x_0} = \lim_{n \rightarrow \infty} \frac{0}{x_n - x_0} = 0.$$

We can also show that $(g \circ f)'(x_0) = 0$, using the sequence definition

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) = \lim_{n \rightarrow \infty} \frac{f(x_n) - f(x_0)}{x_n - x_0} = \lim_{n \rightarrow \infty} \frac{0}{x_n - x_0} = 0.$$

We can also show that $(g \circ f)'(x_0) = 0$, using the sequence definition on the [previous slide](#).

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) = \lim_{n \rightarrow \infty} \frac{f(x_n) - f(x_0)}{x_n - x_0} = \lim_{n \rightarrow \infty} \frac{0}{x_n - x_0} = 0.$$

We can also show that $(g \circ f)'(x_0) = 0$, using the sequence definition on the [previous slide](#). *Try to fill in this last detail*

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) = \lim_{n \rightarrow \infty} \frac{f(x_n) - f(x_0)}{x_n - x_0} = \lim_{n \rightarrow \infty} \frac{0}{x_n - x_0} = 0.$$

We can also show that $(g \circ f)'(x_0) = 0$, using the sequence definition on the [previous slide](#). *Try to fill in this last detail*, or look it up (TBB [§7.3.2, p. 411](#)).

Proof of the chain rule.

- 1 Suppose there is an open interval I , with $x_0 \in I$, and $f(x) \neq f(x_0)$ for all $x \in I \setminus \{x_0\}$. Then we can take the limit $x \rightarrow x_0$ in  and we get the [chain rule](#).
- 2 Next suppose that no open interval like the one hypothesized above exists. Then, in any open interval containing x_0 , there must be at least one point $x \neq x_0$ for which $f(x) = f(x_0)$. Therefore, we can construct a sequence of open intervals I_n , with lengths decreasing to 0, such that each I_n contains x_0 and a point $x_n \neq x_0$ with $f(x_n) = f(x_0)$. Therefore, since $f'(x_0)$ exists, and we recall the [previous slide](#), we can compute $f'(x_0)$ via

$$f'(x_0) = \lim_{n \rightarrow \infty} \frac{f(x_n) - f(x_0)}{x_n - x_0} = \lim_{n \rightarrow \infty} \frac{0}{x_n - x_0} = 0.$$

We can also show that $(g \circ f)'(x_0) = 0$, using the sequence definition on the [previous slide](#). *Try to fill in this last detail*, or look it up (TBB [§7.3.2, p. 411](#)).

Note: TBB's proof leaves out the proof that $f'(x_0) = 0$ in case 2 above. 

More on the derivative

More on the derivative

Theorem (Derivative at local extrema)

More on the derivative

Theorem (Derivative at local extrema)

Let $f : (a, b) \rightarrow \mathbb{R}$.

More on the derivative

Theorem (Derivative at local extrema)

Let $f : (a, b) \rightarrow \mathbb{R}$. If x is a maximum or minimum point of f in (a, b)

More on the derivative

Theorem (Derivative at local extrema)

Let $f : (a, b) \rightarrow \mathbb{R}$. If x is a maximum or minimum point of f in (a, b) , and f is differentiable at x

More on the derivative

Theorem (Derivative at local extrema)

Let $f : (a, b) \rightarrow \mathbb{R}$. If x is a maximum or minimum point of f in (a, b) , and f is differentiable at x , then

More on the derivative

Theorem (Derivative at local extrema)

Let $f : (a, b) \rightarrow \mathbb{R}$. If x is a maximum or minimum point of f in (a, b) , and f is differentiable at x , then $f'(x) = 0$.

More on the derivative

Theorem (Derivative at local extrema)

Let $f : (a, b) \rightarrow \mathbb{R}$. If x is a maximum or minimum point of f in (a, b) , and f is differentiable at x , then $f'(x) = 0$.

Note: f need not be differentiable or even continuous at other points.


More on the derivative

Theorem (Derivative at local extrema)

Let $f : (a, b) \rightarrow \mathbb{R}$. If x is a maximum or minimum point of f in (a, b) , and f is differentiable at x , then $f'(x) = 0$.

Note: f need not be differentiable or even continuous at other points.

Idea for proof:



More on the derivative

More on the derivative

Proof that the derivative vanishes at local extrema.

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$,

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\underline{f(x+h) - f(x)}$$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h}$$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq$$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0$$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq$$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x ,

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x ,

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate the limits

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate the limits as $h \rightarrow 0$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate the limits as $h \rightarrow 0$ to obtain

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate the limits as $h \rightarrow 0$ to obtain

$$f'_+(x)$$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate the limits as $h \rightarrow 0$ to obtain

$$f'_+(x) \leq$$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate the limits as $h \rightarrow 0$ to obtain

$$f'_+(x) \leq 0$$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate the limits as $h \rightarrow 0$ to obtain

$$f'_+(x) \leq 0 \leq f'_-(x).$$

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate the limits as $h \rightarrow 0$ to obtain

$$f'_+(x) \leq 0 \leq f'_-(x).$$

But since f is differentiable at x ,

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate the limits as $h \rightarrow 0$ to obtain

$$f'_+(x) \leq 0 \leq f'_-(x).$$

But since f is differentiable at x , the left and right derivatives must be equal,

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate the limits as $h \rightarrow 0$ to obtain

$$f'_+(x) \leq 0 \leq f'_-(x).$$

But since f is differentiable at x , the left and right derivatives must be equal, hence

More on the derivative

Proof that the derivative vanishes at local extrema.

If f has a local maximum at $x \in (a, b)$, then for sufficiently small $h > 0$ we must have

$$\frac{f(x+h) - f(x)}{h} \leq 0 \leq \frac{f(x) - f(x-h)}{h}$$

Since f is differentiable at x , it is left and right differentiable at x , so we can evaluate the limits as $h \rightarrow 0$ to obtain

$$f'_+(x) \leq 0 \leq f'_-(x).$$

But since f is differentiable at x , the left and right derivatives must be equal, hence $f'(x) = 0$. □



Mathematics
and Statistics

$$\int_M d\omega = \int_{\partial M} \omega$$

Mathematics 3A03 Real Analysis I

Instructor: David Earn

Lecture 4
Differentiation III
Monday 13 January 2025

Announcements

Announcements

- In-Class polls:

Announcements

- In-Class polls: if you do not have a device that enables you to participate in polls in class,

Announcements

- In-Class polls: if you do not have a device that enables you to participate in polls in class, or if any other issue prevents you from participating in polls,

Announcements

- In-Class polls: if you do not have a device that enables you to participate in polls in class, or if any other issue prevents you from participating in polls, please let me know by e-mail.

Announcements

- In-Class polls: if you do not have a device that enables you to participate in polls in class, or if any other issue prevents you from participating in polls, please let me know by e-mail.
- The [online syllabus](#) has been revised to account for the changes in how the course will be evaluated (see [slides from Lecture 3](#)).

Announcements

- In-Class polls: if you do not have a device that enables you to participate in polls in class, or if any other issue prevents you from participating in polls, please let me know by e-mail.
- The [online syllabus](#) has been revised to account for the changes in how the course will be evaluated (see [slides from Lecture 3](#)). In particular, the statement about AI use has been changed to:

Announcements

- In-Class polls: if you do not have a device that enables you to participate in polls in class, or if any other issue prevents you from participating in polls, please let me know by e-mail.
- The [online syllabus](#) has been revised to account for the changes in how the course will be evaluated (see [slides from Lecture 3](#)). In particular, the statement about AI use has been changed to:

Generative AI: Unrestricted Use

Students may use generative AI throughout this course in whatever way enhances their learning; no special documentation or citation is required. Note that access to generative AI will **not** be available during tests or exams.

Announcements

- In-Class polls: if you do not have a device that enables you to participate in polls in class, or if any other issue prevents you from participating in polls, please let me know by e-mail.
- The [online syllabus](#) has been revised to account for the changes in how the course will be evaluated (see [slides from Lecture 3](#)). In particular, the statement about AI use has been changed to:

Generative AI: Unrestricted Use

Students may use generative AI throughout this course in whatever way enhances their learning; no special documentation or citation is required. Note that access to generative AI will **not** be available during tests or exams.

- The [course web site](#) has been updated to reflect the changes in how the course will be evaluated.

Assignment 1

Assignment 1

- [Assignment 1](#) has been posted on the course web site.

Last time...

Last time...

- Discussed algebra of derivatives and chain rule.

Last time...

- Discussed algebra of derivatives and chain rule.
- Proved the chain rule.

Last time...

- Discussed algebra of derivatives and chain rule.
- Proved the chain rule.
- Proved that that derivative is zero at extrema.

The Mean Value Theorem

The Mean Value Theorem

Theorem (Rolle's theorem)

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b)

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies$

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$.

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$.

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b)

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints,

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b .

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$,

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant.

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0$

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0 \forall x \in (a, b)$. $\square$

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0 \forall x \in (a, b)$. $\square$

Theorem (Mean value theorem)

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0 \forall x \in (a, b)$. $\square$

Theorem (Mean value theorem)

If f is continuous on $[a, b]$

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0 \forall x \in (a, b)$. $\square$

Theorem (Mean value theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b)

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0 \forall x \in (a, b)$. $\square$

Theorem (Mean value theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) then

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0 \forall x \in (a, b)$. $\square$

Theorem (Mean value theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) then there exists

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0 \forall x \in (a, b)$. $\square$

Theorem (Mean value theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) then there exists $x \in (a, b)$ such that

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0 \forall x \in (a, b)$. $\square$

Theorem (Mean value theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) then there exists $x \in (a, b)$ such that

$$f'(x) =$$

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0 \forall x \in (a, b)$. $\square$

Theorem (Mean value theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) then there exists $x \in (a, b)$ such that

$$f'(x) = \frac{f(b) - f(a)}{b - a}$$

The Mean Value Theorem

Theorem (Rolle's theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) , and $f(a) = f(b)$, then there exists $x \in (a, b)$ such that $f'(x) = 0$.

Proof.

f continuous on $[a, b] \implies f$ has a max and min value on $[a, b]$. If either a max or min occurs at $x \in (a, b)$ then $f'(x) = 0$. If no max or min occurs in (a, b) then they must both occur at the endpoints, a and b . But $f(a) = f(b)$, so f is constant. Hence $f'(x) = 0 \forall x \in (a, b)$. $\square$

Theorem (Mean value theorem)

If f is continuous on $[a, b]$ and differentiable on (a, b) then there exists $x \in (a, b)$ such that

$$f'(x) = \frac{f(b) - f(a)}{b - a}.$$


The Mean Value Theorem

The Mean Value Theorem

Idea for proof:


The Mean Value Theorem

Idea for proof:



The Mean Value Theorem


Idea for proof:



Proof.

The Mean Value Theorem

Idea for proof:




Proof.

Apply Rolle's theorem to

The Mean Value Theorem

Idea for proof:




Proof.

Apply Rolle's theorem to

$$h(x)$$

The Mean Value Theorem

Idea for proof:




Proof.

Apply Rolle's theorem to

$$h(x) =$$

The Mean Value Theorem

Idea for proof:




Proof.

Apply Rolle's theorem to

$$h(x) = f(x)$$

The Mean Value Theorem

Idea for proof:




Proof.

Apply Rolle's theorem to

$$h(x) = f(x) -$$

The Mean Value Theorem

Idea for proof:




Proof.

Apply Rolle's theorem to

$$h(x) = f(x) - \left[f(a) \right]$$

The Mean Value Theorem

Idea for proof:




Proof.

Apply Rolle's theorem to

$$h(x) = f(x) - \left[f(a) + \right.$$

The Mean Value Theorem

Idea for proof:




Proof.

Apply Rolle's theorem to

$$h(x) = f(x) - \left[f(a) + \left(\frac{f(b) - f(a)}{b - a} \right) (x - a) \right]$$

The Mean Value Theorem

Idea for proof:




Proof.

Apply Rolle's theorem to

$$h(x) = f(x) - \left[f(a) + \left(\frac{f(b) - f(a)}{b - a} \right) (x - a) \right]$$

The Mean Value Theorem

Idea for proof:



Proof.

Apply Rolle's theorem to

$$h(x) = f(x) - \left[f(a) + \left(\frac{f(b) - f(a)}{b - a} \right) (x - a) \right].$$



The Mean Value Theorem

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$.

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$,

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#)

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} =$$

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(x_*).$$

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(x_*).$$

But $x_2 - x_1 > 0$

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(x_*).$$

But $x_2 - x_1 > 0$ and since $x_* \in I$,

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(x_*).$$

But $x_2 - x_1 > 0$ and since $x_* \in I$, we know

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(x_*).$$

But $x_2 - x_1 > 0$ and since $x_* \in I$, we know $f'(x_*) > 0$.

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(x_*).$$

But $x_2 - x_1 > 0$ and since $x_* \in I$, we know $f'(x_*) > 0$.

$\therefore$

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(x_*).$$

But $x_2 - x_1 > 0$ and since $x_* \in I$, we know $f'(x_*) > 0$.

$\therefore f(x_2) - f(x_1) > 0$,

The Mean Value Theorem

Example

$f'(x) > 0$ on an interval $I \implies f$ strictly increasing on I .

Proof:

Suppose $x_1, x_2 \in I$ and $x_1 < x_2$. We must show $f(x_1) < f(x_2)$.

Since $f'(x)$ exists for all $x \in I$, f is certainly differentiable on the closed subinterval $[x_1, x_2]$.

Hence by the [Mean Value Theorem](#) $\exists x_* \in (x_1, x_2)$ such that


$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(x_*).$$

But $x_2 - x_1 > 0$ and since $x_* \in I$, we know $f'(x_*) > 0$.


$\therefore f(x_2) - f(x_1) > 0$, i.e., $f(x_1) < f(x_2)$. □

REMINDER: Intermediate Value Property

REMINDER: Intermediate Value Property




REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have the *intermediate value property (IVP)*


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have the *intermediate value property (IVP)* on I


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have the **intermediate value property (IVP)** on I iff


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have the **intermediate value property (IVP)** on I iff for each $a, b \in I$


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have the **intermediate value property (IVP)** on I iff for each $a, b \in I$ with $f(a) \neq f(b)$


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have the **intermediate value property (IVP)** on I iff for each $a, b \in I$ with $f(a) \neq f(b)$, and for each d


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have the **intermediate value property (IVP)** on I iff for each $a, b \in I$ with $f(a) \neq f(b)$, and for each d between $f(a)$ and $f(b)$,


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have the **intermediate value property (IVP)** on I iff for each $a, b \in I$ with $f(a) \neq f(b)$, and for each d between $f(a)$ and $f(b)$, there exists c


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have the **intermediate value property (IVP)** on I iff for each $a, b \in I$ with $f(a) \neq f(b)$, and for each d between $f(a)$ and $f(b)$, there exists c between a and b


REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have the **intermediate value property (IVP)** on I iff for each $a, b \in I$ with $f(a) \neq f(b)$, and for each d between $f(a)$ and $f(b)$, there exists c between a and b for which

REMINDER: Intermediate Value Property



Definition (Intermediate Value Property (IVP))

A function f defined on an interval I is said to have the **intermediate value property (IVP)** on I iff for each $a, b \in I$ with $f(a) \neq f(b)$, and for each d between $f(a)$ and $f(b)$, there exists c between a and b for which $f(c) = d$.

REMINDER: Intermediate Value Property

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

If f is continuous

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

If f is continuous on an interval I

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

If f is continuous on an interval I then

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

If f is continuous on an interval I then f has

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

*If f is continuous on an interval I then f has the **intermediate value property (IVP)***

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

*If f is continuous on an interval I then f has the **intermediate value property (IVP)** on I .*

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

*If f is continuous on an interval I then f has the **intermediate value property (IVP)** on I .*

Note:

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

*If f is continuous on an interval I then f has the **intermediate value property (IVP)** on I .*

Note: The interval I in the statement of the IVT

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

*If f is continuous on an interval I then f has the **intermediate value property (IVP)** on I .*

Note: The interval I in the statement of the IVT does not have to be closed

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

*If f is continuous on an interval I then f has the **intermediate value property (IVP)** on I .*

Note: The interval I in the statement of the IVT does not have to be closed and it does not have to be bounded.

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

*If f is continuous on an interval I then f has the **intermediate value property (IVP)** on I .*

Note: The interval I in the statement of the IVT does not have to be closed and it does not have to be bounded.

Unlike the **extreme value theorem**,

REMINDER: Intermediate Value Property

Theorem (Intermediate Value Theorem (IVT))

*If f is continuous on an interval I then f has the **intermediate value property (IVP)** on I .*

Note: The interval I in the statement of the IVT does not have to be closed and it does not have to be bounded.

Unlike the **extreme value theorem**, the IVT is not a theorem about functions defined on closed and bounded intervals.

Poll

- Go to
https://www.childsmath.ca/childsa/forms/main_login.php
- Click on **Math 3A03**
- Click on **Take Class Poll**
- Fill in poll **Derivatives: IVP and derivatives**
- .

Intermediate Value Property

Intermediate Value Property

Question:

Intermediate Value Property

Question: If a function has the IVP on an interval I , must it be continuous on I ?

Intermediate Value Property

Question: If a function has the IVP on an interval I , must it be continuous on I ?

Example


$$f(x) = \begin{cases} \sin \frac{1}{x} & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Intermediate Value Property

Question: If a function has the **IVP** on an interval I , must it be **continuous** on I ?

Example

$$f(x) = \begin{cases} \sin \frac{1}{x} & x \neq 0, \\ 0 & x = 0. \end{cases}$$



Intermediate value property for derivatives

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f'

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Notes:

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Notes:

- It is f' ,

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Notes:

- It is f' , not f ,

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Notes:

- It is f' , not f , that is claimed to have the intermediate value property

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Notes:

- It is f' , not f , that is claimed to have the intermediate value property in Darboux's theorem.

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Notes:

- It is f' , not f , that is claimed to have the intermediate value property in Darboux's theorem. This theorem does not

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem**

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Notes:

- It is f' , not f , that is claimed to have the intermediate value property in Darboux's theorem. This theorem does not follow from the standard intermediate value theorem because the derivative f'

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Notes:

- It is f' , not f , that is claimed to have the intermediate value property in Darboux's theorem. This theorem does not follow from the standard intermediate value theorem because the derivative f' is not necessarily

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Notes:

- It is f' , not f , that is claimed to have the intermediate value property in Darboux's theorem. This theorem does not follow from the standard intermediate value theorem because the derivative f' is not necessarily continuous.

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the **intermediate value property** on I

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the **intermediate value property** on I then it is impossible

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the **intermediate value property** on I then it is impossible that it is the derivative

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the **intermediate value property** on I then it is impossible that it is the derivative of any function

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the **intermediate value property** on I then it is impossible that it is the derivative of any function on I .

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the **intermediate value property** on I then it is impossible that it is the derivative of any function on I .
- Darboux's theorem implies

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the **intermediate value property** on I then it is impossible that it is the derivative of any function on I .
- Darboux's theorem implies that a derivative

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the **intermediate value property** on I then it is impossible that it is the derivative of any function on I .
- Darboux's theorem implies that a derivative cannot have

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the **intermediate value property** on I then it is impossible that it is the derivative of any function on I .
- Darboux's theorem implies that a derivative cannot have jump or removable discontinuities.

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

*If f is differentiable on an interval I then its derivative f' has the **intermediate value property** on I .*

Notes:

- It is f' , not f , that is claimed to have the **intermediate value property** in Darboux's theorem. This theorem does not follow from the standard **intermediate value theorem** because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the **intermediate value property** on I then it is impossible that it is the derivative of any function on I .
- Darboux's theorem implies that a derivative cannot have jump or removable discontinuities. Any discontinuity of a derivative

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Notes:

- It is f' , not f , that is claimed to have the intermediate value property in Darboux's theorem. This theorem does not follow from the standard intermediate value theorem because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the intermediate value property on I then it is impossible that it is the derivative of any function on I .
- Darboux's theorem implies that a derivative cannot have jump or removable discontinuities. Any discontinuity of a derivative must be essential.

Intermediate value property for derivatives

Theorem (Darboux's Theorem: IVP for derivatives)

If f is differentiable on an interval I then its derivative f' has the intermediate value property on I .

Notes:

- It is f' , not f , that is claimed to have the intermediate value property in Darboux's theorem. This theorem does not follow from the standard intermediate value theorem because the derivative f' is not necessarily continuous.
- *Equivalent (contrapositive) statement of Darboux's theorem:*
If a function does not have the intermediate value property on I then it is impossible that it is the derivative of any function on I .
- Darboux's theorem implies that a derivative cannot have jump or removable discontinuities. Any discontinuity of a derivative must be essential. Recall example of a discontinuous function with IVP.

Intermediate value property for derivatives

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$,

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the **Extreme Value Theorem** implies that f attains its minimum at some point $x \in [a, b]$.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the **Extreme Value Theorem** implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the **Extreme Value Theorem** implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$). Therefore, $x \in (a, b)$.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the **Extreme Value Theorem** implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$). Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b)

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the [theorem on the derivative at local extrema](#)

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the [theorem on the derivative at local extrema](#), we must have

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the [theorem on the derivative at local extrema](#), we must have $f'(x) = 0$.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the **Extreme Value Theorem** implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the **theorem on the derivative at local extrema**, we must have $f'(x) = 0$. Now suppose more generally that $f'(a) < K < f'(b)$.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the **Extreme Value Theorem** implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the **theorem on the derivative at local extrema**, we must have $f'(x) = 0$.

Now suppose more generally that $f'(a) < K < f'(b)$. Let

$g(x) = f(x) - Kx$.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the [theorem on the derivative at local extrema](#), we must have $f'(x) = 0$.

Now suppose more generally that $f'(a) < K < f'(b)$. Let

$g(x) = f(x) - Kx$. Then g is differentiable on I and $g'(x) = f'(x) - K$ for all $x \in I$.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the [theorem on the derivative at local extrema](#), we must have $f'(x) = 0$.

Now suppose more generally that $f'(a) < K < f'(b)$. Let

$g(x) = f(x) - Kx$. Then g is differentiable on I and $g'(x) = f'(x) - K$ for all $x \in I$. In addition, $g'(a) = f'(a) - K < 0$

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the **Extreme Value Theorem** implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the **theorem on the derivative at local extrema**, we must have $f'(x) = 0$.

Now suppose more generally that $f'(a) < K < f'(b)$. Let

$g(x) = f(x) - Kx$. Then g is differentiable on I and $g'(x) = f'(x) - K$ for all $x \in I$. In addition, $g'(a) = f'(a) - K < 0$ and $g'(b) = f'(b) - K > 0$

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the [theorem on the derivative at local extrema](#), we must have $f'(x) = 0$.

Now suppose more generally that $f'(a) < K < f'(b)$. Let

$g(x) = f(x) - Kx$. Then g is differentiable on I and $g'(x) = f'(x) - K$ for all $x \in I$. In addition, $g'(a) = f'(a) - K < 0$ and $g'(b) = f'(b) - K > 0$, so by the argument above

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the [theorem on the derivative at local extrema](#), we must have $f'(x) = 0$.

Now suppose more generally that $f'(a) < K < f'(b)$. Let

$g(x) = f(x) - Kx$. Then g is differentiable on I and $g'(x) = f'(x) - K$ for all $x \in I$. In addition, $g'(a) = f'(a) - K < 0$ and

$g'(b) = f'(b) - K > 0$, so by the argument above, $\exists x \in (a, b)$ such that

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the [theorem on the derivative at local extrema](#), we must have $f'(x) = 0$.

Now suppose more generally that $f'(a) < K < f'(b)$. Let

$g(x) = f(x) - Kx$. Then g is differentiable on I and $g'(x) = f'(x) - K$ for all $x \in I$. In addition, $g'(a) = f'(a) - K < 0$ and

$g'(b) = f'(b) - K > 0$, so by the argument above, $\exists x \in (a, b)$ such that $g'(x) = 0$

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the [theorem on the derivative at local extrema](#), we must have $f'(x) = 0$.

Now suppose more generally that $f'(a) < K < f'(b)$. Let

$g(x) = f(x) - Kx$. Then g is differentiable on I and $g'(x) = f'(x) - K$ for all $x \in I$. In addition, $g'(a) = f'(a) - K < 0$ and

$g'(b) = f'(b) - K > 0$, so by the argument above, $\exists x \in (a, b)$ such that $g'(x) = 0$, i.e., $f'(x) - K = 0$

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the [Extreme Value Theorem](#) implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the [theorem on the derivative at local extrema](#), we must have $f'(x) = 0$.

Now suppose more generally that $f'(a) < K < f'(b)$. Let

$g(x) = f(x) - Kx$. Then g is differentiable on I and $g'(x) = f'(x) - K$ for all $x \in I$. In addition, $g'(a) = f'(a) - K < 0$ and

$g'(b) = f'(b) - K > 0$, so by the argument above, $\exists x \in (a, b)$ such that $g'(x) = 0$, i.e., $f'(x) - K = 0$, i.e., $f'(x) = K$.

Intermediate value property for derivatives

Proof of Darboux's Theorem.

Consider $a, b \in I$ with $a < b$.

Suppose first that $f'(a) < 0 < f'(b)$. We will show $\exists x \in (a, b)$ such that $f'(x) = 0$. Since f' exists on $[a, b]$, we must have f continuous on $[a, b]$, so the **Extreme Value Theorem** implies that f attains its minimum at some point $x \in [a, b]$. This minimum point cannot be an endpoint of $[a, b]$ ($x \neq a$ because $f'(a) < 0$ and $x \neq b$ because $f'(b) > 0$).

Therefore, $x \in (a, b)$. But f is differentiable everywhere in (a, b) , so, by the **theorem on the derivative at local extrema**, we must have $f'(x) = 0$.

Now suppose more generally that $f'(a) < K < f'(b)$. Let

$g(x) = f(x) - Kx$. Then g is differentiable on I and $g'(x) = f'(x) - K$ for all $x \in I$. In addition, $g'(a) = f'(a) - K < 0$ and

$g'(b) = f'(b) - K > 0$, so by the argument above, $\exists x \in (a, b)$ such that $g'(x) = 0$, i.e., $f'(x) - K = 0$, i.e., $f'(x) = K$.

The case $f'(a) > K > f'(b)$ is similar. □

Intermediate value property for derivatives

Intermediate value property for derivatives

Example ($f'(x) \neq 0 \forall x \in I \implies f \nearrow$ or $\searrow$ on I)

If f is differentiable on an interval I and $f'(x) \neq 0$ for all $x \in I$ then f is either increasing or decreasing on the entire interval I .

Intermediate value property for derivatives

Example $(f'(x) \neq 0 \ \forall x \in I \implies f \nearrow \text{ or } \searrow \text{ on } I)$

If f is differentiable on an interval I and $f'(x) \neq 0$ for all $x \in I$ then f is either increasing or decreasing on the entire interval I .

Proof:

Intermediate value property for derivatives

Example $(f'(x) \neq 0 \forall x \in I \implies f \nearrow \text{ or } \searrow \text{ on } I)$

If f is differentiable on an interval I and $f'(x) \neq 0$ for all $x \in I$ then f is either increasing or decreasing on the entire interval I .

Proof:

Suppose $\exists a, b \in I$ such that $f'(a) < 0$ and $f'(b) > 0$.

Intermediate value property for derivatives

Example ($f'(x) \neq 0 \forall x \in I \implies f \nearrow$ or $\searrow$ on I)

If f is differentiable on an interval I and $f'(x) \neq 0$ for all $x \in I$ then f is either increasing or decreasing on the entire interval I .

Proof:

Suppose $\exists a, b \in I$ such that $f'(a) < 0$ and $f'(b) > 0$.

Then, from [Darboux's theorem](#), $\exists c \in I$ such that $f'(c) = 0$.

Intermediate value property for derivatives

Example ($f'(x) \neq 0 \forall x \in I \implies f \nearrow$ or $\searrow$ on I)

If f is differentiable on an interval I and $f'(x) \neq 0$ for all $x \in I$ then f is either increasing or decreasing on the entire interval I .

Proof:

Suppose $\exists a, b \in I$ such that $f'(a) < 0$ and $f'(b) > 0$.

Then, from [Darboux's theorem](#), $\exists c \in I$ such that $f'(c) = 0$. $\Rightarrow \Leftarrow$

Intermediate value property for derivatives

Example $(f'(x) \neq 0 \forall x \in I \implies f \nearrow \text{ or } \searrow \text{ on } I)$

If f is differentiable on an interval I and $f'(x) \neq 0$ for all $x \in I$ then f is either increasing or decreasing on the entire interval I .

Proof:

Suppose $\exists a, b \in I$ such that $f'(a) < 0$ and $f'(b) > 0$.

Then, from [Darboux's theorem](#), $\exists c \in I$ such that $f'(c) = 0$. $\Rightarrow \Leftarrow$

$\therefore$ Either " $\exists a \in I \nmid f'(a) < 0$ " is FALSE
or " $\exists b \in I \nmid f'(b) > 0$ " is FALSE.

Intermediate value property for derivatives

Example $(f'(x) \neq 0 \ \forall x \in I \implies f \nearrow \text{ or } \searrow \text{ on } I)$

If f is differentiable on an interval I and $f'(x) \neq 0$ for all $x \in I$ then f is either increasing or decreasing on the entire interval I .

Proof:

Suppose $\exists a, b \in I$ such that $f'(a) < 0$ and $f'(b) > 0$.

Then, from [Darboux's theorem](#), $\exists c \in I$ such that $f'(c) = 0$. $\Rightarrow \Leftarrow$

$\therefore$ Either " $\exists a \in I \nmid f'(a) < 0$ " is FALSE
or " $\exists b \in I \nmid f'(b) > 0$ " is FALSE.

$\therefore$ Since we know $f'(x) \neq 0 \ \forall x \in I$, it must be that
either $f'(x) > 0 \ \forall x \in I$ or $f'(x) < 0 \ \forall x \in I$

Intermediate value property for derivatives

Example $(f'(x) \neq 0 \ \forall x \in I \implies f \nearrow \text{ or } \searrow \text{ on } I)$

If f is differentiable on an interval I and $f'(x) \neq 0$ for all $x \in I$ then f is either increasing or decreasing on the entire interval I .

Proof:

Suppose $\exists a, b \in I$ such that $f'(a) < 0$ and $f'(b) > 0$.

Then, from [Darboux's theorem](#), $\exists c \in I$ such that $f'(c) = 0$. $\Rightarrow \Leftarrow$

$\therefore$ Either " $\exists a \in I \nmid f'(a) < 0$ " is FALSE
or " $\exists b \in I \nmid f'(b) > 0$ " is FALSE.

$\therefore$ Since we know $f'(x) \neq 0 \ \forall x \in I$, it must be that
either $f'(x) > 0 \ \forall x \in I$ or $f'(x) < 0 \ \forall x \in I$,
i.e., either f is increasing on I or decreasing on I . □

The Derivative of an Inverse

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

Proof: By definition, the inverse function satisfies

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

Proof: By definition, the inverse function satisfies

$$f(f^{-1}(y)) = y.$$

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

Proof: By definition, the inverse function satisfies

$$f(f^{-1}(y)) = y.$$

Suppose that f^{-1} is differentiable at y .

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

Proof: By definition, the inverse function satisfies

$$f(f^{-1}(y)) = y.$$

Suppose that f^{-1} is differentiable at y . Then, by the [Chain Rule](#),

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

Proof: By definition, the inverse function satisfies

$$f(f^{-1}(y)) = y.$$

Suppose that f^{-1} is differentiable at y . Then, by the [Chain Rule](#),

$$f'(f^{-1}(y)) \cdot (f^{-1})'(y) = 1.$$

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

Proof: By definition, the inverse function satisfies

$$f(f^{-1}(y)) = y.$$

Suppose that f^{-1} is differentiable at y . Then, by the [Chain Rule](#),

$$f'(f^{-1}(y)) \cdot (f^{-1})'(y) = 1.$$

But $f^{-1}(y) = x$,

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

Proof: By definition, the inverse function satisfies

$$f(f^{-1}(y)) = y.$$

Suppose that f^{-1} is differentiable at y . Then, by the [Chain Rule](#),

$$f'(f^{-1}(y)) \cdot (f^{-1})'(y) = 1.$$

But $f^{-1}(y) = x$, and $f'(x) = 0$,

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

Proof: By definition, the inverse function satisfies

$$f(f^{-1}(y)) = y.$$

Suppose that f^{-1} is differentiable at y . Then, by the [Chain Rule](#),

$$f'(f^{-1}(y)) \cdot (f^{-1})'(y) = 1.$$

But $f^{-1}(y) = x$, and $f'(x) = 0$, so

$$0 \cdot (f^{-1})'(y) = 1,$$

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

Proof: By definition, the inverse function satisfies

$$f(f^{-1}(y)) = y.$$

Suppose that f^{-1} is differentiable at y . Then, by the [Chain Rule](#),

$$f'(f^{-1}(y)) \cdot (f^{-1})'(y) = 1.$$

But $f^{-1}(y) = x$, and $f'(x) = 0$, so

$$0 \cdot (f^{-1})'(y) = 1,$$

which is impossible!

The Derivative of an Inverse

Example (Sufficient condition for *non*-differentiable inverse)

Suppose f is continuous and one-to-one on an interval I . If $x \in I$ and $f'(x) = 0$ then f^{-1} is not differentiable at $y = f(x)$.

Proof: By definition, the inverse function satisfies

$$f(f^{-1}(y)) = y.$$

Suppose that f^{-1} is differentiable at y . Then, by the **Chain Rule**,

$$f'(f^{-1}(y)) \cdot (f^{-1})'(y) = 1.$$

But $f^{-1}(y) = x$, and $f'(x) = 0$, so

$$0 \cdot (f^{-1})'(y) = 1,$$

which is impossible! $\Rightarrow \Leftarrow$.



The Derivative of an Inverse

The Derivative of an Inverse

Theorem (Inverse function theorem)

The Derivative of an Inverse

Theorem (Inverse function theorem)

If f is differentiable on an interval I

The Derivative of an Inverse

Theorem (Inverse function theorem)

If f is differentiable on an interval I and $f'(x) \neq 0 \forall x \in I$

The Derivative of an Inverse

Theorem (Inverse function theorem)

If f is differentiable on an interval I and $f'(x) \neq 0 \forall x \in I$, then

The Derivative of an Inverse

Theorem (Inverse function theorem)

If f is differentiable on an interval I and $f'(x) \neq 0 \forall x \in I$, then

- 1** *f is one-to-one on I ;*

The Derivative of an Inverse

Theorem (Inverse function theorem)

If f is differentiable on an interval I and $f'(x) \neq 0 \forall x \in I$, then

- 1** *f is one-to-one on I ;*
- 2** *f^{-1} is differentiable on $J = f(I)$;*

The Derivative of an Inverse

Theorem (Inverse function theorem)

If f is differentiable on an interval I and $f'(x) \neq 0 \forall x \in I$, then

- 1** *f is one-to-one on I ;*
- 2** *f^{-1} is differentiable on $J = f(I)$;*
- 3** $(f^{-1})'(f(x)) = \frac{1}{f'(x)}$ *for all $x \in I$*

The Derivative of an Inverse

Theorem (Inverse function theorem)

If f is differentiable on an interval I and $f'(x) \neq 0 \forall x \in I$, then

- 1** *f is one-to-one on I ;*
- 2** *f^{-1} is differentiable on $J = f(I)$;*
- 3** $(f^{-1})'(f(x)) = \frac{1}{f'(x)}$ for all $x \in I$,
i.e., $(f^{-1})'(y) = \frac{1}{f'((f^{-1}(y)))}$ for all $y \in J$.

(TBB Theorem 7.32, p. 445)

The Derivative of an Inverse

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

- Darboux's theorem $\implies f \nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

- Darboux's theorem $\implies f \nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I
- If $y = f(x)$ and $y_0 = f(x_0)$
then

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

■ Darboux's theorem $\implies f \nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I

■ If $y = f(x)$ and $y_0 = f(x_0)$

then $x = f^{-1}(y)$ and $x_0 = f^{-1}(y_0)$

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

■ Darboux's theorem $\implies f \nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I

■ If $y = f(x)$ and $y_0 = f(x_0)$

then $x = f^{-1}(y)$ and $x_0 = f^{-1}(y_0)$,

$$\text{so } \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} =$$

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

- Darboux's theorem $\implies f \nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I
- If $y = f(x)$ and $y_0 = f(x_0)$

then $x = f^{-1}(y)$ and $x_0 = f^{-1}(y_0)$,

$$\text{so } \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} = \frac{x - x_0}{f(x) - f(x_0)}$$

$$=$$

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

- Darboux's theorem $\implies f \nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I
- If $y = f(x)$ and $y_0 = f(x_0)$

then $x = f^{-1}(y)$ and $x_0 = f^{-1}(y_0)$,

$$\begin{aligned} \text{so } \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} &= \frac{x - x_0}{f(x) - f(x_0)} \\ &= \frac{1}{\frac{f(x) - f(x_0)}{x - x_0}}. \end{aligned}$$

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

- Darboux's theorem $\implies f \nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I
- If $y = f(x)$ and $y_0 = f(x_0)$

then $x = f^{-1}(y)$ and $x_0 = f^{-1}(y_0)$,

$$\begin{aligned} \text{so } \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} &= \frac{x - x_0}{f(x) - f(x_0)} \\ &= \frac{1}{\frac{f(x) - f(x_0)}{x - x_0}}. \end{aligned}$$

- Since f continuous at x_0 , we know $x \rightarrow x_0 \implies y \rightarrow y_0$.

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

- Darboux's theorem $\implies f$ $\nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I
- If $y = f(x)$ and $y_0 = f(x_0)$

then $x = f^{-1}(y)$ and $x_0 = f^{-1}(y_0)$,

$$\begin{aligned} \text{so } \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} &= \frac{x - x_0}{f(x) - f(x_0)} \\ &= \frac{1}{\frac{f(x) - f(x_0)}{x - x_0}}. \end{aligned}$$

- Since f continuous at x_0 , we know $x \rightarrow x_0 \implies y \rightarrow y_0$.
- But we need $y \rightarrow y_0 \implies x \rightarrow x_0$

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

- Darboux's theorem $\implies f$ $\nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I
- If $y = f(x)$ and $y_0 = f(x_0)$

then $x = f^{-1}(y)$ and $x_0 = f^{-1}(y_0)$,

$$\begin{aligned} \text{so } \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} &= \frac{x - x_0}{f(x) - f(x_0)} \\ &= \frac{1}{\frac{f(x) - f(x_0)}{x - x_0}}. \end{aligned}$$

- Since f continuous at x_0 , we know $x \rightarrow x_0 \implies y \rightarrow y_0$.
- But we need $y \rightarrow y_0 \implies x \rightarrow x_0$, i.e., f^{-1} continuous at y_0 .

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

- Darboux's theorem $\implies f \nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I
- If $y = f(x)$ and $y_0 = f(x_0)$

then $x = f^{-1}(y)$ and $x_0 = f^{-1}(y_0)$,

$$\begin{aligned} \text{so } \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} &= \frac{x - x_0}{f(x) - f(x_0)} \\ &= \frac{1}{\frac{f(x) - f(x_0)}{x - x_0}}. \end{aligned}$$

- Since f continuous at x_0 , we know $x \rightarrow x_0 \implies y \rightarrow y_0$.
- But we need $y \rightarrow y_0 \implies x \rightarrow x_0$, i.e., f^{-1} continuous at y_0 .
- In fact, f continuous and either $\nearrow$ or $\searrow$ on $I \implies$

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

- Darboux's theorem $\implies f \nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I
- If $y = f(x)$ and $y_0 = f(x_0)$

then $x = f^{-1}(y)$ and $x_0 = f^{-1}(y_0)$,

$$\begin{aligned} \text{so } \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} &= \frac{x - x_0}{f(x) - f(x_0)} \\ &= \frac{1}{\frac{f(x) - f(x_0)}{x - x_0}}. \end{aligned}$$

- Since f continuous at x_0 , we know $x \rightarrow x_0 \implies y \rightarrow y_0$.
- But we need $y \rightarrow y_0 \implies x \rightarrow x_0$, i.e., f^{-1} continuous at y_0 .
- In fact, f continuous and either $\nearrow$ or $\searrow$ on $I \implies f^{-1}$ continuous on $J = f(I)$.

The Derivative of an Inverse

Key insights for proof of inverse function theorem:

- Darboux's theorem $\implies f \nearrow$ or $\searrow$ on $I \implies f$ is 1 : 1 on I
- If $y = f(x)$ and $y_0 = f(x_0)$

then $x = f^{-1}(y)$ and $x_0 = f^{-1}(y_0)$,

$$\begin{aligned} \text{so } \frac{f^{-1}(y) - f^{-1}(y_0)}{y - y_0} &= \frac{x - x_0}{f(x) - f(x_0)} \\ &= \frac{1}{\frac{f(x) - f(x_0)}{x - x_0}}. \end{aligned}$$

- Since f continuous at x_0 , we know $x \rightarrow x_0 \implies y \rightarrow y_0$.
- But we need $y \rightarrow y_0 \implies x \rightarrow x_0$, i.e., f^{-1} continuous at y_0 .
- In fact, f continuous and either $\nearrow$ or $\searrow$ on $I \implies f^{-1}$ continuous on $J = f(I)$. (more generally, cf. [Invariance of Domain](#) thm)