

On Dialectica and Differentiation

Davide Barbarossa

db2437@bath.ac.uk

<https://davidebarbarossa12.github.io/>


Department of Computer Science



Seminar at University of Sussex, Brighton (UK)

13/11/2024

Differentiation




Differentiation

Cartesian differential categories ('24)

$$\frac{f : A \rightarrow B}{Df : A \times A \rightarrow B}$$

Cartesian tangent categories ('24)

$$\frac{f : A \rightarrow B}{Tf : TA \rightarrow TB}$$




Differentiation

Cartesian differential categories ('24)

$$\frac{f : A \rightarrow B}{Df : A \times A \rightarrow B}$$

Cartesian tangent categories ('24)

$$\frac{f : A \rightarrow B}{Tf : TA \rightarrow TB}$$




Differentiation

Cartesian differential categories ('24)

$$\frac{f : A \rightarrow B}{Df : A \times A \rightarrow B}$$

Cartesian tangent categories ('24)

$$\frac{f : A \rightarrow B}{Tf : TA \rightarrow TB}$$



Differentiation

Cartesian differential categories ('24)

$$\frac{f : A \rightarrow B}{Df : A \times A \rightarrow B}$$

Cartesian tangent categories ('24)


$$\frac{f : A \rightarrow B}{Tf : TA \rightarrow TB}$$

Cartesian reverse diff. categories ('24)

$$\frac{f : A \rightarrow B}{Rf : A \times B \rightarrow A}$$

Cartesian reverse tangent categories ('24)

$$\frac{f : A \rightarrow B}{T^*f : f^*T^*B \rightarrow T^*A}$$



Dialectica (overview)

Source \rightarrow Target

Gödel
(’58)

$$A \in \text{HA} \quad \longmapsto \quad A_D\{w, c\} \in \text{T}$$

such that

$$\vdash_{\text{HA}} A \quad \implies \quad \vdash_{\text{T}} A_D\{M, c\} \text{ for some } M \in \text{T}$$

Dialectica (overview)

	Source \rightarrow Target		
Gödel (’58)	$A \in \mathbf{HA}$ $\vdash_{\mathbf{HA}} A$	\mapsto <i>such that</i> \implies	$A_D\{w, c\} \in \mathbf{T}$ $\vdash_{\mathbf{T}} A_D\{\mathbf{M}, c\}$ for some $\mathbf{M} \in \mathbf{T}$
Pédrot (Diller-Nahm) (’15)	$A \in \Lambda$ $\mathbf{M} \in \Lambda$ $\mathbf{x} : A \vdash_{\Lambda} \mathbf{M} : B$	\mapsto \mapsto <i>such that</i> \implies	$W(A), C(A) \in \mathbf{P}$ $\mathbf{M}^\bullet, \mathbf{M}_x \in \mathbf{P}$ (for \mathbf{x} variable) $\begin{cases} \mathbf{x} : W(A) \vdash_{\mathbf{P}} \mathbf{M}^\bullet : W(B) \\ \mathbf{x} : W(A) \vdash_{\mathbf{P}} \mathbf{M}_x : C(B) \rightarrow \mathcal{M}[C(A)] \end{cases}$

Dialectica (Transformation)

	α	$E \rightarrow F$
W	α_W	$W(E) \rightarrow W(F)$ \times $W(E) \times C(F) \rightarrow \mathcal{M}[C(E)]$
C	α_C	$W(E) \times C(F)$

	x	$\lambda x.M$	PQ
$(_)\bullet$	x	$\left\langle \begin{array}{c} \lambda x.M^\bullet \\ \lambda \pi.(\lambda x.M_x)\pi^1\pi^2 \end{array} \right\rangle$	$P^{\bullet 1}Q^\bullet$
$(_)_y$	$\begin{cases} \lambda \pi.[\pi], & x = y \\ \lambda \pi.0, & y \neq y \end{cases}$	$\lambda \pi.(\lambda x.M_y)\pi^1\pi^2$	$\lambda \pi. \left(\begin{array}{c} P_y\langle Q^\bullet, \pi \rangle \\ + \\ P^{\bullet 2}\langle Q^\bullet, \pi \rangle \gg Q_y \end{array} \right)$

A model $(\mathcal{C}, !)$ of Classical Differential Linear Logic

Arrows in \mathcal{C} : $A \xrightarrow{f} B$ (linear) Arrows in $\mathcal{C}_!$: $A \xrightarrow{f} B := !A \xrightarrow{f} B$ (non-linear)

A model $(\mathcal{C}, !)$ of Classical Differential Linear Logic

Arrows in \mathcal{C} : $A \xrightarrow{f} B$ (linear) Arrows in $\mathcal{C}_!$: $A \xrightarrow{f} B := !A \xrightarrow{f} B$ (non-linear)

Cartesian
+SMC
+Seely

$$\frac{A \quad B}{A \& B} \quad \frac{A \quad B}{A \otimes B} \quad \overline{! \top}$$

$$\frac{}{\text{ev}_{A,B} : [A \multimap B] \otimes A \multimap B}$$

$$\frac{f : A \multimap [E \multimap F]}{\lambda f : A \otimes E \multimap F}$$

A model $(\mathcal{C}, !)$ of Classical Differential Linear Logic

Arrows in \mathcal{C} : $A \xrightarrow{f} B$ (linear) Arrows in $\mathcal{C}_!$: $A \xrightarrow{f} B := !A \multimap B$ (non-linear)

Cartesian

+SMC

+Seely

$$\frac{A \quad B}{A \& B} \quad \frac{A \quad B}{A \otimes B} \quad \overline{! \top}$$

★-autonomous

$$\frac{f : A \multimap B}{f^\perp : B^\perp \multimap A^\perp}$$

$$\overline{\text{ev}_{A,B} : [A \multimap B] \otimes A \multimap B}$$

$$\frac{f : A \multimap [E \multimap F]}{\lambda f : A \otimes E \multimap F}$$

$$\overline{f^{\perp\perp} = f}$$

A model $(\mathcal{C}, !)$ of Classical Differential Linear Logic

Arrows in \mathcal{C} : $A \xrightarrow{f} B$ (linear) Arrows in $\mathcal{C}_!$: $A \xrightarrow{f} B := !A \xrightarrow{f} B$ (non-linear)

Cartesian
+SMC
+Seely

$$\frac{A \quad B}{A \& B} \quad \frac{A \quad B}{A \otimes B} \quad \overline{! \top}$$

★-autonomous
comm.
monoids
enriched

$$\frac{f : A \multimap B}{f^\perp : B^\perp \multimap A^\perp}$$

$$\overline{0_{A,B} : A \multimap B}$$

$$\overline{\text{ev}_{A,B} : [A \multimap B] \otimes A \multimap B}$$

$$\frac{f : A \multimap [E \multimap F]}{\lambda f : A \otimes E \multimap F}$$

$$\overline{f^{\perp\perp} = f}$$

$$\frac{f : A \multimap B \quad g : A \multimap B}{f + g : A \multimap B}$$

A model $(\mathcal{C}, !)$ of Classical Differential Linear Logic

Arrows in \mathcal{C} : $A \xrightarrow{f} B$ (linear) Arrows in $\mathcal{C}_!$: $A \xrightarrow{f} B := !A \xrightarrow{f} B$ (non-linear)

Cartesian
+SMC
+Seely

$$\frac{A \quad B}{A \& B} \quad \frac{A \quad B}{A \otimes B} \quad \overline{! \top}$$

★-autonomous
comm.
monoids
enriched

$$\frac{f : A \multimap B}{f^\perp : B^\perp \multimap A^\perp}$$

$$\overline{0_{A,B} : A \multimap B}$$

DiLL magic

$$\frac{f : A \rightarrow B}{\partial f : A \otimes !A \multimap B}$$

$$\overline{\text{ev}_{A,B} : [A \multimap B] \otimes A \multimap B}$$

$$\frac{f : A \multimap [E \multimap F]}{\lambda f : A \otimes E \multimap F}$$

$$\overline{f^{\perp\perp} = f}$$

$$\frac{f : A \multimap B \quad g : A \multimap B}{f + g : A \multimap B}$$

$$\frac{\frac{f : A \rightarrow B}{f_* : A \rightarrow [A \multimap B]} \quad a : !\top \multimap A}{f_* a : A \multimap B}$$

A model $(\mathcal{C}, !)$ of Classical Differential Linear Logic

Arrows in \mathcal{C} : $A \xrightarrow{f} B$ (linear) Arrows in $\mathcal{C}_!$: $A \xrightarrow{f} B := !A \xrightarrow{f} B$ (non-linear)

Cartesian
+SMC
+Seely

$$\frac{A \quad B}{A \& B} \quad \frac{A \quad B}{A \otimes B} \quad \overline{! \top}$$

★-autonomous
comm.
monoids
enriched

$$\frac{f : A \multimap B}{f^\perp : B^\perp \multimap A^\perp}$$

$$\overline{0_{A,B} : A \multimap B}$$

DiLL magic

$$\frac{f : A \rightarrow B}{\partial f : A \otimes !A \multimap B}$$

$$\overline{\text{ev}_{A,B} : [A \multimap B] \otimes A \multimap B}$$

$$\frac{f : A \multimap [E \multimap F]}{\lambda f : A \otimes E \multimap F}$$

$$\overline{f^{\perp\perp} = f}$$

$$\frac{f : A \multimap B \quad g : A \multimap B}{f + g : A \multimap B}$$

$$\frac{\frac{f : A \rightarrow B}{f_* : A \rightarrow [A \multimap B]} \quad a : !\top \multimap A}{f_* a : A \multimap B}$$

$\mathcal{C}_!$ is a model of differential λ -calculus where we can transpose linear arrows:

$$\frac{f : A \rightarrow B}{Df : A \times A \rightarrow B} \quad (\text{in } \mathcal{C}_!, E \times F := E \& F)$$

Dialectica and (Categorical) Differentiation

$$\sim_B \subseteq \{\vdash_{\mathbf{P}} \mathbf{M} : W(B)\} \times \mathcal{C}_!(\top, B)$$

$$\bowtie_B^A \subseteq \{\vdash_{\mathbf{P}} \mathbf{M} : C(B) \rightarrow \mathcal{M}[C(A)]\} \times \mathcal{C}_!(A, B) \times \mathcal{C}(B^\perp, A^\perp)$$

$\mathbf{M} \sim_{E \rightarrow F} f$	<p>for all $H \sim_E e$, we have $\mathbf{M}^1 H \sim_F f _e : F$</p> <p>and $\lambda \pi. \mathbf{M}^2 \langle H, \pi \rangle \bowtie_F^E \left(\begin{array}{c} \lambda^{-1} f : E \rightarrow F \\ ((\lambda^{-1} f)_* e)^\perp : F^\perp \multimap E^\perp \end{array} \right)$</p>
$\mathbf{M} \bowtie_{E \rightarrow F}^A \left(\begin{array}{c} f \\ g \end{array} \right)$	<p>for all $H \sim_E e$, we have</p> <p>$\lambda \pi. \mathbf{M} \langle H, \pi \rangle \bowtie_F^A \left(\begin{array}{c} f _e : A \rightarrow F \\ g^\perp _e^\perp : F^\perp \multimap A^\perp \end{array} \right)$</p>

The theorem

Let $\mathbf{x} : A \vdash_{\Lambda} \mathbf{M} : B$. Then:

$$1) \quad (\lambda \mathbf{x} . \mathbf{M})^{\bullet} \quad \sim_{A \rightarrow B} \quad \llbracket \lambda \mathbf{x} . \mathbf{M} \rrbracket : [A \rightarrow B]$$

$$2) \quad (\lambda \mathbf{x} . \mathbf{M}_{\mathbf{x}}) \mathbf{N} \quad \bowtie_B^A \quad \left(\begin{array}{ll} \llbracket \mathbf{M} \rrbracket & : \quad A \rightarrow B \\ (\llbracket \mathbf{M} \rrbracket_* a)^{\perp} & : \quad B^{\perp} \multimap A^{\perp} \end{array} \right) \quad \text{for all } \mathbf{N} \sim_A a.$$

Moral:

$$(\lambda \mathbf{x} . \mathbf{M}^{\bullet}, \lambda \mathbf{x} . \mathbf{M}_{\mathbf{x}})$$

“represents” the pair $(\llbracket \mathbf{M} \rrbracket, R[\llbracket \mathbf{M} \rrbracket])$, where

$$R[\llbracket \mathbf{M} \rrbracket] : A \times B^{\perp} \rightarrow A^{\perp}$$

is the reverse differential of $\llbracket \mathbf{M} \rrbracket$.

Reflection moment

Yet, can we say that Dialectica really *is* (reverse) Differentiation?

Reflection moment

Yet, can we say that Dialectica really *is* (reverse) Differentiation?

- All Dialectica transformation/only the transformation of λ -calculus.
What about quantifiers?

Reflection moment

Yet, can we say that Dialectica really *is* (reverse) Differentiation?

- All Dialectica transformation/only the transformation of λ -calculus.
What about quantifiers?
Counterexample (thanks to Thomas Powell), but that's another topic...

Reflection moment

Yet, can we say that Dialectica really *is* (reverse) Differentiation?

- All Dialectica transformation/only the transformation of λ -calculus.
What about quantifiers?
Counterexample (thanks to Thomas Powell), but that's another topic...
- The main feature of Dialectica is that in the target we have (e.g.) binary predicates (non-trivial subobjects in Dialectica categories). Here we don't: aren't we lose something about Dialectica?

Reflection moment

Yet, can we say that Dialectica really *is* (reverse) Differentiation?

- All Dialectica transformation/only the transformation of λ -calculus.
What about quantifiers?
Counterexample (thanks to Thomas Powell), but that's another topic...
- The main feature of Dialectica is that in the target we have (e.g.) binary predicates (non-trivial subobjects in Dialectica categories). Here we don't: aren't we lose something about Dialectica?
Not really, we are just formulating it differently.

Reflection moment

Yet, can we say that Dialectica really *is* (reverse) Differentiation?

- All Dialectica transformation/only the transformation of λ -calculus.
What about quantifiers?
Counterexample (thanks to Thomas Powell), but that's another topic...
- The main feature of Dialectica is that in the target we have (e.g.) binary predicates (non-trivial subobjects in Dialectica categories). Here we don't: aren't we lose something about Dialectica?
Not really, we are just formulating it differently.
- Is this correspondence astonishing/magic? Can we find some “reason” clarifying it?

Reflection moment

Yet, can we say that Dialectica really *is* (reverse) Differentiation?

- All Dialectica transformation/only the transformation of λ -calculus.
What about quantifiers?
Counterexample (thanks to Thomas Powell), but that's another topic...
- The main feature of Dialectica is that in the target we have (e.g.) binary predicates (non-trivial subobjects in Dialectica categories). Here we don't: aren't we lose something about Dialectica?
Not really, we are just formulating it differently.
- Is this correspondence astonishing/magic? Can we find some “reason” clarifying it?
Definitely yes at first sight... but then we can clearly understand its reason by looking at the categorical framework behind it.

Lens Categories

The category $\text{Lens}(\mathcal{L})$ of lenses over \mathcal{L} is defined as follows:

- objects: arrows in \mathcal{L} , which we think as fibre bundles and we write $p : \binom{\alpha}{A}$
- arrows from $p : \binom{\alpha}{A}$ to $q : \binom{\beta}{B}$ are the data of both a $f : A \rightarrow B$ in \mathcal{L} and a span $\alpha \xleftarrow{F} f^*\beta \xrightarrow{\bar{f}} \beta$ in \mathcal{L} , taken from the following pullback diagram:

$$\begin{array}{ccccc}
 \alpha & \xleftarrow{F} & f^*\beta & \xrightarrow{\bar{f}} & \beta \\
 & \searrow p & \downarrow f^*q & \lrcorner & \downarrow q \\
 & & A & \xrightarrow{f} & B
 \end{array}$$

Lens Categories

The category $\text{Lens}(\mathcal{L})$ of lenses over \mathcal{L} is defined as follows:

- objects: arrows in \mathcal{L} , which we think as fibre bundles and we write $p : \binom{\alpha}{A}$
- arrows from $p : \binom{\alpha}{A}$ to $q : \binom{\beta}{B}$ are the data of both a $f : A \rightarrow B$ in \mathcal{L} and a span $\alpha \xleftarrow{F} f^*\beta \xrightarrow{\bar{f}} \beta$ in \mathcal{L} , taken from the following pullback diagram:

$$\begin{array}{ccccc}
 \alpha & \xleftarrow{F} & f^*\beta & \xrightarrow{\bar{f}} & \beta \\
 & \searrow p & \downarrow f^*q & \lrcorner & \downarrow q \\
 & & A & \xrightarrow{f} & B
 \end{array}$$

Let $\mathcal{ELens}(\mathcal{L})$ be the full subcategory of $\text{Lens}(\mathcal{L})$ of trivial bundles, i.e. first projections. Concretely:

- Objects are first projections $\pi_1 : \binom{A \times X}{A}$
- An arrow from $\pi_1 : \binom{A \times X}{A}$ to $\pi_1 : \binom{B \times Y}{B}$ is given by an $f : A \rightarrow B$ and a span $A \times X \xleftarrow{F} A \times Y \xrightarrow{f \times 1} B \times Y$ such that $F; \pi_1^{A,X} = \pi_1^{A,Y}$.

Differentiation through the lens of lenses

Let \mathcal{L} be a Cartesian (closed, if you want λ -calculus) differential category where from the differential Df of a function f (a primitive data in \mathcal{L}) we can define the reverse differential Rf of f . (Think of $\mathcal{L} := \mathcal{C}_!$ of the first part).

Differentiation through the lens of lenses

Let \mathcal{L} be a Cartesian (closed, if you want λ -calculus) differential category where from the differential Df of a function f (a primitive data in \mathcal{L}) we can define the reverse differential Rf of f . (Think of $\mathcal{L} := \mathcal{C}_!$ of the first part).

We have a functor $\mathcal{L} \rightarrow \mathcal{ELens}(\mathcal{L})$ defined by:

$$A \quad \mapsto \quad \pi_1 : \left(A \times A^\perp \right)_A$$

$$A \xrightarrow{f} B \quad \mapsto \quad \left(f, \quad A \times A^\perp \xleftarrow{\langle \pi_1, Rf \rangle} A \times B^\perp \xrightarrow{f \times 1} B \times B^\perp \right).$$

Differentiation through the lens of lenses

Let \mathcal{L} a Cartesian reverse differential category.

Differentiation through the lens of lenses

Let \mathcal{L} a Cartesian reverse differential category.

We have a functor $\mathcal{L} \rightarrow \mathcal{ELens}(\mathcal{L})$ defined by:

$$\begin{aligned} A &\mapsto \pi_1 : \left(\begin{smallmatrix} A \times A \\ A \end{smallmatrix} \right) \\ A \xrightarrow{f} B &\mapsto \left(f, A \times A \xleftarrow{\langle \pi_1, Rf \rangle} A \times B \xrightarrow{f \times 1} B \times B \right) \end{aligned}$$

where $Rf : A \times B \rightarrow A$ is the reverse differential of f (a primitive data in \mathcal{L}).

Differentiation through the lens of lenses

Let \mathcal{L} be a reverse tangent category. This means that \mathcal{L} has a tangent functor T giving tangent bundles $p_A : (T^*_A A)$ of objects A and giving tangent arrows $Tf : T^*_A A \rightarrow T^*_B B$ for arrows $f : A \rightarrow B$, and we can “reverse” T in order to get cotangent bundles $p_A^* : (T^* A)$ and arrows in the pullback diagram below:

$$\begin{array}{ccccc}
 T^* A & \xleftarrow{T^* f} & f^* T^* B & \xrightarrow{\bar{f}} & T^* B \\
 & \searrow p_A^* & \downarrow f^* p_B & \lrcorner & \downarrow p_B^* \\
 & & A & \xrightarrow{f} & B
 \end{array}$$

where $T^* f$ is the diff. geometry formulation of the reverse differential of f .

Differentiation through the lens of lenses

Let \mathcal{L} be a reverse tangent category. This means that \mathcal{L} has a tangent functor T giving tangent bundles $p_A : (T_A^A)$ of objects A and giving tangent arrows $Tf : TA \rightarrow TB$ for arrows $f : A \rightarrow B$, and we can “reverse” T in order to get cotangent bundles $p_A^* : (T^*A)$ and arrows in the pullback diagram below:

$$\begin{array}{ccccc}
 T^*A & \xleftarrow{T^*f} & f^*T^*B & \xrightarrow{\bar{f}} & T^*B \\
 & \searrow p_A^* & \downarrow f^*p_B & \lrcorner & \downarrow p_B^* \\
 & & A & \xrightarrow{f} & B
 \end{array}$$

where T^*f is the diff. geometry formulation of the reverse differential of f .

We have a functor $\mathcal{L} \rightarrow \text{Lens}(\mathcal{L})$ defined by:

$$A \mapsto p_A^* : (T^*A)$$

$$A \xrightarrow{f} B \mapsto (f, T^*A \xleftarrow{T^*f} f^*T^*B \xrightarrow{\bar{f}} T^*B).$$

Expressing Dialectica as a functor

$$\Lambda_{\text{cat}} \rightarrow \mathcal{E}\text{Lens}(\mathbf{P}_{\text{cat}})$$

Expressing Dialectica as a functor

$$\Lambda_{\text{cat}} \rightarrow \mathcal{ELens}(\mathbf{P}_{\text{cat}})$$

- An object A is sent to the typed term $\mathbf{z} : W(A) \times \mathcal{M}[C(A)] \vdash_{\mathbf{P}} \mathbf{z}^1 : W(A)$
- An arrow $\mathbf{z} : A \vdash_{\Lambda} M : B$ in Λ_{cat} from A to B is sent to the arrow in $\mathcal{ELens}(\mathbf{P}_{\text{cat}})$ from $\mathbf{z} : W(A) \times \mathcal{M}[C(A)] \vdash_{\mathbf{P}} \mathbf{z}^1 : W(A)$ to $\mathbf{z} : W(B) \times \mathcal{M}[C(B)] \vdash_{\mathbf{P}} \mathbf{z}^1 : W(B)$ given by the following diagram:

$$\begin{array}{ccccc}
 W(A) \times \mathcal{M}[C(A)] & \xleftarrow{\langle \mathbf{z}^1, \mathbf{z}^2 \rangle \models_{\mathcal{M}(\mathbf{z}^1)}} & W(A) \times \mathcal{M}[C(B)] & \xrightarrow{\langle M^\bullet, \mathbf{z}^2 \rangle} & W(B) \times \mathcal{M}[C(B)] \\
 & \searrow \mathbf{z}^1 & \downarrow \mathbf{z}^1 & \lrcorner & \downarrow \mathbf{z}^1 \\
 & & W(A) & \xrightarrow{M^\bullet} & W(B)
 \end{array}$$

Expressing Dialectica as a functor

$$\Lambda_{\text{cat}} \rightarrow \mathcal{E}\text{Lens}(\mathbf{P}_{\text{cat}})$$

Moral:

The Dialectica transformation of λ -calculus encodes (reverse) Differentiation *because* it is a transformation into a category of Lenses, the latter being the abstract setting for Reverse Differentiation.

Final comments

- I didn't talk about Dialectica categories. I could have said something (ask me if you are interested)
- Explore categorical framework to reverse a Cartesian closed differential category in order to define Cartesian closed reverse differential/tangent categories
- Reverse differential λ -calculus? There is an interesting paper from Ong and Mak.

Final comments

- I didn't talk about Dialectica categories. I could have said something (ask me if you are interested)
- Explore categorical framework to reverse a Cartesian closed differential category in order to define Cartesian closed reverse differential/tangent categories
- Reverse differential λ -calculus? There is an interesting paper from Ong and Mak.

THANK YOU!