

Abelian covers of horocycle flows

Davide Ravotti (U. Lille, CNRS)


joint work with Roberto Castorrini (SNS Pisa)

25th June 2024

The setting


Let

- S be a closed surface with a smooth Riemannian metric of negative curvature,



The setting


Let



- S be a closed surface with a smooth Riemannian metric of negative curvature,
- $M = T^1S$ be the unit tangent bundle of S ,

The setting


Let



- S be a closed surface with a smooth Riemannian metric of negative curvature,
- $M = T^1S$ be the unit tangent bundle of S ,
- $(g_t)_{t \in \mathbb{R}}$ be the geodesic flow on M .

The setting

Let



- S be a closed surface with a smooth Riemannian metric of negative curvature,
- $M = T^1S$ be the unit tangent bundle of S ,
- $(g_t)_{t \in \mathbb{R}}$ be the geodesic flow on M .

The setting

The geodesic flow is an **Anosov flow**: there exist $C \geq 0$, $\lambda > 0$, and a continuous, Dg_t -invariant splitting

$$TM = E_- \oplus \langle X \rangle \oplus E_+,$$

where X is the vector field generating $(g_t)_{t \in \mathbb{R}}$, such that

$$\|Dg_t|_{E_-}\| \leq C \cdot e^{-\lambda t} \quad \text{and} \quad \|Dg_{-t}|_{E_+}\| \leq C \cdot e^{-\lambda t},$$

for all $t \geq 0$.

The setting

The geodesic flow is a **contact flow**: there exists a smooth (invariant) 1-form α such that

$$\alpha(X) = 1 \quad \text{and} \quad \alpha|_{E_- \oplus E_+} = 0.$$

The setting

The geodesic flow is a **contact flow**: there exists a smooth (invariant) 1-form α such that

$$\alpha(X) = 1 \quad \text{and} \quad \alpha|_{E_- \oplus E_+} = 0.$$

Then, the 3-form

$$d\text{vol} = \alpha \wedge d\alpha$$

is never zero and defines a smooth invariant measure for $(g_t)_{t \in \mathbb{R}}$.

The setting

The geodesic flow is a **contact flow**: there exists a smooth (invariant) 1-form α such that

$$\alpha(X) = 1 \quad \text{and} \quad \alpha|_{E_- \oplus E_+} = 0.$$

Then, the 3-form

$$d\text{vol} = \alpha \wedge d\alpha$$

is never zero and defines a smooth invariant measure for $(g_t)_{t \in \mathbb{R}}$.

Theorem (Dolgopyat '98, Liverani '04)

The geodesic flow on (M, vol) is exponentially mixing.

For **hyperbolic surfaces**: Ratner '87.

Horocycle flows

Theorem (Hurder-Katok '90)

- The distributions E_{\pm} are of class $\mathcal{C}^{2-\varepsilon}$, for all $\varepsilon > 0$.
- If E_+ or E_- is of class \mathcal{C}^2 , then S has constant curvature.

Horocycle flows

Theorem (Hurder-Katok '90)

- The distributions E_{\pm} are of class $\mathcal{C}^{2-\varepsilon}$, for all $\varepsilon > 0$.
- If E_+ or E_- is of class \mathcal{C}^2 , then S has constant curvature.
- The distribution E_- integrates to a foliation with 1-dimensional, orientable leaves.

Horocycle flows

Theorem (Hurder-Katok '90)


- The distributions E_{\pm} are of class $\mathcal{C}^{2-\varepsilon}$, for all $\varepsilon > 0$.
 - If E_+ or E_- is of class \mathcal{C}^2 , then S has constant curvature.
-
- The distribution E_- integrates to a foliation with 1-dimensional, orientable leaves.
 - The (stable) horocycle flow $(h_t)_{t \in \mathbb{R}}$ is the unit speed motion along the leaves parallel to E_- .

Horocycle flows

Theorem (Hurder-Katok '90)

- The distributions E_{\pm} are of class $\mathcal{C}^{2-\varepsilon}$, for all $\varepsilon > 0$.
- If E_+ or E_- is of class \mathcal{C}^2 , then S has constant curvature.
- The distribution E_- integrates to a foliation with 1-dimensional, orientable leaves.
- The (stable) horocycle flow $(h_t)_{t \in \mathbb{R}}$ is the unit speed motion along the leaves parallel to E_- .
- The generating vector field $U \in E_-$ is of class $\mathcal{C}^{2-\varepsilon}$, for all $\varepsilon > 0$.

Horocycle flows



Stable horocycle flow $(h_t)_{t \in \mathbb{R}}$ on $T^1\mathbb{D}$

Unique ergodicity

Theorem (Marcus '75, from Margulis '70)

There exists a unique, h_t -invariant probability measure μ .

For **hyperbolic surfaces**: Furstenberg '73.

Unique ergodicity

Theorem (Marcus '75, from Margulis '70)

There exists a unique, h_t -invariant probability measure μ .

For **hyperbolic surfaces**: Furstenberg '73.

Theorem (Adam-Baladi '22)

Under a pinching condition, there exists $\delta > 0$ such that for all $f \in \mathcal{C}^2(M)$ we have

$$\int_0^T f \circ h_t(x) dx = T \cdot \mu(f) + O(T^{1-\delta}).$$

For **hyperbolic surfaces**: Burger '90, Flaminio-Forni '03, Bufetov-Forni '14, R. '23.

The noncompact setting

- Fix $\Gamma_0 \trianglelefteq \Gamma := \pi_1(S)$ so that $\mathcal{G} = \Gamma/\Gamma_0 \simeq \mathbb{Z}^d$, for some $d \geq 1$.

The noncompact setting

- Fix $\Gamma_0 \trianglelefteq \Gamma := \pi_1(S)$ so that $\mathcal{G} = \Gamma/\Gamma_0 \simeq \mathbb{Z}^d$, for some $d \geq 1$.
- The associated cover $p: S_0 \rightarrow S$ has a Galois group isomorphic to \mathbb{Z}^d .


The noncompact setting

- Fix $\Gamma_0 \trianglelefteq \Gamma := \pi_1(S)$ so that $\mathcal{G} = \Gamma/\Gamma_0 \simeq \mathbb{Z}^d$, for some $d \geq 1$.
- The associated cover $p: S_0 \rightarrow S$ has a Galois group isomorphic to \mathbb{Z}^d .
- Since $[\Gamma, \Gamma] \trianglelefteq \Gamma_0$ and $\Gamma/[\Gamma, \Gamma] \simeq H_1(S, \mathbb{Z}) \simeq \mathbb{Z}^{2g}$, there are d linearly independent primitive cohomology classes $[\omega_1], \dots, [\omega_d] \in H_{\text{dR}}^1(S, \mathbb{Z})$ so that $\int_\gamma \omega_i = 0$ for all $[\gamma] \in \Gamma_0/[\Gamma, \Gamma]$ and $p^*[\omega_i] = 0$.

The noncompact setting


- Fix $\Gamma_0 \trianglelefteq \Gamma := \pi_1(S)$ so that $\mathcal{G} = \Gamma/\Gamma_0 \simeq \mathbb{Z}^d$, for some $d \geq 1$.
- The associated cover $p: S_0 \rightarrow S$ has a Galois group isomorphic to \mathbb{Z}^d .
- Since $[\Gamma, \Gamma] \trianglelefteq \Gamma_0$ and $\Gamma/[\Gamma, \Gamma] \simeq H_1(S, \mathbb{Z}) \simeq \mathbb{Z}^{2g}$, there are d linearly independent primitive cohomology classes $[\omega_1], \dots, [\omega_d] \in H_{\text{dR}}^1(S, \mathbb{Z})$ so that $\int_\gamma \omega_i = 0$ for all $[\gamma] \in \Gamma_0/[\Gamma, \Gamma]$ and $p^*[\omega_i] = 0$.
- We can take the forms $\omega_1, \dots, \omega_d$ to be harmonic.

The noncompact setting



- We equip S_0 with the pullback Riemannian metric and measure μ .

The noncompact setting



- We equip S_0 with the pullback Riemannian metric and measure μ .
- We consider the geodesic and horocycle flows $(g_t)_{t \in \mathbb{R}}$ and $(h_t)_{t \in \mathbb{R}}$ on $M_0 = T^1 S_0$.

Ergodic integrals

Some general facts about the ergodic integrals of integrable functions

$$\mathcal{I}_t f(x) := \int_0^t f \circ h_r(x) dr.$$

Ergodic integrals

Some general facts about the ergodic integrals of integrable functions

$$\mathcal{I}_t f(x) := \int_0^t f \circ h_r(x) dr.$$

- Birkhoff Ergodic Theorem: $\mathcal{I}_t f = o(t)$,

Ergodic integrals

Some general facts about the ergodic integrals of integrable functions

$$\mathcal{I}_t f(x) := \int_0^t f \circ h_r(x) dr.$$

- Birkhoff Ergodic Theorem: $\mathcal{I}_t f = o(t)$,
- Aaronson '97: for $f \geq 0$, for any $a(t) > 0$, either

$$\limsup_{t \rightarrow \infty} \frac{\mathcal{I}_t f}{a(t)} = \infty \quad \mu\text{-a.e.}, \quad \text{or} \quad \liminf_{t \rightarrow \infty} \frac{\mathcal{I}_t f}{a(t)} = 0 \quad \mu\text{-a.e.},$$

Ergodic integrals

Some general facts about the ergodic integrals of integrable functions

$$\mathcal{I}_t f(x) := \int_0^t f \circ h_r(x) dr.$$

- Birkhoff Ergodic Theorem: $\mathcal{I}_t f = o(t)$,
- Aaronson '97: for $f \geq 0$, for any $a(t) > 0$, either

$$\limsup_{t \rightarrow \infty} \frac{\mathcal{I}_t f}{a(t)} = \infty \quad \mu\text{-a.e.}, \quad \text{or} \quad \liminf_{t \rightarrow \infty} \frac{\mathcal{I}_t f}{a(t)} = 0 \quad \mu\text{-a.e.},$$

- Hopf '37: for a positive integrable g ,

$$\lim_{t \rightarrow \infty} \frac{\mathcal{I}_t f}{\mathcal{I}_t g} = \frac{\mu(f)}{\mu(g)}.$$

Ergodic integrals

Our goal: describe the ergodic integrals μ -a.e. as

$$\mathcal{I}_t f(x) = a(t) \cdot \mu(f) \cdot \Phi_t(x) \cdot (1 + o(1)).$$

where

Ergodic integrals

Our goal: describe the ergodic integrals μ -a.e. as

$$\mathcal{I}_t f(x) = a(t) \cdot \mu(f) \cdot \Phi_t(x) \cdot (1 + o(1)).$$

where

- $a(t)$ is the “correct” asymptotic rate and

Ergodic integrals

Our goal: describe the ergodic integrals μ -a.e. as

$$\mathcal{I}_t f(x) = a(t) \cdot \mu(f) \cdot \Phi_t(x) \cdot (1 + o(1)).$$

where

- $a(t)$ is the “correct” asymptotic rate and
- $\Phi_t(x)$ is an “oscillating” term, independent of f .

The main result

Theorem (Castorrini-R. '24)

There exist $C_M \geq 1$, $\delta > 0$, and

- a $d \times d$ positive definite symmetric matrix Σ ,
- a function $t_*: M \times [C_M, \infty) \rightarrow \mathbb{R}_{\geq 0}$ satisfying

$$\|T - e^{h_{\text{top}} \cdot t_*(\cdot, T)}\|_{\infty} \leq C_M T^{1-\delta},$$

- a vector-valued function $F_*: M \times \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}^d$ satisfying

$$\frac{F_*(x, t)}{\sqrt{t}} \rightarrow \mathcal{N}(0, \Sigma) \quad \text{in distribution, as } x \sim \text{vol}$$

for which the following holds. (cont'd)

The main result

Theorem (Castorrini-R. '24)

(cont'd)

For every $f \in \mathcal{C}_c^2(M_0)$ and every $x \in M_0$, there exists a constant $C(f, x)$ depending (explicitly) of the \mathcal{C}^2 -norm of f , its support $\text{supp}(f)$, and the distance between x and $\text{supp}(f)$,

such that, denoting $t_* = t_*(p(x), T)$, for all $T \geq C_M$ we have

The main result

Theorem (Castorrini-R. '24)

(cont'd)

For every $f \in \mathcal{C}^2(M_0)$ and every $x \in M_0$, there exists a constant $C(f, x)$ depending (explicitly) of the \mathcal{C}^2 -norm of f , its support $\text{supp}(f)$, and the distance between x and $\text{supp}(f)$,

such that, denoting $t_* = t_*(p(x), T)$, for all $T \geq C_M$ we have

$$\left| \mathcal{I}_T f(x) - \frac{h_{\text{top}}^{\frac{d}{2}}}{(2\pi)^{\frac{d}{2}} \sqrt{\det \Sigma}} \frac{T}{(\log T)^{\frac{d}{2}}} \mu(f) \times \right. \\ \left. \times \exp \left(-\frac{1}{2} \frac{F_*(p(x), t_*) \cdot \Sigma F_*(p(x), t_*)}{t_*} \right) \right| \leq C_M C(f, x) \frac{T \log \log T}{(\log T)^{\frac{d+1}{2}}}.$$

Remarks

- For hyperbolic surfaces, the result without error rates is due to [Ledrappier-Sarig '06](#).


Remarks

- For hyperbolic surfaces, the result without error rates is due to [Ledrappier-Sarig '06](#).
- Results of this type have been obtained for some (very special) classes of \mathbb{Z}^d -covers of translation flows, including [Avila-Doglopyat-Duryev-Sarig '15](#) and [Bruin-Fougeron-R.-Terhesiu '24](#).

Renormalization time

The function t_* is a **renormalization time**, namely is defined by the equality


$$g_{t_*(x,T)} \circ h_T(x) = h_1 \circ g_{t_*(x,T)}(x).$$



Renormalization time

The function t_* is a **renormalization time**, namely is defined by the equality

$$g_{t_*(x,T)} \circ h_T(x) = h_1 \circ g_{t_*(x,T)}(x).$$



In particular, for hyperbolic surfaces, $t_*(x, T) = \log(T)$.

The Frobenius function

- Recall that there are d harmonic 1-forms $\omega_1, \dots, \omega_d$ on S so that their pullback $p^*\omega_1, \dots, p^*\omega_d$ are **exact** on S_0 .

The Frobenius function

- Recall that there are d harmonic 1-forms $\omega_1, \dots, \omega_d$ on S so that their pullback $p^*\omega_1, \dots, p^*\omega_d$ are **exact** on S_0 .
- We still denote by $p^*\omega_i$ their pullback on M_0 under the canonical projection $M_0 = T^1S_0 \rightarrow S_0$.

The Frobenius function

- Recall that there are d harmonic 1-forms $\omega_1, \dots, \omega_d$ on S so that their pullback $p^*\omega_1, \dots, p^*\omega_d$ are **exact** on S_0 .
- We still denote by $p^*\omega_i$ their pullback on M_0 under the canonical projection $M_0 = T^1S_0 \rightarrow S_0$.
- The **Frobenius function/geodesic winding cycle** $F_*(x, t)$ is the d -dimensional vector whose i -th component is

$$\int_x^{g_t(x)} p^*\omega_i = \int_0^t \langle \omega_i, X \rangle \circ g_r(x) dr.$$

The Frobenius function

- Recall that there are d harmonic 1-forms $\omega_1, \dots, \omega_d$ on S so that their pullback $p^*\omega_1, \dots, p^*\omega_d$ are **exact** on S_0 .
- We still denote by $p^*\omega_i$ their pullback on M_0 under the canonical projection $M_0 = T^1S_0 \rightarrow S_0$.
- The **Frobenius function/geodesic winding cycle** $F_*(x, t)$ is the d -dimensional vector whose i -th component is

$$\int_x^{g_t(x)} p^*\omega_i = \int_0^t \langle \omega_i, X \rangle \circ g_r(x) dr.$$

- It is invariant by deck transformations, hence it is a well-defined function on M .

The proof: renormalization

The proof: renormalization

Commutation

There exists a function $\tau: M \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ such that

$$g_t \circ h_s(x) = h_{\tau(s,t,p(x))} \circ g_t(x),$$

for all $x \in M_0$ and $t, s \in \mathbb{R}$.

The proof: renormalization

Commutation

There exists a function $\tau: M \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ such that

$$g_t \circ h_s(x) = h_{\tau(s,t,p(x))} \circ g_t(x),$$

for all $x \in M_0$ and $t, s \in \mathbb{R}$.

For $t \geq 0$, it satisfies

$$C^{-1}e^{h_{\text{top}}t}s \leq \tau(s, -t, x) \leq Ce^{h_{\text{top}}t}s,$$

for some constant $C \geq 1$.

The proof: renormalization

Let

$$J_t(x) = \frac{\partial}{\partial s} \Big|_{s=0} \tau(s, t, x) \quad \text{so that} \quad J_t(h_s(x)) = \frac{\partial}{\partial s} \tau(s, t, x).$$

The proof: renormalization

Let

$$J_t(x) = \frac{\partial}{\partial s} \Big|_{s=0} \tau(s, t, x) \quad \text{so that} \quad J_t(h_s(x)) = \frac{\partial}{\partial s} \tau(s, t, x).$$

Then

$$\int_0^T f \circ h_s(x) \, ds = \int_0^T f \circ g_{-t} \circ g_t \circ h_s(x) \, ds$$

The proof: renormalization

Let

$$J_t(x) = \frac{\partial}{\partial s} \Big|_{s=0} \tau(s, t, x) \quad \text{so that} \quad J_t(h_s(x)) = \frac{\partial}{\partial s} \tau(s, t, x).$$

Then

$$\begin{aligned} \int_0^T f \circ h_s(x) \, ds &= \int_0^T f \circ g_{-t} \circ g_t \circ h_s(x) \, ds \\ &= \int_0^T f \circ g_{-t} \circ h_{\tau(s, t, x)}(g_t(x)) \cdot \frac{\partial}{\partial s} \tau(s, t, x) \cdot J_{-t}(g_t \circ h_s(x)) \, ds \end{aligned}$$

The proof: renormalization

Let

$$J_t(x) = \frac{\partial}{\partial s} \Big|_{s=0} \tau(s, t, x) \quad \text{so that} \quad J_t(h_s(x)) = \frac{\partial}{\partial s} \tau(s, t, x).$$

Then

$$\begin{aligned} \int_0^T f \circ h_s(x) \, ds &= \int_0^T f \circ g_{-t} \circ g_t \circ h_s(x) \, ds \\ &= \int_0^T f \circ g_{-t} \circ h_{\tau(s, t, x)}(g_t(x)) \cdot \frac{\partial}{\partial s} \tau(s, t, x) \cdot J_{-t}(g_t \circ h_s(x)) \, ds \\ &= \int_0^{\tau(T, t, x)} f \circ g_{-t} \circ h_r(g_t(x)) \cdot J_{-t} \circ h_r(g_t(x)) \, dr. \end{aligned}$$

The proof: renormalization

Let

$$J_t(x) = \frac{\partial}{\partial s} \Big|_{s=0} \tau(s, t, x) \quad \text{so that} \quad J_t(h_s(x)) = \frac{\partial}{\partial s} \tau(s, t, x).$$

Then

$$\begin{aligned} \int_0^T f \circ h_s(x) \, ds &= \int_0^T f \circ g_{-t} \circ g_t \circ h_s(x) \, ds \\ &= \int_0^T f \circ g_{-t} \circ h_{\tau(s, t, x)}(g_t(x)) \cdot \frac{\partial}{\partial s} \tau(s, t, x) \cdot J_{-t}(g_t \circ h_s(x)) \, ds \\ &= \int_0^{\tau(T, t, x)} f \circ g_{-t} \circ h_r(g_t(x)) \cdot J_{-t} \circ h_r(g_t(x)) \, dr. \end{aligned}$$

Weighted transfer operators

We are led to consider the operators $\mathcal{L}_t: f \mapsto f \circ g_{-t} \cdot J_{-t}$ on $\mathcal{C}_c^{2-\varepsilon}(M_0)$.

The proof: a Fourier decomposition

We identify Γ_0 with $\Gamma_0/[\Gamma, \Gamma] \leq H_1(S, \mathbb{Z})$.

The proof: a Fourier decomposition

We identify Γ_0 with $\Gamma_0/[\Gamma, \Gamma] \leq H_1(S, \mathbb{Z})$. Let

- Deck = {deck transformations of the cover $p: S_0 \rightarrow S$ },

The proof: a Fourier decomposition

We identify Γ_0 with $\Gamma_0/[\Gamma, \Gamma] \leq H_1(S, \mathbb{Z})$. Let

- Deck = {deck transformations of the cover $p: S_0 \rightarrow S$ },
- $\mathcal{H} = \{\omega \text{ harmonic 1-form} : p^*[\omega] = 0, \omega(\Gamma_0) = 0\} = \langle \omega_i : i = 1, \dots, d \rangle,$

The proof: a Fourier decomposition

We identify Γ_0 with $\Gamma_0/[\Gamma, \Gamma] \leq H_1(S, \mathbb{Z})$. Let

- Deck = {deck transformations of the cover $p: S_0 \rightarrow S$ },
- $\mathcal{H} = \{\omega \text{ harmonic 1-form} : p^*[\omega] = 0, \omega(\Gamma_0) = 0\} = \langle \omega_i : i = 1, \dots, d \rangle$,
- $\mathcal{H}(\mathbb{Z}) = \mathbb{Z}\omega_1 \oplus \dots \oplus \mathbb{Z}\omega_d$, and $\mathbb{T}^d = \mathcal{H}/\mathcal{H}(\mathbb{Z})$.

The proof: a Fourier decomposition

We identify Γ_0 with $\Gamma_0/[\Gamma, \Gamma] \leq H_1(S, \mathbb{Z})$. Let

- Deck = {deck transformations of the cover $p: S_0 \rightarrow S$ },
- $\mathcal{H} = \{\omega \text{ harmonic 1-form} : p^*[\omega] = 0, \omega(\Gamma_0) = 0\} = \langle \omega_i : i = 1, \dots, d \rangle$,
- $\mathcal{H}(\mathbb{Z}) = \mathbb{Z}\omega_1 \oplus \dots \oplus \mathbb{Z}\omega_d$, and $\mathbb{T}^d = \mathcal{H}/\mathcal{H}(\mathbb{Z})$.

Then, every homomorphism $\chi: \text{Deck} \rightarrow U(1)$ is of the form

$$\chi(D) = \chi([\gamma] + \Gamma_0) = \exp\left(2\pi i \int_{[\gamma]} \omega\right),$$

for a unique $\omega \in \mathbb{T}^d = \mathcal{H}/\mathcal{H}(\mathbb{Z})$.

The proof: a Fourier decomposition

Define

$$\mathcal{C}^2(M, \omega) := \{f \in \mathcal{C}^2(M_0) : f \circ D^{-1} = \chi_\omega(D)f\}.$$

The proof: a Fourier decomposition

Define

$$\mathcal{C}^2(M, \omega) := \{f \in \mathcal{C}^2(M_0) : f \circ D^{-1} = \chi_\omega(D)f\}.$$

We have

$$\pi_\omega : \mathcal{C}_c^2(M_0) \rightarrow \mathcal{C}^2(M, \omega), \quad \pi_\omega(f) = \sum_{D \in \text{Deck}} \chi_\omega(D) f \circ D$$

and

$$f = \int_{\mathbb{T}^d} \pi_\omega(f) d\omega.$$

The proof: a Fourier decomposition

Define

$$\Xi_{\omega}: \mathcal{C}^2(M, \eta + \omega) \rightarrow \mathcal{C}^2(M, \eta), \quad \Xi_{\omega}(f)(x) = f(x) \exp \left(2\pi i \int_{x_0}^x p^* \omega \right)$$

The proof: a Fourier decomposition

Define

$$\Xi_{\omega}: \mathcal{C}^2(M, \eta + \omega) \rightarrow \mathcal{C}^2(M, \eta), \quad \Xi_{\omega}(f)(x) = f(x) \exp \left(2\pi i \int_{x_0}^x p^* \omega \right)$$

We have

$$\begin{array}{ccc} \mathcal{C}^2(M, \omega) & \xrightarrow{\mathcal{L}_t} & \mathcal{C}^2(M, \omega) \\ \uparrow \Xi_{-\omega} & & \Xi_{\omega} \downarrow \\ \mathcal{C}^2(M, 0) & & \mathcal{C}^2(M, 0) \end{array}$$

The proof: a Fourier decomposition

Define

$$\Xi_{\omega}: \mathcal{C}^2(M, \eta + \omega) \rightarrow \mathcal{C}^2(M, \eta), \quad \Xi_{\omega}(f)(x) = f(x) \exp \left(2\pi i \int_{x_0}^x p^* \omega \right)$$

We have

$$\begin{array}{ccc} \mathcal{C}^2(M, \omega) & \xrightarrow{\mathcal{L}_t} & \mathcal{C}^2(M, \omega) \\ \uparrow \Xi_{-\omega} & & \Xi_{\omega} \downarrow \\ \mathcal{C}^2(M, 0) & & \mathcal{C}^2(M, 0) \end{array}$$

Twisted (and weighted) transfer operators

We study the family of operators $\mathcal{L}_t^{(\omega)}: \mathcal{C}^{2-\varepsilon}(M) \rightarrow \mathcal{C}^{2-\varepsilon}(M)$ given by

$$\mathcal{L}_t^{(\omega)} f = f \circ g_{-t} \cdot J_{-t} \cdot \exp \left(2\pi i \int_0^t \langle \omega, X \rangle \circ g_{-r} dr \right)$$

The proof: spectral theory

The proof: spectral theory

We consider a pair of Banach spaces $\mathcal{B}_w, \mathcal{B}$ with dense inclusions

$$\mathcal{C}^{2-\varepsilon}(M) \hookrightarrow \mathcal{B} \hookrightarrow \mathcal{B}_w \hookrightarrow \mathcal{C}^{2-\varepsilon}(M)^*,$$

The proof: spectral theory

We consider a pair of Banach spaces $\mathcal{B}_w, \mathcal{B}$ with dense inclusions

$$\mathcal{C}^{2-\varepsilon}(M) \hookrightarrow \mathcal{B} \hookrightarrow \mathcal{B}_w \hookrightarrow \mathcal{C}^{2-\varepsilon}(M)^*,$$

on which the linear functionals

$$L_{x,\varphi}: f \mapsto \int_0^1 f \circ h_s(x) \cdot \varphi(s) ds, \quad \text{for } x \in M \text{ and } \varphi \in \mathcal{C}_c^{2-\varepsilon}((0,1)),$$

defined for $f \in \mathcal{C}^{2-\varepsilon}(M)$, extend by continuity.

The proof: spectral theory

We prove that there exists $\delta > 0$ such that, for all $v \in \mathcal{C}^{2-\varepsilon}(M)$, we have

$$e^{-h_{\text{top}} t} \mathcal{L}_t^{(\omega)} v = e^{z(\omega)t} \Pi_{\omega} v + Q_{\omega,t} v,$$

where

The proof: spectral theory

We prove that there exists $\delta > 0$ such that, for all $v \in \mathcal{C}^{2-\varepsilon}(M)$, we have

$$e^{-h_{\text{top}} t} \mathcal{L}_t^{(\omega)} v = e^{z(\omega)t} \Pi_{\omega} v + Q_{\omega,t} v,$$

where

- Π_{ω} is a projection,

The proof: spectral theory

We prove that there exists $\delta > 0$ such that, for all $v \in \mathcal{C}^{2-\varepsilon}(M)$, we have

$$e^{-h_{\text{top}} t} \mathcal{L}_t^{(\omega)} v = e^{z(\omega)t} \Pi_{\omega} v + Q_{\omega,t} v,$$

where

- Π_{ω} is a projection,
- $\|Q_{\omega,t} v\|_{\mathcal{B}_w} \leq e^{-\delta t} \|v\|_{\mathcal{C}^{2-\varepsilon}},$

and

The proof: spectral theory

We prove that there exists $\delta > 0$ such that, for all $v \in \mathcal{C}^{2-\varepsilon}(M)$, we have

$$e^{-h_{\text{top}} t} \mathcal{L}_t^{(\omega)} v = e^{z(\omega)t} \Pi_{\omega} v + Q_{\omega,t} v,$$

where

- Π_{ω} is a projection,
- $\|Q_{\omega,t} v\|_{\mathcal{B}_w} \leq e^{-\delta t} \|v\|_{\mathcal{C}^{2-\varepsilon}},$

and

- $z(\omega) \leq 0$ and $z(\omega) = 0$ if and only if $\omega = 0$,

The proof: spectral theory

We prove that there exists $\delta > 0$ such that, for all $v \in \mathcal{C}^{2-\varepsilon}(M)$, we have

$$e^{-h_{\text{top}} t} \mathcal{L}_t^{(\omega)} v = e^{z(\omega)t} \Pi_{\omega} v + Q_{\omega,t} v,$$

where

- Π_{ω} is a projection,
- $\|Q_{\omega,t} v\|_{\mathcal{B}_w} \leq e^{-\delta t} \|v\|_{\mathcal{C}^{2-\varepsilon}},$

and

- $z(\omega) \leq 0$ and $z(\omega) = 0$ if and only if $\omega = 0$,
- for all $\omega \in B(0, \delta) \subseteq \mathbb{T}^d$, we have

$$z(\omega) = -4\pi^2 \omega \cdot \Sigma \omega + O(|\omega|^3).$$

The proof: combining the pieces

The proof: combining the pieces

- Step 1: Fourier decomposition

$$\begin{aligned}\mathcal{J}_T f(x) &= \int_{\mathbb{T}^d} \int_0^T \pi_\omega(f) \circ h_s(x) \, ds \, d\omega \\ &= \int_{\mathbb{T}^d} \int_0^T \Xi_{-\omega} \circ \pi_0(f) \circ \Xi_\omega \circ h_s(x) \, ds \, d\omega.\end{aligned}$$

The proof: combining the pieces

- Step 1: Fourier decomposition

$$\mathcal{I}_T f(x) = \int_{\mathbb{T}^d} \int_0^T \Xi_{-\omega}(f_\omega) \circ h_s(x) ds d\omega.$$

- Step 2: renormalization

The proof: combining the pieces

- Step 1: Fourier decomposition

$$\mathcal{I}_T f(x) = \int_{\mathbb{T}^d} \int_0^T \Xi_{-\omega}(f_\omega) \circ h_s(x) ds d\omega.$$

- Step 2: renormalization

$$\begin{aligned} \mathcal{I}_T f(x) &= \int_{\mathbb{T}^d} \int_0^1 \mathcal{L}_{t_*}[\Xi_{-\omega}(f_\omega)] \circ h_s(x) ds d\omega \\ &= \int_{\mathbb{T}^d} \int_0^1 \Xi_{-\omega} \circ \mathcal{L}_{t_*}^{(\omega)}(f_\omega) \circ h_s(g_{t_*}(x)) ds d\omega \\ &\approx \int_{\mathbb{T}^d} e^{-2\pi i \int_x^{g_{t_*}(x)} p^* \omega} L_{g_{t_*}(x), \varphi} \left[\mathcal{L}_{t_*}^{(\omega)}(f_\omega) \right] d\omega \end{aligned}$$

The proof: combining the pieces

- Step 1: Fourier decomposition

$$\mathcal{J}_T f(x) = \int_{\mathbb{T}^d} \int_0^T \Xi_{-\omega}(f_\omega) \circ h_s(x) \, ds \, d\omega.$$

- Step 2: renormalization

$$\mathcal{J}_T f(x) \approx \int_{\mathbb{T}^d} e^{-2\pi i \int_x^{\mathbf{g}_{t_*}(x)} p^* \omega} L_{\mathbf{g}_{t_*}(x), \varphi} \left[\mathcal{L}_{t_*}^{(\omega)}(f_\omega) \right] d\omega$$

- Step 3: spectral theory

The proof: combining the pieces

- Step 1: Fourier decomposition

$$\mathcal{I}_T f(x) = \int_{\mathbb{T}^d} \int_0^T \Xi_{-\omega}(f_\omega) \circ h_s(x) ds d\omega.$$

- Step 2: renormalization

$$\mathcal{I}_T f(x) \approx \int_{\mathbb{T}^d} e^{-2\pi i \int_x^{\mathbf{g}_{t_*}(x)} p^* \omega} L_{\mathbf{g}_{t_*}(x), \varphi} \left[\mathcal{L}_{t_*}^{(\omega)}(f_\omega) \right] d\omega$$

- Step 3: spectral theory

$$\begin{aligned} \mathcal{I}_T f(x) &\approx \int_{\mathbb{T}^d} e^{-2\pi i \int_x^{\mathbf{g}_{t_*}(x)} p^* \omega} e^{h_{\text{top}} t_*} L_{\mathbf{g}_{t_*}(x), \varphi} \left[e^{z(\omega)t} \Pi_\omega f_\omega + Q_{\omega, t} f_\omega \right] d\omega \\ &\approx \int_{B(0, \delta)} e^{-2\pi i \int_x^{\mathbf{g}_{t_*}(x)} p^* \omega} e^{h_{\text{top}} t_*} e^{-4\pi^2 \omega \cdot \Sigma \omega t_*} L_{\mathbf{g}_{t_*}(x), \varphi} [\Pi_\omega f_\omega] d\omega \end{aligned}$$

The proof: combining the pieces

- Step 1: Fourier decomposition

$$\mathcal{I}_T f(x) = \int_{\mathbb{T}^d} \int_0^T \Xi_{-\omega}(f_\omega) \circ h_s(x) ds d\omega.$$

- Step 2: renormalization

$$\mathcal{I}_T f(x) \approx \int_{\mathbb{T}^d} e^{-2\pi i \int_x^{\mathbf{g}_{t_*}(x)} p^* \omega} L_{\mathbf{g}_{t_*}(x), \varphi} \left[\mathcal{L}_{t_*}^{(\omega)}(f_\omega) \right] d\omega$$

- Step 3: spectral theory

$$\mathcal{I}_T f(x) \approx \int_{B(0, \delta)} e^{-2\pi i \int_x^{\mathbf{g}_{t_*}(x)} p^* \omega} e^{h_{\text{top}} t_*} e^{-4\pi^2 \omega \cdot \Sigma \omega t_*} L_{\mathbf{g}_{t_*}(x), \varphi} [\Pi_\omega f_\omega] d\omega$$

- Step 4: computations of exponential integrals (“stationary phase”-type estimates)

Thank you for your attention.

Happy birthday, Giovanni!