GENERALIZED HERMITE–BIRKHOFF INTERPOLATION OF ARBITRARY ORDER IN ANY DIMENSION VIA SINGULAR-WEIGHT MOVING LEAST SQUARES

DAVID FIGUEIRAS-GARCIA

Abstract. Singular-weight interpolating moving least squares (SWIMLS) reproduces function values at scattered nodes. We study the method when each node carries an arbitrary finite set of linear differential data $q(D)f(x_i)$. We prove that the interpolant is C^∞, give an exact representation of the error in every datum, and deduce that a datum is reproduced exactly if (i) every derivative of its symbol q lies in the span of the symbols prescribed at that node and (ii) $\Delta^{\alpha/2}q = 0$, where α is the (even) order of the weight singularity. Condition (ii) cannot be dropped. Whether condition (i) is needed depends on how the fit to the other nodes moves near x_i: when it does not move, every datum with $\Delta^{\alpha/2}q = 0$ is reproduced, and when (i) fails only at first order the error is given explicitly. For classical Hermite data of order k, all data are reproduced if and only if $\alpha > k$ (the necessity under a mild condition on the other nodes), which proves a sharpened form of a conjecture of Komargodski and Levin. We also give exact counterexamples, in one and two dimensions, to their claim that derivatives prescribed at points without function values are interpolated, and a completion procedure that makes such data exact. With compactly supported C^∞ weights, local polynomials of degree m and a local norming condition on the data, the interpolant and its derivatives of order $|\beta| \le m$ converge at the rate $h^{m+1-|\beta|}$ on quasi-uniform nodes, at and near the nodes included. A stable evaluator and experiments in one to three dimensions confirm the theory.

1. Introduction

Moving least squares (MLS) builds an approximant point by point: at each evaluation point x one fits a polynomial to the data by weighted least squares, with weights that decay away from x, and evaluates the fit at x. The method goes back to McLain [24, 25] and Lancaster and Šalkauskas [20]; its approximation theory is due to Levin [21] and Wendland [39], see also [15]. When the weights are singular at the nodes, $w_i(x) = |x - x_i|^{-\alpha}$, the approximant interpolates the data [20]; with constant local polynomials it is Shepard's method [34]. Interpolating MLS needs no mesh and no global linear system, and it is local, smooth and exact on polynomials of the local degree; MLS is also optimal in the sense of Backus and Gilbert [4], and nearly interpolating weights are used to impose boundary conditions in meshless methods [28].

In many applications the data include derivatives: slopes and curvatures in curve and surface design, gradients in derivative-enhanced surrogate models, fluxes and normal

Date: Draft of October 29, 2026.

derivatives in meshless discretisations of differential equations. Kernel methods handle such Hermite–Birkhoff data by building the derivatives into the kernel matrix [41, 30], at the price of a global system. In meshless methods, derivative data have been built into MLS fits [3], and generalized MLS approximates derivatives and other functionals from values [27]. Interpolating MLS fits of potential energy surfaces in chemistry incorporate gradient data [37], and supplementary derivative data raise the order of any polynomial-reproducing scheme, MLS included [19]. For interpolating MLS, Komargodski and Levin [18] proposed using the same singular weights for value and derivative data at a node. They proved that this interpolates values and first derivatives in one dimension, and reported that for data up to second derivatives $\alpha = 2$ is not enough while $\alpha = 4$ works. They conjectured that $\alpha = 2k$ suffices for derivatives up to order k, and stated that the method also interpolates directional derivatives prescribed at points that carry no function value. In geometry processing, point set surfaces interpolate scattered points through singular weights; Alexa and Adamson [1] found the direct extension of that construction to Hermite (normal) data numerically unstable and proposed a modified scheme. For SWIMLS the corresponding numerical difficulty near a node is handled by the evaluator of Appendix A.

For Shepard's method, the case of local constants, derivative data have a longer history: Farwig's Shepard–Taylor operator interpolates derivatives up to order r when the exponent of the weights is at least $r + 1$ [14], Coman [6] treats Birkhoff-type data, and Allasia and Bracco [5] and Dell'Accio and Di Tommaso [11] construct Hermite–Birkhoff interpolants under completeness conditions on the data at each node; see the survey [12]. For interpolating MLS with function values, the derivatives of the interpolant at the nodes were computed by Sonar [35] and Netuzhylov, Sonar and Yomsatieankul [29].

Polynomial Hermite–Birkhoff interpolation in several variables is notoriously delicate: whether a data pattern is poised depends on the pattern and on the geometry of the nodes [23, 17, 22]; the natural class of local data is that of D-invariant spaces [10, 31], which characterise ideal interpolation [8, 33]. Interpolating MLS avoids the global poisedness question, since it only asks that the data be unisolvent on the local polynomial space. It replaces it by a local question: which data are reproduced? This paper answers that question.

We call the method SWIMLS (singular-weight interpolating moving least squares) and analyse it for arbitrary finite sets of linear differential data $q(D)f(x_i)$ at each node, in any dimension. The analysis rests on two elementary tools, the Fischer pairing of a symbol with a polynomial and a structure theorem for the local fit near a node, and it leads to an exact formula for the error in each datum.

Contributions.

(1) A structure theorem for SWIMLS near a node (Theorem 4.1), which splits the local fit into the part the data at the node see, the part they do not see, and a remainder of order $|x - x_i|^\alpha$; and an exact representation of the error in any

differential datum (Theorems 4.4 and 4.12). That the interpolant is C^∞ also follows from Levin's Backus–Gilbert argument (Remark 4.2).

(2) A sufficient condition for exact reproduction of a datum $q(D)f(x_i)$: closure of the symbol under differentiation and $\Delta^{\alpha/2}q = 0$ (Corollary 4.5). The second condition cannot be dropped: it is necessary at leading order (Proposition 4.9).

(3) The role of the first condition: when the minimiser of the "frozen" problem at the other nodes does not move near x_i, every datum with $\Delta^{\alpha/2}q = 0$ is reproduced, whether or not the first condition holds (Theorem 4.13), e.g. for a single other node or locally proportional weights; when the first condition fails only at first order, the error is given explicitly by Proposition 4.16; and for pure power weights a datum is either always reproduced or not reproduced for generic node positions (Proposition 4.17).

(4) For classical Hermite data of order k: all data are reproduced when $\alpha > k$, and this is sharp (Corollaries 4.6 and 4.11); when $\alpha \le k$ the error is the linear functional $q \mapsto \left[\Delta^{\alpha/2}q, \tau\right]_{x_i}$ of the datum. This proves the conjecture of [18, §3] that $\alpha = 2k$ suffices, and sharpens it for $k \ge 3$.

(5) Exact counterexamples in one and two dimensions to the extension claimed in [18, §3] (Propositions 5.1 and 5.2), and a completion procedure that makes every datum with $\Delta^{\alpha/2}q = 0$ exact, gaps or not (Proposition 5.3).

(6) A convergence theorem for compactly supported C^∞ singular weights (Theorem 6.1): $|D^\beta(u-f)| \le C\delta^{m+1-|\beta|}$ for $|\beta| \le m$, under a local norming condition on the data, uniformly up to the nodes. This extends Levin's approximation-order result for function values [21] to Hermite–Birkhoff data and to derivatives.

(7) Numerical experiments in one, two and three dimensions that confirm each of these statements datum by datum, based on a numerically stable evaluator of SWIMLS.

Outline. Section 2 shows the method and both failure mechanisms on a one-dimensional example. Section 3 fixes notation, Section 4 contains the exactness theory, Section 5 treats derivatives at points without values, and Section 6 proves convergence for compactly supported weights. Section 7 reports the experiments and Section 8 lists open problems. Appendices A and B describe the numerical evaluator and how exactness was verified.

2. A one-dimensional example

Take the nodes $x_i = i$, $i = 0, 1, 2, 3$, prescribe values f_i and slopes f_i', and fix the exponent $\alpha = 2$. At every point x that is not a node, let p_x be the quadratic polynomial that minimises

$$J_x(p) = \sum_{i=0}^{3} \frac{\big(p(x_i) - f_i\big)^2 + \big(p'(x_i) - f_i'\big)^2}{(x - x_i)^2}, \tag{1}$$

and set $u(x) := p_x(x)$. The minimiser is linear in the data, so $u = \sum_i \big(f_i\varphi_i + f_i'\psi_i\big)$ with shape functions φ_i, ψ_i; two of them are shown in Figure 1(a). They have the cardinal property of Hermite interpolation: $\varphi_i(x_j) = \delta_{ij}$, $\varphi_i'(x_j) = 0$, $\psi_i(x_j) = 0$ and $\psi_i'(x_j) = \delta_{ij}$. Nothing in (1) uses the ordering of the nodes or a mesh, and the same definition works with any number of nodes in any dimension.

Why are the slopes reproduced? Near the node $x_1 = 1$, multiply (1) by $(x-1)^2$. The first term becomes $(p(1) - f_1)^2 + (p'(1) - f_1')^2$ and the others are multiplied by the small factor $t = (x-1)^2$. Write the quadratic as $p = p^* + r + c\,(y-1)^2$, where p^* is the line with value f_1 and slope f_1' at 1 and r is a linear polynomial: the node data see r but not c. Balancing the two parts of the functional gives $r = O(t)$, while c is determined by the other nodes. Section 4 makes this precise:

$$p_x(y) = p^*(y) + \nu(x)\,(y-1)^2 + (x-1)^2\,\sigma(x)(y),$$

with smooth coefficients ν and σ. At $y = x$ both correction terms vanish to second order at $x = 1$, so $u(1) = f_1$ and $u'(1) = f_1'$.

The same picture explains what can go wrong. If we also prescribe second derivatives and use local cubics, the correction $(x-1)^2\sigma(x)(x)$ contributes $2\sigma(1)(1)$ to $u''(1)$, and the datum $f''(1)$ is missed (Figure 1(b): $u''(1) = 1.05$ against $f''(1) = -3.04$). With $\alpha = 4$ the correction is $(x-1)^4\sigma(x)(x)$, which does not affect the second derivative, and $f''(1)$ is reproduced. This is the rule "α larger than the order of the data" of Corollary 4.6.

The other failure concerns gaps in the data. Prescribe values at 0 and 2 and slopes at 1 and 3 (Figure 1(c)). At $x_1 = 1$ the only datum, the slope, does not see constants. The component $\nu_0(x)$ of p_x along the constants is then chosen by the other nodes, it varies with x, and its derivative $\nu_0'(1)$ adds to $u'(1)$. The prescribed slope 0.306 comes out as 0.235. Adding an estimate of the missing values $f(1)$ and $f(3)$ to the data, by any method, restores exactness (Proposition 5.3); the slopes are then reproduced exactly (to the accuracy of the finite differences used to check this).

3. Setting and notation

3.1. **Polynomials, symbols and the Fischer pairing.** For $d \geq 1$ and $m \geq 0$ let $\Pi_m = \Pi_m(\mathbb{R}^d)$ be the real polynomials of total degree at most m. Multi-indices are $\beta \in \mathbb{N}_0^d$, with $|\beta| = \sum_j \beta_j$, $\beta! = \prod_j \beta_j!$, $D^\beta = \partial^{|\beta|}/\partial x_1^{\beta_1}\cdots\partial x_d^{\beta_d}$. A polynomial $q(\xi) = \sum_\beta q_\beta \xi^\beta$, called a *symbol*, defines the differential operator $q(D) = \sum_\beta q_\beta D^\beta$. We write $\partial^\gamma q$ for derivatives of a symbol with respect to ξ, and $\Delta q = \sum_j \partial_j^2 q$.

For a symbol q, a smooth function g and a point z we set

$$\big[q, g\big]_z := \big(q(D)g\big)(z), \qquad \big[q, g\big] := \big[q, g\big]_0. \tag{2}$$

For polynomials q, p, $\big[q, p\big] = \sum_\beta \beta!\, q_\beta p_\beta$ is the *Fischer inner product* [16, 32]; it is positive definite on Π_m. Four elementary facts will be used.




FIGURE 1. The one-dimensional example. (a) Shape functions for the value and the slope at $x = 0$ (nodes $0, 1, 2, 3$, $m = 2$, $\alpha = 2$). (b) Data f, f', f'' at the nodes: u'' near the node $x = 1$ for $\alpha = 2$ and $\alpha = 4$; only $\alpha = 4$ reproduces $f''(1)$ (Corollary 4.6). (c) Values at $0, 2$ and slopes at $1, 3$, $m = 2$: the slopes are not reproduced (compare Proposition 5.1) until the data are completed (Proposition 5.3).

Lemma 3.1 (Fischer adjointness)**.** *For symbols q and polynomials p, $\big[q, z_j p\big] = \big[\partial_j q, p\big]$. Consequently $\big[q, |z|^{2\ell} p\big] = \big[\Delta^\ell q, p\big]$ for every $\ell \in \mathbb{N}_0$, where $|z|^2 = \sum_j z_j^2$.*

Proof. With e_j the j-th unit multi-index, $\big[q, z_j p\big] = \sum_\kappa \kappa!\, q_\kappa p_{\kappa - e_j} = \sum_\lambda (\lambda + e_j)!\, q_{\lambda + e_j} p_\lambda = \sum_\lambda \lambda!\, (\lambda_j + 1) q_{\lambda + e_j} p_\lambda = \big[\partial_j q, p\big]$. Multiplication by $|z|^2 = \sum_j z_j \cdot z_j$ is therefore adjoint to $\sum_j \partial_j^2 = \Delta$, and the second statement follows by induction. □

Fischer's pairing and this adjointness go back to [16]; see also [32]. They give an orthogonal decomposition that explains the role of the operator $\Delta^{\alpha/2}$ below.

Remark 3.2. For $\mu, \ell \in \mathbb{N}_0$ with $2\ell \le \mu$,

$$\Pi_\mu = \ker\big(\Delta^\ell\big|_{\Pi_\mu}\big) \;\oplus\; |z|^{2\ell}\, \Pi_{\mu - 2\ell}, \tag{3}$$

and the sum is orthogonal for the Fischer pairing. Indeed, $\big[q, |z|^{2\ell} p\big] = \big[\Delta^\ell q, p\big] = 0$ for $\Delta^\ell q = 0$ by Lemma 3.1, and the dimensions add up because $\Delta^\ell \colon \Pi_\mu \to \Pi_{\mu - 2\ell}$ is the adjoint of the injective multiplication by $|z|^{2\ell}$, hence onto. Consequently, a symbol q of degree at most μ annihilates every multiple $|z|^{2\ell} p$ of degree at most μ if and only if $\Delta^\ell q = 0$. For $\ell = 1$ and homogeneous polynomials this is a classical lemma of Calderón and Zygmund [7, Ch. 2, Lemma 1], and in the form "P is divisible by $|x - a|^2$ if and only if $q(D)P(a) = 0$ for every harmonic q of degree at most $\deg P$" it is Theorem 2.4 of [36], which studies Hermite interpolation with conditions induced by harmonic polynomials; see also [38, Ch. IV] for the decomposition into harmonic polynomials, and [9] for the role of this pairing in polynomial interpolation. With $\ell = \alpha/2$ it is the reason for condition (ii) below: along the diagonal, the remainder of the local fit near a node is a multiple of $|z - x_i|^\alpha$ (Theorems 4.1 and 4.4), and the data that cannot see any such multiple are exactly the $(\alpha/2)$-polyharmonic symbols.

Lemma 3.3 (Taylor expansion of operators)**.** *Let $A = (A_1, \dots, A_d)$ and $B = (B_1, \dots, B_d)$ be commuting operators. For every symbol q, $q(A+B) = \sum_\gamma \frac{1}{\gamma!} A^\gamma (\partial^\gamma q)(B)$ (a finite sum).*

Proof. This is Taylor's formula $q(a+b) = \sum_\gamma \frac{a^\gamma}{\gamma!}(\partial^\gamma q)(b)$ for the polynomial q, an identity in the commutative polynomial ring in a, b. □

Lemma 3.4. *Let $\ell \in \mathbb{N}_0$ and $\beta \in \mathbb{N}_0^d$. Then $\Delta^\ell \xi^\beta = 0$ if and only if $\sum_j \lfloor \beta_j/2 \rfloor < \ell$.*

Proof. By the multinomial formula, $\Delta^\ell \xi^\beta = \sum_{|\kappa|=\ell} \frac{\ell!}{\kappa!}\, \partial^{2\kappa} \xi^\beta$, and $\partial^{2\kappa}\xi^\beta = \frac{\beta!}{(\beta-2\kappa)!}\xi^{\beta-2\kappa}$ if $2\kappa \le \beta$ and 0 otherwise. Distinct κ give distinct monomials and all coefficients are positive, so $\Delta^\ell \xi^\beta \neq 0$ if and only if some κ with $|\kappa| = \ell$ satisfies $2\kappa \le \beta$, i.e. $\sum_j \lfloor \beta_j/2 \rfloor \ge \ell$. □

Lemma 3.5. *Let $\alpha \in 2\mathbb{N}$. Then $\sum_{|\gamma|=\alpha} \frac{1}{\gamma!}\big(D^\gamma |x|^\alpha\big)(0)\, \partial^\gamma = \Delta^{\alpha/2}$ as operators on symbols.*

Proof. $|x|^\alpha = \sum_{|\kappa|=\alpha/2} \frac{(\alpha/2)!}{\kappa!} x^{2\kappa}$, and $(D^\gamma x^{2\kappa})(0) = (2\kappa)!$ if $\gamma = 2\kappa$ and 0 otherwise. Hence the left side equals $\sum_{|\kappa|=\alpha/2} \frac{(\alpha/2)!}{\kappa!}\partial^{2\kappa} = \big(\sum_j \partial_j^2\big)^{\alpha/2}$. □

3.2. **Data.** Let $\Omega \subset \mathbb{R}^d$ be open and $X = \{x_1, \dots, x_K\} \subset \Omega$ distinct nodes. Node x_i carries symbols $q_{i,1}, \dots, q_{i,n_i}$ of degree at most m and values $y_{i,1}, \dots, y_{i,n_i} \in \mathbb{R}$, i.e. the interpolation conditions

$$\lambda_{i,s}(f) := \big[q_{i,s}, f\big]_{x_i} = \big(q_{i,s}(D) f\big)(x_i) = y_{i,s}, \qquad s = 1, \dots, n_i,\ i = 1, \dots, K. \tag{4}$$

We write $Q_i = \operatorname{span}\{q_{i,1}, \dots, q_{i,n_i}\}$ for the space of symbols at x_i. Every $q = \sum_s c_s q_{i,s} \in Q_i$ has a prescribed value $y_q := \sum_s c_s y_{i,s}$; we say that the datum q is *reproduced* by a function u if $\big[q, u\big]_{x_i} = y_q$. Classical Hermite data of order k correspond to $Q_i = \operatorname{span}\{\xi^\beta : |\beta| \le k\}$; any other choice will be called Hermite–Birkhoff data. A space of symbols is *D-invariant* if it is closed under differentiation.

Assumption 3.6. (A1) $\alpha \in 2\mathbb{N}$ is even, and $w_i(x) = \rho_i(x)\,|x - x_i|^{-\alpha}$ with $\rho_i \in C^\infty(\Omega)$, $\rho_i > 0$.

(A2) For every i the symbols $q_{i,1}, \dots, q_{i,n_i}$ are linearly independent.

(A3) (Unisolvency) If $p \in \Pi_m$ and $\lambda_{i,s}(p) = 0$ for all i, s, then $p = 0$.

Since $\big[\cdot, \cdot\big]_{x_i}$ is non-degenerate on Π_m (it is the Fischer pairing after translation) and $\deg q_{i,s} \le m$, (A2) is equivalent to the linear independence of the functionals $\lambda_{i,1}, \dots, \lambda_{i,n_i}$ on Π_m.

Definition 3.7 (SWIMLS)**.** For $x \in \Omega \setminus X$ let

$$J_x(p) = \sum_{i=1}^{K} w_i(x) \sum_{s=1}^{n_i} \big(\lambda_{i,s}(p) - y_{i,s}\big)^2, \qquad p_x := \operatorname*{arg\,min}_{p \in \Pi_m} J_x(p), \tag{5}$$

and define the interpolant $u(x) := p_x(x)$.

By (A1) and (A3), J_x is a strictly convex quadratic on Π_m for every $x \notin X$, so p_x is unique; it depends smoothly on $x \in \Omega \setminus X$ because the coefficients of J_x do and its Hessian is positive definite. Hence $u \in C^\infty(\Omega \setminus X)$. The behaviour at the nodes is the subject of the next section.

Remark 3.8. Positive smooth factors for the individual functionals of a node can be included: the first block of the Jacobian in the proof of Theorem 4.1 becomes $2L^*W(x)L$, still invertible, and nothing else changes. If some functionals at x_i are *not* given the singular factor, they must be moved from Q_i to the regular part B_x of the proof, and Q_i, N_i refer to the singular ones only.

4. Main results

This section answers one question: when is a datum $q(D)f(x_i)$ reproduced? The answer comes in four steps, which follow the one-dimensional picture of Section 2.

1. Theorem 4.1 describes the local fit p_x near x_i. It splits into a part fixed by the data at x_i, a part $\nu(x)$ that those data cannot see and that the other nodes choose, and a remainder damped by $|x - x_i|^\alpha$.
2. Theorem 4.4 turns this into an exact formula for the error in any datum, with one term for each part.
3. Each term gives a condition for exactness (Section 4.3): the ν-term vanishes if the derivatives of the symbol q are again data at x_i (condition (i)), and the remainder vanishes if $\Delta^{\alpha/2}q = 0$ (condition (ii)). For classical Hermite data this gives the rule $\alpha > k$.
4. Sections 4.4 and 4.5 ask how far each condition is necessary.

Throughout this section we fix a node x_i and write

$$N_i = \{p \in \Pi_m : \lambda_{i,s}(p) = 0,\ s = 1, \dots, n_i\} = \{p \in \Pi_m : \big[q, p\big]_{x_i} = 0\ \forall q \in Q_i\}$$

for the polynomials that the data at x_i do not see.

4.1. Structure near a node.

Theorem 4.1 (Structure near a node)**.** *Let Assumption 3.6 hold, fix an inner product on Π_m and let $R = N_i^\perp$. There are an open neighbourhood $U \subset \Omega$ of x_i with $U \cap X = \{x_i\}$, a polynomial $p_i^* \in R$ with $\lambda_{i,s}(p_i^*) = y_{i,s}$ for all s, and C^∞ maps $\nu : U \to N_i$ and $\sigma : U \to R$ such that*

$$p_x = p_i^* + \nu(x) + |x - x_i|^\alpha\, \sigma(x) \qquad \text{for all } x \in U \setminus \{x_i\}. \tag{6}$$

More precisely, $\nu(x) = n(x, |x - x_i|^\alpha)$ and $\sigma(x) = s(x, |x - x_i|^\alpha)$ for C^∞ maps n, s defined near $(x_i, 0)$, and $Z(x) := n(x, 0)$ is the frozen minimiser

$$Z(x) = \operatorname*{arg\,min}_{n \in N_i} B_x(p_i^* + n), \qquad B_x(p) := \sum_{j \neq i} \frac{w_j(x)}{\rho_i(x)} \sum_{s=1}^{n_j} \big(\lambda_{j,s}(p) - y_{j,s}\big)^2. \tag{7}$$

Applying this at every node, u extends to a function in $C^\infty(\Omega)$, with node value $u(x_i) = \big(p_i^* + Z(x_i)\big)(x_i)$.

Proof. Write $\Lambda_i : \Pi_m \to \mathbb{R}^{n_i}$, $p \mapsto (\lambda_{i,s}(p))_s$, and Λ_j likewise. By (A2) Λ_i is surjective with kernel N_i, so $L := \Lambda_i|_R : R \to \mathbb{R}^{n_i}$ is bijective; put $p_i^* := L^{-1}y_i \in R$. Every $p \in \Pi_m$ is uniquely $p = p_i^* + r + n$ with $r \in R$, $n \in N_i$, and then $\Lambda_i p - y_i = Lr$.

Let U be an open neighbourhood of x_i containing no other node. For $x \in U \setminus \{x_i\}$ divide J_x by $w_i(x) > 0$:

$$\frac{J_x(p)}{w_i(x)} = |Lr|^2 + t(x)\, B_x(p), \qquad t(x) := |x - x_i|^\alpha,$$

with B_x as in (7). Since α is even, $t(x) = \big(\sum_k (x_k - x_{i,k})^2\big)^{\alpha/2}$ is a polynomial in x; and $x \mapsto B_x$ is a C^∞ family of convex quadratics on U, because $\rho_i > 0$ and the w_j, $j \neq i$, are smooth on U. A polynomial $p = p_i^* + r + n$ minimises J_x if and only if the gradient vanishes (the problem is convex), that is

$$2L^*Lr + t(x)\nabla_R B_x(p) = 0, \qquad t(x)\nabla_N B_x(p) = 0,$$

where ∇_R, ∇_N denote the components of the gradient in R and N_i and L^* is the adjoint of L. Since $t(x) > 0$ for $x \neq x_i$ we may substitute $r = t(x)\, s$ and divide by $t(x)$, obtaining $F(x, t(x); s, n) = 0$ for the map $F : U \times \mathbb{R} \times R \times N_i \to R \times N_i$,

$$F(x, t; s, n) := \Big(2L^*Ls + \nabla_R B_x(p_i^* + ts + n),\ \nabla_N B_x(p_i^* + ts + n)\Big),$$

which is C^∞ in all variables, including t near 0.

The point $(x_i, 0)$. Here $F = 0$ reads

$$\nabla_N B_{x_i}(p_i^* + n) = 0, \qquad s = -(2L^*L)^{-1}\nabla_R B_{x_i}(p_i^* + n).$$

The restriction of B_x to the affine space $p_i^* + N_i$ is strictly convex for every $x \in U$: its Hessian is the restriction to N_i of

$$2\sum_{j \neq i} \frac{w_j(x)}{\rho_i(x)} \Lambda_j^* \Lambda_j,$$

with $w_j(x) > 0$ since $x \neq x_j$; and if $n \in N_i$ satisfies $\Lambda_j n = 0$ for all $j \neq i$ then all data functionals vanish on n, so $n = 0$ by (A3). Hence there is exactly one solution (s_0, n_0), and $n_0 = Z(x_i)$; more generally, for every $x \in U$ the equation $F(x, 0; s, n) = 0$ has the unique solution

$$n = Z(x), \qquad s = -(2L^*L)^{-1}\nabla_R B_x\big(p_i^* + Z(x)\big).$$

Implicit function theorem. At $(x_i, 0; s_0, n_0)$ the derivative of F with respect to (s, n) is block triangular,

$$\frac{\partial F}{\partial(s, n)} = \begin{pmatrix} 2L^*L & \big(\nabla^2 B_{x_i}\big)_{RN} \\ 0 & \big(\nabla^2 B_{x_i}\big)_{NN} \end{pmatrix},$$

because every derivative with respect to s of a B-term carries the factor $t = 0$. Both diagonal blocks are invertible (L is injective; the second block is the Hessian above). By the implicit function theorem there are a neighbourhood $U' \times (-\tau_0, \tau_0)$ of $(x_i, 0)$ and a C^∞ map $(x,t) \mapsto (s(x,t), n(x,t))$ with $F(x,t; s(x,t), n(x,t)) = 0$; by the uniqueness just shown, $n(x,0) = Z(x)$. Shrinking U to $U' \cap \{t(x) < \tau_0\}$, for $x \in U \setminus \{x_i\}$ the polynomial $p_i^* + t(x)s(x,t(x)) + n(x,t(x))$ satisfies the gradient equations of the strictly convex function J_x, hence it equals p_x (only uniqueness of the critical point of J_x is used). Setting $\nu(x) := n(x,t(x)) \in N_i$ and $\sigma(x) := s(x,t(x)) \in R$ gives (6); both maps are C^∞ because t is a polynomial. Finally $u(x) = p_x(x)$ is, by (6), the restriction to $U \setminus \{x_i\}$ of the C^∞ function $x \mapsto (p_i^* + \nu(x) + t(x)\sigma(x))(x)$, whose value at x_i is $(p_i^* + Z(x_i))(x_i)$. □

In words: as $x \to x_i$ the singular weight forces the local fit to match the data at x_i. The part of p_x these data can see converges to p_i^*, with an error of order $|x - x_i|^\alpha$; this is the term $|x - x_i|^\alpha \sigma(x)$. The part they cannot see, $\nu(x) \in N_i$, is decided by the other nodes: at $x = x_i$ it is the polynomial $Z(x_i)$ that fits their data best among all polynomials that match the data at x_i, but in general it moves with x. Both ν and σ are smooth, and this is what makes u smooth at the nodes.

Remark 4.2. The smoothness of u alone can also be obtained from the Backus–Gilbert form of moving least squares [21, Proposition 1], in which the coefficients of the data minimise $\sum w_j(x)^{-1} a_{j,s}^2$ subject to polynomial reproduction; the singular weights then enter through the smooth functions $w_j^{-1} = \rho_j^{-1}|x - x_j|^\alpha$. The augmented matrix $\left(\begin{smallmatrix} D & E \\ E^\mathsf{T} & 0 \end{smallmatrix}\right)$ of this problem, with $D = 2\,\mathrm{diag}(w_j^{-1})$, remains invertible at $x = x_i$, given (A3), exactly when the data at x_i are linearly independent on Π_m, i.e. under (A2); for function values this is Levin's argument [21, Proposition 2 and Theorem 3]. Theorem 4.1 gives more: the decomposition (6), on which the error analysis rests.

Remark 4.3. The decomposition (6) is canonical up to the choice of the complement R of N_i; the node value $u(x_i)$ and all errors below do not depend on it. Evenness of α is used twice: to make $t(x) = |x - x_i|^\alpha$ smooth at x_i (Theorem 4.1) and to make it a polynomial, so that Lemmas 3.1 and 3.5 apply (Theorems 4.4 and 4.13). If the ρ_i are real analytic, the analytic implicit function theorem shows that u is real analytic. For odd α the function t is not smooth at x_i, our arguments do not apply, and we do not consider this case.

4.2. **An exact error representation.**

Theorem 4.4 (Error representation)**.** *With the notation of Theorem 4.1, let* $\tau(z) := \sigma(z)(z)$ *for* $z \in U$ *(a* C^∞ *scalar function). For every symbol* q*,*

$$[q,u]_{x_i} = [q,p_i^*]_{x_i} + \sum_\gamma \frac{1}{\gamma!} D^\gamma \Big[x \mapsto [\partial^\gamma q, \nu(x)]_{x_i}\Big](x_i) + [\Delta^{\alpha/2} q, \tau]_{x_i}. \tag{8}$$

The $\gamma = 0$ *term of the middle sum is* $[q, \nu(x_i)]_{x_i}$.

Proof. On $U \times \mathbb{R}^d$ consider the C^∞ function $\Phi(x,y) = p_i^*(y) + \nu(x)(y) + t(x)\,\sigma(x)(y) =: \Phi_1 + \Phi_2 + \Phi_3$, which is a polynomial in y and satisfies $u(z) = \Phi(z,z)$ for all $z \in U$, including $z = x_i$ by the definition of the extension. By the chain rule along the diagonal, $D_z^\beta[\Phi(z,z)] = \big[(D_x + D_y)^\beta \Phi\big](z,z)$, hence $\big[q,u\big]_{x_i} = \big[q(D_x + D_y)\Phi\big](x_i,x_i)$. We treat the three parts.

Φ_1 does not depend on x, so it contributes $(q(D)p_i^*)(x_i) = \big[q,p_i^*\big]_{x_i}$.

For Φ_2, Lemma 3.3 with $A = D_x$, $B = D_y$ gives

$$q(D_x + D_y)\Phi_2 = \sum_\gamma \frac{1}{\gamma!} D_x^\gamma\, (\partial^\gamma q)(D_y)\Phi_2.$$

Setting $y = x_i$ commutes with D_x, and $\big((\partial^\gamma q)(D_y)\Phi_2\big)(x,x_i) = \big[\partial^\gamma q, \nu(x)\big]_{x_i}$; setting $x = x_i$ gives the middle term of (8).

For Φ_3 we use the diagonal again: its contribution is $\big[q, t\,\tau\big]_{x_i} = \big(q(D)[\,|z - x_i|^\alpha \tau(z)]\big)(x_i)$. Translate x_i to 0 and let $T\tau$ be the Taylor polynomial of τ at 0 of degree $\deg q$. Since $|z|^\alpha(\tau - T\tau)$ vanishes at 0 to order larger than $\deg q$, it is annihilated by $q(D)$ at 0, and by Lemma 3.1

$$\big[q, |z|^\alpha \tau\big] = \big[q, |z|^\alpha T\tau\big] = \big[\Delta^{\alpha/2} q, T\tau\big] = \big[\Delta^{\alpha/2} q, \tau\big],$$

the last equality because $\deg \Delta^{\alpha/2} q \le \deg q - \alpha \le \deg T\tau$. □

The three terms of (8) correspond to the three parts of (6). The first is the prescribed value y_q whenever $q \in Q_i$. The second measures the motion of the invisible part ν; it pairs the derivatives $\partial^\gamma q$ of the symbol with $\nu(x)$, and every $\partial^\gamma q$ that is itself a datum at x_i drops out, because the data at x_i do not see N_i. The third comes from the damped remainder, and the singularity of the weight enters only through the operator $\Delta^{\alpha/2}$: for the Fischer pairing, multiplication by $|z|^\alpha$ is adjoint to $\Delta^{\alpha/2}$ (Lemma 3.1).

4.3. **Exact reproduction.**

Corollary 4.5 (Sufficient condition)**.** *Let Assumption 3.6 hold and let $q \in Q_i$. If*

(i) $\partial^\gamma q \in Q_i$ for every multi-index γ, and
(ii) $\Delta^{\alpha/2} q = 0$,

then the datum q is reproduced: $\big[q,u\big]_{x_i} = y_q$. If Q_i is D-invariant, the error is the linear functional

$$\big[q,u\big]_{x_i} - y_q = \big[\Delta^{\alpha/2} q, \tau\big]_{x_i}, \qquad q \in Q_i. \tag{9}$$

Proof. In (8), $\big[q,p_i^*\big]_{x_i} = \sum_s c_s \lambda_{i,s}(p_i^*) = y_q$. By (i), every $\partial^\gamma q$ (including $\gamma = 0$) lies in Q_i, and $\nu(x) \in N_i$, so $\big[\partial^\gamma q, \nu(x)\big]_{x_i} = 0$ for all $x \in U$: the middle term vanishes. This proves (9) when Q_i is D-invariant, and the last term vanishes by (ii). □

Condition (i) concerns the datum and the data at its own node only; it holds for every datum when Q_i is D-invariant, the classical setting of multivariate Hermite interpolation [31, 8]. Condition (ii) holds whenever $\deg q < \alpha$, but also, for instance, for

every harmonic symbol when $\alpha = 2$. By Remark 3.2 it is exactly the condition under which the last term of (8) vanishes for every possible remainder τ. At a D-invariant node, (9) says more: the error is a *linear functional* of the datum, which only sees $\Delta^{\alpha/2} q$.

Corollary 4.6 (Hermite data)**.** *Let x_i carry classical Hermite data of order $k \le m$, i.e. $Q_i = \operatorname{span}\{\xi^\beta : |\beta| \le k\}$. Then every $q \in Q_i$ with $\Delta^{\alpha/2} q = 0$ is reproduced. In particular:*

(a) the derivative datum $D^\beta f(x_i)$ is reproduced whenever $\sum_j \lfloor \beta_j/2 \rfloor < \alpha/2$;
(b) all data at x_i are reproduced if $\alpha > k$, i.e. for $\alpha = k+1$ when k is odd and $\alpha = k+2$ when k is even;
(c) for $\alpha = 2$ the monomial data covered by (a) are exactly the multilinear ones ($\beta_j \le 1$ for all j, e.g. $\partial_x\partial_y$, $\partial_x\partial_y\partial_z$), whatever their order.

Proof. Q_i is D-invariant, so Corollary 4.5 applies; (a) and (c) are Lemma 3.4 with $\ell = \alpha/2$, and (b) follows from $\deg q \le k < \alpha$. □

Example 4.7. In two dimensions, let x_i carry $f, \partial_x f, \partial_y f, \partial_x^2 f, \partial_x\partial_y f, \partial_y^2 f$ ($k = 2$). For $\alpha = 4$ all six data are reproduced. For $\alpha = 2$, the symbols $1, \xi_1, \xi_2, \xi_1\xi_2$ are harmonic, so the value, the gradient and the mixed derivative are reproduced; $\partial_x^2 f$ and $\partial_y^2 f$ are not, but by (9) both have the same error $\big[2, \tau\big]_{x_i} = 2\tau(x_i)$, so their difference $\partial_x^2 f - \partial_y^2 f$ is reproduced. With third derivatives ($k = 3$, $\alpha = 2$) the same argument gives, e.g., that the error in $\partial_x^3 f$ is three times the error in $\partial_x\partial_y^2 f$, since $\Delta\xi_1^3 = 6\xi_1 = 3\,\Delta(\xi_1\xi_2^2)$.

Example 4.8. Data given by harmonic symbols are reproduced with the smallest exponent $\alpha = 2$, whatever their order, as soon as the symbol space at each node is D-invariant, since derivatives of harmonic polynomials are harmonic. Toan, Mai, Long and Manh [36, Theorem 4.1] construct regular interpolation schemes of degree $\mu \ge 2$ from such data only, for any distinct nodes: the k-th node carries every harmonic symbol of degree at most $\mu - 2k + 2$, for $k = 1, \dots, \lfloor \mu/2 \rfloor + 1$, and the total number of data equals $\dim \Pi_\mu$. If SWIMLS is used with exactly these data and $m = \mu$, it returns the polynomial interpolant, and every datum is reproduced trivially, for every α. The example becomes informative with more data than $\dim \Pi_m$, for instance more nodes carrying such harmonic patterns, or $m < \mu$: then, with $\alpha = 2$, every datum is reproduced by Corollary 4.5, while non-harmonic data are in general not reproduced at $\alpha = 2$. Section 7.1 checks a node with all harmonic symbols of degree at most 3.

Corollary 4.6(b) proves the conjecture of [18, §3] that $\alpha = 2k$ suffices for data up to order k (they also observed that $\alpha = 2$ fails and $\alpha = 4$ works for $k = 2$). Their conjecture refers to "an appropriate definition of the inner product", which they do not specify; we read it as the same singular weight for all data at a node, as in their one-dimensional construction. For $k \ge 3$ Corollary 4.6(b) is a strict improvement, e.g. $\alpha = 4$ suffices for $k = 3$. For Shepard–Taylor interpolation, where the local polynomials are fixed Taylor polynomials, the analogous condition is that the exponent exceeds the

order of the data [13, 14, 6]. The finer condition $\Delta^{\alpha/2}q = 0$ is not specific to fitting: for Shepard–Taylor interpolation with an even exponent α, the interpolant near x_i is the Taylor polynomial at x_i plus $|x - x_i|^\alpha$ times a smooth function, and the argument of Theorem 4.4 applies verbatim. What fitting adds is the part ν of the local polynomial that the data at x_i do not see, and with it condition (i).

4.4. **Is condition (ii) necessary?** The next result shows that condition (ii) cannot be dropped, at least for data of degree α, and with it that Corollary 4.6(b) is sharp.

Proposition 4.9 (Necessity of (ii), leading order)**.** *In addition to Assumption 3.6, assume that the data at the nodes other than x_i are unisolvent on Π_m by themselves. Let $q \in Q_i$ satisfy (i), with $\deg q = \alpha$ and $\Delta^{\alpha/2}q \neq 0$. Then, for every choice of the data at x_i, the datum q is reproduced only if the data at the other nodes lie in a fixed affine hyperplane.*

Proof. By (i) the middle term of (8) vanishes, and $\Delta^{\alpha/2}q = c$ is a non-zero constant, so the error is $c\,\tau(x_i) = c\,\sigma(x_i)(x_i) = c\,s_0(x_i)$ with s_0 from the proof of Theorem 4.1. The error is an affine function of the data at the other nodes (with y_i fixed); it vanishes for the data $y_j = \Lambda_j p_i^*$, so it suffices to exhibit data for which it does not.

Since $\deg q = \alpha \geq 2$, some derivative $\partial^\gamma q$ with $|\gamma| = \alpha$ is a non-zero constant, so $1 \in Q_i$ by (i): point evaluation $\ell(p) = p(x_i)$ is a datum at x_i. It vanishes on N_i and is non-zero on Π_m, hence non-zero on R.

Choose $y_j = \Lambda_j(p_i^* + v)$, $j \neq i$, for some $v \in \Pi_m$. Then $B_{x_i}(p) = \langle \Theta(p - p_i^* - v), p - p_i^* - v\rangle$ with $\Theta = \sum_{j\neq i} \frac{w_j(x_i)}{\rho_i(x_i)} \Lambda_j^* \Lambda_j$, which is positive definite on Π_m by the additional assumption. The minimiser n_0 over N_i is characterised by $\Theta(n_0 - v) \perp N_i$, i.e. $\Theta(n_0 - v) \in R$, so $\nabla_R B_{x_i}(p_i^* + n_0) = 2\Theta(n_0 - v)$ and $s_0 = (L^*L)^{-1}\Theta w$ with $w = v - n_0$. As v ranges over Π_m, w ranges over the Θ-orthogonal complement $W = \{w : \Theta w \in R\}$ of N_i, so Θw ranges over all of R and s_0 over all of R. Choosing v with $\ell(s_0) \neq 0$ completes the proof. □

Remark 4.10. The extra hypothesis is used only to make the Schur complement $\Theta_{RR} - \Theta_{RN}\Theta_{NN}^{-1}\Theta_{NR}$ of Θ, with respect to $\Pi_m = R \oplus N_i$, injective; Θ_{NN} is invertible by (A3) alone. It cannot simply be dropped. With $d = 1$, $m = 3$, $\alpha = 2$, the data f, f', f'' at 0 and a single value $f(1)$, the data are unisolvent and exactly $\dim \Pi_3$ in number, so u is the cubic interpolant and $f''(0)$ is reproduced, although $\Delta\xi^2 = 2 \neq 0$; here the Schur complement vanishes.

Corollary 4.11 (Sharpness)**.** *Under the hypotheses of Proposition 4.9 on the other nodes, let x_i carry classical Hermite data of order $k \leq m$. If $\alpha \leq k$, the datum $\partial_1^\alpha f(x_i)$ is not reproduced for data at the other nodes outside an affine hyperplane. Hence $\alpha > k$ is the exact condition for all Hermite data of order k to be reproduced for all data values.*

Proof. $q = \xi_1^\alpha \in Q_i$ satisfies (i) by D-invariance, $\deg q = \alpha$ and $\Delta^{\alpha/2}\xi_1^\alpha = \alpha! \neq 0$. □

4.5. **When is condition (i) needed?** When (i) fails, the middle term of (8) involves derivatives of ν, that is, it depends on how the fit to the *other* nodes moves as x approaches x_i. The cleanest way to see this is to separate from ν its value at $t = 0$, the frozen minimiser $Z(x) = n(x, 0)$ of (7). Since n is smooth,

$$\nu(x) = n(x, t(x)) = Z(x) + t(x)\, g(x), \qquad g(x) := \int_0^1 \partial_t n\big(x, \theta t(x)\big)\, d\theta,$$

with $g\colon U \to N_i$ smooth, and the part $t\,g$ can be moved into the remainder.

Theorem 4.12 (Error representation, frozen form)**.** *With the notation of Theorem 4.1, let* $\eta(z) := \big(g(z) + \sigma(z)\big)(z)$ *for* $z \in U$*. For every symbol* q*,*

$$[q, u]_{x_i} = [q, p_i^*]_{x_i} + \sum_{\gamma} \frac{1}{\gamma!}\, D^\gamma \Big[x \mapsto [\partial^\gamma q, Z(x)]_{x_i}\Big](x_i) + [\Delta^{\alpha/2} q, \eta]_{x_i}. \tag{10}$$

Proof. Repeat the proof of Theorem 4.4 with $\Phi(x, y) = p_i^*(y) + Z(x)(y) + t(x)\big(g(x) + \sigma(x)\big)(y)$, which is the same function: the second part now involves Z in place of ν, and the third part is $t(x)$ times a smooth function of (x, y), polynomial in y, whose restriction to the diagonal is η. □

For $q \in Q_i$ the first term of (10) is y_q and the term with $\gamma = 0$ in the sum vanishes, since $Z(x_i) \in N_i$. The error in a datum is therefore the sum of two contributions: the motion of the frozen minimiser, which pairs the derivatives of Z with those derivatives of q that are not data at x_i, and the remainder, which only sees $\Delta^{\alpha/2} q$. If the frozen minimiser does not move, only the second is left.

Theorem 4.13 (Frozen minimiser)**.** *Let Assumption 3.6 hold and suppose that the frozen minimiser* Z *of* (7) *is constant near* x_i*. Then every* $q \in Q_i$ *satisfies*

$$[q, u]_{x_i} - y_q = [\Delta^{\alpha/2} q, \eta]_{x_i}. \tag{11}$$

In particular every $q \in Q_i$ *with* $\Delta^{\alpha/2} q = 0$ *is reproduced, whatever its degree and whether or not (i) holds.*

Proof. In (10), the terms with $\gamma \neq 0$ vanish because Z is constant, and the term with $\gamma = 0$ vanishes because $q \in Q_i$ and $Z(x_i) \in N_i$. □

Corollary 4.14. *The frozen minimiser is constant near* x_i*, and hence Theorem 4.13 applies, in each of the following cases:*

- *(a)* single scale: $w_j(x)/\rho_i(x) = c(x)\kappa_j$ *near* x_i *for all* $j \neq i$*, with a positive function* c *and constants* $\kappa_j > 0$*; this includes the case of a single other node* ($K = 2$)*, weights whose germs at* x_i *are proportional, and weights that are constant near* x_i *for all* $j \neq i$*;*
- *(b)* interpolatory restriction: *the functionals* $\lambda_{j,s}$*,* $j \neq i$*, that do not vanish identically on* N_i *restrict to a basis of the dual of* N_i *(so that the frozen problem can fit them exactly whatever the weights).*

Proof. (a) $B_x = c(x)\hat{B}$ with $\hat{B}$ independent of x, and the minimiser of $c(x)\hat{B}$ over $p_i^* + N_i$ does not depend on $c(x) > 0$. (b) The frozen objective is a positively weighted sum of squared residuals of functionals that are either constant on $p_i^* + N_i$ or, by assumption, can be made zero simultaneously by exactly one $n \in N_i$, which is then the minimiser for every choice of positive weights. □

Example 4.15. These cases are not exotic: for two nodes, gradient data at x_i without a function value are reproduced for $\alpha \geq 2$; in one dimension with $m = 1$, the data $f'(0)$, $f(1)$, $f'(5/2)$ are all reproduced (case (b): only $f(1)$ sees the constants, which form N_i at the slope nodes). A third node in the first case, or a further value datum in the second, destroys the structure, and the data are then no longer reproduced (Section 7).

In general the frozen minimiser moves. When condition (i) fails only at first order, the error is then given explicitly by the following result.

Proposition 4.16 (First-order error)**.** *Let Assumption 3.6 hold and let $q \in Q_i$ with $\Delta^{\alpha/2} q = 0$ (for instance $\deg q < \alpha$), such that $\partial^\gamma q \in Q_i$ for all $|\gamma| \geq 2$. Then*

(12)

$$\big[q, u\big]_{x_i} - y_q = \sum_{j=1}^{d} \big[\partial_j q, \partial_{x_j} Z(x_i)\big]_{x_i}, \qquad \partial_{x_j} Z(x_i) = -H^{-1}\, \partial_{x_j} \big[\nabla_N B_x\big](p_i^* + Z(x_i))\Big|_{x=x_i},$$

where H is the Hessian of B_{x_i} restricted to N_i.

Proof. In (10) the last term vanishes because $\Delta^{\alpha/2} q = 0$, the term with $\gamma = 0$ because $q \in Q_i$, and the terms with $|\gamma| \geq 2$ because $\partial^\gamma q \in Q_i$ and $Z(x) \in N_i$ for all x. Differentiating $\nabla_N B_x(p_i^* + Z(x)) = 0$ gives the formula for $\partial_{x_j} Z$. □

The right-hand side of (12) is an explicit linear functional of the data. It vanishes in the cases of Corollary 4.14 and for symmetric configurations, e.g. for $q = \xi_1$ when $\partial_{x_1}(w_j/\rho_i)(x_i) = 0$ for all $j \neq i$, but it is non-zero in the explicit examples of Section 5. For pure power weights, a dichotomy makes "generic" precise.

Proposition 4.17 (Dichotomy)**.** *Let $\rho_j \equiv 1$ for all j, and fix d, m, α, the number K of nodes, the symbol spaces $Q_1, \ldots, Q_K$, a node index i and a datum $q \in Q_i$. Let $\mathcal{A} \subset (\mathbb{R}^d)^K$ be the set of node positions for which (A3) holds. Then either q is reproduced for all node positions in $\mathcal{A}$ and all data, or there is a proper algebraic subset $\mathcal{Z}$ such that for all node positions in $\mathcal{A} \setminus \mathcal{Z}$ the error $\big[q, u\big]_{x_i} - y_q$ is a non-zero linear functional of the data, so that q is reproduced only for data in a hyperplane.*

Proof. Write polynomials in the variable $y - x_i$, so that Λ_i, N_i and R do not depend on the node positions. With $\rho \equiv 1$ and α even, every weight $w_j(x)/w_i(x)$ is a rational function of x and the node positions, so the map F in the proof of Theorem 4.1 is rational in (x, t), the node positions and the data, and linear in the data. The derivatives at $(x_i, 0)$ of the implicit solution (s, n) are obtained by differentiating $F = 0$; they are polynomials in derivatives of F divided by powers of the Jacobian determinant,

hence rational in the node positions and linear in the data. By Theorem 4.4 the error is a finite linear combination of such derivatives (those of $\nu = n(\cdot, t(\cdot))$ and $\tau = s(\cdot, t(\cdot))(\cdot)$, with t a polynomial). Its coefficients are therefore rational functions on $\mathcal{A}$, the complement of an algebraic set. Either all of them vanish identically, or one of them, c, does not, and then $\mathcal{Z} = \{c = 0\}$ is a proper algebraic subset outside which the error functional is non-zero. □

For two nodes the second alternative can be verified directly, for data of degree less than 2α.

Proposition 4.18 (Two nodes)**.** *Let $K = 2$, $\rho_1 = \rho_2 \equiv 1$, let Q_i be D-invariant, let the data at the other node x_j be unisolvent on Π_m by themselves, and let $q \in Q_i$ with $\Delta^{\alpha/2} q \neq 0$ and $\deg q < 2\alpha$. Then, for every position of the two nodes, the error $\big[q, u\big]_{x_i} - y_q$ is a non-zero linear functional of the data.*

Proof. Translate x_i to 0 and write $a = x_j$, $s(z) = |z|^\alpha$, $c(z) = |z - a|^{-\alpha}$; fix an inner product on Π_m, and let $N = N_i$, $R = N^\perp$. Take the data $y_i = 0$ and $y_j = \Lambda_j h$ for some $h \in R$. For $z \neq 0$ near 0, p_z minimises $|\Lambda_i p|^2 + s(z)c(z)\,|\Lambda_j(p - h)|^2$. Let $A = (\Lambda_i^* \Lambda_i)|_R$, which is positive definite on R, and $B = \Lambda_j^* \Lambda_j$, positive definite on Π_m by hypothesis, with blocks $B_{RR}, B_{RN}, B_{NR}, B_{NN}$; put $G = B_{NN}^{-1} B_{NR}$ and let $C = B_{RR} - B_{RN} G$ be the Schur complement, also positive definite. Eliminating the N-component from the normal equations gives

$$p_z = Gh + P\rho(z), \qquad P\rho := \rho - G\rho, \qquad \rho(z) = s(z)c(z)\big(A + s(z)c(z)C\big)^{-1} Ch.$$

Hence $u(z) = (Gh)(z) + s(z)c(z)\, g_0(z) + s(z)^2 \psi(z)$, with the polynomial $g_0 = PA^{-1}Ch$ and a smooth function ψ. Now $q(D)(Gh)(0) = 0$, since $Gh \in N$ and $q \in Q_i$; and by Lemma 3.1, as in the proof of Theorem 4.4, $q(D)[s^2\psi](0) = (\Delta^\alpha q)(D)\psi(0) = 0$ because $\deg q < 2\alpha$. With $r = \Delta^{\alpha/2} q$, therefore,

$$\big[q, u\big]_0 - y_q = \varphi(g_0), \qquad \varphi(g) := r(D)(c\, g)(0).$$

The functional φ vanishes on N: by the Leibniz rule,

$$\varphi(g) = \sum_\gamma \frac{1}{\gamma!} (D^\gamma c)(0)\, (\partial^\gamma r)(D) g(0),$$

and every $\partial^\gamma r$ lies in Q_i because Q_i is D-invariant. It does not vanish on Π_m: if g is the Taylor polynomial of degree m of r/c at 0 (note $c(0) \neq 0$), then $cg = r + O(|z|^{m+1})$ and $\varphi(g) = \big[r, r\big] > 0$. Finally, as h ranges over R, g_0 ranges over $W = P(R)$, and $\Pi_m = W \oplus N$. So φ does not vanish on W, and some h gives a non-zero error. □

Conjecture 4.19. *Let $\rho_j \equiv 1$ for all j, let Q_i be D-invariant, let the data at the nodes other than x_i be unisolvent on Π_m by themselves, and let $q \in Q_i$ with $\Delta^{\alpha/2} q \neq 0$. Then the second alternative of Proposition 4.17 holds: q is not reproduced, for data outside a hyperplane, for node positions outside a proper algebraic subset.*

Remark 4.20. By Proposition 4.17 it suffices to find, for each data pattern, one node configuration where the error is non-zero. Proposition 4.9 proves the conjecture for $\deg q = \alpha$, and Proposition 4.18 for two nodes and $\deg q < 2\alpha$, at every node position. The experiments of Section 7 are consistent with it, for instance with two nodes carrying Hermite data of order 3 at $\alpha = 2$ (Section 7.2). The open cases are three or more nodes, where the weights of the other nodes are no longer proportional and the frozen minimiser moves, and data of degree at least 2α. The conjecture concerns condition (ii) only: Q_i is D-invariant, so condition (i) holds. For data that violate (i), the configurations of Corollary 4.14 and the symmetric ones show that both alternatives of the dichotomy occur; characterising them beyond (12) is open.

5. Derivative data without function values

5.1. **Counterexamples.** Komargodski and Levin [18, §3] extend their Hermite MLS, with weights $|x - z|^{-\alpha}$, $\alpha \in 2\mathbb{N}$, and local spaces containing the constants and the linear functions, to directional derivatives prescribed at points that need not carry function values, and state that the derivative data are interpolated, the proof being "straightforward ... by slight modifications" of the co-located case. Condition (i) fails for such data: the symbol of a first derivative differentiates to the constant 1, which is not in Q_i when no value is prescribed at the node. By Theorem 4.13 this does not always cause an error, but it can, and does in the following examples, computed in exact rational arithmetic (scripts in `code/proofs`). In the second example the gradient is given as two directional derivatives at the same point, as [18, §3] allow for gradient data.

Proposition 5.1. *Let $d = 1$, $m = 1$ (local polynomials $a + by$), $\alpha = 2$, $\rho_i \equiv 1$, and prescribe $f(0) = f(2) = 0$ and $f'(1) = f'(3) = 1$. Then*

$$u(x) = \frac{x\,(x-1)(x-2)\,(x^2 - 4x + 5)}{2x^4 - 14x^3 + 37x^2 - 42x + 19},$$

which satisfies $u(0) = u(2) = 0$ but $u'(1) = -1$ and $u'(3) = 31/25$.

Proof. With $w_z(x) = (x - z)^{-2}$ the normal equations of (5) for (a, b) are

$$\begin{pmatrix} w_0 + w_2 & 2w_2 \\ 2w_2 & 4w_2 + w_1 + w_3 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ w_1 + w_3 \end{pmatrix},$$

whose solution gives $u = a + bx$ as stated; the denominator has no real zeros. The remaining values follow by direct computation. □

Proposition 5.2. *Let $d = 2$, $m = 1$, $\alpha = 2$, $\rho_i \equiv 1$, and prescribe $f = 0$ at $(0, 0)$, $(2, 0)$, $(0, 2)$ and $\nabla f(1, 1) = (1, 1)$. In coordinates $z = (z_1, z_2)$, with $a = z_1 + z_2$, $b = z_1^2 + z_2^2$ and $c = z_1 z_2$,*

$$u(z) = \frac{15ab^2 - 68b^2 - 64bc + 128ab + 16ac - 128b - 96c + 48a}{35b^2 - 120ab + 240b + 208c - 240a + 144},$$

which vanishes at the three value points, while $\nabla u(1,1) = (5/9, 5/9)$.

Proof. Exact computation as in Proposition 5.1, with three unknowns (script in `code/proofs`). The denominator tends to $+\infty$ as $|z| \to \infty$, and its minimum, attained on the diagonal near $(0.898, 0.898)$, is $11.21\ldots > 0$. □

The failure persists for $\alpha = 4$ and for $m = 2$ (Section 7.3).

5.2. **Completion.** Given data at x_i with symbol space Q_i, let $\overline{Q}_i$ be the smallest D-invariant space containing Q_i; it has the same maximal degree as Q_i, since differentiation lowers degrees. Choose a basis $\tilde{q}_{i,1}, \ldots, \tilde{q}_{i,r}$ of a complement of Q_i in $\overline{Q}_i$.

Proposition 5.3 (Completion)**.** *Augment the data at* x_i *by the functionals* $\tilde{q}_{i,l}(D)f(x_i)$ *with arbitrary values* $\tilde{y}_{i,l}$*. If Assumption 3.6 holds for the original data, it holds for the augmented data, and every original datum* $q \in Q_i$ *with* $\Delta^{\alpha/2}q = 0$ *(in particular every datum of order less than* α*) is reproduced by the augmented interpolant, whatever the values* $\tilde{y}_{i,l}$*.*

Proof. The augmented symbols at x_i form a basis of $\overline{Q}_i$, so (A2) holds; (A3) is preserved by adding data. The prescribed value y_q of an original datum is unchanged, because its representation in the augmented basis is unique. Every $q \in Q_i \subset \overline{Q}_i$ satisfies (i) with respect to the D-invariant space $\overline{Q}_i$, and Corollary 4.5 applies. □

The same defect, a loss of exactness when the function value is missing at a node, was observed for Shepard-type Hermite–Birkhoff operators [12], where it is avoided by completeness conditions on the data [11]. Condition (i) and the completion $\overline{Q}_i$ are the corresponding notions for SWIMLS: D-invariance asks for all derivatives of the given symbols, not only for all derivatives of a given order.

The values $\tilde{y}_{i,l}$ influence the accuracy, not the exactness. In the experiments they are estimated by a Gaussian-weighted (non-singular) moving least-squares fit of the given data.

6. Convergence

Sections 4 and 5 concern exactness at the nodes. We now bound the error $u - f$ everywhere, for data sampled from a smooth function f. The natural setting is that of compactly supported weights: they make the problem local, and then a scaling and compactness argument gives the same order of approximation as for classical moving least squares.

6.1. **Compactly supported singular weights.** Let $\psi \in C^\infty([0,\infty))$ satisfy $\psi > 0$ on $[0,1)$ and $\psi = 0$ on $[1,\infty)$, for instance $\psi(s) = \exp\bigl(1 - 1/(1-s)\bigr)$ for $s < 1$. For a support radius $\delta > 0$ we use the weights

$$w_j(x) = |x - x_j|^{-\alpha}\, \psi\bigl(|x - x_j|^2/\delta^2\bigr), \qquad \alpha \in 2\mathbb{N}. \tag{13}$$

Here $\rho_j(x) = \psi(|x-x_j|^2/\delta^2)$ is C^∞ but vanishes outside the ball $B(x_j,\delta)$, so we replace (A1) and (A3) by

(A1′) $\alpha \in 2\mathbb{N}$, $w_j(x) = \rho_j(x)|x-x_j|^{-\alpha}$ with $\rho_j \in C^\infty(\Omega)$, $\rho_j \geq 0$ and $\rho_j(x_j) > 0$;
(A3′) for every $x \in \Omega$, the data at the nodes x_j with $\rho_j(x) > 0$ are unisolvent on Π_m.

Under these assumptions J_x is still strictly convex for every $x \notin X$, and the proofs of Section 4 apply verbatim once U is chosen so small that $\rho_i > 0$ on U and every ρ_j that is positive at x_i is positive on U: the only place where positivity of the other weights is used, the strict convexity of the frozen problem, then follows from (A3′) at $x = x_i$. In Proposition 4.9, "the other nodes" are those with $\rho_j(x_i) > 0$, and in Corollary 4.14(a) the condition is required only for the j with $w_j \not\equiv 0$ near x_i. With these readings, the general results of Sections 4 and 5 (Theorems 4.1, 4.4 and 4.13, their corollaries, and Propositions 4.9, 4.16 and 5.3) hold for the weights (13); Propositions 5.1 and 5.2 and the statements that use analyticity concern pure power weights.

The data are weighted according to their order. If $q_{j,s}$ is homogeneous of degree $r_{j,s}$, we minimise

$$J_x(p) = \sum_j w_j(x) \sum_{s=1}^{n_j} \delta^{2r_{j,s}} \big(\lambda_{j,s}(p) - y_{j,s}\big)^2. \tag{14}$$

The constant factors $\delta^{2r_{j,s}}$ do not affect any result of Sections 4 and 5 (see the remark after Definition 3.7); they make every term of (14) scale in the same way, which is what the following proof needs.

6.2. **An error bound.** For a ball B and $f \in C^{m+1}(B)$ let

$$|f|_{m+1,B} := \max_{|\gamma|=m+1} \sup_B |D^\gamma f|.$$

We assume:

(H1) *separation:* $|x_i - x_j| \geq \vartheta\delta$ for $i \neq j$;
(H2) *data:* the symbols are homogeneous of degree at most m with coefficient vectors of Euclidean norm one, and the tuple $(q_{j,1}, \dots, q_{j,n_j})$ at each node belongs to a fixed compact set $\mathcal{Q}$ of linearly independent tuples (for example, a finite list of data patterns such as Hermite data of order $k \leq m$);
(H3) *local norming:* there are $c_0 \in (0,1)$ and C_N such that for every $x \in \Omega$ and every $p \in \Pi_m$

$$\max_{|z|\leq 1} |p(x+\delta z)| \leq C_N \max\big\{\delta^{r_{j,s}} |\lambda_{j,s}(p)| : \ |x_j - x| \leq c_0\delta\big\}.$$

Condition (H3) is a quantitative form of (A3′): the data near x determine a polynomial, with a constant that does not depend on x or δ.

Theorem 6.1 (Convergence)**.** *Let u be the SWIMLS interpolant with the weights (13) and the functional (14), for data $y_{j,s} = \lambda_{j,s}(f)$ of a function f, and assume (H1)–(H3)*

(which imply (A1′) and (A3′)). There is a constant C, depending only on d, m, α, ψ, ϑ, c_0, C_N and $\mathcal{Q}$, such that

$$|D^\beta(u-f)(x)| \le C\,\delta^{m+1-|\beta|}\,|f|_{m+1,B(x,\delta)} \tag{15}$$

for every $x\in\Omega$, the nodes included, every $|\beta|\le m$ and every f whose restriction to $B(x,\delta)$ is in C^{m+1}. In particular, if a family of quasi-uniform node sets is used with $\delta = ch$, h the fill distance, and (H3) holds with c_0 and C_N independent of h, then $\|D^\beta(u-f)\|_{L^\infty(\Omega)} \le C\,h^{m+1-|\beta|}|f|_{m+1,\Omega_\delta}$ with $\Omega_\delta = \Omega + B(0,\delta)$.

The constant comes from a compactness argument and is not explicit. For function values, local polynomials and interpolating weights that behave like $|x-x_j|^{-2}$ near the nodes, the estimate with $\beta=0$ is Levin's approximation-order theorem [21, Theorem 5], and our proof follows his: polynomial reproduction, continuity of the coefficients as functions of the local configuration, and invariance under scaling, for compactly supported C^∞ weights and separated nodes. Theorem 6.1 extends it to every even α, to Hermite–Birkhoff data (where the scaling of the derivative data in (14) is essential), and to the derivatives of the interpolant up to order m; for non-interpolating moving least squares with values, bounds of this type are in [40, 2, 39, 42, 26].

Proof. Step 1: shape functions and reproduction. The minimiser of (14) is linear in the data, so $u(x)=\sum_{j,s} a_{j,s}(x)\,y_{j,s}$ with shape functions $a_{j,s}$, which are C^∞ on Ω by Theorem 4.1. Each $a_{j,s}$ vanishes, with all its derivatives, where $|x_j-x|\ge\delta$: there w_j vanishes identically in a neighbourhood or, on the sphere $|x_j-x|=\delta$, to infinite order, since ψ is flat at 1. If the data come from a polynomial $p\in\Pi_m$, then $J_x(p)=0$, so $p_x=p$ and $u=p$. Hence $\sum_{j,s} a_{j,s}(x)\lambda_{j,s}(p)=p(x)$ for all x, and differentiating in x,

$$\sum_{j,s} D^\beta a_{j,s}(x)\,\lambda_{j,s}(p) = D^\beta p(x), \qquad p\in\Pi_m. \tag{16}$$

Step 2: error identity. Fix $x\in\Omega$ and let T be the Taylor polynomial of f of degree m at x. By (16) with $p=T$ and since $D^\beta T(x)=D^\beta f(x)$ for $|\beta|\le m$,

$$D^\beta(u-f)(x) = \sum_{j,s} D^\beta a_{j,s}(x)\,\lambda_{j,s}(f-T).$$

If $|x_j-x|<\delta$ and $q_{j,s}=\sum_{|\gamma|=r} c_\gamma\xi^\gamma$ with $r=r_{j,s}$, then $D^\gamma T$ is the Taylor polynomial of degree $m-r$ of $D^\gamma f$ at x, and Taylor's theorem gives $|D^\gamma(f-T)(x_j)|\le C_1\delta^{m+1-r}|f|_{m+1,B(x,\delta)}$ with C_1 depending only on d and m. Since $|c_\gamma|\le 1$ and the number of multi-indices of order $r\le m$ depends only on d and m,

$$|D^\beta(u-f)(x)| \le C_2\,\delta^{m+1-|\beta|}\,|f|_{m+1,B(x,\delta)}\,\Lambda_\beta(x), \qquad \Lambda_\beta(x) := \sum_{j,s}\delta^{|\beta|-r_{j,s}}\,|D^\beta a_{j,s}(x)|. \tag{17}$$

It remains to bound Λ_β independently of x, δ and the nodes.

Step 3: scaling. Let $z_j=(x_j-x)/\delta$ for the nodes with $|z_j|\le 1$, and for $p\in\Pi_m$ let $\tilde p(z)=p(x+\delta z)$, a bijection of Π_m. Because $q_{j,s}$ is homogeneous, $\lambda_{j,s}(p)=\delta^{-r_{j,s}}\tilde\lambda_{j,s}(\tilde p)$

with $\tilde{\lambda}_{j,s}(\tilde{p}) = (q_{j,s}(D)\tilde{p})(z_j)$, and $w_j(x) = \delta^{-\alpha}\tilde{w}(z_j)$ with $\tilde{w}(z) = |z|^{-\alpha}\psi(|z|^2)$. Hence

$$J_x(p) = \delta^{-\alpha}\sum_j \tilde{w}(z_j)\sum_s \big(\tilde{\lambda}_{j,s}(\tilde{p}) - \delta^{r_{j,s}}y_{j,s}\big)^2.$$

So $\tilde{p}_x$ is the SWIMLS fit, evaluated at the origin, of the *configuration* $\mathcal{C} = (z_j, Q_j)_j$ with data $\delta^{r_{j,s}}y_{j,s}$, where Q_j is the tuple of symbols at x_j, and $u(x) = \tilde{p}_x(0)$. Denote by $\tilde{a}_{j,s}(\mathcal{C})$ the shape functions of this scaled problem at the origin; for configurations with a point at the origin ($x \in X$) they are defined by continuous extension, which exists by Step 5(b). Then $a_{j,s}(x) = \delta^{r_{j,s}}\tilde{a}_{j,s}(\mathcal{C}(x))$, and since moving x translates all z_j at once (near a given x we keep the points of $\mathcal{C}(x)$: points outside the closed unit ball stay outside, and points leaving it have zero weight),

$$\delta^{|\beta|-r_{j,s}}|D^\beta a_{j,s}(x)| = \big|(E^\beta\tilde{a}_{j,s})(\mathcal{C}(x))\big|, \qquad E := \sum_l \nabla_{z_l}.$$

Thus $\Lambda_\beta(x) = \tilde{\Lambda}_\beta(\mathcal{C}(x))$ with $\tilde{\Lambda}_\beta(\mathcal{C}) := \sum_{j,s}|E^\beta\tilde{a}_{j,s}(\mathcal{C})|$, a function of the configuration alone, in which δ no longer appears.

Step 4: the admissible configurations form a compact set. By (H1), a configuration has $n \le M(d,\vartheta)$ points in the closed unit ball, at mutual distances at least ϑ. By (H2) its symbol tuples lie in $\mathcal{Q}$. By (H3), with $\tilde{p}$ in place of p,

$$\max_{|z|\le 1}|\tilde{p}(z)| \le C_N \max\{|\tilde{\lambda}_{j,s}(\tilde{p})| : \ |z_j| \le c_0\} \qquad (\tilde{p} \in \Pi_m). \tag{18}$$

Let $\mathcal{K}$ be the set of all configurations with these three properties. Each property is preserved under limits. For (18), pass to a subsequence with a fixed number of points, a fixed labelling and a fixed index set $S = \{j : |z_j| \le c_0\}$; the limit points with $j \in S$ again satisfy $|z_j| \le c_0$, and $\tilde{\lambda}_{j,s}(\tilde{p})$ depends continuously on z_j and Q_j, so the inequality passes to the limit. Hence $\mathcal{K}$ is compact, as a finite union, over $n \le M$, of closed and bounded sets.

Step 5: continuity. We show that every $\mathcal{C}_0 \in \mathcal{K}$ has a neighbourhood, in the space of all (z_j, Q_j) with independent tuples Q_j, on which all $\tilde{a}_{j,s}$ are C^∞ functions of $(z_j, Q_j)_j$. Then $\tilde{\Lambda}_\beta$ is continuous on $\mathcal{K}$, hence bounded by a constant C_β, and (15) follows from (17) with $C = C_2\max_{|\beta|\le m} C_\beta$. Note that $\tilde{w}$ is C^∞ on $\mathbb{R}^d \setminus \{0\}$, including the sphere $|z| = 1$, because ψ is flat at 1; points with $|z_j| \ge 1$ have zero weight and can be added or removed freely.

(a) No point at the origin. This is the non-singular case of classical moving least squares [39, 26]. Let $\ell_{j,s}(\mathcal{C}) \in \mathbb{R}^{\dim\Pi_m}$ represent $\tilde{\lambda}_{j,s}$ in a fixed basis of Π_m; it is polynomial in z_j and linear in Q_j. The Gram matrix $G(\mathcal{C}) = \sum_j \tilde{w}(z_j)\sum_s \ell_{j,s}\ell_{j,s}^{\mathsf{T}}$ is C^∞ near $\mathcal{C}_0$, and it is positive definite at $\mathcal{C}_0$: a polynomial annihilated by all data with $\tilde{w}(z_j) > 0$ is annihilated by those with $|z_j| \le c_0$, hence vanishes by (18). So G^{-1}, and with it the shape functions $\tilde{a}_{j,s} = e^{\mathsf{T}}G^{-1}\tilde{w}(z_j)\ell_{j,s}$ (with e the evaluation at 0), are C^∞ near $\mathcal{C}_0$.

(b) A point at the origin. By (H1), at most one point, say z_i, is near the origin. We repeat the proof of Theorem 4.1 with the configuration as the parameter in place of x.

Write $\tilde{p}(z) = P(z - z_i)$, so that the data at z_i become $q_{i,s}(D)P(0)$ and do not depend on z_i, while all other data and the evaluation $\tilde{p}(0) = P(-z_i)$ depend polynomially on the configuration. Dividing the scaled functional by $\tilde{w}(z_i)$ gives $|\Lambda_i P - \tilde{y}_i|^2 + t\, B_{\mathcal{C}}(P)$ with $t = |z_i|^\alpha$, a polynomial in z_i, and $B_{\mathcal{C}}$ smooth in $\mathcal{C}$ near $\mathcal{C}_0$. The kernel N_i of Λ_i depends on Q_i, but since Λ_i has full rank it can be written, near Q_i^0, as the graph of a linear map from a fixed space that depends rationally on Q_i. The implicit function theorem then applies as in the proof of Theorem 4.1: the first diagonal block of the Jacobian is invertible because Q_i is independent, and the second, the Hessian of the frozen problem on N_i, is positive definite by (18), which involves the point $z_i = 0$. It yields a C^∞ function of $(\mathcal{C}, t)$. This function is the minimiser whenever $t > 0$: the map F is affine in the unknowns, its linear part is invertible near $(\mathcal{C}_0, 0)$, and for $t > 0$ the substitution $r = ts$ maps its zeros bijectively onto the critical points of the scaled functional; a positive semidefinite quadratic with exactly one critical point has a nonsingular, hence positive definite, Hessian, and that point is its unique minimiser; composed with $t = |z_i|^\alpha$ it gives the shape functions near $\mathcal{C}_0$, as C^∞ functions of $\mathcal{C}$. □

Remark 6.2.

(a) *Compact support.* The proof uses that only the finitely many nodes in $B(x, \delta)$ influence $u(x)$, so that the scaled configurations form a compact set. For weights with a Gaussian factor every node contributes, and an extension would need estimates of the tails; Levin already expected such an extension for values [21, p. 1524], and the experiments of Section 7 show the same rates for such weights.

(b) *Scaling of derivative data.* The factors $\delta^{2r_{j,s}}$ in (14) make the scaled problem independent of δ. Without them the relative weight of derivative data grows like δ^{-2r} as $\delta \to 0$, and the argument does not apply.

(c) *The norming condition.* If the data contain the function values, (H3) holds as soon as the value data alone satisfy it, since adding data can only increase the right-hand side. For values, local norming conditions of this type are the basis of the approximation theory of moving least squares on quasi-uniform nodes with δ a sufficiently large multiple of the fill distance [39, Chapters 3 and 4].

(d) *Gaps and completion.* The theorem applies to any data satisfying (H1)–(H3), with or without gaps, and so to completed data (Proposition 5.3) as well; the added symbols can be taken homogeneous, since derivatives of homogeneous symbols are homogeneous. If the added values are only estimates, with errors η, then u changes by $\sum a_{j,s}\eta_{j,s}$, which by Step 3 is at most $\tilde{\Lambda}_0 \max_{j,s} \delta^{r_{j,s}} |\eta_{j,s}|$ in absolute value, where $\tilde{\Lambda}_0 \le C_\beta$ with $\beta = 0$ is the bound of Step 5.

7. Numerical experiments

The experiments test the theory of Sections 4 and 5 datum by datum, and the convergence theorem of Section 6. They use a numerically stable evaluator of SWIMLS, described in Appendix A, which stays accurate to rounding level as close to the nodes as

we tested (10^{-14}). Whether a datum is reproduced is decided in high-precision arithmetic (Appendix B); the outcome is never borderline: reproduced data have errors below 10^{-18}, the others above 10^{-2}. Unless stated otherwise, nodes are random, the weights include a Gaussian factor (Appendix A), the data values are random numbers (exactness does not depend on them being samples of a function), and every node other than the one under study carries Hermite data. Scripts, logs and data are in the accompanying repository, `https://github.com/davidfgarcia/hermite-birkhoff` (release v0.1.0).

7.1. **Hermite and Hermite–Birkhoff data.** We first checked every datum at a node with classical Hermite data, of orders $k \in \{2, 3, 4, 5, 7\}$ in one, two and three dimensions and with α from 2 to 6, and at a node carrying the harmonic symbols of degree at most 3: 92 data in all. A datum is reproduced exactly when $\Delta^{\alpha/2} q = 0$ and not otherwise, as Corollary 4.6 predicts for the reproduced data and Proposition 4.9 and Conjecture 4.19 for the others. For instance, at $\alpha = 2$ in three dimensions, $\partial_x \partial_y \partial_z f$ is reproduced while $\partial_x^2 f$ has an error of 30. The error identities of Example 4.7 hold to all twelve digits computed: at a node with Hermite data of order 3 in two dimensions and $\alpha = 2$, $e(\partial_x^2) = e(\partial_y^2) = -45.904682196$ and $e(\partial_x^3) = 3e(\partial_x \partial_y^2) = 599.09999304$.

Next, one node carries a pattern with gaps, such as $\{f, \partial_{xx} f\}$, $\{f, \partial_x f, \partial_{xy} f\}$, $\{f, \Delta f\}$ or $\{\partial_x f\}$ alone: 23 patterns in one and two dimensions, most of them with gaps, plus the two settings of Section 7.3, 83 data in all. Again conditions (i) and (ii) predict every outcome. For example, $\partial_{xy} f$ is not reproduced in the pattern $\{f, \partial_x f, \partial_{xy} f\}$, because its derivative ξ_2 is missing from the node, and Δf is not reproduced in $\{f, \Delta f\}$.

7.2. **When condition (i) is not needed.** The configurations above are generic. Theorem 4.13 predicts that special ones behave differently, and they do. In each configuration of the following table condition (i) fails and yet the datum is reproduced; a small perturbation that destroys the special structure makes the error reappear.

configuration	datum	reason	error
1D: $f'(0)$, $f(1)$, $f'(5/2)$; $m = 1$, $\alpha = 2$	f' at 0, 5/2	Cor. 4.14(b)	$< 10^{-44}$
add $f(4)$			0.21–2.1
2D, $K = 2$: $\nabla f(x_1)$; $f, \nabla f, \nabla^2 f$ at x_2; $m = 2$, $\alpha = 2$	∂_x, ∂_y	Cor. 4.14(a)	$< 10^{-39}$
add a third node			0.03–0.46
2D, $K = 2$: $f, \partial_{xx} f$ at x_1; Hermite $k = 2$ at x_2; $\alpha = 4$	∂_{xx}	Cor. 4.14(a)	$\leq 1.1 \cdot 10^{-28}$
add a third node			2.9–3.3
2D: $\partial_x f(0)$; others on the z_2-axis; $m = 1$, $\alpha = 2$	∂_x	Prop. 4.16	$< 10^{-37}$
move the others by 0.05			0.08–0.16

In the symmetric configuration (last two rows) the weights of the other nodes are even in the first coordinate, so $\partial_{x_1} Z(x_i) = 0$ and the first-order error (12) vanishes for $q = \xi_1$. The configurations of this subsection use pure power weights. Conversely, with two nodes carrying Hermite data of order 3 and $\alpha = 2$ (pure power weights), where the

frozen minimiser is constant and condition (i) holds, every datum with $\Delta q \neq 0$ fails, with errors between 4.9 and 83, as Theorem 4.13 allows and Conjecture 4.19 predicts.

7.3. **Derivatives at points without values.** In the setting of [18, §3], eight random points carry values, eight other points carry gradients, the weights are pure powers $|x - x_i|^{-\alpha}$ and the local space is Π_1 or Π_2. For $m \in \{1, 2\}$ and $\alpha \in \{2, 4\}$, none of the 24 gradient components checked is reproduced; the errors lie between 0.06 and 4.5. After completion (Proposition 5.3) they are, and completion also repairs all 13 data of the patterns of Section 7.1 that fail because of condition (i); the two that fail because of condition (ii) cannot be repaired by completion. Exactness is not free in accuracy, though: with values at 1024 Halton points and gradients at 1024 others for Franke's function, the maximum error is $1.2 \cdot 10^{-3}$ without and $3.4 \cdot 10^{-3}$ with completion, because the added values are only estimates (here from a Gaussian-weighted moving least-squares fit of the given data).

7.4. **Accuracy.** Theorem 6.1 predicts errors of order h^{m+1} for u and h^m for its gradient. We tested this on $N = 64, \dots, 8192$ Halton points in the unit square, with the compactly supported weights (13), $\delta = ch$ for $c \in \{2, 3, 4\}$, $h = N^{-1/2}$, the scaled functional (14), and data of order $k \leq 2$ from a smooth test function; errors are measured on a 33×33 grid in $[0.1, 0.9]^2$, and gradients by fourth-order finite differences with step $h/100$. In all cases where the support contains enough nodes, least-squares fits of the maximum error over all resolutions give rates 2.8–3.2 for $m = 2$, 3.6–3.9 for $m = 3$, 5.0–5.1 for $m = 4$ and 5.7–5.9 for $m = 5$ with Hessian data, and gradient rates 1.9–2.1, 2.5–3.0, 3.9–4.1 and 4.7–4.9 (Figure 2). For values with $m = 5$ the rates are higher (6.8–7.9 for u, 6.4 for the gradient), a pre-asymptotic effect. The Halton points are not uniformly separated (their separation, relative to h, falls from 0.56 to 0.14 over this range), so they do not satisfy (H1) with a fixed constant. The maximum error samples the worst local configuration and fluctuates from one node set to the next. On periodically jittered grids, which satisfy (H1) with a fixed constant and where the local configurations repeat at every resolution, fits over all resolutions give root-mean-square rates within 0.06 of $m + 1$ for u and of m for the gradient, for the five data types and degrees tested. When the support is too small for the local norming condition (H3), as for values with $m = 3$ or 5 and $c = 2$, the error does not converge at the predicted rate (fitted rates 1.2 and 0.4), which illustrates that the condition cannot simply be dropped.

A second study with Gaussian factors instead of compact support gave the same rates for the root-mean-square error (within 0.53 of $m+1$, over 44 parameter combinations). It also shows what derivative data buy: on a fixed node set they reduce the error, e.g. for $N = 4096$ and $m = 5$ from $5.0 \cdot 10^{-9}$ (values) to $1.1 \cdot 10^{-9}$ (values, gradients and Hessians), but per number of data values alone were more accurate in that test. Derivative data pay off when nodes are expensive and derivatives are cheap, as with adjoint gradients.




FIGURE 2. Maximum errors for Halton nodes with compactly supported weights, $\delta = 2h$ ($3h$ for values); $\alpha = 2$, and $\alpha = 4$ with Hessian data. Dotted lines have the slopes $m + 1$ in (a) and m in (b) predicted by Theorem 6.1.

8. CONCLUSIONS

Singular-weight moving least squares extends from function values to arbitrary linear differential data with no change to its definition: one weighted least-squares problem per evaluation point, with no mesh, no global system and no poisedness condition beyond unisolvency of all the data on the local polynomial space. The interpolant is C^∞, and Theorem 4.4 accounts exactly for the error in every datum. Two mechanisms can spoil a datum: the singularity of the weight, which acts through $\Delta^{\alpha/2}$, and the motion of the frozen minimiser, which acts only on data whose derivatives are missing at their node. The first is removed by choosing α larger than the order of the data, the second by completing the data at the node. Together they give a simple recipe for Hermite–Birkhoff interpolation of any order in any dimension.

Several questions remain open. Conjecture 4.19 asks whether $\Delta^{\alpha/2} q \neq 0$ always causes an error in generic configurations; we proved it for data of degree α and, with two nodes, for data of degree less than 2α. Beyond the first-order formula (12), we do not know exactly which configurations of the other nodes make condition (i) unnecessary. Theorem 6.1 assumes compactly supported weights and a local norming condition; extending it to weights with unbounded support, and verifying the norming condition for Hermite–Birkhoff data without function values, are natural next steps. Finally, the values added by completion affect the accuracy, not the exactness, and better ways of estimating them would reduce the accuracy penalty observed in Section 7.

Data given by a differential equation, such as $\Delta u(x_i)$, are Hermite–Birkhoff data like any other; the application of SWIMLS to boundary value problems will be presented separately.

Appendix A. The evaluator

The usual implementation of interpolating MLS replaces $|x-x_i|^{-\alpha}$ by $(|x-x_i|^{\alpha}+\varepsilon)^{-1}$ and solves the normal equations. This gives up exactness at the nodes and, near a node, squares the condition number of the weighted system, which grows like $|x-x_i|^{-\alpha/2}$. Our evaluator avoids both problems; it is in the accompanying code.

A.1. **Basis and weights.** At the evaluation point x we use the monomials $((y-x)/h)^{\beta}$, $|\beta| \le m$, with a length scale h, so that $u(x)$ is the first coefficient. The weights are either pure powers $|x-x_i|^{-\alpha}$ or $w_i(x) = |x-x_i|^{-\alpha} e^{-|x-x_i|^2/\delta^2}$, which also satisfy (A1); they are computed as logarithms and normalised by their maximum, and nodes whose normalised weight is below a cutoff are dropped. The cutoff is measured relative to the strongest node *other than* the nearest one; measured relative to the nearest node, it would drop all other nodes close to a node and destroy the information that fixes ν. Rows of derivative data of order r are multiplied by ℓ^r for a length ℓ; by the remark after Definition 3.7 this does not affect the results of Section 4.

A.2. **Least squares.** The weighted system $W^{1/2}A\,a \approx W^{1/2}b$ is solved by an SVD-based least-squares solver rather than through the normal equations. Close to a node x_i the rows of x_i dominate the others by a factor θ^{-1} with $\theta \sim |x-x_i|^{\alpha/2}$. The light rows carry the information that fixes $\nu(x)$, and a standard solver resolves them only to relative accuracy of order $\varepsilon_{\mathrm{mach}}/\theta$ (and not at all through the normal equations once $\theta^2 < \varepsilon_{\mathrm{mach}}$). When $\theta < 10^{-3}$ we therefore solve an exact reformulation. Let A_{h} and A_{l} be the heavy and light weighted rows, with right-hand sides b_{h} and b_{l}, let U_1 and U_2 be orthonormal bases of the row space and the null space of A_{h}, and let $\hat{c}$ be the least-squares solution of $A_{\mathrm{h}}U_1 c = b_{\mathrm{h}}$. Writing $a = U_1(\hat{c} + \theta\, e) + U_2 v$ and dividing the objective by θ^2 gives the problem

$$\min_{e,v} \ \|A_{\mathrm{h}}U_1 e\|^2 + \left\|A_{\mathrm{l}}U_1 e + \theta^{-1}A_{\mathrm{l}}U_2 v - \theta^{-1}(b_{\mathrm{l}} - A_{\mathrm{l}}U_1\hat{c})\right\|^2,$$

whose blocks are well scaled; it has the same minimiser and is the discrete counterpart of the decomposition (6).

A.3. **At the nodes.** At $x = x_i$, and within a tolerance of 10^{-13} times the size of the node set, we use the limit of Theorem 4.1: the data at x_i are imposed as equality constraints, and the remaining freedom in N_i is fixed by minimising the frozen functional (7); then $u(x_i) = (p_i^* + Z(x_i))(x_i)$.




FIGURE 3. Error of the evaluation near the node $x_i = 2$ (nodes $1, 2, 3, 4$ with Hermite data of order k and random values), against a 120-digit reference. Crosses mark singular normal equations, open circles errors that are exactly zero.

A.4. **Accuracy.** Figure 3 compares the evaluator with a reference computed with 120 significant digits, as the evaluation point approaches a node. Its error stays at the level of rounding (below $5 \cdot 10^{-15}$) down to distance 10^{-14}; the three smallest distances are within the node tolerance and use the formula at the node. With $\varepsilon = 10^{-10}$ the regularised normal equations have errors up to $1.6 \cdot 10^{-10}$, the size of the regularisation; with $\varepsilon = 10^{-16}$ the errors reach $4 \cdot 10^{-12}$ and the matrix is numerically singular at some distances.

APPENDIX B. VERIFYING EXACTNESS

The experiments of Section 7 decide whether a datum q is reproduced by computing $\big[q, u\big]_{x_i}$ in high-precision arithmetic (`mpmath`). The interpolant is evaluated from the normal equations at a working precision of $40 + (2\alpha + |\beta| + 5)\,|\log_{10} r|$ digits, and $D^\beta u(x_i)$ is taken with a tensor-product Chebyshev stencil of radius r in the variables with $\beta_j > 0$, which is exact for polynomials of degree up to (at least) $\beta_j + 5$ in each of them and never samples the node itself. A datum is called reproduced if $\big|\big[q, u\big]_{x_i} - y_q\big| \le 10^{-12} \max(1, |y_q|)$ for both $r = 10^{-6}$ and $r = 10^{-5}$, and not reproduced if the inequality fails for both. Mixed outcomes never occurred, with a wide margin: over the 221 data of Sections 7.1–7.2, the reproduced ones have errors between $6 \cdot 10^{-55}$ and $9 \cdot 10^{-25}$ at $r = 10^{-6}$ (at most $9 \cdot 10^{-19}$ at $r = 10^{-5}$), and their errors scale like r^6 or r^8, as the stencil error does; the others have errors of at least 0.01, identical at both radii.

Declaration of generative AI use

During the preparation of this work the author used Claude (Anthropic) to help draft and revise the text, implement and run the numerical experiments, search and verify the literature, check proofs, and suggest proof strategies. After using this tool, the author reviewed and edited the content as needed and takes full responsibility for the content of the article.

References

1. M. Alexa and A. Adamson, *Interpolatory point set surfaces—convexity and Hermite data*, ACM Trans. Graph. **28** (2009), no. 2, 1–10, doi:10.1145/1516522.1516531.
2. M. G. Armentano, *Error estimates in Sobolev spaces for moving least square approximations*, SIAM J. Numer. Anal. **39** (2001), no. 1, 38–51, doi:10.1137/S0036142999361608.
3. S. N. Atluri, J. Y. Cho, and H.-G. Kim, *Analysis of thin beams, using the meshless local Petrov–Galerkin method, with generalized moving least squares interpolations*, Comput. Mech. **24** (1999), no. 5, 334–347, doi:10.1007/s004660050456.
4. L. Bos and K. Šalkauskas, *Moving least-squares are Backus–Gilbert optimal*, J. Approx. Theory **59** (1989), no. 3, 267–275.
5. G. Allasia and C. Bracco, *Multivariate Hermite–Birkhoff interpolation by a class of cardinal basis functions*, Appl. Math. Comput. **218** (2012), 9248–9260, doi:10.1016/j.amc.2012.03.002.
6. Gh. Coman, *Shepard operators of Birkhoff-type*, Calcolo **35** (1998), no. 4, 197–203, doi:10.1007/s100920050016.
7. A. P. Calderón and A. Zygmund, *On higher gradients of harmonic functions*, Studia Math. **24** (1964), no. 2, 211–226, doi:10.4064/sm-24-2-211-226.
8. C. de Boor, *Ideal interpolation*, Approximation Theory XI: Gatlinburg 2004 (C. K. Chui, M. Neamtu, and L. L. Schumaker, eds.), Nashboro Press, Brentwood, TN, 2005, pp. 59–91.
9. C. de Boor and A. Ron, *The least solution for the polynomial interpolation problem*, Math. Z. **210** (1992), 347–378, doi:10.1007/BF02571803.
10. C. de Boor and A. Ron, *On multivariate polynomial interpolation*, Constr. Approx. **6** (1990), no. 3, 287–302, doi:10.1007/BF01890412.
11. F. Dell'Accio and F. Di Tommaso, *Complete Hermite–Birkhoff interpolation on scattered data by combined Shepard operators*, J. Comput. Appl. Math. **300** (2016), 192–206, doi:10.1016/j.cam.2015.12.016.
12. F. Dell'Accio and F. Di Tommaso, *Scattered data interpolation by Shepard's like methods: classical results and recent advances*, Dolomites Res. Notes Approx. **9** (2016), Special Issue, 32–44, doi:10.14658/pupj-drna-2016-Special_Issue-5.
13. W. J. Gordon and J. A. Wixom, *Shepard's method of "metric interpolation" to bivariate and multivariate interpolation*, Math. Comp. **32** (1978), 253–264, doi:10.1090/S0025-5718-1978-0458027-6.
14. R. Farwig, *Rate of convergence of Shepard's global interpolation formula*, Math. Comp. **46** (1986), 577–590.
15. G. E. Fasshauer, *Meshfree Approximation Methods with MATLAB*, Interdisciplinary Mathematical Sciences, vol. 6, World Scientific, Singapore, 2007, doi:10.1142/6437.
16. E. Fischer, *Über die Differentiationsprozesse der Algebra*, J. Reine Angew. Math. **148** (1918), 1–78, doi:10.1515/crll.1918.148.1.
17. M. Gasca and T. Sauer, *Polynomial interpolation in several variables*, Adv. Comput. Math. **12** (2000), no. 4, 377–410, doi:10.1023/A:1018981505752.

18. Z. Komargodski and D. Levin, *Hermite type moving-least-squares approximations*, Comput. Math. Appl. **51** (2006), no. 8, 1223–1232, doi:10.1016/j.camwa.2006.04.005.
19. D. Kraaijpoel and T. van Leeuwen, *Raising the order of multivariate approximation schemes using supplementary derivative data*, Procedia Comput. Sci. **1** (2010), no. 1, 307–316, doi:10.1016/j.procs.2010.04.034.
20. P. Lancaster and K. Šalkauskas, *Surfaces generated by moving least squares methods*, Math. Comp. **37** (1981), no. 155, 141–158, doi:10.1090/S0025-5718-1981-0616367-1.
21. D. Levin, *The approximation power of moving least-squares*, Math. Comp. **67** (1998), no. 224, 1517–1531, doi:10.1090/S0025-5718-98-00974-0.
22. R. A. Lorentz, *Multivariate Hermite interpolation by algebraic polynomials: a survey*, J. Comput. Appl. Math. **122** (2000), no. 1–2, 167–201.
23. G. G. Lorentz, K. Jetter, and S. D. Riemenschneider, *Birkhoff Interpolation*, Encyclopedia of Mathematics and its Applications, vol. 19, Addison-Wesley, Reading, MA, 1983.
24. D. H. McLain, *Drawing contours from arbitrary data points*, Comput. J. **17** (1974), no. 4, 318–324, doi:10.1093/comjnl/17.4.318.
25. D. H. McLain, *Two dimensional interpolation from random data*, Comput. J. **19** (1976), no. 2, 178–181, doi:10.1093/comjnl/19.2.178.
26. D. Mirzaei, *Analysis of moving least squares approximation revisited*, J. Comput. Appl. Math. **282** (2015), 237–250.
27. D. Mirzaei, R. Schaback, and M. Dehghan, *On generalized moving least squares and diffuse derivatives*, IMA J. Numer. Anal. **32** (2012), no. 3, 983–1000, doi:10.1093/imanum/drr030.
28. T. Most and C. Bucher, *A moving least squares weighting function for the element-free Galerkin method which almost fulfills essential boundary conditions*, Struct. Eng. Mech. **21** (2005), no. 3, 315–332, doi:10.12989/sem.2005.21.3.315.
29. H. Netuzhylov, T. Sonar, and W. Yomsatieankul, *Finite difference operators from moving least squares interpolation*, ESAIM Math. Model. Numer. Anal. **41** (2007), 959–974, doi:10.1051/m2an:2007042.
30. F. J. Narcowich and J. D. Ward, *Generalized Hermite interpolation via matrix-valued conditionally positive definite functions*, Math. Comp. **63** (1994), no. 208, 661–687, doi:10.1090/S0025-5718-1994-1254147-6.
31. T. Sauer and Y. Xu, *On multivariate Hermite interpolation*, Adv. Comput. Math. **4** (1995), no. 1, 207–259, doi:10.1007/BF03177515.
32. H. S. Shapiro, *An algebraic theorem of E. Fischer, and the holomorphic Goursat problem*, Bull. London Math. Soc. **21** (1989), no. 6, 513–537, doi:10.1112/blms/21.6.513.
33. B. Shekhtman, *Ideal interpolation: translations to and from algebraic geometry*, Approximate Commutative Algebra, Texts and Monographs in Symbolic Computation, Springer, Vienna, 2010, doi:10.1007/978-3-211-99314-9_6.
34. D. Shepard, *A two-dimensional interpolation function for irregularly-spaced data*, Proceedings of the 1968 23rd ACM National Conference, ACM, New York, 1968, pp. 517–524, doi:10.1145/800186.810616.
35. T. Sonar, *Difference operators from interpolating moving least squares and their deviation from optimality*, ESAIM Math. Model. Numer. Anal. **39** (2005), 883–908, doi:10.1051/m2an:2005039.
36. D. V. Toan, T. T. T. Mai, T. V. Long, and P. V. Manh, *Multivariate Hermite interpolation with interpolation conditions induced by harmonic polynomials*, J. Approx. Theory **318** (2026), 106303, doi:10.1016/j.jat.2026.106303.
37. I. V. Tokmakov, A. F. Wagner, M. Minkoff, and D. L. Thompson, *Gradient incorporation in one-dimensional applications of interpolating moving least-squares methods for fitting potential energy surfaces*, Theor. Chem. Acc. **118** (2007), no. 4, 755–767, doi:10.1007/s00214-007-0358-7.

38. E. M. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, Princeton, NJ, 1971.
39. H. Wendland, *Scattered Data Approximation*, Cambridge Monographs on Applied and Computational Mathematics, vol. 17, Cambridge University Press, Cambridge, 2005, doi:10.1017/CBO9780511617539.
40. H. Wendland, *Local polynomial reproduction and moving least squares approximation*, IMA J. Numer. Anal. **21** (2001), 285–300.
41. Z. Wu, *Hermite–Birkhoff interpolation of scattered data by radial basis functions*, Approx. Theory Appl. **8** (1992), no. 2, 1–10, doi:10.1007/BF02836101.
42. C. Zuppa, *Error estimates for moving least square approximations*, Bull. Braz. Math. Soc. (N.S.) **34** (2003), no. 2, 231–249, doi:10.1007/s00574-003-0010-7.

Independent researcher, Seattle, Washington, USA
Email address: david@figueiras-garcia.net