


Fig 1. Consolidated Standards of Reporting Trials diagram. PD, pharmacodynamic (population); PK, pharmacokinetic (population); PP, per protocol (population).

Table 1 Atopic dermatitis. Baseline demographics and clinical characteristics (safety population)

	ASN002 20 mg, n = 9	ASN002 40 mg, n = 9	ASN002 80 mg, n = 9	ASN002 Overall, n = 27	Placebo, n = 9
Baseline demographics					
Age, years	38·2 ± 14·36	42·4 ± 13·88	33·1 ± 10·42	37·9 ± 13·09	29·9 ± 9·33
Sex					
Male	5 (56)	5 (56)	5 (56)	15 (56)	3 (33)
Female	4 (44)	4 (44)	4 (44)	12 (44)	6 (67)
Ethnicity					
White	6 (67)	7 (78)	7 (78)	20 (74)	8 (89)
Black	2 (22)	1 (11)	2 (22)	5 (19)	1 (11)
Asian	0 (0·0)	1 (11)	0 (0)	1 (4)	0 (0)
Other	1 (11)	0 (0)	0 (0)	1 (4)	0 (0)
Weight at screening, kg	76·3 ± 11·65	72·3 ± 11·46	85·4 ± 16·23	78·0 ± 13·94	72·3 ± 15·11
Height at screening, cm	169·1 ± 5·57	165·0 ± 8·45	174·0 ± 6·22	169·4 ± 7·59	165·0 ± 9·82
BMI at screening, kg m^{-2}	26·8 ± 4·92	26·6 ± 3·87	28·2 ± 5·44	27·2 ± 4·66	26·5 ± 4·75
Clinical characteristics					
EASI score	29·0 ± 13·49	21·8 ± 6·21	28·2 ± 11·67	26·3 ± 10·97	21·6 ± 6·24
IGA score					
3 (Moderate)	5 (56)	8 (89)	4 (44)	17 (63)	8 (89)
4 (Severe)	4 (44)	1 (11)	5 (56)	10 (37)	1 (11)
BSA	44·1 ± 20·19	31·4 ± 17·96	36·9 ± 25·88	37·5 ± 21·42	25·1 ± 10·49
Pruritus NRS	5·4 ± 2·26	6·1 ± 2·47	6·0 ± 2·55	5·8 ± 2·36	5·9 ± 2·57

Data are presented as mean ± SD or n (%). BMI, body mass index; BSA, body surface area; EASI, Eczema Area and Severity Index; IGA, Investigator's Global Assessment; NRS, numeric rating scale.

and mean terminal half-life ($t_{1/2}$) ranging from 7·3 to 14·1 at steady state. Maximum observed concentration (C_{max}) and area under the plasma concentration (AUC) parameters showed dose-dependent exposure, and AUC approximately proportional to the increase of dose (Table S1; see Supporting Information). Interpatient variability in C_{max} and AUC was low to moderate, and minimal drug accumulation was measured at steady state.

© 2019 The Authors. *British Journal of Dermatology* published by John Wiley & Sons Ltd on behalf of British Association of Dermatologists.




Fig 1. Consolidated Standards of Reporting Trials diagram. PD, pharmacodynamic (population); PK, pharmacokinetic (population); PP, per protocol (population).

Table 1 Atopic dermatitis. Baseline demographics and clinical characteristics (safety population)

	ASN002 20 mg, n = 9	ASN002 40 mg, n = 9	ASN002 80 mg, n = 9	ASN002 Overall, n = 27	Placebo, n = 9
Baseline demographics					
Age, years	38·2 ± 14·36	42·4 ± 13·88	33·1 ± 10·42	37·9 ± 13·09	29·9 ± 9·33
Sex					
Male	5 (56)	5 (56)	5 (56)	15 (56)	3 (33)
Female	4 (44)	4 (44)	4 (44)	12 (44)	6 (67)
Ethnicity					
White	6 (67)	7 (78)	7 (78)	20 (74)	8 (89)
Black	2 (22)	1 (11)	2 (22)	5 (19)	1 (11)
Asian	0 (0·0)	1 (11)	0 (0)	1 (4)	0 (0)
Other	1 (11)	0 (0)	0 (0)	1 (4)	0 (0)
Weight at screening, kg	76·3 ± 11·65	72·3 ± 11·46	85·4 ± 16·23	78·0 ± 13·94	72·3 ± 15·11
Height at screening, cm	169·1 ± 5·57	165·0 ± 8·45	174·0 ± 6·22	169·4 ± 7·59	165·0 ± 9·82
BMI at screening, kg m^{-2}	26·8 ± 4·92	26·6 ± 3·87	28·2 ± 5·44	27·2 ± 4·66	26·5 ± 4·75
Clinical characteristics					
EASI score	29·0 ± 13·49	21·8 ± 6·21	28·2 ± 11·67	26·3 ± 10·97	21·6 ± 6·24
IGA score					
3 (Moderate)	5 (56)	8 (89)	4 (44)	17 (63)	8 (89)
4 (Severe)	4 (44)	1 (11)	5 (56)	10 (37)	1 (11)
BSA	44·1 ± 20·19	31·4 ± 17·96	36·9 ± 25·88	37·5 ± 21·42	25·1 ± 10·49
Pruritus NRS	5·4 ± 2·26	6·1 ± 2·47	6·0 ± 2·55	5·8 ± 2·36	5·9 ± 2·57

Data are presented as mean ± SD or n (%). BMI, body mass index; BSA, body surface area; EASI, Eczema Area and Severity Index; IGA, Investigator's Global Assessment; NRS, numeric rating scale.

and mean terminal half-life ($t_{1/2}$) ranging from 7·3 to 14·1 at steady state. Maximum observed concentration (C_{max}) and area under the plasma concentration (AUC) parameters showed dose-dependent exposure, and AUC approximately proportional to the increase of dose (Table S1; see Supporting Information). Interpatient variability in C_{max} and AUC was low to moderate, and minimal drug accumulation was measured at steady state.

© 2019 The Authors. *British Journal of Dermatology* published by John Wiley & Sons Ltd on behalf of British Association of Dermatologists.