

Simplicial Sets, Spaces, and Singular Chains

Algebraic Topology I | Fall 2026 | September 21 and 23

Contents

1 Presheaves and Functor Categories	1
1.1 Presheaves and Natural Transformations	1
1.2 Representables and Yoneda	2
2 Simplices and Their Operators	2
2.1 The Topological Simplex	2
2.2 The Simplex Category and Affine Operators	3
2.3 Simplicial Sets	3
3 Geometric Realization: Gluing the Simplices	5
3.1 The Quotient and Its Universal Property	5
3.2 Standard Simplices and Elementary Realizations	5
4 The Singular Functor: All Simplices Mapping into a Space	6
4.1 Singular Simplices and Their Operators	6
4.2 Functoriality and Examples	6
5 The Realization-Singular Adjunction	7
5.1 The Natural Bijection	7
5.2 The Unit and Counit	7
6 The Free Simplicial Abelian Group Functor	8
6.1 First Take the Free Abelian Group on a Set	8
6.2 Apply This Construction in Every Degree	8
7 Alternating Faces and Singular Chains	9
7.1 The Alternating-Face Chain Complex	9
7.2 Degenerate Simplices and the Homology of a Point	10
7.3 The Singular Chain Functor	10
8 Coefficients in Topology	11
8.1 Coefficient Complexes and Universal Coefficient Sequences	11
8.2 The Real Projective Plane	11

1 Presheaves and Functor Categories

1.1 Presheaves and Natural Transformations

Definition 1.1. Let $\mathcal{D}$ be a small category, regarded as a diagram shape, and let $\mathcal{C}$ be a locally small category. The category of $\mathcal{C}$ -valued presheaves on $\mathcal{D}$ is

$$\mathcal{P}(\mathcal{D}; \mathcal{C}) = \text{Fun}(\mathcal{D}^{\text{op}}, \mathcal{C}).$$

Its objects are functors $F : \mathcal{D}^{\text{op}} \rightarrow \mathcal{C}$, equivalently contravariant diagrams on $\mathcal{D}$. Thus F assigns an object $F(d)$ to each $d \in \mathcal{D}$ and a restriction map $f^* = F(f) : F(d') \rightarrow F(d)$ to each $f : d \rightarrow d'$,

with

$$\mathrm{id}_d^* = \mathrm{id}_{F(d)}, \quad (g \circ f)^* = f^* \circ g^*.$$

Its morphisms are natural transformations: a morphism $\eta : F \rightarrow G$ is a family $\eta_d : F(d) \rightarrow G(d)$ satisfying

$$G(f) \circ \eta_{d'} = \eta_d \circ F(f) \quad \text{for every } f : d \rightarrow d'.$$

Identities and composition are componentwise: $(\mathrm{id}_F)_d = \mathrm{id}_{F(d)}$ and $(\theta \circ \eta)_d = \theta_d \circ \eta_d$.

The first argument of $\mathcal{P}(\mathcal{D}; \mathcal{C})$ specifies the diagram shape; the second specifies the category of values. The opposite category $\mathcal{D}^{\mathrm{op}}$ reverses arrows. Here small means that objects and morphisms form sets; locally small means that morphisms between any two fixed objects form a set.

More generally, the functor category $\mathrm{Fun}(\mathcal{I}, \mathcal{C})$ has covariant functors $\mathcal{I} \rightarrow \mathcal{C}$ as objects and natural transformations as morphisms. For $f : i \rightarrow j$, naturality reads $G(f)\eta_i = \eta_j F(f)$; composition is again componentwise. The definition above is the case $\mathcal{I} = \mathcal{D}^{\mathrm{op}}$.

1.2 Representables and Yoneda

Definition 1.2. Specializing to $\mathcal{C} = \mathrm{Set}$, write

$$\mathcal{P}(\mathcal{D}) = \mathcal{P}(\mathcal{D}; \mathrm{Set}) = \mathrm{Fun}(\mathcal{D}^{\mathrm{op}}, \mathrm{Set}).$$

Thus an omitted second argument means sets. By contrast, $\mathcal{P}(\mathcal{D}; \mathrm{Ab})$ consists of contravariant diagrams of abelian groups and homomorphisms.

Each object $d \in \mathcal{D}$ defines the representable presheaf $\mathrm{Mor}_{\mathcal{D}}(-, d)$, whose value at e is $\mathrm{Mor}_{\mathcal{D}}(e, d)$ and whose restriction maps are precomposition. Simplicial sets will be the case $\mathcal{D} = \Delta$.

Proposition 1.3. For $F \in \mathcal{P}(\mathcal{D})$ and $d \in \mathcal{D}$, evaluation at id_d gives a natural bijection

$$\mathrm{Mor}_{\mathcal{P}(\mathcal{D})}(\mathrm{Mor}_{\mathcal{D}}(-, d), F) \cong F(d).$$

Proof. An element $x \in F(d)$ defines $\eta_e(f) = f^*x$ for $f : e \rightarrow d$. Functoriality makes η natural, and naturality forces this formula from $x = \eta_d(\mathrm{id}_d)$. These constructions are inverse and natural. $\square$

2 Simplicies and Their Operators

2.1 The Topological Simplex

Definition 2.1. The standard topological n -simplex is


$$|\Delta^n| = \left\{ (t_0, \dots, t_n) \in \mathbb{R}^{n+1} : t_i \geq 0, \sum_{i=0}^n t_i = 1 \right\}.$$

The realization notation will be justified when we construct geometric realization. Its vertices are the coordinate vectors $e_0, \dots, e_n$. The coordinates t_i are called barycentric coordinates.

Thus $|\Delta^0|$ is a point, $|\Delta^1|$ is an interval, and $|\Delta^2|$ is a filled triangle. The ordering of the vertices determines the signs in its oriented boundary:

$$\partial[e_0, e_1, e_2] = [e_1, e_2] - [e_0, e_2] + [e_0, e_1].$$

The triangle can be displayed as



This is an oriented geometric triangle. In a general simplicial set, its edges are not assumed to be morphisms in a category, and the triangle is not an assertion of equality between composites.

2.2 The Simplex Category and Affine Operators

Definition 2.2. The simplex category Δ has objects $[n] = \{0 < \dots < n\}$ for $n \geq 0$ and order-preserving maps as morphisms. A map $\alpha : [m] \rightarrow [n]$ determines the affine map

$$\alpha_* : |\Delta^m| \rightarrow |\Delta^n|, \quad (\alpha_* t)_j = \sum_{\alpha(i)=j} t_i.$$

An empty sum is zero. In particular, $\alpha_*(e_i) = e_{\alpha(i)}$.

For $\beta : [\ell] \rightarrow [m]$, grouping the coordinates gives $(\alpha\beta)_* = \alpha_*\beta_*$. Thus we obtain a covariant functor $\Delta \rightarrow \mathbf{Top}$ that sends $[n]$ to $|\Delta^n|$.

Two kinds of maps generate Δ :

1. The coface $\delta^i : [n-1] \hookrightarrow [n]$ omits i . Its affine map inserts a zero into barycentric coordinates at position i .
2. The codegeneracy $\sigma^i : [n+1] \twoheadrightarrow [n]$ identifies i and $i+1$. Its affine map replaces the adjacent coordinates t_i, t_{i+1} by their sum.

Every order-preserving map factors uniquely as a surjection followed by an injection. Geometrically, it first collapses repeated vertices and then includes a face.

2.3 Simplicial Sets

Definition 2.3. A simplicial set is a presheaf of sets on Δ , that is, a functor $K : \Delta^{\text{op}} \rightarrow \mathbf{Set}$. Write $K_n = K([n])$ and, for $\alpha : [m] \rightarrow [n]$, write $\alpha^* : K_n \rightarrow K_m$. A simplicial map $f : K \rightarrow L$ is a natural transformation. These objects and maps form the presheaf category $\mathcal{P}(\Delta) = \mathbf{Fun}(\Delta^{\text{op}}, \mathbf{Set})$.

The constructions fit into the following diagram of functors:

$$\begin{array}{ccccc} \mathbf{Top} & \xrightarrow{\text{Sing}} & \mathcal{P}(\Delta) & \xrightarrow{\mathbb{Z}[-]} & \mathcal{P}(\Delta; \mathbf{Ab}) & \xrightarrow{C^{\text{alt}}} & \mathbf{Ch}_{\geq 0}(\mathbf{Ab}). \\ & \xleftarrow{|\cdot|} & & \xleftarrow{U} & & & \end{array}$$

Here $|\cdot| \dashv \text{Sing}$ and $\mathbb{Z}[-] \dashv U$: realization is left adjoint to the singular functor, and the free simplicial abelian group functor is left adjoint to forgetting addition. The composite from left to right is the singular chain functor.

Elements of K_n are called n -simplices. The following theorem gives an equivalent description using only faces and degeneracies.

Theorem 2.4. The data of a simplicial set is equivalently a sequence of sets $K_0, K_1, K_2, \dots$ equipped with face maps and degeneracy maps

$$\begin{aligned} d_i : K_n &\rightarrow K_{n-1} & (n \geq 1, 0 \leq i \leq n), \\ s_i : K_n &\rightarrow K_{n+1} & (n \geq 0, 0 \leq i \leq n), \end{aligned}$$

subject to the simplicial identities

$$\begin{aligned} d_i d_j &= d_{j-1} d_i & (i < j), \\ s_i s_j &= s_{j+1} s_i & (i \leq j), \\ d_i s_j &= \begin{cases} s_{j-1} d_i, & i < j, \\ \text{id}, & i = j \text{ or } i = j + 1, \\ s_j d_{i-1}, & i > j + 1. \end{cases} \end{aligned}$$

The identities are required whenever the indicated composites are defined; composition is read from right to left.

More precisely, every such system extends uniquely to a functor $K : \Delta^{\text{op}} \rightarrow \text{Set}$ with $K([n]) = K_n$, $K(\delta^i) = d_i$, and $K(\sigma^i) = s_i$. Under this correspondence, simplicial maps are exactly families of functions $f_n : K_n \rightarrow L_n$ commuting with every face and degeneracy map.

Proof. A functor gives the stated maps by applying it to cofaces and codegeneracies. Their relations in Δ , with composition reversed, give the displayed identities.

Conversely, every order-preserving map factors uniquely as a surjection followed by an injection. The surjection is specified by its repeated adjacent values, and the injection by its omitted values. Thus each has a canonical expression in codegeneracies or cofaces. The elementary relations move any expression into this form: mixed relations move omissions past repetitions, cancelling when necessary, and the remaining relations order the omissions and repetitions. Consequently these relations present Δ by generators and relations.

Assigning d_i and s_i to these generators therefore defines the pullback along every map of Δ , independently of its expression. The same relations ensure functoriality. Generation gives uniqueness and also shows that commuting with faces and degeneracies is equivalent to naturality. See [3] for the generators-and-relations formulation. $\square$

The faces of an edge e are its target $d_0 e$ and source $d_1 e$. The simplex $s_0 v$ is the degenerate edge at v ; both of its endpoints are v . Geometrically, the first identity says that deleting vertex j and then vertex $i < j$ has the same effect as deleting vertex i and then the renumbered vertex $j - 1$.

A simplex is *degenerate* if it is in the image of some s_i ; otherwise it is *nondegenerate*. A degenerate simplex is still an element of K_n . It is not removed from the simplicial set.

Example 2.5. The simplicial set Δ^n is defined by

$$\Delta_m^n = \text{Mor}_{\Delta}([m], [n]), \quad \beta^*(\alpha) = \alpha \beta.$$

Its nondegenerate m -simplices are precisely the injective order-preserving maps $[m] \hookrightarrow [n]$. Thus Δ^1 has two nondegenerate vertices and one nondegenerate edge, but it has simplices in every degree.

The notations $[n]$, Δ^n , and $|\Delta^n|$ refer respectively to an ordered set, a simplicial set, and a topological space.

Proposition 2.6. There is a natural bijection

$$\text{Mor}_{\mathcal{P}(\Delta)}(\Delta^n, K) \cong K_n.$$

Proof. The standard simplex Δ^n is the representable presheaf $\text{Mor}_{\Delta}(-, [n])$. Apply Yoneda: $x \in K_n$ corresponds to the simplicial map $\Delta^n \rightarrow K$ whose degree- m component is $\text{Mor}_{\Delta}([m], [n]) \rightarrow K_m$, $\alpha \mapsto \alpha^* x$. $\square$

3 Geometric Realization: Gluing the Simplices

3.1 The Quotient and Its Universal Property

Definition 3.1. The geometric realization of a simplicial set K is the quotient space

$$|K| = \left(\coprod_{n \geq 0} K_n \times |\Delta^n| \right) / \sim,$$

where each K_n has the discrete topology and the equivalence relation is generated by

$$(\alpha^*x, t) \sim (x, \alpha_*t) \quad \text{for } \alpha : [m] \rightarrow [n], x \in K_n, t \in |\Delta^m|.$$

Write $[x, t]$ for the equivalence class of (x, t) .

The face relations attach the boundary of one simplex to the specified lower-dimensional simplices. The degeneracy relations collapse a redundant simplex onto its lower-dimensional source. For example,

$$[s_0v, (t_0, t_1)] = [v, (1)],$$

so a degenerate edge becomes a point, not an extra loop.

Every $x \in K_n$ has a characteristic map

$$\chi_x : |\Delta^n| \rightarrow |K|, \quad t \mapsto [x, t].$$

These maps obey the commutative diagrams

$$\begin{array}{ccc} |\Delta^m| & \xrightarrow{\alpha_*} & |\Delta^n| \\ \chi_{\alpha^*x} \downarrow & & \downarrow \chi_x \\ |K| & \xlongequal{\quad} & |K|. \end{array}$$

Define the category of simplices Δ/K to have objects $([n], x)$ with $x \in K_n$; a morphism $([m], y) \rightarrow ([n], x)$ is a map $\alpha : [m] \rightarrow [n]$ with $y = \alpha^*x$. Then

$$|K| \cong \operatorname{colim}_{([n], x) \in \Delta/K} |\Delta^n|.$$

This is a colimit in **Top**: continuous maps $|K| \rightarrow X$ correspond exactly to families of continuous maps $f_x : |\Delta^n| \rightarrow X$ satisfying $f_{\alpha^*x} = f_x \alpha_*$. Indeed, such a family defines a continuous map from the disjoint union, and compatibility is precisely what makes it descend through the quotient.

A simplicial map $f : K \rightarrow L$ induces

$$|f| : |K| \rightarrow |L|, \quad [x, t] \mapsto [f_n(x), t].$$

Compatibility with the relation follows from $f_m(\alpha^*x) = \alpha^*f_n(x)$. This defines a functor $|-| : \mathcal{P}(\Delta) \rightarrow \mathbf{Top}$.

3.2 Standard Simplices and Elementary Realizations

Example 3.2.

1. The realization of the standard simplicial simplex is canonically the topological simplex introduced in Section 2. The map $|\Delta^n| \rightarrow \{t \in \mathbb{R}^{n+1} : t_i \geq 0, \sum_i t_i = 1\}$ is given by $[\alpha, t] \mapsto \alpha_*t$. Its inverse sends u to $[\operatorname{id}_{[n]}, u]$; the defining relation proves that the two composites are identities. We use $|\Delta^n|$ for this identified space.

2. A constant simplicial set on a set S has value S and every structure map is the identity. Its realization is the discrete space S .
3. Let $\partial\Delta^1 \subset \Delta^1$ consist of the two vertices and all their degeneracies. The degreewise quotient

$$S_\Delta^1 = \Delta^1 / \partial\Delta^1$$

has one vertex v , one nondegenerate edge e with $d_0e = d_1e = v$, and no higher nondegenerate simplices. Its realization is the interval with its endpoints identified, hence a circle.

Repeated vertices, loops, and distinct simplices with the same boundary are all allowed in a simplicial set. This flexibility goes beyond an abstract simplicial complex.

4 The Singular Functor: All Simplices Mapping into a Space

4.1 Singular Simplices and Their Operators

Definition 4.1. We work in $\mathbf{Top}$, the category of topological spaces and continuous maps, without separation assumptions. For a topological space X , define

$$(\mathrm{Sing} X)_n = \mathrm{Mor}_{\mathbf{Top}}(|\Delta^n|, X).$$

For $\alpha : [m] \rightarrow [n]$, define $\alpha^*(\tau) = \tau \alpha_*$. The sets of continuous maps here are regarded simply as sets.

Precomposition and $(\alpha\beta)_* = \alpha_*\beta_*$ give the required contravariance. Thus $\mathrm{Sing} X$ is a simplicial set, called the *singular simplicial set* of X .

The first few degrees have direct geometric meanings:

1. $(\mathrm{Sing} X)_0$ is the underlying set of points of X .
2. $(\mathrm{Sing} X)_1$ is the set of continuous paths, with the interval parametrized by $(1-u, u)$ for $0 \leq u \leq 1$. Thus $d_1\gamma = \gamma(0)$ and $d_0\gamma = \gamma(1)$.
3. $(\mathrm{Sing} X)_2$ consists of continuous maps of a filled triangle into X . Its three faces are the restrictions to its three ordered edges.

A singular simplex need not be injective or linear, and its image need not look like a geometric simplex. In particular, an arbitrary space need not come with a chosen triangulation for its singular simplicial set to be defined.

For $\tau : |\Delta^n| \rightarrow X$, the degeneracy $s_i\tau : |\Delta^{n+1}| \rightarrow X$ is

$$(t_0, \dots, t_{n+1}) \mapsto \tau(t_0, \dots, t_{i-1}, t_i + t_{i+1}, t_{i+2}, \dots, t_{n+1}).$$

For $n = 0$, this is a constant path. A path that goes out and comes back is generally *not* a degenerate edge: degeneracy means the specified factorization through a lower-dimensional standard simplex, not merely that endpoints coincide.

4.2 Functoriality and Examples

If $f : X \rightarrow Y$ is continuous, postcomposition defines

$$(\mathrm{Sing} f)_n : (\mathrm{Sing} X)_n \rightarrow (\mathrm{Sing} Y)_n, \quad \tau \mapsto f\tau.$$

Since postcomposition commutes with precomposition, this is simplicial and defines the functor $\mathrm{Sing} : \mathbf{Top} \rightarrow \mathcal{P}(\Delta)$.

Example 4.2. The singular simplicial set of a point is Δ^0 : it has one simplex in every degree. If X is discrete, every map $|\Delta^n| \rightarrow X$ is constant, so $\text{Sing } X$ is the constant simplicial set on X . For $X = [0, 1]$, the simplicial set $\text{Sing } X$ is much larger than Δ^1 : it includes every continuous path and every continuous singular simplex in the interval.

5 The Realization-Singular Adjunction

An adjunction here is a natural bijection between two sets of maps. The relevant bijection says that mapping a glued space into X is the same as assigning compatible singular simplices in X to its pieces.

5.1 The Natural Bijection

Theorem 5.1. For a simplicial set K and a topological space X , there is a natural bijection

$$\text{Mor}_{\text{Top}}(|K|, X) \cong \text{Mor}_{\mathcal{P}(\Delta)}(K, \text{Sing } X).$$

Consequently, geometric realization is left adjoint to the singular functor.

Proof. By the colimit universal property, a continuous map $|K| \rightarrow X$ is a compatible family $f_x : |\Delta^n| \rightarrow X$. Since $(\text{Sing } X)_n = \text{Mor}_{\text{Top}}(|\Delta^n|, X)$, this is exactly a family of functions

$$K_n \rightarrow (\text{Sing } X)_n, \quad x \mapsto f_x,$$

commuting with restriction maps: a morphism of presheaves $K \rightarrow \text{Sing } X$. Explicitly, $f : |K| \rightarrow X$ determines $g : K \rightarrow \text{Sing } X$ with $g_n(x) = f \chi_x$. Conversely, $g : K \rightarrow \text{Sing } X$ determines $f : |K| \rightarrow X$ by $f([x, t]) = g_n(x)(t)$. Both commute with precomposition in K and postcomposition in X , so the bijection is natural. $\square$

5.2 The Unit and Counit

The identity of $|K|$ corresponds to the *unit*

$$(\eta_K)_n : K_n \rightarrow (\text{Sing } |K|)_n, \quad x \mapsto \chi_x.$$

Thus each abstract simplex becomes the singular simplex that parametrizes its own image in the realization.

The identity of $\text{Sing } X$ corresponds to the *counit*, or evaluation map,

$$\epsilon_X : |\text{Sing } X| \rightarrow X, \quad [\tau, t] \mapsto \tau(t).$$

The triangle identities are

$$\epsilon_{|K|} \circ |\eta_K| = \text{id}_{|K|}, \quad \text{Sing}(\epsilon_X) \circ \eta_{\text{Sing } X} = \text{id}_{\text{Sing } X}.$$

For example, the first composite sends $[x, t]$ to $[\chi_x, t]$ and then to $\chi_x(t) = [x, t]$. The second says that evaluating the characteristic simplex associated to τ recovers τ .

Remark 5.2. The map η_{Δ^1} includes the simplices determined by the ordered vertices into all singular simplices of the interval; most continuous paths are not in its image. Likewise, evaluation need not be a homeomorphism. The deeper comparison with homotopy types is a theorem beyond the adjunction itself, and is not needed to construct singular chains.

6 The Free Simplicial Abelian Group Functor

6.1 First Take the Free Abelian Group on a Set

Definition 6.1. For a set S , the free abelian group on S is

$$\mathbb{Z}[S] = \bigoplus_{s \in S} \mathbb{Z}s.$$

Its elements are finite formal sums $\sum_s a_s s$ with $a_s \in \mathbb{Z}$. For a function $f : S \rightarrow T$, set

$$\mathbb{Z}[f] \left(\sum_s a_s s \right) = \sum_s a_s f(s).$$

We write s also for the formal generator labelled by $s \in S$; addition in $\mathbb{Z}[S]$ is addition of formal sums. If f identifies two elements, their coefficients add. For example, if $f(s) = f(s') = t$, then $2s - s'$ maps to t .

For every abelian group B , a function $S \rightarrow U(B)$ extends uniquely to a homomorphism $\mathbb{Z}[S] \rightarrow B$ by linearity. Here $U(B)$ denotes its underlying set. Thus

$$\text{Mor}_{\text{Ab}}(\mathbb{Z}[S], B) \cong \text{Mor}_{\text{Set}}(S, U(B)).$$

6.2 Apply This Construction in Every Degree

Definition 6.2. A simplicial abelian group is a functor $A : \Delta^{\text{op}} \rightarrow \text{Ab}$. Its structure maps, including its faces and degeneracies, are group homomorphisms. Thus simplicial abelian groups form the presheaf category $\mathcal{P}(\Delta; \text{Ab}) = \text{Fun}(\Delta^{\text{op}}, \text{Ab})$. Forgetting the group structures degree-wise defines $U : \mathcal{P}(\Delta; \text{Ab}) \rightarrow \mathcal{P}(\Delta)$.

For a simplicial set K , define

$$\mathbb{Z}[K]_n = \mathbb{Z}[K_n], \quad \alpha^* \left(\sum_x a_x x \right) = \sum_x a_x \alpha^* x.$$

The simplicial identities hold on generators because they hold in K , and therefore hold on all finite sums. In particular,

$$d_i x = d_i x, \quad s_i x = s_i x.$$

A simplicial map $f : K \rightarrow L$ induces the degree-wise homomorphisms $\mathbb{Z}[f_n] : \mathbb{Z}[K_n] \rightarrow \mathbb{Z}[L_n]$ given by $x \mapsto f_n(x)$. We obtain the functor

$$\mathbb{Z}[-] : \mathcal{P}(\Delta) \rightarrow \mathcal{P}(\Delta; \text{Ab}).$$

The group in degree n is free on *all* n -simplices, including degenerate simplices. It is not the free group on the vertices alone.

Proposition 6.3. Let $L : \mathcal{D} \rightleftarrows \mathcal{E} : R$ be an adjunction and $\mathcal{T}$ a small category. Postcomposition induces an adjunction

$$L_* : \text{Fun}(\mathcal{T}, \mathcal{D}) \rightleftarrows \text{Fun}(\mathcal{T}, \mathcal{E}) : R_*.$$

Proof. For $F : \mathcal{T} \rightarrow \mathcal{D}$ and $G : \mathcal{T} \rightarrow \mathcal{E}$, transpose each component $LF(i) \rightarrow G(i)$ under $L \dashv R$. Naturality of the adjunction bijection identifies the naturality equations for these components with those for $F(i) \rightarrow RG(i)$. Thus componentwise transposition gives a natural bijection between the two sets of natural transformations. $\square$

Proposition 6.4. For $K \in \mathcal{P}(\Delta)$ and $A \in \mathcal{P}(\Delta; \mathbf{Ab})$, there is a natural bijection

$$\mathrm{Mor}_{\mathcal{P}(\Delta; \mathbf{Ab})}(\mathbb{Z}[K], A) \cong \mathrm{Mor}_{\mathcal{P}(\Delta)}(K, U(A)).$$

Proof. Apply the pointwise adjunction proposition to $\mathbb{Z}[-] : \mathbf{Set} \rightleftarrows \mathbf{Ab} : U$ with $\mathcal{T} = \Delta^{\mathrm{op}}$. Concretely, $g : K \rightarrow U(A)$ extends uniquely by

$$\bar{g}_n \left(\sum_x a_x x \right) = \sum_x a_x g_n(x).$$

Naturality of g makes these extensions commute with every simplicial operator. Restriction to generators is the inverse. $\square$

The unit in degree n is the function $K_n \rightarrow \mathbb{Z}[K_n]$ sending a simplex to its formal generator. The counit is the simplicial homomorphism $\mathbb{Z}[U(A)] \rightarrow A$ that evaluates each finite formal sum using the addition in A . A generator labelled by a sum in A is distinct from the formal sum of the corresponding generators; the counit sends them to the same element.

Remark 6.5. The functor $\mathbb{Z}[-]$ freely introduces sums and additive inverses while retaining all faces and degeneracies. A simplicial abelian group still has $n + 1$ face maps in degree n . The next construction combines them into one differential.

7 Alternating Faces and Singular Chains

7.1 The Alternating-Face Chain Complex

Definition 7.1. All chain complexes here are homologically graded and concentrated in nonnegative degrees. We retain degenerate simplices. For a simplicial abelian group A , let $C^{\mathrm{alt}}(A)_n = A_n$ for $n \geq 0$, and put

$$\partial_n = \sum_{i=0}^n (-1)^i d_i : A_n \rightarrow A_{n-1} \quad (n \geq 1), \quad \partial_0 : A_0 \rightarrow 0.$$

The first two differentials are $\partial_1 = d_0 - d_1$ and $\partial_2 = d_0 - d_1 + d_2$. Thus the boundary of an edge is target minus source, and the boundary of a triangle is its three edges with alternating signs.

Proposition 7.2. The equation $\partial_{n-1} \partial_n = 0$ holds. A simplicial homomorphism induces a chain map, so this construction defines $C^{\mathrm{alt}} : \mathcal{P}(\Delta; \mathbf{Ab}) \rightarrow \mathbf{Ch}_{\geq 0}(\mathbf{Ab})$.

Proof. For $n \geq 2$, expand the composite as

$$\sum_{i=0}^{n-1} \sum_{j=0}^n (-1)^{i+j} d_i d_j.$$

Pair the term with $i < j$ with the term indexed by $(j-1, i)$. Their maps agree by $d_i d_j = d_{j-1} d_i$, and their signs are opposite because $i+j$ and $j-1+i$ differ by one. This pairs every term exactly once. In degree one the assertion follows from $\partial_0 = 0$.

A simplicial homomorphism commutes with each face map, so it commutes with their signed sum. Identities and composition are preserved degree-wise. $\square$

Each codimension-two face occurs twice, with opposite signs. See also [2].

For a simplicial set K , write

$$C_*(K; \mathbb{Z}) = C^{\mathrm{alt}}(\mathbb{Z}[K]), \quad H_n(K; \mathbb{Z}) = \ker(\partial_n) / \mathrm{im}(\partial_{n+1}).$$

These are the unnormalized simplicial chains and their homology. In positive degrees,

$$\partial x = \sum_{i=0}^n (-1)^i d_i x.$$

7.2 Degenerate Simplices and the Homology of a Point

A degenerate simplex is not the zero element of a free chain group. For instance, the constant edge $a = s_0 v$ has $\partial a = 0$, but a is a nonzero generator of C_1 . It becomes a boundary: if $b = s_0 s_0 v$, then all three faces of b equal a , so $\partial b = a - a + a = a$.

Example 7.3. For $K = \Delta^0$, each chain group is $\mathbb{Z}$, and all face maps are identities. Hence

$$\partial_n = \begin{cases} 0, & n \text{ odd,} \\ \text{id}, & n \geq 2 \text{ even.} \end{cases}$$

In low degrees the complex is

$$\dots \rightarrow \mathbb{Z} \xrightarrow{0} \mathbb{Z} \xrightarrow{\text{id}} \mathbb{Z} \xrightarrow{0} \mathbb{Z} \rightarrow 0,$$

where the rightmost $\mathbb{Z}$ is in degree zero. It has $H_0 \cong \mathbb{Z}$ and $H_n = 0$ for $n > 0$. Keeping the degenerate simplices therefore gives the correct answer without deleting any generators by hand.

7.3 The Singular Chain Functor

Definition 7.4. The singular chain complex of a topological space X is

$$C_\bullet^{\text{sing}}(X; \mathbb{Z}) = C^{\text{alt}}(\mathbb{Z}[\text{Sing } X]).$$

Its degree- n group is freely generated by continuous maps $\tau : |\Delta^n| \rightarrow X$. For $n \geq 1$, its boundary is

$$\partial \tau = \sum_{i=0}^n (-1)^i \tau \delta_*^i.$$

The singular homology of X is $H_n^{\text{sing}}(X; \mathbb{Z}) = H_n(C_\bullet^{\text{sing}}(X; \mathbb{Z}))$.

The construction separates three steps:

1. Sing records all continuous simplices and their restrictions and repetitions.
2. $\mathbb{Z}[-]$ permits finite integral combinations of these simplices.
3. C^{alt} adds the face maps with signs to define a boundary operator.

None of these steps requires choosing a triangulation of X .

For a continuous map $f : X \rightarrow Y$, the induced chain map sends τ to $f\tau$. For every $n \geq 1$, we have the commutative square

$$\begin{array}{ccc} C_n^{\text{sing}}(X; \mathbb{Z}) & \xrightarrow{f_\#} & C_n^{\text{sing}}(Y; \mathbb{Z}) \\ \partial \downarrow & & \downarrow \partial \\ C_{n-1}^{\text{sing}}(X; \mathbb{Z}) & \xrightarrow{f_\#} & C_{n-1}^{\text{sing}}(Y; \mathbb{Z}). \end{array}$$

It follows that singular homology is functorial.

Example 7.5. For a path $\gamma : x \rightarrow y$, its boundary is $y - x$. Consequently,

$$H_0^{\text{sing}}(X; \mathbb{Z}) \cong \mathbb{Z}[\pi_0^{\text{path}} X],$$

where $\pi_0^{\text{path}} X$ is the set of path components, not merely connected components.

For a singular triangle τ with ordered edges $\gamma_{01}, \gamma_{02}, \gamma_{12}$,

$$\partial \tau = \gamma_{12} - \gamma_{02} + \gamma_{01}.$$

Thus the cycle given by the two successive edges minus the long edge is a boundary. This is a relation among chains modulo boundaries; the individual nonclosed edges need not represent classes in H_1 .

8 Coefficients in Topology

8.1 Coefficient Complexes and Universal Coefficient Sequences

Let X be a simplicial set and let $A_\bullet = C_\bullet(X; \mathbb{Z})$. Each A_n is a free abelian group on all n -simplices. Define

$$H_n(X; G) = H_n(A_\bullet \otimes_{\mathbb{Z}} G), \quad H^n(X; G) = H^n(\text{Mor}_{\mathbb{Z}}(A_\bullet, G)).$$

Here G is an abelian group and $H_n(X; \mathbb{Z}) = H_n(A_\bullet)$. Applied to $X = \text{Sing}(Y)$, these are precisely the usual definitions of singular homology and cohomology: the chains are free on singular simplices and the differential is the alternating sum of restrictions to faces.

Remark 8.1. For any chain complex $A_\bullet$ of free abelian groups concentrated in nonnegative degrees, there are natural short exact sequences

$$0 \rightarrow H_n(A) \otimes_{\mathbb{Z}} G \rightarrow H_n(A \otimes_{\mathbb{Z}} G) \rightarrow \text{Tor}_1^{\mathbb{Z}}(H_{n-1}(A), G) \rightarrow 0$$

and

$$0 \rightarrow \text{Ext}_{\mathbb{Z}}^1(H_{n-1}(A), G) \rightarrow H^n(\text{Mor}_{\mathbb{Z}}(A, G)) \rightarrow \text{Mor}_{\mathbb{Z}}(H_n(A), G) \rightarrow 0.$$

Take $H_{-1}(A) = 0$. Both sequences split as sequences of abelian groups, but in general there is no natural choice of splitting. These are the algebraic universal coefficient theorems; we state them here without a full proof.

The algebra behind these statements is that subgroups of free abelian groups are free. Writing $Z_n = \ker \partial_n$ and $B_n = \text{im } \partial_{n+1}$, the sequence

$$0 \rightarrow Z_n \rightarrow A_n \rightarrow B_{n-1} \rightarrow 0$$

splits, whereas $0 \rightarrow B_n \rightarrow Z_n \rightarrow H_n(A) \rightarrow 0$ is a free resolution of $H_n(A)$. Applying tensor or Hom to the latter sequence produces the Tor or Ext correction. The first sequence explains why choices of splittings enter the calculation.

8.2 The Real Projective Plane

Example 8.2. The cellular chain complex of $\mathbb{RP}^2$ over $\mathbb{Z}$ is

$$0 \rightarrow \mathbb{Z} \xrightarrow{2} \mathbb{Z} \xrightarrow{0} \mathbb{Z} \rightarrow 0$$

in degrees 2, 1, 0. Thus $H_1(\mathbb{RP}^2; \mathbb{Z}) = \mathbb{Z}/2$ and $H_2(\mathbb{RP}^2; \mathbb{Z}) = 0$. The cellular comparison theorem identifies these groups with singular homology.

For coefficients $\mathbb{Z}/2$, the multiplication-by-two differential vanishes, giving

$$H_2(\mathbb{RP}^2; \mathbb{Z}/2) \cong \mathbb{Z}/2.$$

This class is invisible in $H_2(\mathbb{RP}^2; \mathbb{Z}) \otimes \mathbb{Z}/2$, which is zero. It is supplied by

$$\mathrm{Tor}_1^{\mathbb{Z}}(H_1(\mathbb{RP}^2; \mathbb{Z}), \mathbb{Z}/2) = \mathrm{Tor}_1^{\mathbb{Z}}(\mathbb{Z}/2, \mathbb{Z}/2) \cong \mathbb{Z}/2.$$

Likewise, applying $\mathrm{Mor}_{\mathbb{Z}}(-, \mathbb{Z})$ to the cellular complex gives $H^2(\mathbb{RP}^2; \mathbb{Z}) \cong \mathbb{Z}/2$. This is the Ext contribution

$$\mathrm{Ext}_{\mathbb{Z}}^1(H_1(\mathbb{RP}^2; \mathbb{Z}), \mathbb{Z}) \cong \mathbb{Z}/2,$$

even though $\mathrm{Mor}_{\mathbb{Z}}(H_2(\mathbb{RP}^2; \mathbb{Z}), \mathbb{Z}) = 0$.

A simplicial chain complex is free in each degree, but it is usually not a resolution of its degree-zero homology: it may have higher homology. To compute Tor of a single module, use an actual resolution. To compute homology with coefficients of a space, retain the full chain complex of that space.

References

- [1] Kerodon, Section 1.2.3, *The Geometric Realization of a Simplicial Set*, especially Proposition 1.2.3.4 and Corollary 1.2.3.5. <https://kerodon.net/tag/0021>. The realization-singular bijection in this lecture is proved explicitly in Section 5.
- [2] The Stacks Project, Section 14.23, *Simplicial Objects and Chain Complexes*. <https://stacks.math.columbia.edu/tag/0194>. Our notation $C^{\mathrm{alt}}(A)$ denotes the alternating-face complex called $s(A)$ there.
- [3] The Stacks Project, Lemmas 14.2.4 and 14.3.2, *The Category of Finite Ordered Sets and Simplicial Objects*. <https://stacks.math.columbia.edu/tag/0169>.