

Derived Functors: Tensor Products, Tor, and Ext

Math 615 Algebraic Topology

A presentation records generators and relations. Tensoring a presentation computes a tensor product. Continuing the presentation to a resolution records relations among relations; tensoring that resolution computes Tor. Applying the morphism functor to the same resolution reverses the arrows and produces the cochain complex computing Ext.

All rings have an identity and all modules are unital. In Sections 1–3, R is commutative. In Sections 4–5, R is an arbitrary ring and R -modules are left modules. Write $\text{Mor}_R(M, N)$ for the abelian group of R -linear maps. Unless explicitly specified, modules and presentations need not be finitely generated.

Contents

1	Tensor Products and Their Computation	1
2	Resolutions and Relations Among Relations	4
3	Tor and the Failure of Exactness	6
4	Cochain Complexes over an Arbitrary Ring	8
5	Ext and a Noncommutative Computation	10

1 Tensor Products and Their Computation

1.1 The Universal Bilinear Map

Fix a commutative ring R . An R -bilinear map $b : M \times N \rightarrow L$ is linear in each variable separately. The tensor product turns such a map into a linear map with a single input.

Definition 1.1 (Tensor Product). The tensor product of M and N is an R -module $M \otimes_R N$ with a bilinear map $\tau : M \times N \rightarrow M \otimes_R N$ such that, for every bilinear $b : M \times N \rightarrow L$, there is a unique linear $\tilde{b}$ with $\tilde{b}\tau = b$. Write $\tau(m, n) = m \otimes n$.

$$\begin{array}{ccc} M \times N & \xrightarrow{b} & L \\ \tau \downarrow & \nearrow \tilde{b} & \\ M \otimes_R N & & \end{array}$$

Proposition 1.2 (Construction for Arbitrary Modules). Let $F = \bigoplus_{(m,n) \in M \times N} R$ be the free R -module with basis symbols $[m, n]$. Let $U \subseteq F$ be generated by

$$\begin{aligned} [m + m', n] - [m, n] - [m', n], & \quad [m, n + n'] - [m, n] - [m, n'], \\ [rm, n] - r[m, n], & \quad [m, rn] - r[m, n] \end{aligned}$$

for all indicated elements. Then F/U , with $m \otimes n = [m, n] + U$, is a tensor product. It is unique up to a unique isomorphism compatible with τ .

Proof of 1.2. The relations make τ bilinear. A function $(m, n) \mapsto b(m, n)$ extends uniquely to a linear map $F \rightarrow L$, and that map kills U exactly when b is bilinear. It therefore factors uniquely through F/U . For

uniqueness, the universal properties of two tensor products give maps in both directions compatible with τ ; their composites are identities by uniqueness. $\square$

Used in: Propositions 1.3 and 1.4.

Every tensor is a finite sum of elementary tensors. Such an expression need not be unique, and a tensor need not be elementary. For example, over a field k , the tensor $e_1 \otimes e_1 + e_2 \otimes e_2$ in $k^2 \otimes_k k^2$ corresponds to a rank-two matrix, whereas an elementary tensor corresponds to a matrix of rank at most one.

Linear maps $f : M \rightarrow M'$ and $g : N \rightarrow N'$ induce $f \otimes g$ by $m \otimes n \mapsto f(m) \otimes g(n)$. The universal property also gives the natural isomorphisms

$$R \otimes_R N \cong N, \quad M \otimes_R N \cong N \otimes_R M, \quad \left(\bigoplus_{i \in I} R \right) \otimes_R N \cong \bigoplus_{i \in I} N.$$

The first map sends $r \otimes n$ to rn , the second interchanges the factors, and the third sends $e_i \otimes n$ to n in the i th summand. Their inverse maps are specified on the same generators. Each vector in a direct sum has finite support. For a finite indexing set I , we write R^I and N^I .

Proposition 1.3 (Tensor–Morphism Adjunction and Right Exactness). There is a natural isomorphism

$$\text{Mor}_R(M \otimes_R N, L) \cong \text{Mor}_R(M, \text{Mor}_R(N, L)).$$

For fixed N , tensoring preserves cokernels and direct sums. In particular, if $M' \xrightarrow{u} M \xrightarrow{v} M'' \rightarrow 0$ is exact, then

$$M' \otimes_R N \xrightarrow{u \otimes \text{id}} M \otimes_R N \xrightarrow{v \otimes \text{id}} M'' \otimes_R N \rightarrow 0$$

is exact.

Proof of 1.3. Commutativity makes $\text{Mor}_R(N, L)$ an R -module by $(r\phi)(n) = r\phi(n)$. Sending b to the map $m \mapsto (n \mapsto b(m, n))$ identifies bilinear maps with the right-hand side. The universal property identifies them with the left-hand side. Thus $-\otimes_R N$ is a left adjoint, so it preserves colimits, including cokernels and direct sums. Explicitly, a bilinear map on $M \times N$ vanishing on $u(M') \times N$ factors uniquely through $M'' \times N$; this is exactly the asserted cokernel property. $\square$

Used in: Proposition 1.4 and Definition 3.1.

1.2 Presentations Give a General Computation

A presentation is an exact sequence $\bigoplus_{j \in J} R \xrightarrow{A} \bigoplus_{i \in I} R \rightarrow M \rightarrow 0$. Its basis elements indexed by I give generators m_i of M , and its columns give relations among those generators. Such a presentation always exists: take a free module surjecting onto M , then a free module surjecting onto the kernel.

Proposition 1.4 (One Presentation or Two). For an arbitrary presentation of M and an arbitrary module N ,

$$M \otimes_R N \cong \text{coker}(A_N : \bigoplus_{j \in J} N \rightarrow \bigoplus_{i \in I} N). \quad (1)$$

If $A(e_j) = \sum_i a_{ij}e_i$, then

$$A_N((n_j)_{j \in J}) = \left(\sum_{j \in J} a_{ij}n_j \right)_{i \in I}.$$

Each column of A has finite support, and each input has finite support, so this formula is defined even for infinite sets.

If also $\bigoplus_{j \in J'} R \xrightarrow{B} \bigoplus_{i \in I'} R \rightarrow N \rightarrow 0$ is a presentation, then

$$M \otimes_R N \cong \frac{\bigoplus_{(i,i') \in I \times I'} R}{\text{im}(A \otimes \text{id}) + \text{im}(\text{id} \otimes B)}. \quad (2)$$

The two maps in the denominator have domains $\bigoplus_{(j,i') \in J \times I'} R$ and $\bigoplus_{(i,j') \in I \times J'} R$, respectively, and target $\bigoplus_{(i,i') \in I \times I'} R$.

Proof of 1.4. Apply Proposition 1.3 to the presentation of M and identify the tensors of free modules with direct sums of N . This proves (1). For (2), the generator indexed by (i, i') maps to $m_i \otimes n_{i'}$. The relations from A impose every relation among the m_i for each fixed $n_{i'}$, and those from B impose every relation among the $n_{i'}$ for each fixed m_i . A linear map out of this quotient is therefore precisely a bilinear map $M \times N \rightarrow L$. Apply Definition 1.1. $\square$

Used in: Examples 1.6, 1.7, and 3.3.

Remark 1.5 (What the Formula Computes). For finite presentations $A : R^a \rightarrow R^b$ and $B : R^c \rightarrow R^d$, formula (2) is the cokernel of the block matrix

$$[A \otimes I_d \mid I_b \otimes B] : R^{ad+bc} \rightarrow R^{bd},$$

using the basis order specified by the tensor factors. Over $\mathbb{Z}$ one can reduce finite matrices by Smith normal form. Over an arbitrary ring, the cokernel presentation is still a complete algebraic description, but a finite effective algorithm for simplifying it requires additional input about the ring. Arbitrary modules need not admit finite presentations.

Example 1.6 (Quotients and Cyclic Modules). For any ideal $I \subseteq R$,

$$(R/I) \otimes_R N \cong N/IN.$$

The map sends $(r + I) \otimes n$ to $rn + IN$, with inverse induced by $n \mapsto (1 + I) \otimes n$. Here IN consists of finite sums of elements an with $a \in I$. Hence

$$(R/I) \otimes_R (R/J) \cong R/(I + J).$$

For positive integers a, b , this gives $\mathbb{Z}/a\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/b\mathbb{Z} \cong \mathbb{Z}/\gcd(a, b)\mathbb{Z}$. In particular, nonzero modules can have zero tensor product.

Example 1.7 (A Presentation with Two Generators). Let $M = \text{coker } A$ over $\mathbb{Z}$, where

$$A = \begin{pmatrix} 2 & 0 \\ 4 & 6 \end{pmatrix} : \mathbb{Z}^2 \rightarrow \mathbb{Z}^2, \quad N = \mathbb{Z}/4\mathbb{Z}.$$

Tensoring replaces A by its action on N^2 , which is the diagonal matrix with entries 2, 2. Therefore

$$M \otimes_{\mathbb{Z}} N \cong N^2 / (2N \oplus 2N) \cong (\mathbb{Z}/2\mathbb{Z})^2.$$

In coordinates, the map on $(\mathbb{Z}/4\mathbb{Z})^2$ is

$$A_N(x, y) = (2x, 4x + 6y) = (2x, 2y).$$

Its image consists of $(0, 0), (2, 0), (0, 2), (2, 2)$. The map $(x, y) \mapsto (x \bmod 2, y \bmod 2)$ is surjective with precisely this kernel, so it gives the displayed quotient isomorphism. For comparison, the integer row operation $\text{row}_2 \leftarrow \text{row}_2 - 2\text{row}_1$ changes A to $\text{diag}(2, 6)$. Thus $M \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/6\mathbb{Z}$, and tensoring the two summands with $\mathbb{Z}/4\mathbb{Z}$ gives the same two copies of $\mathbb{Z}/2\mathbb{Z}$.

2 Resolutions and Relations Among Relations

2.1 Free and Projective Resolutions

The preceding construction computes degree zero. To measure the exactness lost by tensoring, we extend a presentation to an exact sequence. The constructions in this section work for left modules over any ring; we continue with commutative R until Section 4.

Definition 2.1 (Projective Module). A module P is projective if, for every surjection $q : E \rightarrow L$ and linear map $f : P \rightarrow L$, there is a linear $\tilde{f} : P \rightarrow E$ with $q\tilde{f} = f$. Equivalently, $\text{Mor}_R(P, -)$ is exact. Free modules are projective: lift the images of their basis elements. A projective module is a direct summand of a free module, since a free surjection onto it splits. Conversely, any direct summand of a free module has the lifting property.

Definition 2.2 (Injective Module). A left R -module I is injective if, for every injection $j : A \rightarrow B$ and map $f : A \rightarrow I$, there is a map $\tilde{f} : B \rightarrow I$ with $\tilde{f}j = f$. Equivalently, $\text{Mor}_R(-, I)$ takes short exact sequences to short exact sequences of abelian groups with the arrows reversed. The extension property supplies the surjection at the right; exactness at the other two terms follows from the kernel universal property.

Definition 2.3 (Enough Projectives and Enough Injectives). An abelian category has enough projectives if every object admits an epimorphism from a projective object. It has enough injectives if every object admits a monomorphism into an injective object. Module categories have both properties. Free modules supply the first; the construction below supplies the second.

Proposition 2.4 (Embedding in an Injective Module). Every left module over a unital ring embeds in an injective left module.

Proof of 2.4. First, a divisible abelian group D is injective. Given a map from a subgroup A of an abelian group B , use Zorn's lemma to choose a maximal extension to a subgroup C . If $b \notin C$, the set of integers n with $nb \in C$ is an ideal. If it is zero, choose any image of b . Otherwise it is $d\mathbb{Z}$ for $d > 0$; choose $y \in D$ with $dy = f(db)$ and set $f(c + nb) = f(c) + ny$. The ideal description proves this is well defined. Both cases extend the map to $C + \mathbb{Z}b$, contradicting maximality. Hence $C = B$.

The group $\mathbb{Q}/\mathbb{Z}$ is divisible, and its characters separate points of any abelian group M : a nonzero element has a cyclic subgroup admitting a character nonzero on that element, which extends by injectivity. Therefore M embeds by evaluation into $D = (\mathbb{Q}/\mathbb{Z})^{\text{Mor}_{\mathbb{Z}}(M, \mathbb{Q}/\mathbb{Z})}$. This product is injective because each component of a map extends.

For a left R -module M , take such an embedding $j : M \hookrightarrow D$ of underlying abelian groups. Put $E = \text{Mor}_{\mathbb{Z}}(R, D)$ and define $(r\phi)(s) = \phi(sr)$. The bijection

$$\text{Mor}_R(X, E) \cong \text{Mor}_{\mathbb{Z}}(X, D), \quad f \mapsto (x \mapsto f(x)(1)),$$

has inverse $g \mapsto (x \mapsto (s \mapsto g(sx)))$. Since forgetting the R -action is exact and D is injective, E is injective. The map $M \rightarrow E$ given by $m \mapsto (s \mapsto j(sm))$ is R -linear and injective, as evaluation at 1 recovers j . $\square$

Example 2.5 (Projective and Injective Are Different). The abelian group $\mathbb{Z}$ is free and therefore projective. It is not injective: the map $2\mathbb{Z} \rightarrow \mathbb{Z}$ sending 2 to 1 cannot extend to $\mathbb{Z}$. Both $\mathbb{Q}$ and $\mathbb{Q}/\mathbb{Z}$ are injective by divisibility. The nonzero group $\mathbb{Q}/\mathbb{Z}$ is not projective: projective abelian groups are summands of free abelian groups and hence torsion-free, whereas $\mathbb{Q}/\mathbb{Z}$ is torsion. Thus

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Q} \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow 0$$

is an injective resolution of $\mathbb{Z}$, with $\mathbb{Q}$ in cochain degree zero.

Definition 2.6 (Projective Resolution). A projective resolution of M is an exact augmented sequence

$$\cdots \rightarrow P_2 \xrightarrow{d_2} P_1 \xrightarrow{d_1} P_0 \xrightarrow{\epsilon} M \rightarrow 0 \quad (3)$$

with every P_i projective. The associated chain complex $P_\bullet$ has $P_i = 0$ for $i < 0$, differential $d_0 = 0$, $H_0(P_\bullet) \cong M$, and $H_i(P_\bullet) = 0$ for $i > 0$. The augmentation is not part of the unaugmented complex.

Every module has a free resolution. Choose a free surjection $P_0 \rightarrow M$; choose a free surjection $P_1 \rightarrow \ker \epsilon$; choose a free surjection $P_2 \rightarrow \ker d_1$; and continue. The successive kernels are called syzygies. For a completely general choice, use a free module on the underlying set of each kernel. The process may require infinite rank and may continue indefinitely.

Remark 2.7 (A Presentation Is Only the Beginning). A presentation supplies $d_1 : P_1 \rightarrow P_0$, but does not assert that d_1 is injective. To compute degree-one derived functors one generally also needs d_2 , which records the relations among the original relations. To compute degree i , one needs the resolution through $P_{i+1} \rightarrow P_i \rightarrow P_{i-1}$.

2.2 Why the Choice of Resolution Does Not Matter

Theorem 2.8 (Comparison of Resolutions). Let $P_\bullet \rightarrow M$ and $Q_\bullet \rightarrow L$ be projective resolutions. Every map $f : M \rightarrow L$ lifts to a chain map $\tilde{f} : P_\bullet \rightarrow Q_\bullet$. Any two lifts of the same f are chain homotopic. In particular, any two projective resolutions of M are chain homotopy equivalent by maps lifting id_M .

Proof of 2.8. Lift $f \epsilon_P$ through $Q_0 \rightarrow L$ using projectivity of P_0 . Having constructed the lift through degree $n-1$, the map $\tilde{f}_{n-1} d_n$ lands in the kernel of the next differential (or of the augmentation when $n=1$). Exactness identifies this kernel with the image of Q_n . Projectivity of P_n gives $\tilde{f}_n$.

For two lifts u, v , construct $h_n : P_n \rightarrow Q_{n+1}$ satisfying

$$u_n - v_n = d_{n+1}^Q h_n + h_{n-1} d_n^P, \quad h_{-1} = 0.$$

In degree zero, $u_0 - v_0$ is killed by the augmentation, so lift it through d_1^Q . Inductively, the chain-map identities and the formula already established in degree $n-1$ show that

$$d_n^Q(u_n - v_n - h_{n-1} d_n^P) = 0.$$

Exactness and projectivity give the required h_n . Finally, lifts of id_M in both directions have composites homotopic to the appropriate identities, by the uniqueness statement. $\square$

Used in: Definitions 3.1 and 5.1.

2.3 Left and Right Derived Functors

Let $\mathcal{A}, \mathcal{B}$ be abelian categories and let all functors in this subsection be additive. The constructions apply to objects of $\mathcal{A}$; complexes of objects require a further construction.

Definition 2.9 (Derived Functors on Objects). If $\mathcal{A}$ has enough projectives and $F : \mathcal{A} \rightarrow \mathcal{B}$ is right exact, choose a projective resolution $P_\bullet \rightarrow M$ and set

$$L_n F(M) = H_n(F(P_\bullet)), \quad n \geq 0.$$

If $\mathcal{A}$ has enough injectives and $G : \mathcal{A} \rightarrow \mathcal{B}$ is left exact, construct an exact sequence $0 \rightarrow M \rightarrow I^0 \rightarrow I^1 \rightarrow \cdots$ with each I^q injective, by successively embedding the cokernels. Set

$$R^n G(M) = H^n(G(I^\bullet)), \quad n \geq 0.$$

For a cochain complex, H^n means the kernel of the differential out of degree n modulo the image of the differential into degree n .

Proposition 2.10 (Independence, Degree Zero, and Vanishing). These constructions give functors independent of the resolutions up to canonical natural isomorphism. There are natural isomorphisms $L_0 F \cong F$ and $R^0 G \cong G$. For $n > 0$, $L_n F(P) = 0$ for projective P and $R^n G(I) = 0$ for injective I .

Proof of 2.10. The comparison theorem and its dual lift maps of objects to maps of resolutions uniquely up to chain or cochain homotopy. Additive functors preserve the homotopy equations, so their induced homology maps are unique and respect identities and composition. Comparing identity lifts between two resolutions gives mutually inverse isomorphisms; uniqueness shows their compatibility with a third resolution. Right exactness identifies the cokernel of $F(P_1) \rightarrow F(P_0)$ with $F(M)$. Left exactness identifies the kernel of $G(I^0) \rightarrow G(I^1)$ with $G(M)$. Finally use the resolution concentrated in degree zero when the object is itself projective or injective, respectively. $\square$

Remark 2.11 (Long Exact Sequences and Dimension Shifting). The horseshoe construction resolves a short exact sequence by a degreewise split short exact sequence of projective resolutions. Additivity preserves each splitting, so applying F and taking homology gives

$$\cdots \rightarrow L_1 F(M) \rightarrow L_1 F(M'') \rightarrow F(M') \rightarrow F(M) \rightarrow F(M'') \rightarrow 0.$$

The dual construction with injectives gives

$$0 \rightarrow G(M') \rightarrow G(M) \rightarrow G(M'') \rightarrow R^1 G(M') \rightarrow R^1 G(M) \rightarrow \cdots.$$

In particular, an exact sequence $0 \rightarrow K \rightarrow P \rightarrow M \rightarrow 0$ with P projective gives $L_{n+1} F(M) \cong L_n F(K)$ for $n \geq 1$; an exact sequence $0 \rightarrow M \rightarrow I \rightarrow C \rightarrow 0$ with I injective gives $R^{n+1} G(M) \cong R^n G(C)$ for $n \geq 1$. These are the dimension-shifting isomorphisms.

3 Tor and the Failure of Exactness

3.1 Definition and a Computation Procedure

Again let R be commutative. Tensoring the injection $2 : \mathbb{Z} \rightarrow \mathbb{Z}$ with $\mathbb{Z}/2\mathbb{Z}$ gives the zero map $\mathbb{Z}/2\mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z}$, which is not injective. Tensor products preserve cokernels but can fail to preserve kernels.

Definition 3.1 (Tor). Choose a projective resolution $P_\bullet \rightarrow M$. Define

$$\mathrm{Tor}_i^R(M, N) = H_i(P_\bullet \otimes_R N) = \frac{\ker(d_i \otimes \mathrm{id}_N)}{\mathrm{im}(d_{i+1} \otimes \mathrm{id}_N)} \quad (i \geq 0),$$

where the differential out of degree zero is zero. These are R -modules. Right exactness gives

$$\mathrm{Tor}_0^R(M, N) \cong M \otimes_R N.$$

The comparison theorem makes these groups independent of the resolution up to canonical isomorphism: a chain homotopy remains a chain homotopy after tensoring. Lifting maps of M gives functoriality in the first variable; tensoring maps of N gives functoriality in the second. Thus Tor is covariant in both variables.

For a free resolution $P_i = \bigoplus_{j \in I_i} R$, the computation is completely explicit:

$$\cdots \rightarrow \bigoplus_{j \in I_2} N \xrightarrow{(d_2)_N} \bigoplus_{j \in I_1} N \xrightarrow{(d_1)_N} \bigoplus_{j \in I_0} N \rightarrow 0.$$

Use the same coefficient matrices as in the free resolution, now acting on direct sums of N ; take kernels modulo images. A presentation by itself computes $\text{Tor}_0^R(M, N)$, while $\text{Tor}_1^R(M, N)$ generally requires the next map.

Example 3.2 (A Non-Zero-Divisor). Suppose multiplication by $a \in R$ is injective on R . Then $0 \rightarrow R \xrightarrow{a} R \rightarrow R/(a) \rightarrow 0$ is a free resolution. Writing $N[a] = \{n \in N : an = 0\}$, we obtain

$$\text{Tor}_0^R(R/(a), N) = N/aN, \quad \text{Tor}_1^R(R/(a), N) = N[a], \quad \text{Tor}_i^R(R/(a), N) = 0 \quad (i \geq 2).$$

In particular, for $a, b > 0$, $\text{Tor}_1^{\mathbb{Z}}(\mathbb{Z}/a\mathbb{Z}, \mathbb{Z}/b\mathbb{Z}) \cong \mathbb{Z}/\gcd(a, b)\mathbb{Z}$. The kernel description is natural; an identification with a standard cyclic group involves a choice of generator.

Example 3.3 (Continuing the Matrix Computation). For the matrix in Example 1.7, $\det A = 12 \neq 0$, so $A : \mathbb{Z}^2 \rightarrow \mathbb{Z}^2$ is injective. It therefore gives a length-one free resolution of M . Modulo four, A acts by $(x, y) \mapsto (2x, 2y)$, whose kernel is $\{0, 2\}^2$. Thus

$$\text{Tor}_1^{\mathbb{Z}}(M, \mathbb{Z}/4\mathbb{Z}) \cong (\mathbb{Z}/2\mathbb{Z})^2, \quad \text{Tor}_i^{\mathbb{Z}}(M, \mathbb{Z}/4\mathbb{Z}) = 0 \quad (i \geq 2).$$

Indeed, $A(x, y) = 0$ over $\mathbb{Z}$ implies $2x = 0$, hence $x = 0$, and then $6y = 0$, hence $y = 0$. The tensor complex is $0 \rightarrow (\mathbb{Z}/4\mathbb{Z})^2 \xrightarrow{(2x, 2y)} (\mathbb{Z}/4\mathbb{Z})^2 \rightarrow 0$, in degrees one and zero. A degree-one cycle has each coordinate either zero or two. There is no degree-two term and hence no nonzero degree-one boundary. The map $(\bar{a}, \bar{b}) \mapsto (2a, 2b)$ identifies $(\mathbb{Z}/2\mathbb{Z})^2$ with this kernel. In degree zero the cokernel was computed in Example 1.7.

Example 3.4 (A Resolution That Does Not Terminate). Let $R = k[\varepsilon]/(\varepsilon^2)$ for a field k , and let $K = R/(\varepsilon)$. Since the kernel and image of multiplication by ε are both (ε) ,

$$\cdots \xrightarrow{\varepsilon} R \xrightarrow{\varepsilon} R \xrightarrow{\varepsilon} R \rightarrow K \rightarrow 0$$

is a free resolution. To check this directly, write an element of R uniquely as $a + b\varepsilon$. Multiplication by ε sends it to $a\varepsilon$, so its kernel and image are both $k\varepsilon$. The augmentation has the same kernel. Since $\varepsilon K = 0$, tensoring gives one copy of k in each nonnegative degree and zero differential. Every vector is a cycle and the only boundary is zero, so $\text{Tor}_i^R(K, K) \cong k$ for every $i \geq 0$. The non-zero-divisor hypothesis in Example 3.2 cannot be dropped.

3.2 Flatness and Long Exact Sequences

Definition 3.5 (Flat Module). A module F is flat if $- \otimes_R F$ is exact. Since tensoring is right exact, this means that it preserves injections. Free modules are flat because tensoring with a free module is a direct sum of copies of the original sequence; direct sums of exact sequences of modules are exact. Direct summands of flat modules are flat, so projective modules are flat.

Theorem 3.6 (Either Variable Can Be Resolved). If $P_\bullet \rightarrow M$ and $Q_\bullet \rightarrow N$ are projective resolutions, there are natural isomorphisms

$$H_i(P_\bullet \otimes_R N) \cong H_i(M \otimes_R Q_\bullet).$$

Consequently $\text{Tor}_i^R(M, N) \cong \text{Tor}_i^R(N, M)$ naturally.

Proof of 3.6. Form the first-quadrant double complex $P_p \otimes_R Q_q$ and its total complex

$$T_n = \bigoplus_{p+q=n} P_p \otimes_R Q_q, \quad d(x \otimes y) = d_P x \otimes y + (-1)^p x \otimes d_Q y.$$

The sign makes the two mixed terms in d^2 cancel. Since P_p is flat, each augmented column is exact, with degree-zero homology $P_p \otimes_R N$. Similarly, each augmented row is exact, with degree-zero homology $M \otimes_R Q_q$.

The two augmentations $T \rightarrow P_\bullet \otimes_R N$ and $T \rightarrow M \otimes_R Q_\bullet$ are quasi-isomorphisms. One elementary way to see this is to take the augmented double complex for either map: its columns, respectively rows, are exact. In a total cycle, remove the component with largest column, respectively row, index by writing it as a boundary in that exact column or row. Subtract its total boundary and repeat. Each subtraction introduces terms only at smaller indices. There are finitely many components on each total diagonal, so this procedure terminates and proves that the augmented total complex is acyclic. Taking homology gives the first assertion. The factor-interchange isomorphism then gives symmetry. $\square$

Used in: Theorem 3.7.

Theorem 3.7 (Exact Sequences and the Flatness Criterion). For a short exact sequence $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$, there is a natural long exact sequence

$$\begin{aligned} \cdots \rightarrow \operatorname{Tor}_i^R(M, N') \rightarrow \operatorname{Tor}_i^R(M, N) \rightarrow \operatorname{Tor}_i^R(M, N'') \xrightarrow{\partial} \operatorname{Tor}_{i-1}^R(M, N') \rightarrow \cdots \\ \cdots \rightarrow \operatorname{Tor}_1^R(M, N'') \xrightarrow{\partial} M \otimes_R N' \rightarrow M \otimes_R N \rightarrow M \otimes_R N'' \rightarrow 0. \end{aligned}$$

There is also a long exact sequence in the first variable. A module F is flat if and only if $\operatorname{Tor}_1^R(M, F) = 0$ for every M ; in that case all $\operatorname{Tor}_i^R(M, F)$ vanish for $i > 0$.

Proof of 3.7. Each term of a projective resolution of M is flat, so tensoring the short exact sequence gives a degreewise short exact sequence of chain complexes. Its homology long exact sequence is the displayed one. Theorem 3.6 gives the sequence in the first variable.

If F is flat, tensoring the augmented resolution of M preserves exactness, so all positive homology vanishes. Conversely, apply the long exact sequence in the first variable to $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$. Vanishing of $\operatorname{Tor}_1^R(C, F)$ implies that $A \otimes_R F \rightarrow B \otimes_R F$ is injective. This is flatness. $\square$

Remark 3.8 (The Precise Obstruction). For a particular short exact sequence, the kernel of $M \otimes_R N' \rightarrow M \otimes_R N$ is the image of the connecting map from $\operatorname{Tor}_1^R(M, N'')$. That connecting map need not be injective. Thus the kernel is a quotient of this Tor group, not generally the whole group.

4 Cochain Complexes over an Arbitrary Ring

4.1 Cohomological Indexing

From now on, R is an arbitrary unital ring and all modules are left R -modules. A cochain complex uses an increasing differential. No commutativity is needed for its definition or for the usual exact-sequence arguments in R -modules.

Definition 4.1 (Cochain Complex and Cohomology). A cochain complex $C^\bullet$ consists of maps $d_C^n : C^n \rightarrow C^{n+1}$ with $d_C^{n+1} d_C^n = 0$. Its cocycles, coboundaries, and cohomology are

$$Z^n(C) = \ker d_C^n, \quad B^n(C) = \operatorname{im} d_C^{n-1}, \quad H^n(C) = Z^n(C)/B^n(C).$$

A cochain map $f : C \rightarrow D$ satisfies $d_D^n f^n = f^{n+1} d_C^n$. It induces a map on cohomology. It is a quasi-isomorphism when those maps are isomorphisms for all n .

Reindexing a chain complex by $C^n = C_{-n}$ and $d^n = d_{-n}$ gives a cochain complex, with $H^n(C^\bullet) = H_{-n}(C_\bullet)$. By contrast, applying the contravariant morphism functor to a chain complex reverses its arrows and gives

cochains in nonnegative degrees; that is the construction used for Ext.

Definition 4.2 (Cochain Homotopy, Shift, and Cone). A cochain homotopy from g to f is a family $h^n : C^n \rightarrow D^{n-1}$ with

$$f^n - g^n = d_D^{n-1} h^n + h^{n+1} d_C^n.$$

On cocycles the difference is a coboundary, so homotopic maps induce the same cohomology maps. The shift is $C[r]^n = C^{n+r}$ with differential $(-1)^r d_C^{n+r}$. In particular, $H^n(C[r]) = H^{n+r}(C)$.

For a cochain map $f : C \rightarrow D$, its cone is

$$\text{Cone}(f)^n = D^n \oplus C^{n+1}, \quad d^n = \begin{pmatrix} d_D^n & f^{n+1} \\ 0 & -d_C^{n+1} \end{pmatrix}.$$

The chain-map equation makes this differential square to zero. Inclusion of the first summand and projection onto the second give $0 \rightarrow D \rightarrow \text{Cone}(f) \rightarrow C[1] \rightarrow 0$.

Proposition 4.3 (The Cohomology Long Exact Sequence). A degreewise short exact sequence $0 \rightarrow A \xrightarrow{\iota} B \xrightarrow{\rho} C \rightarrow 0$ of cochain complexes gives

$$\cdots \rightarrow H^n(A) \rightarrow H^n(B) \rightarrow H^n(C) \xrightarrow{\partial} H^{n+1}(A) \rightarrow \cdots.$$

For a cocycle $c \in C^n$, choose $b \in B^n$ with $\rho b = c$ and write $d_B b = \iota a$. Then $\partial[c] = [a]$.

Proof of 4.3. The equation $\rho d_B b = d_C c = 0$ gives a , and $\iota d_A a = d_B^2 b = 0$ makes it a cocycle. Changing b by ιx changes a by $d_A x$; changing c by a coboundary and lifting its primitive does not change the resulting cohomology class. The usual kernel–image chase, or the snake lemma on kernels and cokernels of consecutive differentials, gives exactness. Equivalently, reindex all three complexes as chain complexes and apply the homology long exact sequence; its connecting map is exactly this lift-and-differentiate formula. $\square$

Used in: Proposition 5.2.

4.2 Applying the Morphism Functor

Definition 4.4 (The Cochain Complex Associated to a Resolution). For a chain complex $P_\bullet$ in nonnegative degrees and a module N , put

$$\text{Mor}_R(P_\bullet, N)^n = \text{Mor}_R(P_n, N), \quad \delta^n(\phi) = \phi \circ d_{n+1}.$$

Thus the arrows run as follows:

$$0 \rightarrow \text{Mor}_R(P_0, N) \xrightarrow{\delta^0} \text{Mor}_R(P_1, N) \xrightarrow{\delta^1} \text{Mor}_R(P_2, N) \rightarrow \cdots.$$

The relation $\delta^{n+1} \delta^n = 0$ follows from $d_{n+1} d_{n+2} = 0$. For general R , this is a cochain complex of abelian groups, canonically of modules over the center $Z(R)$. It is not generally a complex of left R -modules: multiplying the values of an R -linear map by a noncentral element need not give an R -linear map.

Remark 4.5 (The General Morphism Complex and Its Sign). For cochain complexes C, D , define the abelian-group complex

$$\mathcal{M}^n(C, D) = \prod_p \text{Mor}_R(C^p, D^{p+n}), \quad (\delta f)^p = d_D^{p+n} f^p - (-1)^n f^{p+1} d_C^p.$$

Its degree-zero cocycles are cochain maps, and its degree-zero coboundaries are nullhomotopic maps. If $C^{-n} = P_n$ and $D = N$ is concentrated in degree zero, this convention gives $\delta \phi = (-1)^{n+1} \phi d_{n+1}$. Multiplication in degree n by $(-1)^{n(n+1)/2}$ identifies the unsigned complex in Definition 4.4 with this signed convention. We use the unsigned complex for the Ext computations below.

5 Ext and a Noncommutative Computation

5.1 Definition, Variance, and Computation

The ring R remains arbitrary. Projective resolutions of left R -modules exist by Section 2.

Definition 5.1 (Ext). For left modules M, N , choose a projective resolution $P_\bullet \rightarrow M$ and define

$$\text{Ext}_R^n(M, N) = H^n(\text{Mor}_R(P_\bullet, N)) = \frac{\ker(\delta^n)}{\text{im}(\delta^{n-1})} \quad (n \geq 0),$$

where the image in degree zero is zero. These are abelian groups with a natural $Z(R)$ -module structure; if R is an algebra over a field k , they are k -vector spaces. When R is commutative, they are naturally R -modules.

The comparison theorem again proves independence of resolutions. More explicitly, if $u, v : P_\bullet \rightarrow Q_\bullet$ satisfy $u - v = d_Q h + h d_P$, precomposition gives cochain-homotopic maps: on $\phi : Q_n \rightarrow N$ use the homotopy $\phi \mapsto \phi h_{n-1}$. Thus Ext is contravariant in M and covariant in N . In degree zero, a map $P_0 \rightarrow N$ lies in $\ker \delta^0$ exactly when it kills $\text{im } d_1$, so it factors through M and

$$\text{Ext}_R^0(M, N) \cong \text{Mor}_R(M, N).$$

For an arbitrary free resolution $P_n = \bigoplus_{i \in I_n} R$, note the essential change from tensor products:

$$\text{Mor}_R\left(\bigoplus_{i \in I_n} R, N\right) \cong \prod_{i \in I_n} N.$$

Maps out of a free direct sum may have arbitrary values on its basis, so this is a product, not a direct sum. If $d_{n+1}(e_j) = \sum_i r_{ij} e_i$ with coefficients on the left, then

$$\delta^n((n_i)_i) = \left(\sum_i r_{ij} n_i \right)_j.$$

Every sum is finite because each column of the differential has finite support. Compute Ext by taking the kernel of this map modulo the image of the preceding map. Over a commutative ring with finite free terms, these are the transposed differential matrices acting on finite powers of N . For a noncommutative ring, the displayed order of the coefficients specifies the maps unambiguously.

Proposition 5.2 (Exactness and Projectivity). For $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$, there is a natural long exact sequence

$$\begin{aligned} 0 \rightarrow \text{Mor}_R(M, N') \rightarrow \text{Mor}_R(M, N) \rightarrow \text{Mor}_R(M, N'') \\ \xrightarrow{\partial} \text{Ext}_R^1(M, N') \rightarrow \text{Ext}_R^1(M, N) \rightarrow \text{Ext}_R^1(M, N'') \xrightarrow{\partial} \text{Ext}_R^2(M, N') \rightarrow \dots \end{aligned}$$

If P is projective, $\text{Ext}_R^n(P, N) = 0$ for $n > 0$. Conversely, if $\text{Ext}_R^1(M, N) = 0$ for every N , then M is projective.

Proof of 5.2. Apply $\text{Mor}_R(P_i, -)$ to the short exact sequence. Each resulting sequence is exact by projectivity, so Proposition 4.3, applied in abelian groups, gives the displayed sequence. A projective P has a resolution concentrated in degree zero, proving its higher vanishing. For the converse, take a free surjection $F \rightarrow M$ with kernel K . The exact sequence makes $\text{Mor}_R(M, F) \rightarrow \text{Mor}_R(M, M)$ surjective because $\text{Ext}_R^1(M, K) = 0$. A preimage of id_M splits $F \rightarrow M$, so M is projective. $\square$

There is also a long exact sequence in the first, contravariant variable. For $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$, its beginning is

$$\begin{aligned} 0 \rightarrow \text{Mor}_R(M'', N) \rightarrow \text{Mor}_R(M, N) \rightarrow \text{Mor}_R(M', N) \\ \rightarrow \text{Ext}_R^1(M'', N) \rightarrow \text{Ext}_R^1(M, N) \rightarrow \text{Ext}_R^1(M', N) \rightarrow \dots \end{aligned}$$

One constructs compatible projective resolutions with $P_i = P'_i \oplus P''_i$ by successively lifting the maps from P''_i ; exactness of the sequence of successive kernels permits the next lift. This is the horseshoe construction. Applying $\text{Mor}_R(-, N)$ gives a degree-wise split short exact sequence of cochain complexes in the reverse order, and hence the displayed sequence.

5.2 Extensions as Degree-One Classes

Theorem 5.3 (The Meaning of Ext in Degree One). The group $\text{Ext}_R^1(M, N)$ classifies short exact sequences of left R -modules

$$0 \rightarrow N \xrightarrow{\iota} E \xrightarrow{\rho} M \rightarrow 0 \quad (4)$$

up to isomorphisms of sequences that are the identity on N and M . The zero class consists precisely of the split extensions.

Proof of 5.3. Choose $0 \rightarrow K \xrightarrow{j} P_0 \rightarrow M \rightarrow 0$ with P_0 projective. Extend it to a projective resolution. A map $P_1 \rightarrow N$ is a degree-one cocycle exactly when it kills $\text{im } d_2 = \ker(P_1 \rightarrow K)$, hence factors through K . Degree-one coboundaries are restrictions of maps $P_0 \rightarrow N$. Therefore

$$\text{Ext}_R^1(M, N) \cong \text{coker}(\text{Mor}_R(P_0, N) \xrightarrow{j^*} \text{Mor}_R(K, N)). \quad (5)$$

Given $\alpha : K \rightarrow N$, define

$$E_\alpha = (N \oplus P_0) / \{(\alpha(k), -j(k)) : k \in K\}.$$

The maps $n \mapsto [(n, 0)]$ and $[(n, p)] \mapsto \epsilon(p)$ give an extension: the first is injective because j is injective, and its image is the kernel of the second. If $\alpha' = \alpha + \beta j$, the map $(n, p) \mapsto (n - \beta(p), p)$ descends to an isomorphism $E_\alpha \rightarrow E_{\alpha'}$ of extensions.

Conversely, lift $P_0 \rightarrow M$ through $E \rightarrow M$. The restriction of a lift s to K lands in N , giving a map α with $sj = \iota\alpha$. The map $(n, p) \mapsto \iota(n) + s(p)$ identifies E_α with E . Changing s changes α by a restriction from P_0 . These constructions are inverse on equivalence classes. Finally, $E_0 \cong N \oplus M$, and a splitting of an extension allows a lift whose restriction to K is zero. Thus its class vanishes exactly when it splits. $\square$

Used in: Example 5.4.

The addition transported by this classification is the Baer sum: take the direct sum of two extensions, pull back along the diagonal $M \rightarrow M \oplus M$, and push out along addition $N \oplus N \rightarrow N$. In the description above, these operations replace the two maps $\alpha, \beta : K \rightarrow N$ by $\alpha + \beta$.

5.3 An Explicit Noncommutative Ring

Example 5.4 (Upper-Triangular Matrices). Let k be a field and

$$R = \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} : a, b, c \in k \right\}, \quad e_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad u = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

This ring is noncommutative: $e_1 u = u$ whereas $u e_1 = 0$. Define left modules $S_1 = S_2 = k$ as vector spaces, with actions

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \cdot x = ax \quad \text{on } S_1, \quad \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \cdot x = cx \quad \text{on } S_2.$$

The left ideals $P_1 = R e_1$ and $P_2 = R e_2$ are projective because $R = R e_1 \oplus R e_2$. They have bases e_1 and u, e_2 , respectively. There is a projective resolution

$$0 \rightarrow P_1 \xrightarrow{d} P_2 \xrightarrow{\rho} S_2 \rightarrow 0, \quad d(x) = xu, \quad \rho(bu + ce_2) = c. \quad (6)$$

Right multiplication by u is a map of left modules. It sends e_1 to u , so it is injective with image $ku = \ker \rho$.

For any left R -module N , evaluation at e_i gives

$$\text{Mor}_R(Re_i, N) \cong e_i N.$$

Indeed, $f(e_i) = e_i f(e_i)$, and conversely $n \in e_i N$ determines $f_n(re_i) = rn$. Under these identifications, precomposition with d is

$$e_2 N \rightarrow e_1 N, \quad n \mapsto un,$$

because $d(e_1) = u = ue_2$. Thus the entire Ext computation is the two-term cochain complex

$$e_2 N \xrightarrow{u} e_1 N \quad \text{in degrees } 0, 1.$$

Consequently

$$\text{Mor}_R(S_2, N) \cong \ker(u : e_2 N \rightarrow e_1 N),$$

$$\text{Ext}_R^1(S_2, N) \cong e_1 N / u(e_2 N),$$

$$\text{Ext}_R^n(S_2, N) = 0 \quad (n \geq 2).$$

For $N = S_1$, the actions give $e_2 S_1 = 0$ and $e_1 S_1 = k$. Thus the cochain complex is $0 \rightarrow 0 \rightarrow k \rightarrow 0$, in degrees zero and one. Its degree-zero cohomology is zero, and its degree-one cohomology is $k/0 = k$. Hence

$$\text{Ext}_R^1(S_2, S_1) \cong k.$$

For $N = S_2$, the complex instead is $0 \rightarrow k \rightarrow 0 \rightarrow 0$, so $\text{Mor}_R(S_2, S_2) = k$ and $\text{Ext}_R^1(S_2, S_2) = 0$. For $N = P_2$, we have $e_2 P_2 = ke_2$ and $e_1 P_2 = ku$; the differential $ce_2 \mapsto cu$ is an isomorphism, so both cohomology groups vanish. By contrast, $S_1 \cong P_1$ is projective, so $\text{Ext}_R^1(S_1, S_2) = 0$. Unlike Tor over a commutative ring, Ext is not symmetric in its variables.

The resolution (6) itself, after identifying P_1 with S_1 , is a nonsplit extension of S_2 by S_1 . If it had a section, the image v of $1 \in S_2$ would satisfy $\rho(v) = 1$, so $v = bu + e_2$ for some $b \in k$. But u acts trivially on S_2 , while $uv = u \neq 0$, contradicting linearity of the section.

More explicitly, the class $\lambda \in k$ corresponds to $E_\lambda = kx \oplus ky$ with

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} x = ax, \quad \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} y = b\lambda x + cy.$$

The sequence $0 \rightarrow S_1 \xrightarrow{1 \mapsto x} E_\lambda \xrightarrow{x \mapsto 0, y \mapsto 1} S_2 \rightarrow 0$ has class λ under (5). It splits exactly when $\lambda = 0$; $E_1 \cong P_2$. Nonzero values can give isomorphic middle modules while representing different extension classes, because equivalence of extensions must fix both end modules.

5.4 Cyclic Groups: Every Kernel and Cokernel

Example 5.5 (The Pair $\mathbb{Z}/12\mathbb{Z}$ and $\mathbb{Z}/8\mathbb{Z}$). The free resolution $0 \rightarrow \mathbb{Z} \xrightarrow{12} \mathbb{Z} \rightarrow \mathbb{Z}/12\mathbb{Z} \rightarrow 0$ is exact because multiplication by 12 is injective and its cokernel is $\mathbb{Z}/12\mathbb{Z}$. Set $N = \mathbb{Z}/8\mathbb{Z}$. Tensoring gives

$$0 \rightarrow N \xrightarrow{4} N \rightarrow 0 \quad \text{in chain degrees } 1, 0.$$

The image is $\{0, 4\}$, and the kernel is $\{0, 2, 4, 6\}$. The quotient map $N \rightarrow \mathbb{Z}/4\mathbb{Z}$ has kernel $\{0, 4\}$; hence $\mathbb{Z}/12\mathbb{Z} \otimes_{\mathbb{Z}} N \cong \mathbb{Z}/4\mathbb{Z}$. The map $\mathbb{Z}/4\mathbb{Z} \rightarrow N$, $[t]_4 \mapsto [2t]_8$, is injective with image $\{0, 2, 4, 6\}$. There is no incoming degree-two boundary, so it identifies $\text{Tor}_1^{\mathbb{Z}}(\mathbb{Z}/12\mathbb{Z}, N)$ with $\mathbb{Z}/4\mathbb{Z}$. Higher Tor vanishes.

Applying $\text{Mor}_{\mathbb{Z}}(-, N)$ reverses the arrows and gives

$$0 \rightarrow N \xrightarrow{4} N \rightarrow 0 \quad \text{in cochain degrees } 0, 1.$$

Thus $\text{Mor}_{\mathbb{Z}}(\mathbb{Z}/12\mathbb{Z}, N) = \{0, 2, 4, 6\}$, while $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}/12\mathbb{Z}, N) = N/\{0, 4\} \cong \mathbb{Z}/4\mathbb{Z}$. Higher Ext vanishes. The same map is being used, but homological and cohomological grading put its kernel and cokernel in different degrees.

For general $a, b > 0$, put $g = \gcd(a, b)$, $a = ga'$, and $b = gb'$. The image of multiplication by a on $\mathbb{Z}/b\mathbb{Z}$ equals the subgroup generated by g , by Bézout's identity. Its cokernel is therefore $\mathbb{Z}/g\mathbb{Z}$. For the kernel, $b \mid ax$ is equivalent to $b' \mid x$, since $\gcd(a', b') = 1$. Its elements are the g multiples of b' modulo b . The map $[t]_g \mapsto [b't]_b$ gives its cyclic-group identification. This computes tensor, Tor, morphism groups, and Ext in the preceding manner.

Example 5.6 (Ext for the Two-Generator Presentation). For $M = \text{coker} \begin{pmatrix} 2 & 0 \\ 4 & 6 \end{pmatrix}$ and $N = \mathbb{Z}/4\mathbb{Z}$, a homomorphism $\mathbb{Z}^2 \rightarrow N$ is specified by a row (x, y) . Precomposition with A sends it to $(2x + 4y, 6y)$. Equivalently, on column coordinates the cochain differential is A^T . Modulo four this is $(x, y) \mapsto (2x, 2y)$. Its kernel is $\{0, 2\}^2$, and its image is also $\{0, 2\}^2$ in the next copy of N^2 . Consequently

$$\text{Mor}_{\mathbb{Z}}(M, N) \cong (\mathbb{Z}/2\mathbb{Z})^2, \quad \text{Ext}_{\mathbb{Z}}^1(M, N) \cong (\mathbb{Z}/2\mathbb{Z})^2,$$

and all higher groups vanish. The map defining Ext is the transposed map, even though in this particular example reduction modulo four makes it coincide with the map used for Tor.

5.5 Computing Ext with Injective Resolutions

For a fixed left R -module M , the functor $\text{Mor}_R(M, -)$ is left exact. Its right derived functors are the Ext groups defined above.

Proposition 5.7 (The Two Resolutions Compute the Same Ext). If $P_{\bullet} \rightarrow M$ is a projective resolution and $0 \rightarrow N \rightarrow I^{\bullet}$ is an injective resolution, there are natural isomorphisms

$$H^n \text{Mor}_R(P_{\bullet}, N) \cong H^n \text{Mor}_R(M, I^{\bullet}) = R^n \text{Mor}_R(M, -)(N).$$

Proof of 5.7. Use the first-quadrant double complex $C^{p,q} = \text{Mor}_R(P_p, I^q)$. Its horizontal map is precomposition with d_{p+1} , and its vertical map is postcomposition with the differential of I . These maps commute; on the total complex take $D = h + (-1)^p v$. Thus $D^2 = 0$. The total degree- n group is $\bigoplus_{p+q=n} C^{p,q}$, a finite sum.

For fixed p , exactness of $\text{Mor}_R(P_p, -)$ makes the augmented column $0 \rightarrow \text{Mor}_R(P_p, N) \rightarrow C^{p,0} \rightarrow C^{p,1} \rightarrow \dots$ exact. For fixed q , injectivity of I^q makes the augmented row $0 \rightarrow \text{Mor}_R(M, I^q) \rightarrow C^{0,q} \rightarrow C^{1,q} \rightarrow \dots$ exact. The augmentation maps from both single complexes to the total complex are quasi-isomorphisms. Here is the elementary total-complex argument: in a total cocycle, eliminate its component with largest positive vertical degree using exactness of the corresponding column, subtracting the total coboundary of a preimage. Repeating leaves only vertical degree zero, where the kernel is the column augmentation. The same elimination applied to a total primitive shows that a class from the augmentation is a boundary only if it already was one in the single complex. There are finitely many components in each total degree, so this process terminates. Interchanging rows and columns proves the other augmentation is a quasi-isomorphism. This also proves naturality by the comparison maps of resolutions. $\square$

Example 5.8 (An Injective Calculation over $\mathbb{Z}$). To compute $\text{Ext}_{\mathbb{Z}}^n(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z})$ for $m \geq 2$, use $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Q} \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow 0$ from Example 2.5. A homomorphism $\mathbb{Z}/m\mathbb{Z} \rightarrow \mathbb{Q}$ must send 1 to an element killed by m ; the only such rational number is zero. A homomorphism to $\mathbb{Q}/\mathbb{Z}$ is instead determined by one of

$$0, \quad \frac{1}{m} + \mathbb{Z}, \quad \frac{2}{m} + \mathbb{Z}, \quad \dots, \quad \frac{m-1}{m} + \mathbb{Z}.$$

The resulting cochain complex is $0 \rightarrow 0 \rightarrow (\frac{1}{m}\mathbb{Z})/\mathbb{Z} \rightarrow 0$, with its two terms in degrees zero and one. Hence

$$\text{Mor}_{\mathbb{Z}}(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z}) = 0, \quad \text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z}) \cong \mathbb{Z}/m\mathbb{Z},$$

and higher Ext vanishes. The projective-resolution calculation gives $0 \rightarrow \mathbb{Z} \xrightarrow{m} \mathbb{Z} \rightarrow 0$, with the same kernel zero and cokernel $\mathbb{Z}/m\mathbb{Z}$. This checks the comparison by two distinct resolutions.

5.6 Comparing the Computations

For comparison with Section 3, suppose R is commutative and a is a non-zero-divisor. The same resolution used there now gives

$$0 \rightarrow N \xrightarrow{a} N \rightarrow 0 \quad \text{in cochain degrees } 0, 1.$$

Hence $\text{Mor}_R(R/(a), N) = N[a]$, $\text{Ext}_R^1(R/(a), N) = N/aN$, and higher Ext vanishes. The kernel and cokernel appear in the opposite degrees from the tensor computation. For the dual-number example, applying $\text{Mor}_R(-, K)$ makes every differential zero, so $\text{Ext}_R^n(K, K) \cong k$ for every $n \geq 0$.

The two procedures can be compared directly:

	Tensor and Tor, R commutative	Morphism Groups and Ext, R arbitrary
Resolution	$P_{\bullet} \twoheadrightarrow M$	$P_{\bullet} \twoheadrightarrow M$
Complex	$P_{\bullet} \otimes_R N$	$\text{Mor}_R(P_{\bullet}, N)$
Free term $\bigoplus_{i \in I} R$	$\bigoplus_{i \in I} N$	$\prod_I N$
Degree zero	$M \otimes_R N$	$\text{Mor}_R(M, N)$
Higher degrees	$H_i = \text{Tor}_i^R(M, N)$	$H^i = \text{Ext}_R^i(M, N)$
Variance	covariant in both	contravariant, then covariant

In either computation, the central work is to construct enough of a resolution and then compute a kernel modulo an image. A single presentation suffices for a tensor product; its higher relations control Tor and Ext.

References

[Sta26] The Stacks Project Authors, *The Stacks Project*, 2026. Further reading: Tensor Products, Tag 00CV, and Tor Groups and Flatness, Tag 00LY.

[May, n.d.] J. Peter May, *Notes on Tor and Ext*. University of Chicago lecture notes.