

Multimorbidity and Access to Social Care: exploiting emerging administrative data sources in Scotland

David Alexander Gunn Henderson

Submitted in partial fulfilment of the requirements of the degree of ...

Month Year

College of Social Sciences

University of Glasgow

Abstract

Blad de Blah blah blah. I may play about with centering and italicised styles here

Contents

Abstract	i
List of Tables	vi
List of Figures	vii
Acknowledgements	viii
Declaration	ix
1 Introduction	1
1.1 Aims and Objectives	2
1.2 Scientific contribution	2
1.3 Conventions	3
2 Literature Review	4
2.1 Introduction	4
2.2 Access to Social Care	6
2.2.1 Definitions	6
2.2.2 International models of social care	9
2.2.3 Access to Social Care - Social Theory of Eligibility	13
2.2.4 Access to Social Care - Eligibility in the UK	15
2.2.5 Access to Social Care - Social Theory of Resource allocation	19
2.2.6 Acces to Social Care - Resource Allocation in Scotland	22
2.2.7 Health inequalities	24
2.2.8 Summary	26
2.3 Health and Social Care Interaction	27
2.3.1 Public Policy	27
2.3.2 Health and Social Care Integration	27
2.3.3 Research on Health and Social Care Interaction	28
2.3.4 Summary	28

2.4	Multimorbidity	28
2.4.1	Why focus on Multimorbidity?	28
2.4.2	Definitions	28
2.4.3	Measurement	30
2.4.4	Epidemiology	32
2.4.5	Summary	33
2.5	Conclusion	33
3	Methods	35
3.1	Overview	35
3.2	Administrative data research	35
3.2.1	Overview	35
3.2.2	Information Governance	35
3.3	Data Sources	36
3.4	Data linkage	36
3.5	Timeline	36
4	Measuring Multimorbidity	37
4.1	Introduction	37
4.2	Background	38
4.3	Methods	40
4.3.1	Data	40
4.3.2	Analysis	40
4.4	Results	42
4.4.1	Pairwise correlation	42
4.4.2	Grouping diseases	42
4.4.3	Grouping individuals	44
4.4.4	Sociodemographic breakdown of latent groups	48
4.5	Discussion	51
4.5.1	Limitations	53
4.5.2	Future research	54
4.6	Conclusion	54
5	Renfrewshire Council Pilot Project	56
5.1	Introduction	56

5.1.1	Aims	56
5.1.2	Research Questions	56
5.2	Background	57
5.3	Methods	57
5.4	Results	57
5.5	Discussion	57
5.6	Conclusion	57
6	Linked Health and Social Care Data: Quality Assessment	58
6.1	Intro	58
6.2	Background	58
6.3	Methods	58
6.4	Results	58
6.5	Discussion	58
6.6	Conclusion	58
7	Access to social care	59
7.1	Introduction	59
7.1.1	Aims	59
7.1.2	Research questions	59
7.2	Background	60
7.3	Methods	60
7.4	Results	60
7.5	Discussion	60
7.6	Conclusion	60
8	Social care & Health outcomes	61
8.1	Introduction	61
8.1.1	Aims	61
8.1.2	Research Questions	62
8.2	Background	62
8.3	Methods	62
8.4	Results	62
8.5	Discussion	62
8.6	Conclusion	62

Table of Contents

9 Conclusion	63
A Title A	64
B Title B	65
Bibliography	66

List of Tables

2.1	Models of social care in OECD countries adapted from Colombo et al (2011)	10
2.2	Seven stage model of candidacy	15
2.3	Six channels of middle class advantage	20
4.1	Overlapping groups of diseases derived from "clumping" procedure . . .	43
4.2	Non-overlapping groups of diseases derived from Ng algorithm	44

List of Figures

2.1	Older recipients of long-term care services as a share of the over 65 population, 2008	12
4.1	Significant Pairwise Correlations	42
4.2	Item response probabilities for nine latent group model	46
4.3	Latent Group Proportions	47
4.4	Histograms of Age, by Latent Group	48
4.5	Count of Sec, by Latent Group - with Proportions	49
4.6	Count of Carstairs Decile, by Latent Group	50

Acknowledgements

Again maybe play about with centering and layout.

How about a nice quotation at the end???

Declaration

I declare, except where explicit reference is made to the contribution of others, that this thesis is the result of my own work and has not been submitted for any other degree at the University of Glasgow or any other institution.

Printed Name: David Henderson

Signature:

Chapter 1

Introduction

Integration of health and social care became law in Scotland on 1st April 2016. Reflects patterns across the developing world to restructure health services to cope with demands of an ageing population.

Social Care of increasing policy (and political) importance. Link to healthcare (and demands on health services) becoming increasingly apparent (increase delayed discharge etc).

This, in part, due to long-term conditions now major burden of global disease (replacing infectious diseases). Large proportions of population have multimorbidity (OECD) which has a number of negative outcomes including mortality and health care use.

Association of multimorbidity and social care use is unknown.

PhD funding from Scottish Government to assess these topics. (2020 vision and other policy link)

Important part of the funding to link administrative data sources in order to identify the benefit of this process. *Measurement of social (or LTC) is important for a number of reasons - OECD(2013) page 18, Care co-ordination (integration) is not measured well pp 76, administrative databases potential to help these problems plus ideas for outcome measures pp 76 & 79, pp81 obstacles to data collection (overcome by data linkage in Scotland??*

(Need WHO policy outlines and other suitable high-level policy docs in this section)

1.1 Aims and Objectives

The thesis has both substantive and methodological aims. Substantively, it aims to contribute to the debate surrounding health and social care integration by looking specifically at a group that are likely to be regular users of both health and social care services, i.e. those with multimorbidity. Methodologically, the thesis aims to contribute to efforts to improve the exploitation of administrative data as a means to analyse public service performance and effectiveness.

Aims of the project are:-

1. Describe and compare social inequalities in the use of social care services using linked health and social care data.
2. Explore the effects of social care use for those with multimorbidity on
 - (a) unscheduled health care use and
 - (b) mortality.

The objectives of the project are:-

1. To assess how access to social care services varies for people with multimorbidity, especially by socioeconomic status.
2. To assess the impacts of social care service use on health service use and health outcomes for people with multimorbidity, where possible exploiting geographic differences in social care as "natural experiments".
3. To make recommendations for policy on the future of integration of health and social care services based on these results.
4. To assess tho what extant measures of multimorbidity and of health and social care service use can be operationalised using existing linked health and social care administrative data.
5. To make recommendations to policy makers on administrative data collections.

1.2 Scientific contribution

Explicit description of what thesis adds to knowledge

1.3 Conventions

Outline definitions

- Social care refers to Adult social care (with link to subsection [2.2.1](#))
- Multimorbidity and morbidity burden as opposed to comorbidity (with link to subsection [2.4.2](#))

Chapter 2

Literature Review

2.1 Introduction

Introductory paragraph with link from Thesis introduction - also may require explanation of lack of structured lit search

Many countries, including the United Kingdom (UK), have recently seen policies implemented that aim to integrate the provision of health and social care services [Burgess, 2012, Stewart et al., 2003, Glasby, 2017]. In addition to reducing variations in the provision of care across geographic areas, these policies hope to save public money by reducing unplanned admissions and delayed discharges from hospital whilst also improving the quality of services for individuals [Burgess, 2016, Scottish-Government, 2016b, OECD/EU, 2013]. The World Health Organisation [2015] cites relative inequalities in improvements of health and life expectancy, within and between countries, as justification for recommended structural change to healthcare. The main recommendation is for governmental policy to shift healthcare focus from acute hospital care to outpatient and community care [WHO, 2015]. The paradigm shift in the method of service delivery is suggested in response to increasing long-term, chronic conditions forming the major burden of care worldwide. Acute secondary care is expensive, thus the shift in care also reflects the need to find affordable models of healthcare in an age of diminishing budgets worldwide [WHO, 2015]. Integrating health and social care services and increasing primary care spend are cited as two potential ways of facilitating this shift in focus [WHO, 2015]. Policies introduced that facilitate integration of services

have been implemented despite little evidence to suggest they will have the desired effect [Petch, 2009, 2012, Burgess, 2012, Robson, 2013, Weatherly et al., 2010, Stewart et al., 2003, Kaehne et al., 2017, Damery et al., 2016]. The continued drive to integrate services does, however, implicitly acknowledge that health and social care services are linked. How these services interact at the individual level and whether differing levels of provision in each service affects the other is not well understood [Glasby et al., 2011, Bardsley et al., 2012b, OECD/EU, 2013].

Internationally, provision of social care has become one of the most important issues for policy makers in recent years [OECD/EU, 2013, Humphries et al., 2016]. In Scotland, approximately two-thirds of individuals receiving social care services are over the age of 65 [Scottish-Government, 2016a] whilst approximately two-thirds of all those over the age of 65 have multimorbidity (two or more health conditions) [Barnett et al., 2012]. It would seem intuitive that a large proportion of those receiving social care (if not all) have multimorbidity. However, no single data source exists that allows this comparison to be made. Nevertheless, guidelines exist for healthcare professionals to assist in assessing the social care needs of older people with multiple long term conditions [NICE, 2015]. Multimorbidity is associated with a number of negative outcomes including increased health care usage [NICE, 2016]. Whether multimorbidity increases use of social care services is unknown but this could have an important role in informing policy decisions regarding social care provision.

Levels of multimorbidity in the Scottish population follow a stark socioeconomic profile with those of lower socioeconomic position having higher levels of multiple conditions and more complex care needs [Barnett et al., 2012, Mercer et al., 2009]. This inequality in outcome is compounded by the fact that primary care provision in areas of higher socioeconomic disadvantage, and thus areas of higher need, receive the same or less funding as other more affluent areas. This inequity in provision of service demonstrates existence of the inverse care law in primary care services [Tudor-Hart, 1971, Mercer et al., 2012, McLean et al., 2015] which states, “The availability of good medical care tends to vary inversely with the need for it in the population served [Tudor-Hart, 1971].” Little is known about how access to social care differs across socioeconomic and geographic strata. Health services are free to access at the point of need whereas social care is provided by Local Authorities and access is regulated by eligibility criteria. In an age of austerity, the question of whether an inverse *social* care law exists remains unanswered.

Until recently many local authorities had attempted to protect front-line services, such as social care, from austerity cuts [Hastings et al., 2015]. However, given continued year-on-year reductions and a further 7.2% cut to local authority spending in 2016/2017 [Audit-Scotland, 2016a], the ability to protect social care from reductions in spend becomes less likely. Decreased local government budgets across the UK and Scotland since 2010 have affected those living in the poorest areas hardest [Hastings et al., 2015, Gannon et al., 2016]. If social care budgets decrease further, the question of whether the most deprived areas will feel these cuts most is of grave importance.

The overall aim of the PhD project is to empirically explore these issues. This chapter will specifically review literature regarding the three main themes outlined above namely; social care, the interaction of health and social care services, and multimorbidity. Section 2.2 reviews definitions, international models, and variation in access to social care. Section 2.3 outlines the policy framework regarding health and social care services, how these services are funded and delivered, and why they are linked. It then describes the legislation that made health and social care integration law in Scotland before reviewing empirical evidence of the nature of the interaction between health and social care services. Finally, Section 2.4 describes why multimorbidity is important in the context of health and social care integration and then provides an overview of academic literature and policy documents regarding multimorbidity and its definitions, measurement, and epidemiology.

2.2 Access to Social Care

2.2.1 Definitions

As in the case of multimorbidity, discussed in section 2.4.2, there is no internationally (or nationally) accepted definition of social care. Indeed, the difference between what is social care and what is health care has no clear line of demarcation resulting in local variation in provision of services [McDonald, 2006]. The Organisation for Economic Co-operation and Development (OECD) and the European Union (EU) jointly published a report on Long Term Care (LTC) for older people discussing much of what may be described in the UK as social care. In the report, LTC is defined as,

"... a range of services required by persons with a reduced degree

of functional capacity, physical or cognitive, and who are consequently dependent for an extended period of time on help with basic activities of daily living (ADL). This "personal care" component is frequently provided in combination with help with basic medical services such as "nursing care" (wound dressing, pain management, medication, health monitoring), as well as prevention, rehabilitation or palliative care. Long-term care services can also be combined with lower level care related to "domestic help" or help with instrumental activities of daily living (IADL)."

[OECD/EU, 2013, pp38]

A recent NICE guideline [2015] addressing social care needs for older people with multiple chronic conditions, used a definition provided in the UK Health and Social Care Act [2012]:-

““Adult social care”— (a) includes all forms of personal care and other practical assistance provided for individuals who, by reason of age, illness, disability, pregnancy, childbirth, dependence on alcohol or drugs, or any other similar circumstances, are in need of such care or other assistance, but (b) does not include anything provided by an establishment or agency for which Her Majesty’s Chief Inspector of Education, Children’s Services and Skills is the registration authority under section 5 of the Care Standards Act 2000.” (The Health and Social Care Act 2012 c7, Part 3, Chapter 1, Section 65, Subsection 4)

The NICE guideline [2015] advises that social care planning for people with multimorbidity should include holistic assessment of biopsychosocial factors including sexual, spiritual, cultural, and communication needs as well as considering access to leisure and social activities incorporating issues regarding mobility and transport. Specifically, the guideline cites; self-care, taking medicines, learning, volunteering, maintaining a home, financial management, employment, socialising with friends and hobbies as activities that all patients should be able to take part in should they wish and social care assessment should assess the ability of the individual to achieve this.

A more succinct definition of social care is used in a report to the Minister for Care Services at the UK Department of Health, :-

"The group of services that provide personal care and support to people

in social situations – such as family; the community; a communal setting; to help them achieve independence and to promote their positive contribution as citizens." Platt

[2007, pp. 4]

Huxley et al. [2007] are critical of this service-based definition and argue that social care is intended to improve general well-being for those that are in need. As quality of life is an important factor of well-being, Huxley et al. [2007] argue that wider issues regarding environment and the quality of public and private services also play an important role in social care. Indeed, Daly and Lewis [2000, pp.287] argue that social care is "... an activity and set of relations lying at the intersection of state, market, family (and voluntary sector) relations."

This view is reflected in an aspirational constitution for social care published by an independent, cross-party think-tank [Bartlett and Guglielmi, 2009]. The authors argue that all citizens should have an equal ability to live and control a full and active life. Where this is not possible the state should have a duty to provide the necessary help, in whatever form that is required, to individuals who require it.

A more clearly defined concept is that of *personal care* which has been provided for free in Scotland since 2002. The legislation introduced by the then Scottish Executive necessitated a clear definition and constitutes six dimensions [Cavaye, 2006, pp.256].

- Personal Hygiene: washing etc.
- Personal Assistance: help with dressings, prostheses etc.
- Continence Management: toileting, catheter management etc.
- Food and Diet: help with eating, food preparation etc.
- Problems of Immobility:
- Simple Treatments: help with medicines, creams, oxygen therapy etc.

Personal care is, however, only one aspect of social care provision and clear definitions of other services provided to individuals are lacking. Nevertheless, the definitions of social (or long-term) care above all highlight services that are required to aid with an individual's functional or cognitive needs.

A final definition provided by Colombo et al [2011] will be used for the purposes of this thesis,

- a) a group of services such as; skilled nursing care, social work, personal care, medical equipment & technologies, and therapies. Delivered by,
- b) a range of professionals such as; nurses, low-skilled carers, or allied health professionals. In,
- c) various locations such as; at home, in an institution, or via community care."

This definition clearly captures the broad range of services that can be associated with social care that are only partially provided in other definitions. It acknowledges that social care can include an number of components including personal, nursing care and help with other domestic activities, and articulates the variety of settings where this can take place. Whilst it is common in Europe to describe “Long-term care” in relation to these services, this thesis will refer to “social care” as this is the most commonly used term in the UK. Furthermore, unless stated otherwise, reference to social care in this thesis will be with regards to care received by adults over the age of 65.

2.2.2 International models of social care

There are four ways in which social care can be provided to those in need; informally via family or community, formally via voluntary non-profit organisations, formally via the state, or formally via a for-profit organisations [Munday, 2003]. In Europe, increasing demand from users has led to many welfare systems being unable to adequately provide care [Pavolini and Ranci, 2008, Colombo et al., 2011]. Changes in demography, the labour market, democracy, and values have all contributed to the increasing pressure on care services [OECD/EU, 2013, Anttonen, 2005, Colombo et al., 2011]. There is wide consensus that lower birth rates and higher proportions of older people mean that a gap has emerged in the number of adult children able to provide informal care to their parents [Anttonen, 2005, Deusdad et al., 2016, Pavolini and Ranci, 2008, Munday, 2003, Robertson et al., 2014, Colombo et al., 2011]. Traditionally, informal care was provided by women. As gender equality improves, more women are employed outwith domestic circumstances which also reduces the pool of informal social care available [Anttonen, 2005]. Anttonen [2005] also cites changes in societal attitudes from “familism” to “individualism” as having an impact on informal care resources. These combined factors mean that formal care services are increasingly required to provide social care. Pressures on these services has seen increased discussion and comparison of models of

Table 2.1: Models of social care in OECD countries adapted from Colombo et al (2011)

Model	Countries where employed
Universal coverage	
a) tax based	Norway, Sweden, Denmark, Finland
b) public long-term insurance	Germany, Japan, South Korea, Netherlands, Luxembourg
c) health system	Belgium
Mixed systems	
a) parallel universal schemes	Scotland, Italy, Czech Republic, Poland
b) income-related universal benefit or subsidy	Ireland, Australia, Austria, France
c) mix of universal and means-tested (or no) benefit	Switzerland, New Zealand, some Canadian Provinces, Spain, and Greece ¹
Means-tested safety net	England, USA

¹ Spain and Greece have less well developed formal care services

care across Europe over the last 20 years [Anttonen and Sipilä, 1996, Fernández-Alonso and Jaime-Castillo, 2016, Munday, 2003, Sapir, 2006].

In a report for the OECD, Colombo et al [2011] categorised the varying models of social care into three main groups with subdivisions as shown in Table 2.1.

Universal models of social care are characterised by; a) a single system being in place for delivery of services, b) nursing *and* personal care are provided for all individuals meeting defined eligibility criteria, and c) some form of charge is levied on service-users [Colombo et al., 2011]. Three subdivisions of universal coverage social care models are identified in this classification.

The tax-based universal models, also known as the Nordic model of care, have strong local-government influence in both the raising of funds and delivery of services [Colombo et al., 2011, Fernández-Alonso and Jaime-Castillo, 2016, Munday, 2003, Sapir, 2006]. Overall responsibility remains with national government which also contributes funds, often dependent on local population need and structure [Colombo et al., 2011]. Public long-term insurance schemes are mandatory in the countries that employ them, although the age at which citizens begin to contribute varies (e.g. only over 40s contribute in Japan) [Colombo et al., 2011, Forder and Fernández, 2011, Eleftheriades and Wittenberg, 2013]. These models have found it increasingly difficult to fund provision and have either increased user-payments or decreased coverage in recent years [Robertson et al., 2014, Forder and Fernández, 2011, Eleftheriades and Wittenberg, 2013]. In Belgium, social care is viewed as a health risk and is financed through the health system with a cap on user-contribution. This results in a medical (rather than social) model of care delivery, mostly provided by nurses [Colombo et al., 2011].

In contrast to universal models described above where a single system is in place for social care delivery, mixed social care systems employ either; a) universal coverage via different sources/sectors or b) a mixture of universal and means-tested benefits [Colombo et al., 2011]. Whilst acknowledging the wide variation in systems, Colombo et al [2011] identify three subdivisions of mixed system models of social care as shown in table 2.1.

Parallel universal schemes provide social care via two or more sectors (e.g. nursing care via health provider and personal care from a non-healthcare source) [Colombo et al., 2011]. Major differences exist between countries in how this is delivered in practice with varying degrees of coverage. Income related universal benefits provide needs-based universal coverage of social care but benefits received vary according to income [Colombo et al., 2011]. France provides an good example of this system where cash benefits are needs *and* income based - benefits are paid at six levels of dependency, those with higher incomes are paid proportionally less and top-up care costs themselves [Colombo et al., 2011, Glendinning and Bell, 2008]. The final subdivision of mixed social care models has some universal entitlements provided alongside means-tested services e.g. free provision of nursing care with means-testing for personal care.

Colombo et al [2011] include Spain and Greece in this final subdivision of mixed social care models but note these countries have formal care systems that are much less developed than elsewhere in the OECD. Sometimes referred to as the Mediterranean model, informal care from family and other sources constitutes the main form of social care in these countries [Fernández-Alonso and Jaime-Castillo, 2016, Munday, 2003, Sapir, 2006]. As this role is traditionally carried out by women, the Mediterranean model has attracted criticism from a feminist perspective [Munday, 2003].

The final model in Colombo et al's [2011] classification of social care in OECD countries is the Means-tested safety net model (Table 2.1). In this model only those below a pre-determined threshold are entitles to state-provided social care. Despite the free provision of health care and some non-means-tested benefits, England is included in this category in the report. Presumably this is due to the fact that state support in a residential home in England is provided only after an individual has depleted both income *and* assets below a threshold [Colombo et al., 2011], a system not seen in other countries ¹ [Glendinning and Bell, 2008].

¹A cap of approximately £72,000 total contribution was to be introduced in 2016 (Eleftheriades et

Of the three models of social care described in this section, universal systems have a number of advantages over the other two [Colombo et al., 2011, Glendinning and Bell, 2008]. Sharing the burden of social care provision among the entire population, either via general taxation or mandatory insurance schemes, results in a reliable, predictable source of finance enabling states to plan more effectively [Colombo et al., 2011, Glendinning and Bell, 2008, Forder and Fernández, 2011]. Mixed systems of social care can still result in considerable costs for those that require care, whereas the costs to individuals in the means-tested system can be “catastrophic” [Colombo et al., 2011, pp240]. The means-tested system can also be perceived as unfair for those that need to sell assets (such as a house) to pay for care, particularly when there is variation in eligibility criteria within countries [Colombo et al., 2011, Glendinning and Bell, 2008, Charlesworth and Thorlby, 2012, Eleftheriades and Wittenberg, 2013, Forder and Fernández, 2011]. A particular advantage of a universal social care systems is that it,

“...generally reduces utilisation of more expensive health care services and professionals (e.g. hospital care, doctors) for long-term care needs, for example by making “social hospitalisation” of frail people with [social care] needs more difficult.”

[Colombo et al., 2011, pp.222]


Given the pressures on state budgets and the desire to adequately fund care services, this seems a particularly useful benefit of the universal coverage model of social care.

Disadvantages of universal systems are that they, generally, are more expensive to the state than other models of social care and can reduce the amount of informal care provided by relatives for elderly family [Colombo et al., 2011, Eleftheriades and Wittenberg, 2013]. The comprehensiveness of coverage can be over-burdensome for the state with a number of countries recently having to cut service or increase user-contributions to compensate for increasing demand [Robertson et al., 2014, Charlesworth and Thorlby, 2012, Eleftheriades and Wittenberg, 2013, Forder and Fernández, 2011].

A recent examination of the effects of the 2008 financial crisis on the way social care is delivered across Europe suggests that the distinctions between social care models is beginning to blur [Deusdad et al., 2016]. There is evidence those with more comprehensive coverage are reducing levels of care whilst those with less coverage are


al., 2013)

Figure 2.1: Older recipients of long-term care services as a share of the over 65 population, 2008



Note: LTC recipients aged over 65 years. Recipients refer both to home and institutional users. Data for Australia, Belgium, Canada, Denmark, Luxembourg and the Netherlands refer to 2007; data for Spain refer to 2009; data for Sweden and Japan refer to 2006. Data for Japan underestimate the number of recipients in institutions because many elderly people receive long-term care in hospitals. According to Campbell et al. (2009), Japan provides public benefits to 13.5% of its population age over 65 years. Austrian data represent recipients of cash allowances.

Source: OECD Health Data 2010.

StatLink  <http://dx.doi.org/10.1787/888932401577>

increasing provision [Colombo et al., 2011].

Given the marked variations in how social care is funded and delivered across countries, it may be expected that there are also marked variations in levels of access and utilisation. Colombo et al [2011] produced a chart (shown in figure 2.1) derived from OECD data that shows the proportion of over 65s receiving some form of social care across countries for which data is provided. The chart shows that having a universal or mixed system of social care provision (as described above) does not absolutely influence the number of people receiving care. For example, South Korea employs a universal (insurance-based) system and has one of the lowest proportions of older people receiving care, whereas Switzerland has one of the highest whilst employing a mixed system involving some universal and some means-tested benefits. This suggests that allocation of resources and eligibility criteria set within countries are likely to be more important in determining access to social care than any particular system of care delivery.

2.2.3 Access to Social Care - Social Theory of Eligibility

"...the *criteria* under which a given individual is eligible for publicly funded support for long-term care, and for how much support the individual is eligible, and the *processes* involved in selecting from the general population those who receive this support and determining for how much support each person is eligible"

[*Eleftheriades and Wittenberg, 2013, pp.2*]

Two social theories on how eligibility for public services are determined will be discussed in this subsection; Street-level bureaucracy and candidacy.

The term Street-level bureaucracy is generally credited to Michael Lipsky and his book of the same name [1979], along with its more recent revision [Lipsky, 2010]. The theory investigates the extent to which front-line public service workers exercise discretion in which individuals of the general public are eligible to access any given service - doing so from a position of authority [Lipsky, 2010]. As a result, Street-level bureaucrats control, "... the nature, amount, and quality of benefits and sanctions provided by their agencies." [Lipsky, 2010, pp.13]. Using the term, "Street-level" suggests that the theory is concerned with power, where it resides, and who wields it [Hupe et al., 2016]. Street level bureaucrats can be teachers, police officers, nurses, social workers, or any other worker providing a public service [Hupe et al., 2016] and their , "... decisions... , the routines they establish, and the devices they invent to cope with uncertainties and work pressures effectively *become* the public policies they carry out" [Lipsky, 2010, pp.xiii].

Evans[2011] and Ellis [2010] both provide a critique of street-level bureaucracy that suggests the theory needs to be augmented to take into account new structures of public services, particularly in social work, that place greater autonomy with managers than front-line workers. They argue that these new structures mean workers who deal face-to-face with public service users have less discretion about eligibility criteria and

are more likely to have to justify eligibility decisions to senior members of staff who now wield more power in terms of service access.

Empirically exploring the effects of Street-level bureaucracy poses an number of methodological problems [Hupe et al., 2016]. The use of large sample surveys of public sector workers investigating their views and how they make decisions is one way (e.g. [Tummers and Bekkers, 2014]). However Lipsky [2010] and Hupe et al [2016] agree that qualitative interview techniques with public sector wokers in their own workplace is the best way to fully understand how street-level bureaucracy impacts on front-line services.

The term “candidacy” was first used by Dixon-Woods et al [2005; 2006] to form a concept of how vulnerable individuals identified themselves as being a eligible for a particular health service or intervention given for particular illnesses or health conditions. The theory was further explored and augmented by Mackenzie et al [2013; 2015] in relation to access and utilization of all public services as a way to explore concerns about inequal distibution of uptake.

The theory argues that there are a number of social and cultural factors that contribute to an individual’s interpretation of their eligibility for a given service and is based on a seven stage model as shown in Table 2.2

Table 2.2: Seven stage model of candidacy

Stages of candidacy	Description of stage
Self-identification of candidacy	Process by which individuals come to view themselves as legitimate candidates for particular services
The availability and accessibility of services	Knowing how to make contact with appropriate services in relation to identified candidacy
Permeability of services	Includes the level of explicit and implicit gate-keeping within a service and the complexity of its referral systems; in addition it refers the "cultural alignment" between users and services
Appearing at services and asserting candidacy	The work that an individual must do to assert their candidacy in an interaction with a service professional
Professional decision making	Candidacy as expressed by service user is validated or otherwise by service professional. This influences future offers of service
Offers of and resistance to services	Service may be appropriately or inappropriately offered by a professional and this may, or may not, be acted on by service user
Operating conditions and local production of candidacy	Incorporates factors that influence decision about future service provision (e.g. resources) and the relationship that develops between service users and professionals over a number of encounters

This is a much broader theory than that of street-level bureaucracy and focusses on the barriers and enablers individuals face/use when accessing services. It could be argued

that candidacy includes the concept of street level bureaucracy in the fifth and sixth stages of the model shown in table 2.2. “Professional decision making” and “Offers of and resistance to services” are areas where an interface between a service user and service professional takes place - much like the interactions described by Lipsky.

Investigating candidacy as a theory empirically again appears to be best served using qualitative methods. The complex and dynamic nature of candidacy suggests identifying meaningful quantitative measures are unlikely. Both Dixon-Woods et al [2005; 2006] and Mackenzie et al [2013; 2015] employed critical interpretive synthesis in their studies.

Both of the social theories described in this subsection deal with the concept of eligibility and how, in the case of street-level bureaucracy, professionals exercise discretion on access to service and, in the case of candidacy, how service-users identify whether they are eligible and offer themselves for a service in the first place. Both theories describe complex interactions between individuals across an eligibility criteria barrier. The next section describes in detail this eligibility barrier in relation to social care in the UK, firstly from a Scottish perspective and then more broadly in the wider UK.

2.2.4 Access to Social Care - Eligibility in the UK

"It is through the eligibility criteria that resources are rationed, that is "need" is equated with "resources available". This mechanism severely limited the idea that provision could be determined either by need or by the right to services."

[Sharkey, 2006, pp.10]

In Scotland, access to social care is needs-tested via assessment carried out by a social worker. The criteria for social care delivery, therefore, has a very important part to play in how services are accessed.

In 2010 the Scottish Government published a report identifying a strategy for the policy of self-directed support [Scottish-Government, 2010]. The report was written in conjunction with the Convention of Scottish Local Authorities (COSLA) and included the

recommendation that the National Eligibility Framework developed by the Sutherland review into free personal and nursing care [2008] should be applied across all social care services. The framework has four criteria for assessing risk in relation to a person's care needs: Critical, Substantial, Moderate and Low [Scottish-Government, 2015]. The Critical and Substantial levels of risk indicate social care needs should be addressed immediately or imminently, whereas a moderate level of risk may indicate either some or no services being required. There is *no explicit description* of severity or which care needs fall into each category and in practice each Local Authority sets the criteria and decided at which level of risk they would provide social care [Scottish-Government, 2014].

Equity of access to services is directly influenced by an eligibility framework. Indeed, the Strategy for Self-Directed Support [Scottish-Government, 2010, pp.20] acknowledges this and states that such a framework "... can result in resources being narrowly focused on individuals with acute needs." However, the report goes on to state that growing demand and finite resources requires some form of eligibility assessment but this should not have a disproportionate effect on any one group of people requiring care.

The eligibility framework allows each Local Authority to set thresholds for access to care in line with local priorities and resources. This has the effect that access to services varies across differing council areas. The potential for regional variation is again acknowledged by the Strategy for Self-Directed Support [Scottish-Government, 2010, pp.20] which states that, "... further work will be undertaken by the Scottish Government and COSLA to assess whether there is merit in establishing national thresholds for access to formal support across all client groups."

Acknowledgment of problems with eligibility criteria and the promise of "further work" to be undertaken by the Scottish Government and COSLA is repeated in practitioner guidance on Self-Directed Support published in 2014 [Scottish-Government, 2014, pp.19] and that , "... it remains the case that local authorities should operate eligibility criteria to determine whether or not an individual assessed as having social care needs can access formal support and if so, which of their needs are to be met by that support."

Data is not available on levels of care provided by LAs for each of the National Eligibility Framework criteria or for the threshold that each LA provides care at. The Scottish Government collects an annual report of eligibility and waiting times for the 1st quarter of the year. A recent report [Scottish-Government, 2015] provides information on the

time individuals had to wait to receive assessment and the time individuals had to wait to receive care in the period January-March for the preceding 5 years. However, no absolute numbers of people in each category is provided.

The Scottish National Eligibility Framework has striking similarities to that formerly used in England and described in Fair Access to Care Services (FACS) produced by the Social Care Institute for Excellence (SCIE) [2013]. Exactly the same nomenclature is used to describe the eligibility categories of need. Newton and Browne [2008] critiqued a previous version the FACS guidance and found similar issues to those raised above regarding regional variations in service and concentration of services on those with the highest need. Their paper describes further issues with access to social care in the context of social theory described by Lipsky [1979] and “Street level bureaucracy” (discussed in section 2.2.3) where intentional and unintentional judgment of entitlement by social care workers have an impact on whether an individual receives care or not. Newton and Browne [2008] also make the assertion that health and social care has never been accessed equitably by arguing that those with a greater ability to articulate needs and negotiate access are more likely to gain access to services. Although no citation is provided to back-up this argument it has certainly been described elsewhere [Matthews and Hastings, 2013] and sits well in the broader discussion of inequitable access to services [Hastings et al., 2014, Le Grand, 1989, Tudor-Hart, 1971] (discussed further in section 2.2.5).

In England, the Care Act [2014] aimed to remove regional variations in eligibility in access to social care by imposing national minimum thresholds that Local Authorities would have a statutory obligation to provide. The Care Act also aimed to ensure Local Authorities provided care, “...as early as possible to help maintain well-being and independence, and potentially delay a situation where longer-term care and support might be required.” [SCIE, 2015, pp.2]. The minimum criteria for being eligible for care involves an individual having needs that impairs their ability to meet 2 or more of a designated list of outcomes (e.g. managing and maintaining nutrition or maintaining hygiene) [SCIE, 2015] and is set by the Secretary of State for Health [Abrahams et al., 2014].

In practice, the most likely outcome is that the minimum threshold that Local Authorities will have to provide care will be similar to the “Critical” level of the National Eligibility Framework previously used in the FACS guidance [Burchardt et al., 2015, Abrahams

[et al., 2014](#)] (and similar to that used in Scotland). This will legalise a shift that has already been occurring in England where fewer numbers of LAs are providing care for those with “moderate” needs and only providing care for those with “critical” needs [[Burchardt et al., 2015](#), [Abrahams et al., 2014](#)]. Burchardt et al. [2015] state that only 2% of English LAs will have to widen their care threshold whereas 12% could now, legally, tighten care provision as a result of the Care Act. This situation is not new and has been gradually worsening over the past decade and has profound impacts on the quality of, and access to, social care [[Sharkey, 2006](#)].

A recent report by the House of Commons Communities and Local Government Committee [2017] confirmed reductions in the absolute number of people receiving care, the concentration of services in those with highest needs only, reduction in quality of care provided, and the resulting pressures this caused to the health service through increased emergency admissions and delayed discharges. The report highlights the perilous state of social care provision in England and urges immediate attention from the government to address funding shortfalls.

Burchardt et al. [2015] and Abrahams et al. [2014] recognise some positive changes to Social Care policy through the Social Care Act but are damning about past UK Government Social Care policy in England and Wales. They cite chronic underfunding and cuts for over ten years resulting in fewer numbers of people receiving care at a time when demand is sharply increasing due to demographic change. The “intensification” of services on those with the most acute needs is cited by both sets of authors as counterproductive – ignoring those with moderate care needs completely derails one of the main purposes of the Care Act, preventative services. Indeed,

“As well as lacking in moral sense, such an approach is economically unsound. Waiting for people to have high needs before providing care means that care will be more expensive, as well as pushing more older people into an already pressurised NHS. ”

[[Abrahams et al., 2014](#), pp.5]

A similar picture has been seen in Scotland. Absolute numbers of people receiving home care has steadily fallen over the last 10 years whilst the number of hours of care provided has increased [[Scottish-Government, 2016a](#)]. There are wide variations in the number of hours of home care provided per population across Local Authorities [[Audit-Scotland](#),

2012, [Scottish-Government, 2016a](#)]. This may reflect different demographic make-up of each Local Authority although reductions in ratios per population can be seen in almost all Local Authorities [[Scottish-Government, 2016a](#)]. Audit Scotland [2012] also highlighted that intensifying services is likely to be a short-term solution with negative long-term impacts and suggests comparison of performance across Scotland would be beneficial in identifying good practice.

In a report profiling the care at home sector in Scotland. MacLeod and Mair [2015] describe large decreases in absolute numbers of people receiving care at home over the ten years to 2013. There have also been significant reductions in the number of people receiving non-personal care (so called “mopping and shopping”). The increase in the number of hours of home care delivered by all services reflects a focus on smaller numbers of individuals with higher care needs. This means those with moderate or low personal care needs and those requiring “mopping and shopping” services are now less likely to receive publicly funded care. Echoing the views of Burchardt et al. [2015] and Abrahams et al. [2014], Macleod and Mair [2015] highlight the potential false economy of this situation – home care services are likely to reduce the need for costly emergency admissions to hospital and delay the requirement for more intensive home care packages.

2.2.5 Access to Social Care - Social Theory of Resource allocation

"Almost all public expenditure on the social services in Britain benefits the better off to a greater extent than the poor"

[Le Grand, 1989, pp.3]

In his seminal book, “The strategy of equality”, Julian Le Grand [1989] investigated whether social and economic equality had been achieved since the introduction of post-war welfare spending. The book compares the distribution of public expenditure and outcomes across health, education, housing, and transport. It concludes, as highlighted in the quote above, that those with higher socioeconomic position benefited disproportionately from government social services spending across all sectors. Indeed ,

“... there persist substantial inequalities in public expenditure, in use, in opportunity, in access and in outcomes”[Le Grand, 1989, pp.4].

Criticism of Le Grand’s conclusions cites subsequent research that shows empirical evidence indicating a reduction in inequalities and questions the assumption that the sole purpose of the welfare state is to achieve equality [Powell, 1995]. More recent research [Bramley et al., 2005, cited in; Hastings et al., 2014],[Hastings, 2007, Bramley and Evans, 2000] has shown that when comparing distribution of resources at neighbourhood level (rather than national level) there is higher spending in less affluent areas. However some service were found to be “pro-rich” (education, pensions) and others “pro-poor” (parks, environmental services) [Bramley et al., 2005, cited in; Hastings et al., 2014]. Whether a service is more likely to be used by more or less affluent citizens is important in terms of resource allocation - particularly when services are being cut as shown by Gannon et al [2016] and discussed further in section 2.2.6.

Understanding why there are differences in resource allocation for different types of service has led to the investigation of “Middle Class-capture” of services and how it is obtained [Gal, 1998, Matthews and Hastings, 2013, Hastings et al., 2014]. An adapted version of Gal’s [1998] six channel framework of middle class advantage described by Hastings et al [2014] is shown in table 2.3

Table 2.3: Six channels of middle class advantage

Channel	Description of channel ¹
Electoral	Large middle class more likely to vote thus political policies influencing welfare services more likely to be geared toward them.
Organisational	Unions and professional associations representing middle-class occupations have strong influence on welfare policy
Knowledge	Resources of education and access to information possessed by middle class mean they have better understanding of "how the system works" and therefore can better exploit it
Mass Media	Middle class has dominant role in media and can thus exert influence over how policy is covered. Also able to access and influence those that produce mass media more easily
Exit	Ability of middle class to leave public provision for private alternative influences public policy in order to avoid this happening.
Bureaucratic	Public services "run" by the middle classes therefore exert influence over how it is accessed and by whom.

¹ Adapted by Hastings et al [2014] from Gal [1998]

These six “channels” conceptualise the modes of how and why welfare spending in certain areas appears to benefit more affluent groups. In their study investigating street-cleansing services, Hastings et al [2014] observed the influence of middle class capture and some of the channels of advantage described in table 2.3 suggesting the theories of Le Grand [1989], described above, and Tudor-Hart, described below, should not be discounted.

"The availability of good medical care tends to vary inversely with the need for it in the population served

[Tudor-Hart, 1971]. In a similar fashion as Le Grand’s work, “The inverse care law” has informed much research since first coined in the early 1970s. As discussed in section 2.2.7, people living in more deprived areas have lower life expectancy and higher morbidity figures and therefore greater health needs [Baker et al., 2015]. However, the poorest neighbourhoods in England have been reported to have 62.5 General Practitioners (GP) per 100,000 population whereas the most affluent neighbourhoods have 76.2 per 100,000 [CfWI, 2014] which suggests health provision does not match need. Recent planned changes in policy to distribute primary care funding based on population age are likely to exacerbate this situation [Mercer et al., 2012]. Indeed, increases in workload with deteriorating proportions of budgets has lead the King’s Fund to describe the situation in primary care in England and Wales as, “in crisis” [Baird et al., 2016, pp.3]. In Scotland, the even distribution of GP workforce among the population means GP practices in the most deprived areas need to provide more consultations, for people with greater needs, at the same funding level as practices with fewer resource demands [McLean et al., 2015, Mercer and Watt, 2007]. Poorer access to primary health care is associated with greater demand for unnecessary admission to hospital [Rosano et al., 2013, Weston et al., 2016] which is responsible for high proportions of healthcare expenditure.

There has been no research, to the author’s knowledge (is this ok??), of whether the inverse care law is observable in social care - a service delivered, like primary care, in a community setting and also likely to have an impact on secondary health care use. Nor has any research specifically investigated variations in the distribution of social care services by socioeconomic position at the local level. Such research would add useful evidence to the debate regarding the strategy of equality and middle class capture.

The next section describes how social care is funded in Scotland and how cuts to services post-2008 may adversely impact less affluent members of the public.

2.2.6 Acces to Social Care - Resource Allocation in Scotland

Local authorities in Scotland have a statutory obligation to provide social care to individuals they have assessed as eligible for care [[Audit-Scotland, 2012](#)]. All local authority funding is provided by the Scottish Government via a block General Revenue Grant made up of a number of components [[Scottish-Government, 2013](#), [King et al., 2007](#)]. The majority of this grant is calculated via a formula know as the Grant Aided Expenditures (GAE) which accounts for over 80% of the General Revenue Grant [[Scottish-Government, 2016c](#)]. The formula for GAE is calculated using what is called a “client group approach” and is based on 89 services provided by local authorities [[Scottish-Government, 2016c](#)]. A national figure for each service is set and each local authority receives a percentage of that figure based on estimates of the number of people that use that service (a capitation) and other secondary indicators such as area deprivation or rurality [[Scottish-Government, 2016c](#), [Smith, 2003](#)]. For example, funding for primary school teachers is based on the number of children in primary education (primary indicator) and adjusted to take into account the percentage of pupils in small schools (secondary indicator) [[Scottish-Government, 2016c](#)].

The use of formulae to allocate public expenditure has potential to improve efficiency in spending and equity of distribution [[Smith, 2003](#)]. Equity of distribution is achieved via the explicit nature of a formula framework with transparent methodology that can be debated and amended [[Smith, 2003](#)] (The formula for the “Green Book” settlement was agreed with the Convention of Scottish Local Authorities (COSLA) [[Scottish-Government, 2013](#)]). However, as King et al [[2007](#)] note, GAE grants to local authorities are estimates of relative, rather than absolute, spending needs in that area. The “Green book” reporting the annual settlement for local authorities in Scotland stresses that the values allocated for different services are not budgets or targets and that local authorities are free to spend resources (other than ring-fenced monies) as they see fit [[Scottish-Government, 2016c](#)]. In effect, “. . . the capitation payments seek to offer comparable public sector organisations the opportunity to deliver some average level of service, assuming average responses to social and economic circumstances, and an average level of efficiency” [[Smith, 2003](#), pp.309].

The “Green Book” outlines seven main areas of local authority expenditure from which the 89 services mentioned above are derived; Education, Social Work, Roads & Transport, Leisure & Recreation, Cleansing & Environment, Elections & Taxation, and

Other Services [[Scottish-Government, 2016c](#)]. Spending allocation for social care comes under the Social Work heading which is subdivided into 23 subcategories of services, Of these, nine are directly related to social care as defined for the purposes of this thesis (the others being based on e.g. children's services);

- Service for Home Based Elderly
- Residential Accomodation for the Elderly
- Casework and related Administration: Elderly
- Services for People with Disabilities
- Casework and related Administration: People with Disabilities
- Independent Living Fund
- Carers Support and Respite Services
- Care Home Fees
- Personal and Nursing Care for Older People

The expenditure for the first three items in this list as well as Carers Support & Respite Services and Care Home Fees are calculated using population weighted indices for each local authority calculated from; the standardised mortality ratio, census data on self-report long term illness & people living alone, as well as pension credit data (Service for Home Based Elderly) or Council Tax Data (Residential Accomodation for the elderly). Services relevant to People with Disabilities and the Independent Living Fund are calculated depending on the number of people aged 16-64 in each local authority. Expenditure allocation for Personal and Nursing Care for older people is derived from formulae calculated in the Scottish Government Health Directorate Distribution [[Scottish-Government, 2016c](#)].

The GAE formula has been in place for some time (initially outlined in 1992 [[Scottish-Government, 2016c](#)]). The more recent issue facing local authorities in terms of finance has been cuts following the 2008 financial crash. In the financial year 2016/17 the overall grant to Scottish local authorities was cut in real-terms by 5% which added to a cumulative real-terms cut of 11% since 2010/11 [[Audit-Scotland, 2016c](#)]. Authorities have been managing this pressure by reducing spending in all areas of their budgets - with the exception of social work [[Audit-Scotland, 2016c](#)]. £3.1 billion was spent on social work by Scottish local authorities in 2014/15 - an increase of 3% since 2010/11 and a third of all council spending [[Audit-Scotland, 2016b](#)]. However, given the 5% decrease planned for 2016/17 Audit Scotland [[2016c](#)] warn that social work, and specifically

social care, budgets are now likely to be cut, which will likely result in a decrease in the quality of service [Audit-Scotland, 2016b].

These budgetary pressures are difficult for local authorities to manage, but what is the outcome on service users? Using the “pro-rich/pro poor” nomenclature initially used by [Bramley et al., 2005] (and discussed in section 2.2.5), Gannon et al [2016] investigated the social impact of spending cuts in Scotland. The report found that the vast majority of local authority spending is on services that are “pro-poor” i.e. services that are disproportionately used by people with lower socioeconomic position. As a result, despite attempts to protect these services, the cuts to local authority spending have a disproportionate effect on this societal group. Councils with higher numbers of the most deprived citizens are having to make the biggest percentage cuts in services defined as “very pro-poor” (e.g. social work for children and families or citizen’s advice). These findings echoed an earlier report from the project looking at cuts across the UK as a whole [Hastings et al., 2015].

Gannon et al’s report [2016] assigns older persons social work as “pro-poor” along with local authority public transport but does not distinguish between the two in analysis. It is therefore difficult to dis-aggregate the specific effect of cuts on social care from the report particularly, as shown above, as there was an increase in spending between 2010/11 and 2014/15. Nevertheless, cuts expected to social care budgets from 2016/17 [Audit-Scotland, 2016c] are also likely to have a disproportionate effect on those with lower socioeconomic position.

Cuts to services reduce the potential for access to such services. If these cuts are disproportionately affecting more deprived communities it is likely unequal outcomes for these communities will be exacerbated. Given the close link of social care to health care, the question of whether social care influences health inequalities is important. The next section discusses presents and overview of literature on health inequalities.

2.2.7 Health inequalities

In the UK, poverty remains the largest predictor of relative ill health and has associations with increased morbidity, multimorbidity, and decreased life expectancy [Baker et al., 2015]. People living in deprived areas are more likely to engage in unhealthy lifestyle behaviours, experience multimorbidity at a younger age, and live in overcrowded or

unsuitable housing [Baker et al., 2015, Shaw et al., 2006].

The influential Marmot review into health inequalities found that those in the most deprived areas of England die, on average, 7 years earlier than their most affluent peers [Marmot et al., 2010] with the gap in life expectancy increasing between 1995 and 2008 [National-Audit-Office, 2010]. Subsequent research by the King's Fund suggests the gap in life expectancy reduced between the periods 1999-2003 and 2006-2010 [Buck and Maguire, 2017]. The report warns that this improvement may be due to the spending and policy decisions of the New Labour Government of the early 2000s and that recent austerity measures in the UK may undermine the progress made [Buck and Maguire, 2017]. Indeed, the most recent analysis released by the Office for National Statistics [2016] suggests the gap in male life expectancy in England is now 9.1 years. In Northern Ireland, the male life expectancy gap is the lowest of all 4 UK nations, however those in the poorest neighbourhoods die, on average, 4 years earlier than those in the most affluent areas [ONS, 2016]. In Wales the gap is slightly larger at 4.2 years [ONS, 2016]. There is a gap of 7 years in life expectancy at birth in Scottish males - those born in East Dunbartonshire can expect to live to 80.5 years, whereas those in Glasgow city can expect to live 73.4 years [ONS, 2016].

The Scottish Government reports statistics on healthy life expectancy which is defined as the number of years people can expect to live in good health [Scottish-Government, 2017]. The most recent figures suggest men and women in the most deprived areas can expect to become ill 25.1 and 22.1 years earlier than their most affluent peers respectively [Scottish-Government, 2017] meaning Scotland has the highest levels of health inequalities in western and central Europe [Mackenbach et al., 2008, Popham and Boyle, 2010].

There are many theories as to why inequalities in health exist across socioeconomic position [Asthana and Halliday, 2006, Nettleton, 2006]. Some of these, such as statistical artefact and biological reasons, were rejected as being implausible by the Black report [Macintyre, 1997]. To a large extent, epidemiological evidence and theoretical argument has agreed with that view [Asthana and Halliday, 2006, Mackenbach, 2012, McCartney et al., 2013, Nettleton, 2006].

There have been many critiques of other theories proposed in the last 35 years which focus on differing numbers of proposals [Asthana and Halliday, 2006, Nettleton, 2006, Peckham and Meerabeau, 2007, Smith et al., 2016, Mackenbach, 2012]. Whilst arguments over

which theory is most plausible to explain the cause of health inequality, most researchers agree on ways to remedy disparities in health outcome namely the redistribution of income, wealth and political power [Smith et al., 2016, Katikireddi et al., 2013, Asthana and Halliday, 2006, Nettleton, 2006]. Although health services have an important role to play, it is the “upstream” policies of redistribution that will make the biggest impacts in improving health outcomes across society [Smith et al., 2016, Katikireddi et al., 2013, Asthana and Halliday, 2006, Scottish-Parliament, 2015]. Whilst this has been known for some time, government policies in the UK to date have not addressed these issues and have thus failed to make meaningful improvements in health inequalities [Frank et al., 2015, Mackenbach, 2010, Peckham and Meerabeau, 2007].

(Might expand this a little more?)

2.2.8 Summary

Either here or at the very end of the chapter??

There is no agreed standard definition of social care, a term often used synonymously is long-term care. The boundry between what is health care and what is social care is often blurry. The definition chosen for this thesis provided by Colombo et al [Colombo et al., 2011] encapsulates the wide number of services that make-up social care including nursing, personal, equipment, and technological. The definition also identifies that social care can be provided not only at home, but also in institutions or other community settings.

Three broad models of social care are seen internationally; universal, mixed, and means-tested schemes. Within each of these models there are many different methods of delivery across countries and no easy comparison can be made identifying differences in outcomes across countries. It does appear that universal systems spread the risk of the costs of social care more equitably among the populations where it is employed. Importantly, every model of social care involves some user-contribution towards costs.

Eligibility for social care is determined via pre-specified criteria in all cases. How these criteria are set varies greatly across and within countries. In UK terms eligibility criteria are set by local authorities and have been greatly tightened in recent years as a response to budgetary constraint. Also observed is the process of “intensification” where greater

hours of social care are being delivered to smaller numbers of people with higher needs. This has potential to erode an important function of social care - preventing expensive unscheduled health care use.

Eligibility for social care can also be affected by the individual in need, and those applying the pre-determined criteria. Social theories regarding this include “street-level bureaucracy” and “candidacy”. Both theories describe difficulties that may exist in individuals attempting to access public provided services, the latter in more detail and including aspects of the former. Both theories are best suited to being investigated with qualitative methods.

Allocation of resources for social care in Scotland are decided by local authorities. The monies they receive are dependent on a grant from the Scottish Government which is calculated via the GAE formula. The GAE formula allocates money for social care services according based on a mixture of data from each local authority including, mortality and morbidity ratios, the amount of people living alone, and the ratio of people paying certain level of tax or receiving certain benefits. This formula has been in place for over 20 years and was agreed with COSLA.

Social theories regarding allocation of resources for public service suggest those with higher socioeconomic position are more likely to benefit from public spending than their less affluent peers. Empirical analysis of; “The strategy of equality”, “The inverse care law”, and “Middle-class capture” all suggest more affluent groups are better at accessing public services.

There have been significant cuts to local authority budgets across the UK since 2008. Savings have been made whilst trying to protect front-line services but current and future cuts are likely to impact these services. Most local authority spending is on services used by those from lower socioeconomic positions thus cuts will disproportionately affect these people. Whether this will have an impact on already wide health inequalities in Scotland is unknown.

2.3 Health and Social Care Interaction

2.3.1 Public Policy

Section discussing public policy of health and social care. Two silos. Short description of NHS funding and compared with LA funding described in subsection ??.

Discuss why these services are linked - hosp discharge etc.

Highlight current policy such as 2020 vision in Scotland, Health and Social Care Act (2012)(UK) **Christie Commision** etc.

Discuss Why poor co-ordination between services?

2.3.2 Health and Social Care Integration

Overview of health and social care integration. What it aims to do. Why etc.

Policy - The Act - structure such as “Lead Agency” or “Joint Body” models.

Evidence of benefits of integration - (not much!)

2.3.3 Research on Health and Social Care Interaction

Section discussing the few studies that have aimed to assess the interaction between health and social care services.

Bardsley et al[2012a] a good starting point - also studies identified by mini lit search in 1st year review plus update.

2.3.4 Summary

Summary of literature presented in section [2.3](#)

2.4 Multimorbidity

2.4.1 Why focus on Multimorbidity?

Important subsection. Brief definition - Identify high use of health care and association with poor outcomes (with citations). Then make case that use of social care likely to be associated with MM (though no evidence to back this up - hence need for thesis).

Discuss whether access to social care likely to be more or less “fair” for those with MM - ref to Inverse Care Law and PC literature.

Will Health and Social Care integration results in better or worse outcomes for MM? (Should be better - main aim of integration..)

Link to following sections as background of current MM literature to help inform understanding of concept

2.4.2 Definitions

Despite the increasing importance of multimorbidity on health care systems, there is no internationally agreed definition of the term or concept [[Almirall and Fortin, 2013](#), [Lefevre et al., 2014](#)]. Van den Akker et al [[1996](#)] first made the distinction between the terms comorbidity and multimorbidity. Comorbidity was originally described by Feinstein [[Feinstein, 1970](#), pp.467] who stated, “In a patient with a particular index disease, the term co-morbidity refers to any additional co-existing ailment.” Van Den Akker et al. [[1996](#), pp.65] used the term multimorbidity to describe, “...any co-occurrence of medical conditions within a person.” In this sense, multimorbidity does not rely on the presence of a primary, or index, disease but refers to the overall state of multiple illnesses.

Further development of definitions is provided by Valderas et al. [[Valderas et al., 2009](#)] who characterise the construct of the term comorbidity found in the literature in four main groups; (a) comorbidity – additional diseases in the context of an index disease, (b) multimorbidity – more than one disease within an individual (without reference to an index disease), (c) morbidity burden – total impact of physiological dysfunction linked to patient outcomes and (d) patient complexity – the effect of non-health characteristics

(e.g. deprivation, culture, environment) on morbidity burden.

Valderas et al. [Valderas et al., 2009] discuss these four constructs of comorbidity further in relation to three main research areas; clinical care, epidemiology and public health, and health service planning. It is suggested that comorbidity may be a more valid definition for use in specialist clinical care, whereas multimorbidity and morbidity burden would be more appropriate in primary care research. In epidemiological and public health research, the definitions of either comorbidity or multimorbidity would be of use depending on the origin of the diseases being studied and the particular research questions being investigated. Morbidity burden and patient complexity are, according to Valderas et al. [Valderas et al., 2009], the most appropriate definitions for research exploring healthcare use and costs.

A further definition of multimorbidity is offered by the European General Practice Research Network (EGPRN) who report findings of a systematic review in the construction of their definition. Citing over 100 different definitions for multimorbidity in academic research the EGPRN [Le Reste et al., 2013, pp.1] aimed to clarify the concept of multimorbidity and define the term as

"...any combination of chronic disease with at least one other disease (acute or chronic) biopsychosocial factor (associated or not) or somatic risk factor."

This definition goes some way to capture the complexity of the concept of multimorbidity as explained by Valderas et al. [2009] but has not ended debate on the matter.

More recently, a systematic review focused on which diseases, risk factors and symptoms are included in varying definitions of multimorbidity [Willadsen et al., 2016]. Whilst the majority of included studies in the review indicated multimorbidity as the presence of two or more conditions, Willadsen et al [2016] found the total number of diseases, risk factors, and symptoms used varied from 4 to 147. Of the 167 included articles in the review, 115 different ways of defining multimorbidity were identified [Willadsen et al., 2016].

In a recently published guideline, the National Institute for Health and Care Excellence (NICE) [NICE, 2016] acknowledge the complexity of defining multimorbidity. NICE agree with other commentators [Mercer et al., 2009] that basing the definition of multimorbidity on 2 or more health conditions only does not fully capture a clinically

meaningful picture of the concept. The guideline highlights the fact that many people defined as multimorbid in this way may not be ill and have excellent quality of life requiring little or no health care input [NICE, 2016]. For this reason the guideline is aimed at people with more than 1 long-term condition with any of the following:-

- Difficulty managing treatments or day-to-day activities.
- Care from multiple services and requiring care from a new service.
- Both long-term physical and mental health conditions.
- Frailty.
- Frequent use of unplanned or emergency care.
- Prescription of multiple, regular medicines.

[NICE, 2016]

Although multimorbidity may seem to be an intuitive thing to understand, defining a useful concept of the term is much more difficult [Guthrie et al., 2011]. Whilst the co-occurrence of 2 or more long-term conditions is the most commonly used definition of multimorbidity, there are wide variations in the number of conditions from which this definition can be based. This lack of clarity means that identifying useful ways to measure Multimorbidity is also problematic with marked variations in approaches.

2.4.3 Measurement

The findings of three recent systematic reviews have highlighted the myriad ways researchers have approached the measurement of multimorbidity [de Groot et al., 2004, Diederichs et al., 2011, Huntley et al., 2012]. Each review aimed to collate evidence of measurement tools in comorbidity or multimorbidity but from different perspectives: De Groot et al [2004] searched for comorbidity indices to inform research into Multiple Sclerosis, Diederichs et al [2011] specifically searched for multimorbidity measurement indices, whereas Huntley et al [2012] searched for measures of multimorbidity used only in primary care research. The systematic reviews found 13, 39 and 17 exclusive ways of measuring multimorbidity or comorbidity respectively. The number of medical conditions included in these measurements varied from 4 to 102 [2011]. Most indices are developed from secondary care populations but many have been adapted for other populations including primary care [Diederichs et al., 2011, Huntley et al., 2012].

There are two main ways of measuring multimorbidity: simple disease counts or using an index which applies weights to either prescribed medications or medical conditions and other factors in an attempt to explain severity of illness [de Groot et al., 2004, Diederichs et al., 2011, Huntley et al., 2012]. In primary care research, the most frequently used measurement is simple disease counts [Huntley et al., 2012]. This may be because of the ease with which it can be administered compared to more complex indices such as the Charlson index [Charlson et al., 1987] or Chronic Disease Score [Von Korff et al., 1992] and their variations. Despite the large number of multimorbidity indices available, Huntley et al [2012] cite evidence that suggests simple counts of diseases or medications are almost as effective as the more complex indices at predicting mortality or health care use in the primary care setting. However, when aiming to predict mortality in Primary Care, Huntley et al [2012] recommend the best measurement of multimorbidity to be provided by the Charlson index [Charlson et al., 1987] and its variations. Measurement of multimorbidity in relation to primary care healthcare use can be predicted with equivalence by either; the Adjusted Clinical Group system [Starfield et al., 1991], the Charlson index [Charlson et al., 1987], or disease counts [Huntley et al., 2012]. Disease counts were also found by Huntley et al [2012] to have good evidence to suggest they provide a robust measure of multimorbidity in relation to Quality of Life, as does the Charlson index [Charlson et al., 1987]. A count of medicines was found to be a good predictor of primary care use and mortality in a more recent paper [Brilleman and Salisbury, 2013]. In their paper, Perkins et al [2004] argue that indices developed in the secondary care setting, such as the Charlson index, should be used with caution in other settings despite adaptations. More recently, Wallace et al [2016] found little difference between simple (count) and complex (index) measures when predicting hospital admission but noted that all measures of multimorbidity alone were poor predictors of the outcome.

An emerging method of measuring multimorbidity is to identify clusters of medical conditions that co-exist in individuals at rates higher than would be expected - or non-random prevalence. Recent research and academic discussion suggests identification of disease clusters may enable clearer answers to clinically relevant research questions than currently employed measures [Le Reste et al., 2015; RN109; Prados-Torres et al., 2014; Sinnige et al., 2013; Holden et al., 2011; Marengoni et al., 2011; Valderas et al., 2009]. Statistical techniques employed in attempts to identify such clusters include: factor analysis, cluster analysis, the observed-to-expected ratio, multiple correspondence

analysis [Prados-Torres et al., 2014, Clerencia-Sierra et al., 2015], principal component analysis, latent class analysis [Islam et al., 2014, Larsen et al., 2017], and machine learning techniques [Schiltz et al., 2017]. In their systematic review of clustering methods, Prados-Torres et al [2014] found wide variations in approaches to clustering and characteristics of populations studied. As opposed to many of the studies included in the review, they recommend future attempts at clustering diseases use: population-sized datasets, statistical techniques that are suited to the dichotomous nature of diagnostic variables, and large numbers of conditions from which to form clusters [Prados-Torres et al., 2014]. Prados-Torres et al [2014] identified three groups of patterns common to all included studies in their review despite marked heterogeneity namely; cardiovascular and metabolic diseases, mental health conditions, and musculoskeletal disorders. Whilst identification of groups may have some benefit in terms of identifying causal mechanisms between diseases, whether they are useful or meaningful in clinical terms is a matter of debate.

2.4.4 Epidemiology

Sections 2.4.2 and 2.4.3 describe the wide variations in definitions and measures of multimorbidity. It is, therefore, unsurprising that there is marked heterogeneity in reports of multimorbidity prevalence. Fortin et al [2012] illustrate this by reporting variations in the prevalence of multimorbidity from 3.5% to 98.5% across 21 studies included in their systematic review. The variation in findings is explained by the vastly different populations, settings, data collection techniques, and definitions of multimorbidity used by included studies. A more recent systematic review concentrating on primary care populations and aiming to describe prevalence, causes and patterns of multimorbidity [Violan et al., 2014] found reports of multimorbidity prevalence between 12.9% and 95.1%. Similar variations in definitions, measures and populations were found. The number of conditions used to estimate multimorbidity prevalence varied between 5 and 335 [Violan et al., 2014]. In an attempt to standardise conditions to be considered using international disease classification labels, a more recent paper included 60 conditions [Calderón-Larrañaga et al., 2016]. Van den Akker et al [-RN91] highlighted the complications that can arise when attempting to measure prevalence of multimorbidity and suggest that certain decisions made in study design will depend on the specific question being interrogated by researchers (e.g. the number of diseases

to include in the measure of multimorbidity or the age-range of the sample). The systematic reviews of Violan et al [2014] and Fortin et al [2012] may reflect the varying decisions made by research teams in study design. Despite the difficulties in synthesizing evidence on heterogeneous studies, Violan et al [2014] found strong relationships between multimorbidity and: age, female gender, low socioeconomic status, and mental health across studies in their review.

2.4.5 Summary

Multimorbidity is most commonly defined as the presence (or co-occurrence) of 2 or more long-term conditions in an individual. Debate continues as to the type and number of long-term conditions that should be included to provide a meaningful concept for individuals, clinicians and healthcare organisations. The lack of a standard definition is mirrored in the myriad ways of measuring multimorbidity with various counts, indices, and clusters. Despite this, evidence suggests multimorbidity is increasing in prevalence and has a strong socioeconomic pattern. As a result, policy needs to be tailored to account for the complex needs of the increasing numbers of people with multimorbidity.

2.5 Conclusion

Conclusion of chapter 2

Access to social care is influenced in 2 main ways - allocation of resources *to* providers and distribution of services *within* providers. Theory and evidence suggest inequitable access possible.

Major policy shift to integrate health and social care services internationally. One of main drivers increase in older population - services currently disjointed for this group. Two-thirds over 65s have multimorbidity and associated poorer outcomes - with social gradient.

Understanding how this population access social care, therefore, vitally important and under-researched.

Does an inverse *social* care law exist? i.e. Does the allocation of resources (via funding

formulae) to Local Authorities negatively impact on those areas with higher need?

Furthermore, does access to social care vary across Local Authorities - is there a “postcode lottery” in terms of service provision i.e. does application of eligibility criteria depend on where you live?

Is multimorbidity status associated with levels of social care provided within *and* across local authorities?

Important to understand how access to social care influences health care use and mortality - do those with multimorbidity and social care have different outcomes from those with multimorbidity and no social care?

Research questions:-

In people over the age of 65 in Scotland:

1.
 - a. What are the socioeconomic, demographic, and geographic patterns in the use of social care?
 - b. Is there an association between multimorbidity status and the amount and type of social care use over time? Does this vary by the patterns described in 1(a)?
2.
 - a. Is there an association in the use of social care services, multimorbidity status and unscheduled health care use?
 - b. Do multimorbidity status and social care use predict mortality?

Chapter 3

Methods

3.1 Overview

- Discussion of methodological approach. Quant v Qual etc.
- Reference Lit review and describe how Research Questions more suitable for Quant approach

3.2 Administrative data research

3.2.1 Overview

- Broad discussion of admin data research

3.2.2 Information Governance

- Sensitive data - how protected
- eDRIS and SG policy docs

3.3 Data Sources

- List and describe data sources

3.4 Data linkage

- Describe linkage process

3.5 Timeline

Outline time frames of:-

- Approvals process
- DSA
- Linkage

for both Renfrew and linked data

Discuss barriers and setbacks.

Chapter 4

Measuring Multimorbidity

4.1 Introduction

As Chapter 2 showed, multimorbidity can be defined as the presence of two or more chronic health conditions within an individual [NICE, 2016]. It is associated with higher mortality [Gijssen et al., 2001], increased use of health care [Gijssen et al., 2001, Salisbury et al., 2011], psychological distress [Fortin et al., 2006], worse quality of life [Fortin et al., 2004, 2005], and worse functional status [Kadam and Croft, 2007]. It not only affects older people but has been observed in greater absolute numbers in those under the age of 65 and affects those with lower socioeconomic status disproportionately [Barnett et al., 2012]. As the proportion of older people in the population increases, multimorbidity is expected to affect increasing numbers of people in the future [Guthrie et al., 2011, Imison, 2012].

Many of the negative outcomes associated with multimorbidity are due to the structure of healthcare delivery which tends to concentrate treatment goals towards single diseases [Guthrie et al., 2011, Starfield et al. [2005]]. As a result healthcare for individuals with multimorbidity can, at best, preferentially treat one disease to the detriment of others or, at worst, cause harm and affect patient safety e.g. through medication interactions [May et al., 2009].

Epidemiological research into multimorbidity has tended to count numbers or report proportions of health conditions. There is, however, marked heterogeneity in the population considered and the number of diseases included in measurement. As a result,

prevalence rates of multimorbidity have been shown by systematic reviews to vary widely in the general population from 13.1% - 71% [Fortin et al., 2012] and 12.9% - 95.1% [Violan et al., 2014]. There has been little research into the prevalence of combinations of diseases and non-random, co-occurrence of diseases partly due to the high number of theoretical combinations meaning large samples and complex calculations are required [van den Akker et al., 2001, Cornell et al., 2009]. More recent research has applied a variety of statistical techniques to overcome this problem [Prados-Torres et al., 2014].

Identification of non-random co-occurrence of health conditions is important for a number of reasons; a) to gain a better understanding of the complicated nature of multimorbidity, b) to help assess the impact multimorbidity has on health outcomes, c) to help assess the sociodemographic differences in prevalence of multimorbidity which could have implications for health policy and the delivery of health services, and d) unexpected non-random associations could prompt further research into possible causal mechanisms for the association.

The objective of this chapter was to apply a novel two-way clustering framework to a large dataset based on the Scottish population, The framework aims to identify clusters of the most significant non-random co-occurrence of health conditions and multimorbidity patterns among individuals [Ng, 2015]. The specific aims were to a) Identify non-random associations of health conditions from the dataset, b) identify if meaningful, homogeneous sub-groups of individuals according to groups on non-random conditions can be formed, c) assess the sociodemographic make-up of identified clusters.

Add para describing purpose of this chapter to overall thesis

4.2 Background

Previous clustering attempts in the field of multimorbidity have employed a number of different statistical techniques including; factor analysis, cluster analysis, the observed to expected ratio, multiple correspondence analysis [Prados-Torres et al., 2014], principal component analysis, and latent class analysis [Islam et al., 2014]. Each of these techniques is a variant of latent variable modelling where clusters are deemed unobserved, or latent, variables that can be measured indirectly depending on the values of two or more observed variables [Collins and Lanza, 2013]. In their systematic review of

clustering techniques, Prados-Torres et al [2014] found marked heterogeneity in the numbers of diseases included in analyses, populations studied, and resulting clusters of conditions. The authors recommended that future attempts at clustering should be conducted using large numbers of diseases and in population-based datasets as opposed to sub-groups of populations.

An important decision in any clustering research is which statistical technique to apply to a given dataset. Collins & Lanza [2013] argue that when the latent variable (cluster) and the observed variables used to identify the latent variable are categorical in nature, Latent Class Analysis (LCA), a version of a finite mixture model, is an appropriate statistical technique to employ. Other techniques such as factor or cluster analysis rely on latent and/or observed variables being continuous in nature [Collins and Lanza, 2013]. The aim of this chapter is to identify whether homogeneous sub-groups of individuals can be identified from a dataset of diseases represented by binary, categorical data. Such sub-groups would also be categorical in nature suggesting LCA is an appropriate technique to apply.

The limitation to LCA is that as the number of observed variables increases it becomes more difficult for the LCA model to be well identified [Collins and Lanza, 2013]. This is because the first step of LCA is to create a contingency table of all possible combinations of outcomes. For example, in a simple dataset with two binary Yes/No variables there are $2^2 = 4$ potential response outcomes; No/No, Yes/No, No/Yes, and Yes/Yes. In the example of the dataset used in the current chapter with 40 health conditions recorded as binary variables the number of cells in the contingency table will contain 2^{40} (over 1 trillion) cells. This makes good model identification highly unlikely. Ideally the ratio of n/W where n denotes the sample size and W denotes the number of potential response outcomes should be as high as possible [Collins and Lanza, 2013]. When this ratio is very small there are only two options to overcome the problem, "... increase the amount of known information or reduce the amount of unknown information." [Collins and Lanza, 2013, :93]

In response to the "high-dimensional" problem created when trying to identify latent variables from health datasets with large numbers of diseases, Ng [2015] proposed a two-way model to identify multimorbidity clusters. The first step of this method is to "clump" diseases in the dataset into statistically correlated groups of conditions thereby reducing the number of variables and therefore the size of contingency table of

potential responses. This first step is described in more detail in Ng et al [2012]. The second step of the method then aims to identify latent groups of individuals based on response patterns to these “clumped” groups with a finite mixture model similar to LCA [Ng, 2015]. This technique was applied to an Australian national health survey which contained self-reported responses of the presence of 24 physical and mental health conditions with full results detailed in Ng [2015]. The aim of this chapter is to identify whether this technique is valid in a much larger dataset of administrative health data.

4.3 Methods

4.3.1 Data

In Scotland, much multimorbidity research has been informed by the Scottish Programme for Improving Clinical Effectiveness in Primary Care (SPICE-PC) dataset [Agur et al., 2016, Barnett et al., 2012, Elder et al., 2007, McLean et al., 2015]. Diagnostic data from the year 2007 is available on 1,754,133 people in Scotland drawn from 314 general practices. The anonymised dataset has information on the presence of 32 physical and 8 mental health conditions (?? add box with list of diseases??) in addition to age, gender and deprivation index data. Diagnostic data is derived from codes entered into IT systems in General Practices and prescription data. A full description of the 40 included diseases and the methods used to classify them is available as supplementary information in Barnett et al [2012]. The analysis for this chapter was restricted to adults over the age of 18 resulting in n=1,426,196. (10% sample of this so far!!) Ethical approval for secondary analysis of the SPICE-PC dataset using LCA was granted by the Research Ethics Committee of the College of Social Sciences at the University of Glasgow on 29/04/2016.

4.3.2 Analysis

The two-way method proposed by Ng [2015] was applied to the above dataset of 40 health conditions. Firstly groups of conditions that co-occur are identified and then latent groups of individuals are identified using a mixture model approach. To identify the groups of conditions that co-occur the method aims to calculate the significant

pairwise multimorbidity between all conditions. The asymmetric Somer’s D statistic was used to quantify the degree of random co-morbidity with all pairs of health conditions. Significance of the Somer’s D statistic for each pair of conditions was calculated using the Benjamini-Hochberg procedure [Benjamini and Hochberg, 1995] to control the false discovery rate (FDR) with $\alpha = 0.001$. Diseases were then clustered into overlapping groups using a “clumping” procedure based on the technique described by Jardin & Robson [1968] and fully described as equation 11 in Ng et al [2012]. The strength of multimorbidity in each cluster was calculated using the average pairwise Somer’s D statistics of disease within the group. From these overlapping groups, non-overlapping groups of diseases were created using an amended version of the algorithm specified in Ng [2015]:-

1. Name the cluster with the highest strength as the first group and then remove its member health conditions in all subsequent clusters with smaller strength. Each member in a cluster must have the condition with which the pairwise concordance statistic is maximum in the same cluster;
2. Repeat (1) for the next cluster and name it as a group if it is not ‘singular’ (singular cluster is defined as a cluster consisting of a single health condition);
3. If a group is formed in (2), remove its member health conditions in all subsequent clusters with smaller strength or singular clusters;
4. Repeat (2) and (3) until all clusters are visited;
5. Put the condition in a singular cluster into a predefined group where more than half of the member conditions are significantly comorbid with the condition;
6. Name those remaining singular clusters as a singular group.

A matrix was then created assigning a score to individual observations for each identified non-overlapping group. Scores were assigned as:-

- 1 for having no diseases within the group
- 2 for having one of the diseases in the group
- 3 for having 2 or more diseases within the group.

In the second part of the analysis a mixture-model of multivariate generalised Bernoulli distributions was applied to the matrix of these scores in order to identify latent groups of individuals according to response patterns to the score matrix. The most appropriate number of latent groups to fit the dataset was determined by the lowest Bayesian

Information Criterion (BIC) and substantive theoretical analysis. Models with 3 to 9 latent groups were compared to find best model fit. Each observation was then assigned to the latent group for which the posterior probability was highest. Descriptive analysis of sociodemographic difference in latent group was then employed to identify patterns in groupings.

All statistical analysis was completed using R version 3.3.3 [R-Core-Team, 2017]. Two-way clustering was conducted with amended code provided by Ng[2015]. Data manipulation and visualisation was conducted using tidyverse packages [Wickham, 2017] and the corrplot package.

4.4 Results

4.4.1 Pairwise correlation

With α controlled at 0.001, 143 of a possible 780 pairwise correlations proved to be significant. A matrix showing all statistically significant correlations is shown in Figure 4.1. The number of expected false positives for 143 pairs is less than 1.

4.4.2 Grouping diseases

Using the “clumping” procedure detailed in Ng [2015], Thirty-eight overlapping groups of conditions were found as shown in Table 4.1

Six diseases; Epilepsy, Learning Disability, Sinusitis, Crohns, Anorexia, Psoriasis/Eczema did not have strong enough pairwise correlations to be included in any of the 38 groups. Thirteen non-overlapping groups were derived from the results of the clumping method using the algorithm described above and named according to the characteristics of the member disease in the groups as shown in Table 4.2.

A further four diseases; Diverticular disease, Prostate, IBS, and Dyspepsia were excluded from the non-overlapping groups. These four conditions did not appear in any groups with the diseases with which they had strongest pairwise correlation, a condition that had to be met in the first stage of the algorithm described above.




Figure 4.1: Significant Pairwise Correlations

Table 4.1: Overlapping groups of diseases derived from "clumping" procedure

Group	Diseases Included	Strength
1	CHD, CKD, AtrialFib, Hypertension, HeartFailure, Rheu-Arthritis, PVD	0.2833
2	CHD, CKD, AtrialFib, Diabetes, TIA-Stroke, Hypertension, HeartFailure	0.2806
3	CHD, CKD, Hypertension, HeartFailure, Rheu-Arthritis, Pain	0.2653
4	CHD, Hypertension, Prostate	0.2622
5	CHD, CKD, Diabetes, TIA-Stroke, Hypertension, HeartFailure, Pain	0.259
6	CHD, TIA-Stroke, Hypertension, HeartFailure, Laxat-Constip, Pain	0.2525
7	Mental-Alcohol, LiverDisease, Depression, Anxiety	0.2512
8	CHD, Hypertension, HeartFailure, Bronchitis, Pain	0.2487
9	CHD, CKD, Diabetes, TIA-Stroke, Hypertension, PVD, Pain	0.2348
10	CHD, CKD, Hypertension, PVD, Rheu-Arthritis, Pain	0.2331
11	Mental-Alcohol, Mental-Psycho, ViralHepatitis, Depression, Anxiety	0.2324
12	Mental-Psycho, Depression, Schiz-Bipolar, Anxiety	0.2304
13	CHD, Hypertension, Diverticular, Laxat-Constip, Pain	0.2294
14	Depression, Migraine, Pain	0.2278
15	CKD, ThyroidDisorders, Hypertension, Pain	0.2262
16	CHD, Blindness, Glaucoma, Hypertension	0.2236
17	Depression, Dyspepsia, Laxat-Constip, Pain, Anxiety	0.218
18	MS, Depression, Laxat-Constip, Pain	0.2178
19	CHD, Diabetes, TIA-Stroke, Blindness, Hypertension, Pain	0.2171
20	ActiveAsthma, Bronchitis, Bronchiectasis	0.2166
21	CHD, TIS-Stroke, Blindness, Hypertension, Laxat-Constip, Pain	0.2138
22	CHD, Hypertension, PVD, Bronchitis, Pain	0.2114
23	CHD, Dementia, Parkinsons, TIA-Stroke, Hypertension, Laxat-Constip, Pain	0.2068
24	Dementia, Parkinsons, Depression, Laxat-Constip, Pain, Anxiety	0.2068
25	CHD, Dementia, Hypertension, Rheu-Arthritis, Pain	0.204
26	Diabetes, Hypertension, LiverDisease, Pain	0.1964
27	Dementia, Mental-Psycho, Depression, Pain, Anxiety	0.1939
28	Hypertension, Diverticular, Dyspepsia, Laxat-Constip, Pain	0.1938
29	CHD, Dementia, Parkinsons, Hypertension, Laxat-Constip, Pain, Anxiety	0.1922
30	Hypertension, Bronchitis, Bronchiectasis, Pain	0.1917
31	Blindness, HearingLoss, Hypertension	0.179
32	Dementia, Parkinsons, TIA-Stroke, Depression, Laxat-Constip, Pain	0.173
33	HearingLoss, Hypertension, Prostate	0.1729
34	AnyCancer-Last5Yrs, Hypertension, Pain	0.1707
35	Hypertension, LiverDisease, Dyspepsia, Pain, Anxiety	0.1593
36	Depression, IBS, Pain	0.1563
37	CHD, Dementia, Mental-Psycho, Pain, Anxiety	0.1385
38	Diverticular, IBS, Pain	0.1226

Table 4.2: Non-overlapping groups of diseases derived from Ng algorithm

Disease group name	Diseases included
<i>Cardiovascular</i>	CHD, CKD, AtrialFib, Hypertension, HeartFailure, Rheu-Arthritis, PVD
<i>Diabetes/Stroke</i>	Diabetes, Stroke
<i>Pain/MS</i>	Laxatives/Constipation, Pain, MS
<i>Mental Health/Liver</i>	Mental/Alcohol, Liver Diseases, Depression, Anxiety
<i>Psychosis</i>	Mental-Psycho, Viral Hepatitis
<i>Sciz/Bipolar</i>	Sciz-Bipolar
<i>Migraine</i>	Migraine
<i>Thyroid</i>	Thyroid disorders
<i>Blindness</i>	Blindness, Glaucoma
<i>Respiratory</i>	ActiveAsthma, Bronchitis, Bronchiectasis
<i>Neurodegenerative</i>	Dementia, Parkinson's
<i>Hearing Loss</i>	Hearing Loss
<i>Cancer</i>	Any Cancer in Last 5 years

4.4.3 Grouping individuals

Individuals were assigned a score depending on the number of diseases they had in each of the non-overlapping groups according to the criteria described above. The finite mixture model of multivariate generalised Bernoulli distributions was then applied to this dataset in order to identify latent groups of individuals depending on their scores for each of the 13 groups, based on presence of 30 diseases. BIC scores suggested the model for nine latent groups had best fit to the data. The nine-group model was compared with the model with second-lowest value of BIC, that for eight latent groups. However, despite offering a more parsimonious solution, the eight-group model did not provide a more substantive theoretical fit to the data. Thus, the nine-group model was deemed best fit and analysed further.

Item-response probabilities for the nine-group model are shown in Figure 4.2.

Each latent group was labelled according to these item response probabilities as follows;

- Latent Group 1. **Well**. High probabilities of having no diseases in any of the groups.
- Latent Group 2. **Cardiovascular only**. Item response in this Latent group have more than 50% chance of having at least 1 condition from the Cardiovascular group. High probabilities of having no diseases in any of the other groups.
- Latent Group 3. **Cancer**. Individuals in this group have almost 100% probability



Figure 4.2: Item response probabilities for nine latent group model

of having had cancer in last 5 years. Weak probabilities across some other groups.

- Latent Group 4. **Mental Health/Pain** High probability of having at least one diseases from Pain/MS group and at least one disease from Mental Health/Liver group.
- Latent Group 5. **Mental and Physical Multimorbidity** High probabilities of having at least 1 cardiovascular, diabetes/stroke, Pain/MS, and Mental health/Liver disease. This group also has the highest probability across latent groups for having 1 respiratory disease.
- Latent Group 6 **Physical Multimorbidity** High probability of a least one cardiovascular disease and one of Diabetes/Stroke.
- Latent Group 7. **Physical multimorbidity (Strong Cardio)**. Similar to Latent group 6 but individuals in this group are more likely to have 2 or more cardiovascular diseases.
- Latent Group 8. **Dementia with Mental/Physical MM** Individuals in this latent group have almost 100% probability of having dementia or Parkinson's. Also highly likely to have at least one Cardiovascular disease and mildly raised probability of having mental health or psychosis diseases.
- Latent Group 9. **Mental Health only**. High probability of having a least 1 of the Mental Health/Liver group diseases. Also mildly raised probabilities in the Psychosis and Sciz_Bipolar groups.

Proportionate size of each of the assigned latent variable groups is shown in Figure 4.3.

The “Well” latent group was by far the largest group with almost 70% of individuals within it. The other groups formed much smaller proportions of the data.

4.4.4 Sociodemographic breakdown of latent groups

Each of the Histograms of Age for each of the latent groups shown in Figure 4.4 show variation in distribution. The “Well” and “Mental Health only” groups have a much younger age distribution compared to the other groups. The “Dementia” group has a much older distribution.

Figure 4.5 shows higher proportions of males in the “Physical MM” latent group. The “Well” and “Physical MM high Cardio” groups have even splits by gender. All other groups have higher proportions of females.




Figure 4.3: Latent Group Proportions




Figure 4.4: Histograms of Age, by Latent Group

Figure 4.6 shows distribution of deprivation across latent groups by Carstairs Decile. There are clear gradients in the “Mental Health & Pain”, “Ment/Phys MM”, and “Mental Health only” groups with much higher numbers of people in Decile 10 (most deprived) areas.




Figure 4.5: Count of Sec, by Latent Group - with Proportions




Figure 4.6: Count of Carstairs Decile, by Latent Group

4.5 Discussion

When proposing the two-way clustering method, Ng [2015] applied the technique to Australian survey data of 24 self-reported diseases from $n=8841$ respondents. These were “clumped” into nine non-overlapping groups of diseases with four latent groups of individuals being identified by the mixture model. Given the different classification of diseases, the nature of how data were collected, and the different populations, it is unsurprising that results from the present study are markedly different. What is of interest is whether the technique provided a meaningful representation of the data which can be used to improve the understanding of multimorbidity within this population.

The first step of the two-way clustering technique resulted in 13 non-overlapping groups being formed. As a result of this process, ten of the diseases recorded in the dataset were excluded from the final groups. Six diseases; Epilepsy, Learning Disability, Sinusitis, Crohns, Anorexia, and Psoriasis/Eczema did not have strong enough pairwise Somer’s D correlation statistics with any other disease to be included in the non-overlapping “clumping” procedure. A further four diseases; Diverticular, Prostate, IBS, and Dyspepsia did not appear in any non-overlapping group with the disease with which they had the strongest Somer’s D correlation. Identifying which diseases do not co-occur at non-random levels is a particularly interesting finding and adds to the debate about which diseases should be included in any multimorbidity measure.[be explicit – do you think this suggests we shouldn’t count these 10 in any MM measure and, if so, why? -NB] Of the 40 diseases present in the SPICE-PC dataset, in this analysis, only 30 diseases co-occurred with enough statistical power to be added to non-overlapping disease groups and considered in the mixture model.

Past research has highlighted associations between; mental health and thyroid problems, mental health and pain, and diseases associated with the metabolic syndrome [Prados-Torres et al., 2014]. As shown in figure 1, there is no pairwise correlation between any mental health diseases and thyroid disorders in the present study. Pain is associated with a large number of conditions, including mental health diseases. Diseases associated with metabolic syndrome such as, hypertension, coronary heart disease, diabetes, stroke, and heart failure all show correlations with each other to varying degrees. Prados-Torres et al [2014] also identified associations between Chronic Obstructive Pulmonary Disease (COPD) and Gastroesophageal reflux disease (GORD) with mental health diseases. No associations with between any respiratory disease and mental health diseases is apparent

in figure 1. The SPICE-PC dataset does not record GORD, however Dyspepsia may be considered a similar diagnosis and has pairwise correlations with both Depression and Anxiety.

The non-overlapping disease groups were named according to the characteristics of the diseases present in each, however many diseases appeared in groups that may not come from similar aetiology. The groups reflect diseases that commonly co-occur and therefore do not always fit into clinically recognisable groups. Individuals are scored as to the number of diseases they have from each group. This results in a loss of some information but the advantage of this method is that it reduces the number of variables in the dataset making the likelihood of identifying a good mixture model much more likely.

Applying a mixture model of multivariate generalised Bernoulli distributions to these 13 non-overlapping disease groups identified nine latent groups as described above. These groups are clinically recognisable with some showing presence of diseases from only one disease group and others illustrating a more multimorbid population. Clear distinctions between mental and physical disease are made enabling individuals to have diseases from both groups. In their systematic review of clustering studies of multimorbidity, Prados-Torres et al [2014] found that included studies reported between three and twenty diseases clusters. From these they found three most common groups of diseases; cardiovascular, mental health, and musculoskeletal. Latent groups in the current study also contain cardiovascular and mental health elements, although they do not reflect any groups that could be identified as musculoskeletal.

Clear sociodemographic patterns were identified in the latent groups. Those classified as belonging to a latent group with mental health involvement such as; mental health/pain, mental and physical multimorbidity, or mental health only, were more likely to be female and from lower deprivation deciles. Two similar latent groups; physical multimorbidity and physical multimorbidity high cardiovascular, had clear age differences with the former more likely to have younger male individuals assigned to it. These findings are similar to simple analyses of the same dataset which identified those in the most deprived areas being more likely to develop physical and mental health multimorbidity 10-15 years earlier than their more affluent peers [Barnett et al., 2012].

4.5.1 Limitations

The two-step method proposed by Ng [2015] enables reduction of dimensions in datasets with large numbers of disease variables making model good identification of mixture models more likely. This, however, results in a loss of detail making interpretation of results more difficult. Report of the item-response probabilities shown in figure 4 identifies that individuals have a probability of having none, one, or two of the diseases in any group. It is impossible to identify exactly which diseases. Hypertension with 13.4% of individuals is the highest reported disease in the SPICE-PC . To what degree this accounts for individuals having one disease in the cardiovascular group across latent classes is unknown.

This is particularly a problem as the non-overlapping groups formed by the two-way clustering method do not always follow recognised clinical groupings. For example, Rheumatoid arthritis is found in the cardiovascular group despite being a connective tissue disorder. It is the only disease in the group that does not directly affect the cardiovascular system. The reason it is found in this group may be due to the association of increased NSAID use with cardiovascular outcomes (ref) resulting in frequent co-occurrence. As some latent groups are classified as “Cardiovascular only”, there is a possibility that small numbers of people with only Rheumatoid Arthritis are misclassified. [This is could be down to nomenclature and I maybe need to revisit group names. It does not get rid of the problem that for the group with high probs for the cardio/rheum disease we can’t distinguish which disease the individual has]

When clustering individuals, fitting a finite mixture model of multivariate generalised Bernoulli distributions makes the assumption that the indicator variables used to identify latent groups are independent. In a dataset containing medical conditions such as e.g. Hypertension and Coronary Heart Disease, or Atrial Fibrillation and Stroke, this assumption is clearly violated. However, ignoring the independence assumption for multivariate categorical variables often results in better fit than when more complicated techniques are applied to account for non-independence of indicator variables [Ng, 2015, Hand and Yu, 2001, Topchy et al., 2005].

Results presented here account for 10% of the SPICE-PC dataset due to the heavy computational requirements of fitting the finite mixture model. Confirmation of results on the whole dataset is required.

4.5.2 Future research

Sociodemographic comparison in the current study was limited to visualisation of histograms. More detailed reports of measures of central tendency and comparison of latent groups with sociodemographic variables such as Carstairs decile with logistic regression is warranted. SPICE-PC data also contains variables on lifestyle factors such as smoking status and alcohol intake. These variables should be included in further comparisons.

A further calculation offered in the two-way clustering technique [Ng, 2015] is the calculation of a multimorbidity score to each latent group and to each individual in the dataset based on their disease profile and posterior probabilities. Such a score would be continuous in nature and would offer the benefit of comparison with sociodemographic variables such as Carstairs score or Age. Such comparisons would be amenable to well-recognised regression techniques.

Part 1 of the two-step method involves reducing the dimensions of the dataset to be amenable to mixture modelling. Using the outcome of this first step, the non-overlapping groups, Ng's technique for clustering individuals should be compared to Latent Class Analysis which is also a suitable technique for the nature of the data [Collins and Lanza, 2013]. In supplementary material, NG [2015] compared his two-step method with Latent Profile Analysis and found it inferior to the two-step method. Latent Profile Analysis is a mixture model for continuous indicator variables [Collins and Lanza, 2013] and would not be suitable for the SPICE-PC dataset.

4.6 Conclusion

A novel two-step method for clustering health conditions and individuals was applied to large, population representative, dataset of administrative primary care data. The method found 10 of the 40 conditions in the dataset did not co-occur with other diseases strongly enough to be included in further analysis. Thirty conditions were “clumped” into 13 groups of commonly co-occurring conditions and individuals in the dataset were assigned a score depending on the number of diseases within each group they had. From this information nine latent groups of individuals were identified with a finite mixture model. These groups reflected varying degrees of physical, mental and physical/mental

morbidity and showed sociodemographic patterning.

Chapter 5

Renfrewshire Council Pilot Project

5.1 Introduction

Pilot project to get first look at social care data to determine structure and definitions. Main linkage project to use cross-sectional social care data - is this reflective of social care provision throughout the year? Also able to attach datazone and demographic variables - is there variation in amounts of care across these variables?

5.1.1 Aims

- Describe variation in social care provision in Renfrewshire Council area by Age, Gender, and socioeconomic position.
- Describe the temporal variation in levels of care within financial years
- Assess the quality of social care data for research purposes.

5.1.2 Research Questions

1. Is there variation in the amount of social care delivered in the Renfrewshire Council area by Age, Gender, and Socioeconomic position?
2. How does social care use vary within and across financial years 2006/07 - 2015/16?

5.2 Background

Link to SCS and census methods. Any relevant lit of social care analysis (Not much). Funders interested in these questions etc.

5.3 Methods

Link to approval process described in chapter 3 section 3.5. Also link to use of safe haven described in section 3.2.2. Re-mention ethics.

Describe manipulation of variables to produce summary stats.

Describe which methods used to summarise variation in care

5.4 Results

Tables and graphs of results with description

5.5 Discussion

Implications - particular focus on variation in care throughout the year and whether care provided in a census week is a robust measure for research purposes (based on Renfrew results)

Limitations - One council only. Different collection and reporting methods across LAs etc etc.

5.6 Conclusion

Chapter 6

Linked Health and Social Care Data: Quality Assessment

Results chapter describing quality of data (in particular SCS)

??????? Still going to include this??

6.1 Intro

6.2 Background

6.3 Methods

6.4 Results

6.5 Discussion

6.6 Conclusion

Chapter 7

Access to social care

Results chapter for 1st research question

7.1 Introduction

Link to chapter 2 re access to social care. Linked dataset to assess national variation in social care provision

7.1.1 Aims

- Assess how access to social care services varies for people with multimorbidity, especially by socioeconomic status.

7.1.2 Research questions

In people over the age of 65 in Scotland:-

1. What are the socioeconomic, demographic, and geographic patterns in the use of social care across Scotland?
2. Is there an association between multimorbidity status and the amount and type of social care use over time? Does this vary by the patterns described in question 1?

7.2 Background

Mention previous linkages of health and social care identified in section [2.3.3](#).

7.3 Methods

- Link to chapter [3](#)
- Describe social care variables
- Describe PIS variables and summaries
- Describe socioeconomic and geographic sources

7.4 Results

7.5 Discussion

7.6 Conclusion

Chapter 8

Social care & Health outcomes

Results chapter for 2nd research question.

Need to find new title - must convey effect of sc **AND** mm on USC **AND** mortality

8.1 Introduction

- Social care believed to impact on use of unscheduled care
- New models of care important to find
- Understanding interaction between services essential to identify strategies to improve services
- Impact of social care on mortality never studied - multimorbidity an important confounder

8.1.1 Aims

- Assess the impacts of social care service use on unscheduled care use and mortality for people with multimorbidity, accounting for age, gender, and socioeconomic position.

8.1.2 Research Questions

1. Is there an association in the use of social care services, multimorbidity status and unscheduled health care use?
2. Do multimorbidity status and social care use predict mortality?

8.2 Background

8.3 Methods

8.4 Results

8.5 Discussion

8.6 Conclusion

Chapter 9

Conclusion

Reccomendations :-

- Standardised score reflecting need/frailty/vulnerability required to accurately assess access to care (OECD pp181)
- disaggregate SCS a little - distinguishing reablement and intermediate care from LTC important!
- Measure outcomes relevent to social care (falls etc OECD 2013 pp59) OR link to outcome data e.g. preventable admission data, hip fracture data??
- Consider other forms of admin data to help with this e.g. Attendance Allowance??
 - Problem being coverage

Appendix A

Title A

Testing, testing. One-two, one-two.

Appendix B

Title B

Let's see what this looks like

Bibliography

- Caroline Abrahams, Marcus Green, and Jill Mortimer. Care in crisis 2014. Report, Age UK, 2014. URL http://www.ageuk.org.uk/Documents/EN-GB/Campaigns/CIC/Care_in_Crisis_report_2014.pdf?epslanguage=en-GB&dtrk%3Dtrue.
- Karolina Agur, Gary McLean, Kate Hunt, Bruce Guthrie, and Stewart W Mercer. How does sex influence multimorbidity? secondary analysis of a large nationally representative dataset. *International journal of environmental research and public health*, 13(4):391, 2016. URL <http://www.mdpi.com/1660-4601/13/4/391/htm>.
- José Almirall and Martin Fortin. The coexistence of terms to describe the presence of multiple concurrent diseases. *Journal of Comorbidity*, 3(1):4–9, 2013. ISSN 2235-042X.
- Anneli Anttonen. *Empowering social policy: The role of social care services in modern welfare states*, pages 88–117. Palgrave MacMillan UK, London, 2005. URL https://books.google.co.uk/books?hl=en&lr=&id=BOmGDAAQBAJ&oi=fnd&pg=PA88&dq=The+Role+of+Social+Care+Services+in+Modern+Welfare+States%E2%80%9D.&ots=vGRWIM6_y1&sig=E_L4_LhYgdzD18gV0kiSazJSxTY#v=onepage&q=The%20Role%20of%20Social%20Care%20Services%20in%20Modern%20Welfare%20States%E2%80%9D.&f=false.
- Anneli Anttonen and Jorma Sipilä. European social care services: is it possible to identify models? *Journal of European social policy*, 6(2):87–100, 1996. ISSN 0958-9287. URL <http://journals.sagepub.com/doi/abs/10.1177/095892879600600201>.
- Sheena Asthana and Joyce Halliday. *What works in tackling health inequalities?: pathways, policies and practice through the lifecourse*. Policy Press, 2006. ISBN 1861346743.

- Audit-Scotland. Commisioning social care. Report, 2012. URL http://www.audit-scotland.gov.uk/docs/health/2012/nr_120301_social_care.pdf.
- Audit-Scotland. Changing models of health and social care. Report, 2016a. URL http://www.audit-scotland.gov.uk/uploads/docs/report/2016/nr_160310_changing_models_care.pdf.
- Audit-Scotland. Social work in scotland. Report, 2016b. URL http://www.audit-scotland.gov.uk/uploads/docs/report/2016/nr_160922_social_work.pdf.
- Audit-Scotland. An overview of local government in scotland 2016. Report, 2016c. URL http://www.audit-scotland.gov.uk/uploads/docs/report/2016/nr_160317_local_government_overview.pdf.
- Beccy Baird, Anna Charles, Matthew Honeyman, David Maguire, and Preety Das. *Understanding pressures in general practice*. King’S Fund, 2016. ISBN 1909029610.
- M. Baker, R. Mawby, and J. Ware. Health inequalites. Report, Royal College of General Practitioners, 2015. URL <http://www.rcgp.org.uk/Policy/RCGP-policy-areas/~media/Files/Policy/A-Z-policy/2015/Health%20Inequalities.ashx>.
- Martin Bardsley, Theo Georghiou, Ludovic Chassin, Geraint Lewis, Adam Steventon, and Jennifer Dixon. Overlap of hospital use and social care in older people in england. *Journal of Health Services Research and Policy*, 17(3):133–139, 2012a. ISSN 1355-8196. URL <http://hsr.sagepub.com/content/17/3/133.full.pdf>.
- Martin Bardsley, Theo Georghiou, Ludovic Chassin, Geraint Lewis, Adam Steventon, and Jennifer Dixon. Overlap of hospital use and social care in older people in england. *Journal of Health Services Research and Policy*, 17(3):133–139, 2012b. ISSN 1355-8196. doi: 10.1258/jhsrp.2011.010171. URL <http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=104486875&site=ehost-livehttp://hsr.sagepub.com/content/17/3/133.full.pdf>.
- K. Barnett, S. W. Mercer, M. Norbury, G. Watt, S. Wyke, and B. Guthrie. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet*, 380(9836):37–43, 2012. ISSN 0140-6736. doi: 10.1016/S0140-6736(12)60240-2. URL <http://www.sciencedirect.com/science/article/pii/S0140673612602402>.

- J. Bartlett and S. Guglielmi. A constitution for social care. Report, DEMOS, 2009. URL http://demos.co.uk/files/Demos_Social_Care_Constitution_online.pdf.
- Yoav Benjamini and Yosef Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the royal statistical society. Series B (Methodological)*, pages 289–300, 1995. ISSN 0035-9246. URL <http://www.stat.purdue.edu/~doerge/BIOINFORM.D/FALL06/Benjamini%20and%20Y%20FDR.pdf>.
- G Bramley, M Evans, and M Noble. Mainstream public services and their impact on neighbourhood deprivation. *Neighbourhood Renewal Unit. London: Office of the Deputy Prime Minister*, 2005.
- Glen Bramley and Martin Evans. Getting the smaller picture: Small-area analysis of public expenditure incidence and deprivation in three english cities. *Fiscal Studies*, 21(2):231–267, 2000. ISSN 1475-5890.
- Samuel L Brilleman and Chris Salisbury. Comparing measures of multimorbidity to predict outcomes in primary care: a cross sectional study. *Family practice*, 30(2): 172–178, 2013. ISSN 0263-2136. URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3604888/?tool=pmcentrez>.
- David Buck and David Maguire. Inequalities in life expectancy: changes over time and implications for policy. Report, 2017. URL http://www.thehealthwell.info/search-results/inequalities-life-expectancy-changes-over-time-and-implications-policy?source=relatedblock&content=resource&member=572160&catalogue=none&collection=Health%20Inequalities%20&tokens_complete=true.
- T. Burchardt, P. Obolenskaya, and P. Vizard. The coalition’s record on adult social care: Policy spending and outcomes 2010-2015. Report, LSE, 2015. URL <http://sticerd.lse.ac.uk/dps/case/spcc/WP17.pdf>.
- L. Burgess. Spice briefing. integration of health and social care: International comparisons. Report, 2012. URL http://www.parliament.scot/ResearchBriefingsAndFactsheets/Factsheets/SB_12-48.pdf.
- L. Burgess. Spice briefing: Integration of health and social care. Report, 2016. URL http://www.parliament.scot/ResearchBriefingsAndFactsheets/S5/SB_16-70_Integration_of_Health_and_Social_Care.pdf.

Amaia Calderón-Larrañaga, Davide L Vetrano, Graziano Onder, Luis A Gimeno-Feliu, Carlos Coscollar-Santaliestra, Angelo Carfí, Maria S Pisciotta, Sara Angleman, René JF Melis, and Giola Santoni. Assessing and measuring chronic multimorbidity in the older population: A proposal for its operationalization. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, page glw233, 2016. ISSN 1079-5006. URL <https://academic.oup.com/biomedgerontology/article/2731241/>.

Joyce Cavaye. *The Care of Older People in Scotland*. Kynoch and Blaney, 2006. ISBN 0955397308.

CfWI. In-depth review of the general practitioner workforce. Report, 2014. URL <https://www.gov.uk/government/publications/in-depth-review-of-the-general-practitioner-workforce>.

Anita Charlesworth and Ruth Thorlby. Reforming social care: options for funding. Report, 2012. URL <https://www.nuffieldtrust.org.uk/files/2017-01/reforming-social-care-options-funding-web-final.pdf>.

Mary E Charlson, Peter Pompei, Kathy L Ales, and C Ronald MacKenzie. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *Journal of chronic diseases*, 40(5):373–383, 1987. ISSN 0021-9681. URL <http://www.sciencedirect.com/science/article/pii/0021968187901718>.

Mercedes Clerencia-Sierra, Amaia Calderón-Larrañaga, Nicolás Martínez-Velilla, Itziar Vergara-Mitxelorena, Pablo Aldaz-Herce, Beatriz Poblador-Plou, Mónica Machón-Sobrado, Nerea Egüés-Olazabal, Gabor Abellán-van Kan, and Alexandra Prados-Torres. Multimorbidity patterns in hospitalized older patients: Associations among chronic diseases and geriatric syndromes. *PloS one*, 10(7):e0132909, 2015. ISSN 1932-6203. URL <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0132909>.

Linda M Collins and Stephanie T Lanza. *Latent class and latent transition analysis: With applications in the social, behavioral, and health sciences*. John Wiley and Sons, New Jersey, 2013. ISBN 111821076X.

Francesca Colombo, Ana Llena-Nozal, Jérôme Mercier, and Fritz Tjadens. *Help wanted? Providing and paying for long-term care: providing and paying for long-term care*,

- volume 2011. OECD Publishing, Paris, 2011. ISBN 9264097759. URL <http://www.oecd.org/els/health-systems/help-wanted-9789264097759-en.htm>.
- John E Cornell, Jacqueline A Pugh, John W Williams Jr, Lewis Kazis, Austin FS Lee, Michael L Parchman, John Zeber, Thomas Pederson, Kelly A Montgomery, and Polly Hitchcock Noël. Multimorbidity clusters: clustering binary data from multimorbidity clusters: clustering binary data from a large administrative medical database. *Applied Multivariate Research*, 12(3):163–182, 2009. ISSN 1918-1108.
- Mary Daly and Jane Lewis. The concept of social care and the analysis of contemporary welfare states. *The British journal of sociology*, 51(2):281–298, 2000. ISSN 1468-4446. URL <http://onlinelibrary.wiley.com/store/10.1111/j.1468-4446.2000.00281.x/asset/j.1468-4446.2000.00281.x.pdf?v=1&t=ihfdbp2&s=4a5f29e4d44e79043d43cdd3134f4761faf78b2f>.
- Sarah Damery, Sarah Flanagan, and Gill Combes. Does integrated care reduce hospital activity for patients with chronic diseases? an umbrella review of systematic reviews. *BMJ open*, 6(11):e011952, 2016. ISSN 2044-6055. URL <http://bmjopen.bmj.com/content/6/11/e011952>.
- V. de Groot, H. Beckerman, G. Lankhorst, and L. Bouter. How to measure comorbidity: a critical review of available methods. *Journal of Clinical Epidemiology*, 57(3):323–323, 2004. ISSN 0895-4356. doi: 10.1016/j.jclinepi.2003.09.002. URL <http://WOS:000220983500015>.
- Blanca A Deusdad, Charles Pace, and Anneli Anttonen. Facing the challenges in the development of long-term care for older people in europe in the context of an economic crisis. *Journal of Social Service Research*, 42(2):144–150, 2016. ISSN 0148-8376. URL <http://www.tandfonline.com/doi/full/10.1080/01488376.2015.1133147>.
- C. Diederichs, K. Berger, and D. B. Bartels. The measurement of multiple chronic diseases-a systematic review on existing multimorbidity indices. *Journals of Gerontology Series a-Biological Sciences and Medical Sciences*, 66(3):301–311, 2011. ISSN 1079-5006. doi: 10.1093/gerona/glq208. URL <http://WOS:000288415800007http://biomedgerontology.oxfordjournals.org/content/66A/3/301.full.pdf>.
- Mary Dixon-Woods, Ms Deborah Kirk, Ms Shona Agarwal, Ellen Annandale, Tony Arthur, Janet Harvey, Ronald Hsu, Savita Katbamna, Richard Olsen, and Lucy

- Smith. Vulnerable groups and access to health care: a critical interpretive review. *National Coordinating Centre NHS Service Delivery Organ RD (NCCSDO)* Retrieved May, 27:2012, 2005.
- Mary Dixon-Woods, Debbie Cavers, Shona Agarwal, Ellen Annandale, Antony Arthur, Janet Harvey, Ron Hsu, Savita Katbamna, Richard Olsen, and Lucy Smith. Conducting a critical interpretive synthesis of the literature on access to healthcare by vulnerable groups. *BMC medical research methodology*, 6(1):35, 2006. ISSN 1471-2288. URL <https://bmcmedresmethodol.biomedcentral.com/articles/10.1186/1471-2288-6-35>.
- R. Elder, M. Kirkpatrick, W. Sutton, Macleod. M., Guthrie B., M. Sutton, and G. Watt. Measuring quality in primary medical services using data from spice. Report, 2007. URL http://www.em-online.com/download/medical_article/36074_spice_report_july_2007.pdf.
- Christopher Eleftheriades and Raphael Wittenberg. A critical review of international practice on assessment and eligibility in adult social care: Lessons for england. Report, 2013. URL <https://www.chseo.org.uk/downloads/report5-adultsocialcare.pdf>.
- Kathryn Ellis. Street-level bureaucracy revisited: The changing face of frontline discretion in adult social care in england. *Social Policy and Administration*, 45(3): 221–244, 2011. ISSN 1467-9515. URL <http://onlinelibrary.wiley.com/doi/10.1111/j.1467-9515.2011.00766.x/full>.
- Tony Evans. Professionals, managers and discretion: Critiquing street-level bureaucracy. *The British Journal of Social Work*, 41(2):368–386, 2010. ISSN 0045-3102. URL <https://academic.oup.com/bjsw/article/41/2/368/1625696/Professionals-Managers-and-Discretion-Critiquing>.
- Alvan R Feinstein. The pre-therapeutic classification of co-morbidity in chronic disease. *Journal of chronic diseases*, 23(7):455–468, 1970. ISSN 0021-9681. URL <http://www.sciencedirect.com/science/article/pii/0021968170900548>.
- Mercedes Fernández-Alonso and Antonio M Jaime-Castillo. Welfare state and individual expectations of economic support: A comparison of norway and spain. *International Sociology*, 31(1):37–56, 2016. ISSN 0268-5809. URL <http://journals.sagepub.com/doi/abs/10.1177/0268580915613192>.

- Julien Forder and José-Luis Fernández. What works abroad? evaluating the funding of long-term care: International perspectives, report commissioned by bupa care services. Report, PSSRU, 2011. URL <http://www.pssru.ac.uk/pdf/dp2794.pdf>.
- Martin Fortin, Lise Lapointe, Catherine Hudon, Alain Vanasse, Antoine L Ntetu, and Danielle Maltais. Multimorbidity and quality of life in primary care: a systematic review. *Health and Quality of life Outcomes*, 2(1):51, 2004. ISSN 1477-7525. URL <https://hqlo.biomedcentral.com/articles/10.1186/1477-7525-2-51>.
- Martin Fortin, Catherine Hudon, Marie-France Dubois, José Almirall, Lise Lapointe, and Hassan Soubhi. Comparative assessment of three different indices of multimorbidity for studies on health-related quality of life. *Health and Quality of Life Outcomes*, 3(1):74, 2005. ISSN 1477-7525. URL <https://hqlo.biomedcentral.com/articles/10.1186/1477-7525-3-74>.
- Martin Fortin, Gina Bravo, Catherine Hudon, Lise Lapointe, Marie-France Dubois, and José Almirall. Psychological distress and multimorbidity in primary care. *The Annals of Family Medicine*, 4(5):417–422, 2006. ISSN 1544-1709. URL <http://www.annfammed.org/content/4/5/417.short>.
- Martin Fortin, Moira Stewart, Marie-Eve Poitras, José Almirall, and Heather Maddocks. A systematic review of prevalence studies on multimorbidity: toward a more uniform methodology. *The Annals of Family Medicine*, 10(2):142–151, 2012. ISSN 1544-1709. URL <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3315131/pdf/0100142.pdf>.
- John Frank, Catherine Bromley, Larry Doi, Michelle Estrade, Ruth Jepson, John McAteer, Tony Robertson, Morag Treanor, and Andrew Williams. Seven key investments for health equity across the lifecourse: Scotland versus the rest of the uk. *Social Science and Medicine*, 140:136–146, 2015. ISSN 0277-9536. URL <http://www.sciencedirect.com/science/article/pii/S027795361530023X>.
- John Gal. Formulating the matthew principle: on the role of the middle classes in the welfare state. *International Journal of Social Welfare*, 7(1):42–55, 1998. ISSN 1468-2397. URL <http://onlinelibrary.wiley.com/doi/10.1111/j.1468-2397.1998.tb00274.x/full>.
- Maria Gannon, Nick Bailey, Annette Hastings, and Glen Bramley. Financial scrutiny unit briefing: The social impact of the 2016-17 local government budget. Report, 2016.

URL http://www.parliament.scot/ResearchBriefingsAndFactsheets/S5/SB_16-84_The_social_impact_of_the_2016-17_local_government_budget.pdf.

Ronald Gijzen, Nancy Hoeymans, François G Schellevis, Dirk Ruwaard, William A Satariano, and Geertrudis AM van den Bos. Causes and consequences of comorbidity: a review. *Journal of clinical epidemiology*, 54(7):661–674, 2001. ISSN 0895-4356. URL http://ac.els-cdn.com/S0895435600003632/1-s2.0-S0895435600003632-main.pdf?_tid=108170ce-6dc1-11e5-8a3e-00000aacb35f&acdnat=1444311347_f32307220b48c8d4a7d7674af6026a1b.

Jon Glasby. The holy grail of health and social care integration. *British Medical Journal*, 2017. ISSN 0959-8138. doi: 10.1136/bmj.j801. URL <http://www.bmj.com/content/356/bmj.j801?hwoasp=authn%3A1488014287%3A4130222%3A1455071724%3A0%3A0%3AMot%2FocEKT3xDHIhVF9ZjA%3D%3D>.

Jon Glasby, Helen Dickinson, and Robin Miller. All in this together? making best use of health and social care resources in an era of austerity. Report, 2011. URL <http://epapers.bham.ac.uk/760/>.

Caroline Glendinning and David Bell. Rethinking social care and support: What can england learn from other countries. Report, 2008. URL <http://www.jrf.org.uk/sites/default/files/jrf/migrated/files/2335.pdf>.

Bruce Guthrie, Sally Wyke, Jane Gunn, Jan van den Akker, and SW Mercer. *Multi-morbidity: the impact on health systems and their development*. OECD Publishing, 2011. ISBN 9264122303. URL <http://dx.doi.org/10.1787/9789264122314-9-en>.

David J Hand and Keming Yu. Idiot’s bayes—not so stupid after all? *International statistical review*, 69(3):385–398, 2001. ISSN 1751-5823. URL <http://onlinelibrary.wiley.com/doi/10.1111/j.1751-5823.2001.tb00465.x/full>.

Annette Hastings. Territorial justice and neighbourhood environmental services: a comparison of provision to deprived and better-off neighbourhoods in the uk. *Environment and Planning C: Government and Policy*, 25(6):896–917, 2007. ISSN 0263-774X. URL <http://journals.sagepub.com/doi/abs/10.1068/c0657>.

Annette Hastings, Nick Bailey, Glen Bramley, Robert Croudace, and David Watkins. ‘managing’the middle classes: Urban managers, public services and the response

- to middle-class capture. *Local Government Studies*, 40(2):203–223, 2014. ISSN 0300-3930.
- Annette Hastings, Nick Bailey, Glen Bramley, Maria Gannon, and David Watkins. The cost of the cuts: The impact on local government and poorer communities. *Joseph Rowntree Foundation: York, UK*, 2015.
- HM-Government. The health and social care act. Report, 2012. URL <http://www.legislation.gov.uk/ukpga/2012/7/contents/enacted>.
- HM-Government. The care act. Report, 2014. URL <http://www.legislation.gov.uk/ukpga/2014/23/contents/enacted/data.htm>.
- Libby Holden, Paul A Scuffham, Michael F Hilton, Alexander Muspratt, Shu-Kay Ng, and Harvey A Whiteford. Patterns of multimorbidity in working australians. *Popul Health Metr*, 9(1):15, 2011. URL <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3123553/pdf/1478-7954-9-15.pdf>.
- R.; Humphries, R.; Thorlby, H.; Holder, P.; Hall, and A. Charles. Social care for older people: Home truths. Report, 2016. URL https://www.kingsfund.org.uk/sites/files/kf/field/field_publication_file/Social_care_older_people_Kings_Fund_Sep_2016.pdf.
- Alyson L Huntley, Rachel Johnson, Sarah Purdy, Jose M Valderas, and Chris Salisbury. Measures of multimorbidity and morbidity burden for use in primary care and community settings: a systematic review and guide. *The Annals of Family Medicine*, 10(2):134–141, 2012. ISSN 1544-1709. URL <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3315139/pdf/0100134.pdf><http://www.annfammed.org/content/10/2/134.full.pdf>.
- Peter Hupe, Michael Hill, and Aurélien Buffat. *Introduction*. Policy Press, 2016. ISBN 1447313275.
- P. Huxley, S. Evans, and T. Maegusku-Hewitt. ‘social care’, the ‘care ethic’ and ‘carework’: new definitions - new directions? *Research Policy and Planning*, 25(1): 3–11, 2007.
- C. Imison. Future trends: Overview. Report, 2012. URL https://www.kingsfund.org.uk/sites/files/kf/field/field_publication_summary/future-trends-overview.pdf.

- M Mofizul Islam, Jose M Valderas, Laurann Yen, Paresh Dawda, Tanisha Jowsey, and Ian S McRae. Multimorbidity and comorbidity of chronic diseases among the senior australians: prevalence and patterns. *PloS one*, 9(1):e83783, 2014. ISSN 1932-6203. URL <http://www.plosone.org/article/fetchObject.action?uri=info:doi/10.1371/journal.pone.0083783&representation=PDF>.
- Nicholas Jardine and Robin Sibson. The construction of hierarchic and non-hierarchic classifications. *The Computer Journal*, 11(2):177–184, 1968. ISSN 0010-4620. URL http://biocomparison.ucoz.ru/_ld/0/60_jardine_constru.pdf.
- UT Kadam and PR Croft. Clinical multimorbidity and physical function in older adults: a record and health status linkage study in general practice. *Family practice*, 24(5): 412–419, 2007. ISSN 0263-2136. URL <http://fampra.oxfordjournals.org/content/24/5/412.full.pdf>.
- Axel Kaehne, Axel Kaehne, Derek Birrell, Derek Birrell, Robin Miller, Robin Miller, Alison Petch, and Alison Petch. Bringing integration home: policy on health and social care integration in the four nations of the uk. *Journal of Integrated Care*, 25(2): 84–98, 2017. ISSN 1476-9018. URL <http://www.emeraldinsight.com/toc/jica/25/2>.
- Srinivasa Vittal Katikireddi, Martin Higgins, Katherine Elizabeth Smith, and Gareth Williams. Health inequalities: the need to move beyond bad behaviours. *Journal of Epidemiology and Community Health*, 2013. ISSN 0143-005X. URL <http://jech.bmj.com/content/early/2013/03/12/jech-2012-202064.short>.
- David King, Matthew Pashley, and Rob Ball. Scotland’s social services spending needs: an english view. *Environment and Planning C: Government and Policy*, 25(6):918–940, 2007. ISSN 0263-774X. URL <http://journals.sagepub.com/doi/abs/10.1068/c0626>.
- Finn Breinholt Larsen, Marie Hauge Pedersen, Karina Friis, Charlotte Glümer, and Mathias Lasgaard. A latent class analysis of multimorbidity and the relationship to socio-demographic factors and health-related quality of life. a national population-based study of 162,283 danish adults. *PloS one*, 12(1):e0169426, 2017. ISSN 1932-6203. URL <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0169426>.
- Julian Le Grand. *The strategy of equality*. Allen and Unwin, London, 1989. ISBN 0043360750.

J. Le Reste, P. Nabbe, H. Lingner, D. K. Lazic, R. Assenova, M. Munoz, A. Sowinska, C. Lygidakis, C. Doerr, S. Czachowski, S. Argyriadou, J. Valderas, B. Le Floch, J. Deriennic, T. Jan, E. Melot, P. Barraine, M. Odorico, C. Lietard, P. Van Royen, and H. Van Marwijk. What research agenda could be generated from the european general practice research network concept of multimorbidity in family practice? *Bmc Family Practice*, 16, 2015. doi: 10.1186/s12875-015-0337-3. URL <http://www.biomedcentral.com/content/pdf/s12875-015-0337-3.pdf>.

Jean Yves Le Reste, Patrice Nabbe, Benedicte Manceau, Charilaos Lygidakis, Christa Doerr, Heidrun Lingner, Slawomir Czachowski, Miguel Munoz, Stella Argyriadou, and Ana Claveria. The european general practice research network presents a comprehensive definition of multimorbidity in family medicine and long term care, following a systematic review of relevant literature. *Journal of the American Medical Directors Association*, 14(5):319–325, 2013. ISSN 1525-8610. URL <http://www.sciencedirect.com/science/article/pii/S1525861013000029>.

T. Lefevre, J. F. d’Ivernois, V. De Andrade, C. Crozet, P. Lombrail, and R. Gagnayre. What do we mean by multimorbidity? an analysis of the literature on multimorbidity measures, associated factors, and impact on health services organization. *Revue D Epidemiologie Et De Sante Publique*, 62(5):305–314, 2014. doi: 10.1016/j.respe.2014.09.002. URL http://ac.els-cdn.com/S0398762014006816/1-s2.0-S0398762014006816-main.pdf?_tid=2c292592-6ecf-11e5-9c03-00000aab0f6b&acdnat=1444427358_5b2e8eb9c02f2cf4d459dca7df76c8bf.

Michael Lipsky. *Street level bureaucracy*, volume 198. Russell Sage Foundation, New York, 1979.

Michael Lipsky. *Street-level bureaucracy, 30th ann. Ed.: dilemmas of the individual in public service*. Russell Sage Foundation, New York, 2010. ISBN 1610446631.

Lord-Sutherland. Independent review of free personal and nursing care. Report, 2008. URL <http://www.gov.scot/Publications/2008/04/25105036/0>.

Sally Macintyre. The black report and beyond what are the issues? *Social science and medicine*, 44(6):723–745, 1997. ISSN 0277-9536. URL http://ac.els-cdn.com/S0277953696001839/1-s2.0-S0277953696001839-main.pdf?_tid=aea008c0-0fb8-11e7-bfeb-00000aacb361&acdnat=1490267333_5da96aa30db817930caf1e5222f0f3d9.

- Johan P Mackenbach. Has the english strategy to reduce health inequalities failed? *Social science and medicine*, 71(7):1249–1253, 2010. ISSN 0277-9536. URL http://www.nusap.net/spe/mackenbach_strategy%20failure_ssm%202010.pdf.
- Johan P Mackenbach. The persistence of health inequalities in modern welfare states: the explanation of a paradox. *Social science and medicine*, 75(4):761–769, 2012. ISSN 0277-9536.
- Johan P Mackenbach, Irina Stirbu, Albert-Jan R Roskam, Maartje M Schaap, Gwenn Menvielle, Mall Leinsalu, and Anton E Kunst. Socioeconomic inequalities in health in 22 european countries. *New England Journal of Medicine*, 358(23):2468–2481, 2008. ISSN 0028-4793. URL <http://www.nejm.org/doi/full/10.1056/NEJMsa0707519#t=article>.
- Mhairi Mackenzie, Ellie Conway, Annette Hastings, Moira Munro, and Catherine O’Donnell. Is ‘candidacy’ a useful concept for understanding journeys through public services? a critical interpretive literature synthesis. *Social Policy and Administration*, 47(7):806–825, 2013. ISSN 1467-9515.
- Mhairi Mackenzie, Ellie Conway, Annette Hastings, Moira Munro, and Catherine A O’Donnell. Intersections and multiple ‘candidacies’: exploring connections between two theoretical perspectives on domestic abuse and their implications for practicing policy. *Social Policy and Society*, 14(1):43–62, 2015. ISSN 1474-7464.
- B. Macleod and M. Mair. Home delivery: a profile of the care at home sector in scotland 2015. Report, Scottish Care, 2015. URL http://www.scottishcare.org/docs/037_280__schomedeliveryreport2015_1433756203.pdf.
- Alessandra Marengoni, Sara Angleman, René Melis, Francesca Mangialasche, Anita Karp, Annika Garmen, Bettina Meinow, and Laura Fratiglioni. Aging with multimorbidity: a systematic review of the literature. *Ageing research reviews*, 10(4): 430–439, 2011. ISSN 1568-1637. URL http://ac.els-cdn.com/S1568163711000249/1-s2.0-S1568163711000249-main.pdf?_tid=229ff4b2-65e5-11e5-91c1-00000aab0f02&acdnat=1443447230_e0ed43a6740c2eaed9ed7fc6f34cc679.
- Michael G Marmot, Jessica Allen, Peter Goldblatt, Tammy Boyce, Di McNeish, Mike Grady, and Ilaria Geddes. Fair society, healthy lives: Strategic review of health inequal-

ities in england post-2010. Report, 2010. URL <http://www.instituteofhealthequity.org/projects/fair-society-healthy-lives-the-marmot-review>.

Peter Matthews and Annette Hastings. Middle-class political activism and middle-class advantage in relation to public services: A realist synthesis of the evidence base. *Social Policy and Administration*, 47(1):72–92, 2013. ISSN 1467-9515.

Carl May, Victor M Montori, and Frances S Mair. We need minimally disruptive medicine. *BMJ: British Medical Journal (Online)*, 339, 2009. ISSN 1756-1833. URL <http://search.proquest.com/openview/548fdf667b7583537cd7c76e7f48cb8a/1?pq-origsite=gscholar&cbl=2043523>.

Gerry McCartney, Chik Collins, and Mhairi Mackenzie. What (or who) causes health inequalities: Theories, evidence and implications? *Health Policy*, 113(3):221–227, 2013. ISSN 0168-8510. doi: <http://dx.doi.org/10.1016/j.healthpol.2013.05.021>. URL <http://www.sciencedirect.com/science/article/pii/S0168851013001590>.

Ann McDonald. *Understanding community care: a guide for social workers*. Palgrave Macmillan, Basingstoke, 2nd edition, 2006. ISBN 1137055707.

Gary McLean, Bruce Guthrie, Stewart W Mercer, and Graham CM Watt. General practice funding underpins the persistence of the inverse care law: cross-sectional study in scotland. *British Journal of General Practice*, 65(641):e799–e805, 2015. doi: 10.3399/bjgp15X687829. URL <http://bjgp.org/bjgp/65/641/e799.full.pdf>.

Stewart W Mercer and Graham CM Watt. The inverse care law: clinical primary care encounters in deprived and affluent areas of scotland. *The Annals of Family Medicine*, 5(6):503–510, 2007. ISSN 1544-1709. URL <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2094031/pdf/0050503.pdf>.

Stewart W Mercer, Susan M Smith, Sally Wyke, Tom O’Dowd, and Graham CM Watt. Multimorbidity in primary care: developing the research agenda. *Family practice*, 26(2):79–80, 2009. ISSN 0263-2136.

Stewart W Mercer, Bruce Guthrie, John Furler, Graham CM Watt, and Julian Tudor Hart. Multimorbidity and the inverse care law in primary care. *BMJ*, 344, 2012. ISSN 1756-1833.

- Brian Munday. European social services: A map of characteristics and trends. Report, 2003. URL http://www.hiproweb.org/fileadmin/cdroms/Kit_Formation_Services/documents/Additional/C-Community_services_for_persons_with_disabilities/C-11_Social_Services_EU_map_Munday_B.doc.
- National-Audit-Office. Tackling inequalities in life expectancy in areas with the worst health and deprivation. Report, 2010. URL <https://www.nao.org.uk/report/tackling-inequalities-in-life-expectancy-in-areas-with-the-worst-health-and-deprivation/>.
- Sarah Nettleton. *The sociology of health and illness*. Polity, 2006. ISBN 0745628281.
- Jennifer Newton and Leanne Browne. How fair is fair access to care? *Practice: Social Work in Action*, 20(4):235–249, 2008. ISSN 0950-3153.
- Shu Kay Ng, Libby Holden, and Jing Sun. Identifying comorbidity patterns of health conditions via cluster analysis of pairwise concordance statistics. *Statistics in medicine*, 31(27):3393–3405, 2012. ISSN 1097-0258.
- SK Ng. A two-way clustering framework to identify disparities in multimorbidity patterns of mental and physical health conditions among australians. *Statistics in medicine*, 34(26):3444–3460, 2015. ISSN 1097-0258.
- NICE. Ng 22. older people with social care needs and multiple long-term conditions. Report, 2015. URL www.nice.org.uk/guidance/ng22.
- NICE. Multimorbidity: clinical assessment and management. multimorbidity: assessment, prioritisation and management of care for people with commonly occurring multimorbidity: Nice guideline ng56:. Report, 2016. URL <https://www.nice.org.uk/guidance/ng56>.
- OECD/EU. A good life in old age? monitoring and improving quality in long-term care. Report, 2013. URL <http://dx.doi.org/10.1787/9789264194564-en>.
- House of Commons. Adult social care. Report, 2017. URL <https://www.publications.parliament.uk/pa/cm201617/cmselect/cmcomloc/1103/1103.pdf>.
- ONS. Health state life expectancies, uk: 2013 to 2015. Report, 2016. URL <file:///C:/Users/hedgi/Downloads/Health%20state%20life%20expectancies,%20UK%202013%20to%202015.pdf>.

- Emmanuele Pavolini and Costanzo Ranci. Restructuring the welfare state: reforms in long-term care in western european countries. *Journal of European Social Policy*, 18 (3):246–259, 2008. ISSN 0958-9287. URL <http://journals.sagepub.com/doi/abs/10.1177/0958928708091058>.
- Stephen Peckham and Elizabeth Meerabeau. *Social policy for nurses and the helping professions*. McGraw-Hill Education., Maidenhead, 2nd edition, 2007. ISBN 0335219624.
- Anthony J Perkins, Kurt Kroenke, Jürgen Unützer, Wayne Katon, John W Williams, Carol Hope, and Christopher M Callahan. Common comorbidity scales were similar in their ability to predict health care costs and mortality. *Journal of clinical epidemiology*, 57(10):1040–1048, 2004. ISSN 0895-4356. URL http://ac.els-cdn.com/S0895435604000812/1-s2.0-S0895435604000812-main.pdf?_tid=20def0d8-6db5-11e5-9bd1-00000aacb35f&acdnt=1444306221_bc95cb86521c040e032fdf541abb0c37.
- Alison Petch. The evidence base for integrated care. *Journal of Integrated Care*, 17(3): 23–25, 2009. ISSN 1476-9018. URL <http://www.emeraldinsight.com/toc/jica/17/3>.
- Alison Petch. Tectonic plates: aligning evidence, policy and practice in health and social care integration. *Journal of Integrated Care*, 20(2):77–88, 2012. ISSN 1476-9018. URL <http://www.emeraldinsight.com/doi/full/10.1108/14769011211220481>.
- D. Platt. The status of social care, a review 2007, report to the minister of state for care services. Report, 2007. URL https://www.housinglin.org.uk/_assets/Resources/Housing/Support_materials/Other_reports_and_guidance/The_state_of_social_care_in_England_2007-08.pdf.
- Frank Popham and Paul Boyle. Assessing socio-economic inequalities in mortality and other health outcomes at the scottish national level. Report, 2010. URL http://calls.ac.uk/wp-content/uploads/final_report_-_assessing_socio-economic_inequalities_in_mortality_and_other_health_outcomes_at_the_scottish_national_level..pdf.
- Martin Powell. The strategy of equality revisited. *Journal of Social Policy*, 24(02): 163–185, 1995. ISSN 1469-7823.

- Alexandra Prados-Torres, Amaia Calderon-Larranaga, Jorge Hanco-Saavedra, Beatriz Poblador-Plou, and Marjan van den Akker. Multimorbidity patterns: a systematic review. *Journal of Clinical Epidemiology*, 67(3):254–266, 2014. ISSN 0895-4356. doi: 10.1016/j.jclinepi.2013.09.021. URL <http://www.sciencedirect.com/science/article/pii/S0895435613004368>.
- R-Core-Team. R: A language and environment for statistical computing. 2011, vienna, austria: R foundation for statistical computing. Report, 2017. URL <http://www.R-project.org>.
- Ruth Robertson, Sarah Gregory, and Joni Jabbal. The social care and health systems of nine countries. Report, 2014. URL http://www.commed.vcu.edu/IntroPH/Community_Assessment/2014/commission-background-paper-social-care-health-system-other-countries.pdf.
- K. Robson. Spice briefing: Public bodies(joint working) (scotland) bill. Report, 2013. URL http://www.parliament.scot/ResearchBriefingsAndFactsheets/S4/SB_13-50.pdf.
- A. Rosano, C. A. Loha, R. Falvo, J. van der Zee, W. Ricciardi, G. Guasticchi, and A. G. de Belvis. The relationship between avoidable hospitalization and accessibility to primary care: a systematic review. *European Journal of Public Health*, 23(3):356–360, 2013. ISSN 1101-1262. doi: 10.1093/eurpub/cks053. URL <http://eurpub.oxfordjournals.org/content/eurpub/23/3/356.full.pdf>.
- Chris Salisbury, Leigh Johnson, Sarah Purdy, Jose M Valderas, and Alan A Montgomery. Epidemiology and impact of multimorbidity in primary care: a retrospective cohort study. *British Journal of General Practice*, 61(582):e12–e21, 2011. ISSN 0960-1643. URL <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3020068/pdf/bjgp61-e12.pdf>.
- André Sapir. Globalization and the reform of european social models. *Journal of Common Market Studies*, 44(2):369–390, 2006. ISSN 1468-5965. URL <http://onlinelibrary.wiley.com/doi/10.1111/j.1468-5965.2006.00627.x/full>.
- Nicholas K Schiltz, David F Warner, Jiayang Sun, Paul M Bakaki, Avi Dor, Charles W Given, Kurt C Stange, and Siran M Koroukian. Identifying specific combinations of multimorbidity that contribute to health care resource

- utilization: An analytic approach. *Medical Care*, 2017. ISSN 0025-7079. URL http://journals.lww.com/lww-medicalcare/Citation/2017/03000/Identifying_Specific_Combinations_of.11.aspx.
- SCIE. Scie guide 33: Fair access to cares services (facs): proritising eligibility for care and support. Report, 2013. URL <http://www.scie.org.uk/publications/guides/guide33/files/guide33.pdf>.
- SCIE. Eligibility dermination for the care act 2014. Report, 2015. URL <http://www.scie.org.uk/care-act-2014/assessment-and-eligibility/eligibility/>.
- Scottish-Government. Self-directed support: A national strategy for scotland. Report, 2010. URL <http://www.gov.scot/Publications/2010/11/05120810/0>.
- Scottish-Government. Local authorities revenue block grant funding. Report, 2013. URL <http://www.gov.scot/Topics/Government/local-government/17999/CoreRevenueFunding>.
- Scottish-Government. Self-directed support: Practitioner guidance. Report, 2014. URL <http://www.gov.scot/Publications/2014/08/5212>.
- Scottish-Government. Data on eligibility criteria and waiting times. Report, 2015. URL <http://www.gov.scot/Topics/Statistics/Browse/Health/Data/QuarterlySurvey/QECWTJUL2015>.
- Scottish-Government. Social care services, scotland, 2016. Report, 2016a. URL <https://www.gov.uk/government/statistics/social-care-services-scotland-2016>.
- Scottish-Government. Health and social care delivery plan. Report, 2016b. URL <http://www.gov.scot/Resource/0051/00511950.pdf>.
- Scottish-Government. 2016-17 settlement. grant aided expenditure green book. Report, 2016c. URL <http://www.gov.scot/Resource/0049/00499184.pdf>.
- Scottish-Government. Long term monitoring of health inequalities. Report, 2017. URL <http://www.gov.scot/Publications/2017/03/8751/downloads>.
- Scottish-Parliament. Health and sport committee: Report on health inequalities. Report, 2015. URL http://www.parliament.scot/S4_HealthandSportCommittee/Reports/her-15-01w-rev.pdf.

- Mr Peter Sharkey. *Essentials of Community Care*. Palgrave Macmillan, Basingstoke, 2nd edition, 2006. ISBN 1137165367.
- Mary Shaw, Danny Dorling, and George Davey Smith. *Poverty, social exclusion, and minorities*. Oxford University Press, Oxford, 2nd edition, 2006.
- Judith Sinnige, Joze Braspenning, Francois Schellevis, Irina Stirbu-Wagner, Gert Westert, and Joke Korevaar. The prevalence of disease clusters in older adults with multiple chronic diseases - a systematic literature review. *Plos One*, 8(11), 2013. ISSN 1932-6203. doi: 10.1371/journal.pone.0079641. URL <http://www.plosone.org/article/doi/10.1371/journal.pone.0079641?representation=PDF>.
- Katherine E Smith, Clare Bambra, and Sarah Hill. *Background and introduction: UK experiences of health inequalities*. Oxford University Press, Oxford, 2016. ISBN 0191008486.
- Peter C Smith. Formula funding of public services: an economic analysis. *Oxford Review of Economic Policy*, 19(2):301–322, 2003. ISSN 1460-2121. URL <https://academic.oup.com/oxrep/article/19/2/301/465429>.
- Barbara Starfield, Jonathan Weiner, Laura Mumford, and Donald Steinwachs. Ambulatory care groups: a categorization of diagnoses for research and management. *Health services research*, 26(1):53, 1991. URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1069810/>.
- Barbara Starfield, Leiyu Shi, and James Macinko. Contribution of primary care to health systems and health. *Milbank quarterly*, 83(3):457–502, 2005. ISSN 1468-0009. URL <http://onlinelibrary.wiley.com/doi/10.1111/j.1468-0009.2005.00409.x/full>.
- Ailsa Stewart, Alison Petch, and Lisa Curtice. Moving towards integrated working in health and social care in scotland: from maze to matrix. *Journal of Interprofessional Care*, 17(4):335–350, 2003. ISSN 1356-1820. URL [MovingtowardsintegratedworkinginhealthandsocialcareinScotland:frommasetomatrix](http://www.tandf.co.uk/journals/1356-1820/MovingtowardsintegratedworkinginhealthandsocialcareinScotland:frommasetomatrix).
- Alexander Topchy, Anil K Jain, and William Punch. Clustering ensembles: Models of consensus and weak partitions. *IEEE Transactions on pattern analysis and machine*

- intelligence*, 27(12):1866–1881, 2005. ISSN 0162-8828. URL <http://ieeexplore.ieee.org/stamp/stamp.jsp?arnumber=1524981>.
- Julian Tudor-Hart. The inverse care law. *The Lancet*, 297(7696):405–412, 1971. ISSN 0140-6736.
- Lars Tummers and Victor Bekkers. Policy implementation, street-level bureaucracy, and the importance of discretion. *Public Management Review*, 16(4):527–547, 2014. ISSN 1471-9037. URL <http://www.tandfonline.com/doi/full/10.1080/14719037.2013.841978?scroll=top&needAccess=true&instName=University+of+Glasgow>.
- Jose M Valderas, Barbara Starfield, Bonnie Sibbald, Chris Salisbury, and Martin Roland. Defining comorbidity: implications for understanding health and health services. *The Annals of Family Medicine*, 7(4):357–363, 2009. ISSN 1544-1709. URL <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2713155/pdf/0060357.pdf>.
- Marjan van den Akker, Frank Buntinx, and J. André Knottnerus. Comorbidity or multimorbidity. *European Journal of General Practice*, 2(2):65–70, 1996. ISSN 1381-4788. doi: 10.3109/13814789609162146. URL <http://www.tandfonline.com/doi/abs/10.3109/13814789609162146>.
- Marjan van den Akker, Frank Buntinx, Sjeff Roos, and J André Knottnerus. Problems in determining occurrence rates of multimorbidity. *Journal of clinical epidemiology*, 54(7):675–679, 2001. ISSN 0895-4356.
- C. Violan, Q. Foguet-Boreu, G. Flores-Mateo, C. Salisbury, J. Blom, M. Freitag, L. Glynn, C. Muth, and J. M. Valderas. Prevalence, determinants and patterns of multimorbidity in primary care: A systematic review of observational studies. *Plos One*, 9(7), 2014. ISSN 1932-6203. doi: 10.1371/journal.pone.0102149. URL <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0102149>.
- Michael Von Korff, Edward H Wagner, and Kathleen Saunders. A chronic disease score from automated pharmacy data. *Journal of clinical epidemiology*, 45(2):197–203, 1992. ISSN 0895-4356. URL <http://www.sciencedirect.com/science/article/pii/089543569290016G>.
- Emma Wallace, Ronald McDowell, Kathleen Bennett, Tom Fahey, and Susan M Smith. Comparison of count-based multimorbidity measures in predicting emergency

admission and functional decline in older community-dwelling adults: a prospective cohort study. *BMJ open*, 6(9):e013089, 2016. ISSN 2044-6055. URL <http://bmjopen.bmj.com/content/6/9/e013089.abstract>.

Helen Weatherly, Anne Mason, Maria Goddard, and K Wright. Financial integration across health and social care: evidence review. Report, 2010. URL <https://www.york.ac.uk/che/pdf/Financial%20Integration%20Across%20Health%20and%20Social%20Care%20-%20Research%20Findings.pdf>.

Charlotte Weston, Alexander Gilkes, Stevo Durbaba, Peter Schofield, Patrick White, and Mark Ashworth. Long term condition morbidity in english general practice: a cross-sectional study using three composite morbidity measures. *BMC Family Practice*, 17(1):166, 2016. ISSN 1471-2296. URL <https://bmcfampract.biomedcentral.com/articles/10.1186/s12875-016-0563-3>.

WHO. Who global strategy on people-centred and integrated health services: interim report. Report, 2015. URL http://apps.who.int/iris/bitstream/10665/155002/1/WHO_HIS_SDS_2015.6_eng.pdf.

H Wickham. tidyverse: Easily install and load ‘tidyverse’packages [software]. Report, 2017. URL <https://cran.r-project.org/web/packages/tidyverse/index.html>.

Tora Grauers Willadsen, Anna Bebe, Rasmus Køster-Rasmussen, Dorte Ejg Jarbøl, Ann Dorrit Guassora, Frans Boch Waldorff, Susanne Reventlow, and Niels de Fine Olivarius. The role of diseases, risk factors and symptoms in the definition of multimorbidity—a systematic review. *Scandinavian journal of primary health care*, 34(2):112–121, 2016. ISSN 0281-3432. URL <http://www.tandfonline.com/doi/abs/10.3109/02813432.2016.1153242>.