Prediction Card:

Laboratory and Solar System Tests of SCT Modified Gravity

David Alfyorov

March 2026

Credits: Aliaksandr Samatyia contributed research-assistance and workflow support.

STATUS: COMPLETE

Primary bound: $\Lambda > 2.565 \times 10^{-3}$ eV (95% CL, Eöt-Wash).
SCT passes all laboratory and solar system tests.
85 tests PASS, 5 publication figures.

Contents

1 Introduction **2**

2 SCT Modified Potential **2**
2.1 Key Properties . . . 2

3 Yukawa Parametrization **3**

4 Torsion Balance Constraints **3**
4.1 Lee et al. (2020) . . . 3
4.2 Kapner et al. (2007) . . . 3
4.3 Consistency with PPN-1 . . . 3

5 Casimir Effect **4**

6 Atom Interferometry **4**

7 Solar System Tests **4**
7.1 Lunar Laser Ranging (LLR) . . . 4
7.2 Gravity Probe B (GP-B) . . . 4
7.3 MICROSCOPE . . . 4
7.4 Cassini . . . 5

8 Unified Exclusion Plot **5**

9 ξ-Dependence **5**

10 Comparison with Competing Theories **6**

11 Summary of Experimental Constraints **7**

12 Conclusion **7**

13 Verification Status **8**

1 Introduction

Spectral Causal Theory (SCT) predicts a modified gravitational potential at short distances, arising from nonlocal form factors generated by the spectral action. At distances much larger than the inverse spectral scale $1/\Lambda$, the potential reduces to the Newtonian $1/r$ law; at distances comparable to or smaller than $1/\Lambda$, Yukawa-type corrections become significant.

This prediction card confronts the SCT modified potential with all available laboratory and solar system experiments that probe departures from Newtonian gravity at sub-millimeter to astronomical scales. The key result is that the Eöt-Wash torsion balance experiment provides the only meaningful constraint on the SCT spectral scale, setting $\Lambda > 2.565 \times 10^{-3}$ eV at 95% confidence level. All other experiments—Casimir, atom interferometry, Lunar Laser Ranging, Gravity Probe B, MICROSCOPE, and Cassini—are either exponentially suppressed or insensitive to the SCT coupling strengths.

The upstream derivations are:

- NT-1b (SM form factors): provides $\alpha_C = 13/120$ and $\alpha_R(\xi) = 2(\xi - 1/6)^2$.
- NT-4a (linearized field equations): provides the propagator denominators $\Pi_{\mathrm{TT}}(z)$ and $\Pi_{\mathrm{s}}(z, \xi)$, the effective masses m_2 and m_0, and the modified potential.
- PPN-1 (post-Newtonian parameters): provides the solar system constraint $\Lambda \geq 2.38 \times 10^{-3}$ eV.

2 SCT Modified Potential

The linearized SCT field equations, derived in NT-4a from the spectral action with Standard Model matter content, yield a modified Newtonian potential in the local Yukawa approximation:

$$\boxed{\frac{V(r)}{V_N(r)} = 1 - \frac{4}{3}\, e^{-m_2 r} + \frac{1}{3}\, e^{-m_0 r}} \tag{1}$$

where $V_N(r) = -Gm_1m_2/r$ is the Newtonian potential and the effective masses are

$$m_2 = \Lambda\sqrt{\frac{60}{13}} \approx 2.148\,\Lambda, \tag{2}$$

$$m_0(\xi) = \frac{\Lambda}{\sqrt{6\left(\xi - \frac{1}{6}\right)^2}} \quad (\xi \neq \tfrac{1}{6}). \tag{3}$$

The Yukawa amplitudes are **parameter-free**, fixed entirely by the Standard Model field content ($N_s = 4$, $N_f = 45$, $N_v = 12$):

$$\alpha_1 = -\frac{4}{3}, \qquad \alpha_2 = +\frac{1}{3}. \tag{4}$$

2.1 Key Properties

(a) **Finite potential at origin:** $V(0)/V_N(0) = 1 - 4/3 + 1/3 = 0$ (exact). At $\xi = 1/6$, $V(0)/V_N(0) = 1 - 4/3 = -1/3$.

(b) **Newtonian recovery:** $V(r)/V_N(r) \to 1$ as $r \to \infty$ (exponentially fast).

(c) **Monotonicity:** $V'(r) > 0$ for all $r > 0$. Proof: $V'(r) = 0$ requires $4m_2/m_0 = \exp((m_2 - m_0)r)$. Since $m_2 < m_0$ (at $\xi = 0$), the right-hand side is < 1 for $r > 0$, while the left-hand side is $4\sqrt{10/13} \approx 3.508 > 1$. No zero exists.

(d) **Mass ratio:** $m_2/m_0 = \sqrt{10/13} \approx 0.8771$ at $\xi = 0$ (parameter-free).

(e) **Scalar decoupling:** At conformal coupling $\xi = 1/6$, the scalar mass $m_0 \to \infty$ and the second Yukawa term vanishes, leaving only the spin-2 contribution.

3 Yukawa Parametrization

The standard parametrization for short-range gravity experiments writes the potential deviation as

$$V(r) = V_N(r)\big(1 + \alpha\, e^{-r/\lambda}\big). \tag{5}$$

The SCT potential (1) contains *two* Yukawa terms. Mapping to the standard form:

$$\text{Spin-2:} \quad \alpha_1 = -\frac{4}{3}, \quad \lambda_1 = \frac{\hbar c}{m_2 c^2} = \frac{\hbar c}{\Lambda\sqrt{60/13}}, \tag{6}$$

$$\text{Scalar:} \quad \alpha_2 = +\frac{1}{3}, \quad \lambda_2 = \frac{\hbar c}{m_0 c^2} = \frac{\hbar c\sqrt{6(\xi - 1/6)^2}}{\Lambda}. \tag{7}$$

The spin-2 coupling $|\alpha_1| = 4/3$ dominates the scalar coupling $|\alpha_2| = 1/3$ at all distance scales. Experimental exclusion curves in the $(\lambda, |\alpha|)$ plane therefore intersect the SCT spin-2 line first, giving the primary constraint.

At the boundary value $\Lambda = 2.565 \times 10^{-3}$ eV ($\xi = 0$):

$$\lambda_1 = 35.8\ \mu\text{m}, \qquad \lambda_2 = 31.4\ \mu\text{m}, \qquad m_2 = 5.509 \times 10^{-3}\ \text{eV}, \qquad m_0 = 6.282 \times 10^{-3}\ \text{eV}. \tag{8}$$

4 Torsion Balance Constraints

Torsion balance experiments by the Eöt-Wash group at the University of Washington provide the most sensitive tests of the gravitational inverse-square law at sub-millimeter distances.

4.1 Lee et al. (2020)

The most recent result (Lee et al., PRL **124**, 101101, 2020) extends the exclusion region to $\lambda \geq 38.6\ \mu$m at $|\alpha| = 1$. Since the SCT spin-2 coupling $|\alpha_1| = 4/3 > 1$, the SCT exclusion curve crosses the experimental bound at a *shorter* Yukawa range:

$$\lambda_{1,\text{cross}} = 35.8\ \mu\text{m}, \qquad \boxed{\Lambda_{\text{min}} = 2.565 \times 10^{-3}\ \text{eV}} \quad (95\%\ \text{CL}). \tag{9}$$

This is the primary laboratory constraint on the SCT spectral scale.

4.2 Kapner et al. (2007)

The earlier result (Kapner et al., PRL **98**, 021101, 2007) provides the exclusion curve at $\lambda > 200\ \mu$m that is included in the composite exclusion analysis. At these larger ranges, the bound is weaker than the Lee (2020) result.

4.3 Consistency with PPN-1

The PPN-1 analysis derived $\Lambda \geq 2.38 \times 10^{-3}$ eV using the conservative single-Yukawa approximation with $|\alpha| = 1$ crossing at 38.6 μm. The LT-3d bound is **7.8% tighter** because it correctly uses the SCT spin-2 coupling $|\alpha_1| = 4/3$ (crossing at 35.8 μm).

5 Casimir Effect

Casimir force measurements (Decca et al., 2007; Chen et al., PRL **116**, 221102, 2016) probe the gravitational potential at sub-micrometer distances. However, the experimental exclusion curves at these scales require $|\alpha|_{95\%} \gtrsim 3000$, which is far above the SCT couplings $|\alpha_1| = 4/3$ and $|\alpha_2| = 1/3$.

The SCT gravitational correction at Casimir-relevant distances is exponentially suppressed. At $r = 0.1\ \mu$m with $\Lambda = 2.565 \times 10^{-3}$ eV:

$$\left| \frac{V(r)}{V_N(r)} - 1 \right| \approx 0.003 \quad \text{(0.3\% deviation from Newton).} \tag{10}$$

This is many orders of magnitude below the Casimir experimental sensitivity. Casimir experiments therefore provide **no constraint** on the SCT spectral scale (see Fig. 3).

6 Atom Interferometry

Atom interferometry experiments (Sabulsky et al., PRL **123**, 2019; MAGIS-100 concept) probe gravity at centimeter-to-meter scales. At these distances, the exclusion limits are $|\alpha|_{95\%} \sim 10^8$, which is far above the SCT couplings.

Furthermore, the SCT Yukawa range at the boundary value $\lambda_1 = 35.8\ \mu$m is many orders of magnitude shorter than the effective baseline of atom interferometers ($\sim$ 0.01–100 m). The SCT correction at these scales is:

$$\text{At } r = 1 \text{ cm}: \quad e^{-m_2 r} \sim e^{-280} \approx 0. \tag{11}$$

Atom interferometry provides **no constraint** on SCT.

7 Solar System Tests

All solar system tests are exponentially suppressed because the SCT Yukawa range $\lambda_1 \sim 36\ \mu$m is negligible compared to astronomical distances.

7.1 Lunar Laser Ranging (LLR)

At the Earth–Moon distance $r_{\text{Moon}} = 3.844 \times 10^8$ m:

$$m_2 r_{\text{Moon}} = 1.073 \times 10^{13}, \qquad e^{-m_2 r_{\text{Moon}}} \approx 0 \quad \text{(to any representable precision).} \tag{12}$$

LLR provides **zero constraint** on SCT (Williams et al., PRL 2004).

7.2 Gravity Probe B (GP-B)

At the GP-B orbital altitude $r \approx 7000$ km:

$$m_2 r = 1.954 \times 10^{11}, \qquad e^{-m_2 r} \approx 0. \tag{13}$$

GP-B provides **zero constraint** on SCT (Everitt et al., PRL **106**, 221101, 2011).

7.3 MICROSCOPE

The MICROSCOPE satellite (Touboul et al., PRL **129**, 121102, 2022) tests the weak equivalence principle to $|\eta| < 1.5 \times 10^{-15}$. The SCT Yukawa forces couple universally to mass (through the trace of the stress-energy tensor), so a composition-independent force does not violate the WEP. MICROSCOPE provides **no constraint** on the SCT spectral scale.

7.4 Cassini

The Cassini spacecraft (Bertotti et al., Nature **425**, 374, 2003) constrains $|\gamma_{\rm PPN} - 1| < 2.3 \times 10^{-5}$ at 1 AU. At this distance, the SCT correction is $\sim e^{-10^{15}} \approx 0$, giving $\Lambda_{\rm min} \sim 6 \times 10^{-18}$ eV—a factor of 10^{14} weaker than the Eöt-Wash bound. Cassini is entirely superseded.

8 Unified Exclusion Plot

Figure 1 presents the unified exclusion plot combining all experimental channels in the $(\lambda, |\alpha|)$ plane. The SCT prediction appears as two horizontal lines at $|\alpha_1| = 4/3$ and $|\alpha_2| = 1/3$. The crossing of the $|\alpha_1| = 4/3$ line with the Eöt-Wash exclusion curve defines the primary bound.



Figure 1: Unified exclusion plot in the $(\lambda, |\alpha|)$ plane. The horizontal dashed lines show the SCT Yukawa couplings $|\alpha_1| = 4/3$ (spin-2) and $|\alpha_2| = 1/3$ (scalar). The crossing of the spin-2 line with the Eöt-Wash exclusion curve gives $\lambda_1 < 35.8\ \mu$m, corresponding to $\Lambda > 2.565 \times 10^{-3}$ eV (95% CL).

9 ξ-Dependence

The Higgs non-minimal coupling parameter ξ affects only the scalar sector:

$$m_0(\xi) = \frac{\Lambda}{\sqrt{6(\xi - 1/6)^2}}, \qquad \alpha_2 = +\frac{1}{3}. \tag{14}$$

The spin-2 sector ($\alpha_1 = -4/3$, $m_2 = \Lambda\sqrt{60/13}$) is entirely ξ-independent.

Since the primary bound is set by the spin-2 coupling, $\Lambda_{\rm min}$ **is constant across all values of** ξ. This is confirmed numerically: a scan over $\xi \in [0, 0.5]$ (with $\xi = 1/6$ excluded due to scalar decoupling) yields $\Lambda_{\rm min} = 2.565 \times 10^{-3}$ eV to machine precision at every point.



Figure 2: Ratio $V(r)/V_N(r)$ as a function of distance for several values of Λ. The potential is identically zero at the origin and monotonically approaches the Newtonian value.

At $\xi = 1/6$ (conformal coupling), the scalar mode decouples entirely: $m_0 \to \infty$, and the potential reduces to

$$\left.\frac{V(r)}{V_N(r)}\right|_{\xi=1/6} = 1 - \frac{4}{3}\, e^{-m_2 r}. \tag{15}$$

The bound remains $\Lambda > 2.565 \times 10^{-3}$ eV.

10 Comparison with Competing Theories

Theory	Potential type	$\lvert\alpha\rvert$	Free params	Ghost status
SCT (this work)	Two Yukawa	4/3, 1/3	1 (Λ)	Spin-2 (fakeon prescription)
Stelle gravity	Two Yukawa	4/3, 1/3	2 (m_0, m_2)	Spin-2 (Ostrogradsky)
Ghost-free IDG	Error function	—	1 (M_s)	None (entire propagator)
ADD ($n = 2$)	Power-law	—	1 (R)	None
$f(R)$ gravity	One Yukawa	1/3	2 (n, λ_C)	None (chameleon screened)

Key distinguishing features of SCT:

- The Yukawa amplitudes $\alpha_1 = -4/3$ and $\alpha_2 = +1/3$ are *parameter-free*—they are fixed by the Standard Model field content and cannot be adjusted.
- The mass ratio $m_2/m_0 = \sqrt{10/13} \approx 0.8771$ (at $\xi = 0$) is a unique prediction.
- Stelle gravity has the same functional form but treats m_0 and m_2 as free parameters.
- Ghost-free infinite-derivative gravity (IDG) has a Gaussian modification ($V/V_N \sim \mathrm{Erf}(M_s r/2)$), not Yukawa, and is ghost-free by construction.



Figure 3: The SCT gravitational correction at Casimir-relevant distances. The correction is exponentially suppressed and far below the experimental sensitivity of Casimir force measurements.

- $f(R)$ gravity has only a scalar Yukawa with $|\alpha| = 1/3 < 4/3$, and the chameleon mechanism can screen the modification in dense environments.

11 Summary of Experimental Constraints

Experiment	Constrains SCT?	Λ_{min} (eV)	Notes
Eöt-Wash (Lee 2020)	**YES**	$\mathbf{2.565 \times 10^{-3}}$	Only constraining experiment
Casimir (Chen 2016)	No	—	$\lvert\alpha\rvert_{95\%} \gg 4/3$ at sub-μm
Atom interferometry	No	—	$\lvert\alpha\rvert_{95\%} \sim 10^{8}$ at cm–m
LLR (Williams 2004)	No	—	$e^{-1.07\times10^{13}} = 0$
GP-B (Everitt 2011)	No	—	$e^{-1.95\times10^{11}} = 0$
MICROSCOPE (Touboul 2022)	No	—	Universal coupling: no WEP violati
Cassini (Bertotti 2003)	No	6×10^{-18}	$10^{14}\times$ weaker than Eöt-Wash

12 Conclusion

Spectral Causal Theory passes all current laboratory and solar system tests of modified gravity. The primary constraint on the spectral scale is

$$\boxed{\Lambda > 2.565 \times 10^{-3}\ \text{eV}} \quad (95\%\ \text{CL}), \tag{16}$$

corresponding to a spin-2 Yukawa range $\lambda_1 < 35.8\ \mu$m, derived from the Eöt-Wash torsion balance experiment (Lee et al., 2020).



Figure 4: $\Lambda_{\min}$ as a function of ξ. The bound is constant because it is set by the ξ-independent spin-2 Yukawa coupling $|\alpha_1| = 4/3$.

This bound is:

- **7.8% tighter** than the PPN-1 bound ($\Lambda \geq 2.38 \times 10^{-3}$ eV), which used the conservative $|\alpha| = 1$ crossing.
- **ξ-independent**, set entirely by the spin-2 Yukawa coupling $|\alpha_1| = 4/3$.
- **Parameter-free**: the SCT couplings $\alpha_1 = -4/3$, $\alpha_2 = +1/3$ are fixed by the SM field content and cannot be tuned.
- **Falsifiable**: improvements in torsion balance sensitivity at $\lambda \sim 20$–40 μm will either detect the SCT signal or tighten the bound on Λ.

No other experiment currently constrains the SCT spectral scale. Casimir measurements, atom interferometry, and all solar system tests are either exponentially suppressed or insensitive to the SCT coupling strengths.

13 Verification Status

Verification status (March 2026). The LT-3d derivation has been verified through systematic multi-method checks with **85 independent tests** (8 parameter checks, 9 unit conversions, 4 scalar mass, 7 potential, 7 Eöt-Wash, 3 LLR, 3 GP-B, 2 MICROSCOPE, 3 atom interferometry, 5 Casimir, 5 ξ-dependence, 5 dimensional analysis, 3 competing theories, 4 results generation, 4 figures, 7 consistency, 6 edge cases). All key values confirmed unanimously across independent methods. Six independent spot-checks confirm canonical results. No errors found in the code under test.

References

[1] J. G. Lee et al., "New test of the gravitational $1/r^2$ law at separations down to 52 μm," PRL **124**, 101101 (2020).



Figure 5: Independent exclusion analysis from the dual derivation, confirming the crossing point and primary bound.

[2] D. J. Kapner et al., "Tests of the gravitational inverse-square law below the dark-energy length scale," PRL **98**, 021101 (2007).

[3] Y.-J. Chen et al., "Stronger limits on hypothetical Yukawa interactions in the 30–8000 nm range," PRL **116**, 221102 (2016).

[4] R. S. Decca et al., "Tests of new physics from precise Casimir force measurements," PRD **75**, 077101 (2007).

[5] I. Sabulsky et al., "Experiment to detect dark energy forces using atom interferometry," PRL **123**, 061102 (2019).

[6] J. G. Williams, S. G. Turyshev, and D. H. Boggs, "Progress in lunar laser ranging tests of relativistic gravity," PRL **93**, 261101 (2004).

[7] C. W. F. Everitt et al., "Gravity Probe B: Final results of a space experiment to test general relativity," PRL **106**, 221101 (2011).

[8] P. Touboul et al., "MICROSCOPE Mission: Final results of the test of the equivalence principle," PRL **129**, 121102 (2022).

[9] B. Bertotti, L. Iess, and P. Tortora, "A test of general relativity using radio links with the Cassini spacecraft," Nature **425**, 374 (2003).

[10] C. M. Will, "The confrontation between general relativity and experiment," Living Rev. Relativity **17**, 4 (2014); arXiv:1403.7377.

[11] K. S. Stelle, "Renormalization of higher-derivative quantum gravity," PRD **16**, 953 (1977).

[12] L. Modesto, "Super-renormalizable quantum gravity," PRD **86**, 044005 (2012); arXiv:1107.2403.

[13] J. Edholm, A. S. Koshelev, and A. Mazumdar, "Behavior of the Newtonian potential for ghost-free gravity and singularity-free gravity," PRD **94**, 104033 (2016); arXiv:1604.01989.

[14] H. Murata and M. Tanaka, "A review of short-range gravity experiments in the LHC era," Class. Quantum Grav. **32**, 033001 (2015); arXiv:1408.3588.

[15] E. G. Adelberger, B. R. Heckel, and A. E. Nelson, "Tests of the gravitational inverse-square law," Ann. Rev. Nucl. Part. Sci. **53**, 77 (2003).

[16] W.-H. Tan et al., "Improvement for testing the gravitational inverse-square law at the submillimeter range," PRL **124**, 051301 (2020).