OT: One-Loop Optical Theorem with Fakeon Prescription in SCT

Spectral Positivity, Central Charge, and
the Ghost Exclusion Mechanism

David Alfyorov

March 2026

Credits: Aliaksandr Samatyia contributed research-assistance and workflow support.

Abstract

We verify unitarity of the Spectral Causal Theory (SCT) graviton propagator at one loop by establishing the optical theorem with the fakeon prescription. The key result is that the dressed graviton propagator satisfies spectral positivity: $\mathrm{Im}[G_{\mathrm{dressed}}(s)] > 0$ for all $s > 0$. This follows from three structural properties: (1) the propagator denominator $\Pi_{\mathrm{TT}}(z)$ is an entire function with real Taylor coefficients, hence real for real z; (2) the one-loop self-energy absorptive part from Standard Model matter is strictly positive, $\mathrm{Im}[\Sigma_{\mathrm{matter}}(s)] > 0$; and (3) the fakeon prescription excludes the ghost from the physical spectrum. We verify the Anselmi central charge $C_m = 283/120$ for the SM content, reconcile it with the HLZ $N_{\mathrm{eff}} = 143.5$, and show that Type C complex poles contribute $< 0.2\%$ corrections. All results are cross-checked against the A3 (ghost width), GP (dressed propagator), and FK (fakeon convergence) analyses: 67 new tests pass, 2825 total regression tests pass.

Contents

1 Introduction **2**

2 Anselmi Absorptive Polynomials **2**

3 Central Charge and N_{eff} Reconciliation **2**
- 3.1 Anselmi central charge . . . 2
- 3.2 HLZ N_{eff} . . . 2
- 3.3 Reconciliation . . . 3

4 Spectral Positivity with Fakeon Prescription **3**

5 One-Loop Optical Theorem **3**

6 Fakeon vs Feynman Comparison **4**

7 Stelle Gravity Comparison **5**

8 N-Pole Convergence **5**

9 Cross-Checks **7**

10 Verification Summary **7**

11 Conclusion **8**

1 Introduction

The SCT graviton propagator in the transverse-traceless sector is

$$G_{\mathrm{TT}}(k^2) = \frac{1}{k^2\,\Pi_{\mathrm{TT}}(-k^2/\Lambda^2)}\,, \tag{1}$$

where

$$\Pi_{\mathrm{TT}}(z) = 1 + \frac{13}{60}\,z\,\hat{F}_1(z) \tag{2}$$

is an entire function of order 1 (NT-2 verified). The zeros of Π_{TT} produce poles in the propagator:

- $z_L = -1.2807$ (Lorentzian, timelike $k^2 > 0$, residue $R_L = -0.538$)
- $z_0 = +2.4148$ (Euclidean, spacelike $k^2 < 0$, residue $R_0 = -0.493$)
- Type C pairs at $|z| \approx 34, 59, 85, \ldots$ (complex, $|R| < 0.01$)

The question addressed here: does the one-loop graviton scattering amplitude satisfy the optical theorem when the ghost poles are treated as fakeons (purely virtual particles)?

2 Anselmi Absorptive Polynomials

Following Anselmi and Piva [1], the absorptive part of the graviton self-energy from matter fields takes the form

$$\Gamma^{\Phi}_{\mathrm{abs}} = \frac{i}{16\pi}\int \sqrt{-g}\,R_{\mu\nu}\,\theta(r_\Phi)\,\theta(1-r_\Phi)\,\sqrt{1-r_\Phi}\,P_\Phi(r_\Phi)\Big(R^{\mu\nu} - \tfrac{1}{3}g^{\mu\nu}R\Big), \tag{3}$$

where $r_\Phi = -4m_\Phi^2/\Box$. The spin-projected polynomials are:

Species	$P_\Phi(r)$	$Q_\Phi(r)$
Scalar	$(1-r)^2/120$	$(4\eta_s - r)^2/576$
Dirac	$(3 - r - 2r^2)/60$	$r(1-r)/144$
Vector	$1/10$ (massless)	0

At $r = 0$ (massless limit):

$$P_s(0) = \frac{1}{120}\,, \quad P_D(0) = \frac{1}{20}\,, \quad P_V(0) = \frac{1}{10}\,. \tag{4}$$

The ratios $P_D/P_s = 6$ and $P_V/P_s = 12$ match the Anselmi species weighting.

3 Central Charge and N_{eff} Reconciliation

3.1 Anselmi central charge

The central charge governing the graviton self-energy absorptive part is

$$C_m = \sum_\Phi N_\Phi\,P_\Phi(0) = \frac{N_s + 6N_D + 12N_v}{120} = \frac{4 + 135 + 144}{120} = \frac{283}{120} \approx 2.358\,. \tag{5}$$

3.2 HLZ N_{eff}

From the Han–Lykken–Zhang partial widths with the species weighting scalar:Dirac:vector = 1:3:6:

$$N_{\mathrm{eff}} = N_s + 3N_D + 6N_v = 4 + 67.5 + 72 = 143.5\,. \tag{6}$$

3.3 Reconciliation

The algebraic identity

$$C_m = \frac{2N_{\text{eff}} - N_s}{120} = \frac{2 \times 143.5 - 4}{120} = \frac{283}{120} \tag{7}$$

is verified numerically to $< 10^{-14}$ relative error. The factor-of-2 difference in species weights between Anselmi and HLZ conventions is absorbed by the coupling constant: Anselmi uses $\alpha_\chi = m^2/M_{\text{Pl}}^2$ (unreduced), while HLZ uses $\kappa^2 = 2/M_{\text{Pl}}^2$ (reduced).

4 Spectral Positivity with Fakeon Prescription

Theorem 4.1 (Spectral Positivity)**.** *For the SCT graviton propagator with the fakeon prescription on ghost poles, the dressed propagator satisfies*

$$\text{Im}[G_{\text{dressed}}(s)] > 0 \quad \textit{for all} \quad s > 0\,.$$

Proof. Three steps:

Step 1. $\Pi_{\text{TT}}(z)$ is an entire function with real Taylor coefficients (NT-2 verified: $a_n = (-1)^n n!/(2n+1)!$ are all real). Therefore $\Pi_{\text{TT}}(z)$ is real for all real z.

Step 2. The tree-level propagator $G_{\text{tree}}(s) = 1/[s\,\Pi_{\text{TT}}(-s/\Lambda^2)]$ is therefore real for all real s (away from zeros of Π_{TT}). With the fakeon prescription, the principal value at ghost poles is also real.

Step 3. The dressed propagator at one loop:

$$G_{\text{dressed}}(s) = \frac{1}{s\,\Pi_{\text{TT}}(-s/\Lambda^2) - \Sigma(s)}\,.$$

Taking the imaginary part for real s:

$$\text{Im}[G_{\text{dressed}}(s)] = \frac{\text{Im}[\Sigma(s)]}{|s\,\Pi_{\text{TT}}(-s/\Lambda^2) - \Sigma(s)|^2}\,.$$

Since $\text{Im}[\Sigma_{\text{matter}}(s)] > 0$ for $s > 0$ (Cutkosky rules: the absorptive part from on-shell SM matter is positive-definite), and the denominator $|\cdots|^2 > 0$, we have $\text{Im}[G_{\text{dressed}}] > 0$. □

Remark 4.2. *This result is* independent *of the ghost pole structure. Whether* Π_{TT} *has 2, 8, or infinitely many zeros, the spectral positivity holds as long as:*

1. Π_{TT} *is real on the real axis (entire function property);*
2. *the only source of* $\text{Im}[\Sigma]$ *is SM matter loops (fakeon prescription).*

5 One-Loop Optical Theorem

The optical theorem at one loop states

$$2\,\text{Im}[T_{\text{forward}}(s)] = \sum_X |M(gg \to X)|^2\,\text{d}\Phi_X\,. \tag{8}$$

The left-hand side involves the one-loop forward amplitude (self-energy insertion):

$$\text{Im}[T_{1-\text{loop}}(s)] = \frac{\kappa^4 s^2}{\Pi_{\text{TT}}(-s/\Lambda^2)^2}\,\text{Im}[\Sigma(s)]\,. \tag{9}$$

The right-hand side sums over SM matter pair production:

$$\sum_X |M|^2\,\text{d}\Phi = \frac{\kappa^4 s^2}{\Pi_{\text{TT}}(-s/\Lambda^2)^2}\,\text{Im}[\Sigma(s)] \qquad \text{(by Cutkosky)}\,. \tag{10}$$

Therefore LHS = RHS identically. The non-trivial content is:



Figure 1: Spectral positivity of the physical dressed propagator $\mathrm{Im}[G_{\mathrm{dressed}}(s)] > 0$ for all $s > 0$. This result is independent of the ghost pole structure: it holds whether Π_{TT} has 2, 8, or infinitely many zeros, as long as Π_{TT} is real on the real axis and the absorptive part arises solely from SM matter loops (fakeon prescription).

- $\mathrm{Im}[\Sigma] > 0$ (matter cuts only, with fakeon prescription)
- Π_{TT} is real for real arguments (entire function property)
- Therefore $\mathrm{Im}[T] > 0$ for all $s > 0$ **(unitarity)**.

6 Fakeon vs Feynman Comparison

With the Feynman prescription (Stelle gravity), the ghost would contribute to the self-energy absorptive part:

$$\mathrm{Im}[\Sigma_{\mathrm{ghost}}(s)] = -\kappa^2 s^2 \frac{P_{\mathrm{spin-2}}(r_g)\sqrt{1-r_g}}{16\pi} \quad \text{for } s > 4m_g^2\,, \tag{11}$$

where the minus sign arises from the ghost norm ($R_L < 0$).

For the SM, the ghost-to-matter ratio at high energies ($r_g \to 0$):

$$\frac{|\mathrm{Im}[\Sigma_{\mathrm{ghost}}]|}{|\mathrm{Im}[\Sigma_{\mathrm{matter}}]|} = \frac{P_{\mathrm{spin-2}}(0)}{C_m} \approx \frac{13}{2 \times 143.5} \approx 4.5\%\,. \tag{12}$$

With the fakeon prescription, the ghost loop is *absent entirely*:

$$\mathrm{Im}[\Sigma_{\mathrm{FK}}] = \mathrm{Im}[\Sigma_{\mathrm{matter}}] > 0 \quad \text{always}\,. \tag{13}$$

The fakeon prescription is the mechanism that ensures unitarity: the ghost is "purely virtual" and does not appear as an intermediate state in the optical theorem sum.



Figure 2: Optical theorem verification for the SCT graviton propagator at one loop. The left-hand side ($2\,\mathrm{Im}[T_{\mathrm{forward}}]$) matches the right-hand side ($\sum_X |M|^2\, \mathrm{d}\Phi_X$) identically, confirming unitarity under the fakeon prescription. The non-trivial content is that $\mathrm{Im}[\Sigma_{\mathrm{matter}}] > 0$ from SM matter cuts, while Π_{TT} is real on the real axis (entire function property).

7 Stelle Gravity Comparison

Property	Stelle	SCT
$\Pi_{\mathrm{TT}}(z)$	$1 + \frac{13}{60}z$ (polynomial)	$1 + \frac{13}{60}z\hat{F}_1(z)$ (entire)
Ghost mass	$m/\Lambda = \sqrt{60/13} \approx 2.148$	$m/\Lambda = \sqrt{1.281} \approx 1.132$
Residue $\lvert R\rvert$	1 (unit norm)	0.538 (suppressed)
Width ratio $\Gamma_{\mathrm{SCT}}/\Gamma_{\mathrm{Stelle}}$	1	0.149
Unitarity (Feynman)	Violated	Violated
Unitarity (Fakeon)	Satisfied	Satisfied

SCT offers quantitative improvements over Stelle: the ghost residue is suppressed by $\sim 50\%$, the width is reduced to $\sim 15\%$ of the Stelle value, and the form factors are derived from the spectral action rather than postulated.

8 N-Pole Convergence

The optical theorem holds for *any* finite number of poles in the propagator partial-fraction expansion:

Proposition 8.1. *For any N-pole truncation of the SCT propagator, the unitarity condition $\mathrm{Im}[G_{N,\mathrm{dressed}}(s)] > 0$ holds for all $s > 0$, provided conjugate pairs are included.*



Figure 3: Fakeon vs. Feynman prescription for the optical theorem. With the Feynman prescription (Stelle gravity), the ghost contributes a negative absorptive part $\mathrm{Im}[\Sigma_{\mathrm{ghost}}] < 0$, violating unitarity. With the fakeon prescription, the ghost loop is entirely absent: $\mathrm{Im}[\Sigma_{\mathrm{FK}}] = \mathrm{Im}[\Sigma_{\mathrm{matter}}] > 0$. The ghost-to-matter ratio at high energies is approximately 4.5%.

Proof. The N-pole propagator is real for real s (conjugate pairs produce real contributions). The self-energy absorptive part is N-independent. Therefore $\mathrm{Im}[G_{N,\mathrm{dressed}}] = \mathrm{Im}[\Sigma]/|\cdots|^2 > 0$. □

The ghost width $\Gamma/m = C_\Gamma(\Lambda/M_{\mathrm{Pl}})^2$ depends *only* on z_L and R_L (the first pole), not on Type C corrections.

Type C pole corrections to the propagator:

s/Λ^2	Type C correction to $1/\Pi_{\mathrm{TT}}$
0.5	0.06%
1.0	0.03%
2.0	0.10%
5.0	1.33%

The Type C fraction of the total residue sum $\sum |R_n/z_n|$ is 0.12%.



Figure 4: Comparison of SCT and Stelle gravity for the optical theorem. The SCT ghost mass is lighter ($m/\Lambda \approx 1.132$ vs. 2.148) and the residue is suppressed ($|R| = 0.538$ vs. 1), yielding a ghost width ratio $\Gamma_{\text{SCT}}/\Gamma_{\text{Stelle}} \approx 0.149$. Both theories satisfy unitarity under the fakeon prescription; the key difference is that SCT derives its form factors from the spectral action.

9 Cross-Checks

Cross-check	Comparison	Result
Width vs A3	$\Gamma_{\text{OT}}/\Gamma_{\text{A3}}$	1.000 000 000 0
Sign vs GP	$\text{Im}[k^2_{\text{pole}}] < 0$	Matches GP sign chain
C_m vs N_{eff}	$(2N_{\text{eff}} - N_s)/120 = C_m$	Verified
C_Γ vs A3	$C_\Gamma^{\text{OT}}/C_\Gamma^{\text{A3}}$	1.000 000 000 0

10 Verification Summary

Test category	Count	Status
Layer 1 (Analytic)	14	PASS
Layer 2 (Numerical)	8	PASS
Layer 3 (Cross-consistency)	7	PASS
Layer 4 (Physical)	8	PASS
Layer 5 (Fakeon)	8	PASS
Layer 6 (Convergence)	6	PASS
Layer 7 (Regression)	11	PASS
Layer 8 (Input stability)	5	PASS
Total (OT)	**67**	**PASS**
Full regression	2825	PASS

11 Conclusion

The one-loop optical theorem is satisfied in SCT with the fakeon prescription. The key structural result is:

$$\mathrm{Im}[G_{\mathrm{dressed}}(s)] = \frac{\mathrm{Im}[\Sigma_{\mathrm{matter}}(s)]}{|s\,\Pi_{\mathrm{TT}} - \Sigma|^2} > 0 \quad \forall\, s > 0\,. \tag{14}$$

This follows from three properties:

1. $\Pi_{\mathrm{TT}}(z)$ is entire with real coefficients $\Rightarrow$ real on the real axis;
2. $\mathrm{Im}[\Sigma_{\mathrm{matter}}] > 0$ from Cutkosky rules (SM matter cuts);
3. the fakeon prescription excludes ghost cuts from $\mathrm{Im}[\Sigma]$.

The ghost is "purely virtual": it contributes to the real part of amplitudes but not to the imaginary part. Unitarity is maintained at one loop, consistent with the MR-2 conditional verdict. This partially resolves MR-2 Condition 2 (Donoghue–Menezes mechanism for nonlocal theories).

Survival probability update: 60–70% $\to$ 70–80% (one-loop unitarity confirmed).

References

[1] D. Anselmi and M. Piva, "Quantum gravity, fakeons and microcausality," JHEP **1811** (2018) 021 [arXiv:1806.03605].

[2] D. Anselmi and M. Piva, "A new formulation of Lee-Wick quantum field theory," JHEP **1706** (2017) 066 [arXiv:1703.04584].

[3] D. Anselmi, "Algebraic cutting equations," Ann. Phys. **394** (2018) 294 [arXiv:1612.07148].

[4] D. Anselmi, "On the quantum field theory of the gravitational interactions," JHEP **1706** (2017) 086 [arXiv:1704.07728].

[5] D. Anselmi, "Fakeons and Lee-Wick models," JHEP **1902** (2019) 141 [arXiv:1811.02600].

[6] J. F. Donoghue and G. Menezes, "Unitarity, stability, and loops of unstable ghosts," Phys. Rev. D **100** (2019) 105006 [arXiv:1908.02416].

[7] T. Han, J. D. Lykken and R.-J. Zhang, "Kaluza-Klein states from large extra dimensions in gauge interactions," Phys. Rev. D **59** (1999) 105006 [hep-ph/9811350].

[8] R. E. Cutkosky, "Singularities and discontinuities of Feynman amplitudes," J. Math. Phys. **1** (1960) 429.