

# Hilbert's Nullstellensatz: Computation and Proof

David Snider

Directed Reading Program  
UNC Department of Mathematics

April 2022


# About the Talk

- ▶ Why Give This Talk?
  - ▶ Expose undergraduates to an area of modern research
  - ▶ Gain an appreciation for algebra's applications



# About the Talk

- ▶ Why Give This Talk?
  - ▶ Expose undergraduates to an area of modern research
  - ▶ Gain an appreciation for algebra's applications
- ▶ Agenda
  - ▶ Motivation / Basic Terms (V and I)
  - ▶ Computing  $I(P)$
  - ▶ Proof of the Nullstellensatz



## Motivation



## Motivation





## Motivation




$$x^2 + y^2 = 1, \text{ or}$$

## Motivation




$$x^2 + y^2 = 1, \text{ or}$$
$$x^2 + y^2 - 1 = 0$$

## Motivation






# Motivation




$$z^2 + x^2 - y^2 = 0$$

# Motivation




$$\begin{aligned} z^2 + x^2 - y^2 &= 0 \\ z - 1 &= 0 \end{aligned}$$

# Motivation



$$\{z^2 + x^2 - y^2 = 0\} \cap \{z - 1 = 0\}$$

# Motivation



$$V(z^2 + x^2 - y^2, z - 1)$$

## V and I

For  $k$  a field,  $k[X_1, \dots, X_n]$  is a ring.

## V and I

For  $k$  a field,  $k[X_1, \dots, X_n]$  is a ring.

Eg.  $p \in \mathbb{R}[X] \implies p(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n, a_i \in \mathbb{R}$

## V and I

For  $k$  a field,  $k[X_1, \dots, X_n]$  is a ring.

Eg.  $p \in \mathbb{R}[X] \implies p(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n, a_i \in \mathbb{R}$

$p \in \mathbb{R}[X, Y] \implies p(x, y) = \sum_{i,j} a_{ij}x^i y^j$  (finite sum)

## V and I

In  $k[X_1, \dots, X_n]$ , ideals are generated by a finite number of polynomials.



## V and I

In  $k[X_1, \dots, X_n]$ , ideals are generated by a finite number of polynomials.

Eg.  $(x^2 + y^2 - 1) \subset \mathbb{R}[X, Y]$

## V and I

In  $k[X_1, \dots, X_n]$ , ideals are generated by a finite number of polynomials.

Eg.  $(x^2 + y^2 - 1) \subset \mathbb{R}[X, Y]$

►  $\{(x^2 + y^2 - 1)r : r \in \mathbb{R}[X, Y]\}$

## V and I

In  $k[X_1, \dots, X_n]$ , ideals are generated by a finite number of polynomials.

Eg.  $(x^2 + y^2 - 1) \subset \mathbb{R}[X, Y]$

►  $\{(x^2 + y^2 - 1)r : r \in \mathbb{R}[X, Y]\}$

Eg.  $(x^2 + y^2 - 1, x - y) \subset \mathbb{R}[X, Y]$

## V and I

In  $k[X_1, \dots, X_n]$ , ideals are generated by a finite number of polynomials.

Eg.  $(x^2 + y^2 - 1) \subset \mathbb{R}[X, Y]$

►  $\{(x^2 + y^2 - 1)r : r \in \mathbb{R}[X, Y]\}$

Eg.  $(x^2 + y^2 - 1, x - y) \subset \mathbb{R}[X, Y]$

►  $\{(x^2 + y^2 - 1)r_1 + (x - y)r_2 : r_1, r_2 \in \mathbb{R}[X, Y]\}$

## V and I

### Definition

Let  $I$  be an ideal in  $k[X_1, \dots, X_n] =: R$ .


## V and I

### Definition

Let  $I$  be an ideal in  $k[X_1, \dots, X_n] =: R$ .


Define  $V(I) := \{x \in k^n : f(x) = 0, \forall f \in I\}$

V and I



$$V(x^2 + y^2 - 1)$$


V and I



$$V(z^2 + x^2 - y^2)$$
$$V(z - 1)$$



V and I



$$V(z^2 + x^2, z - 1)$$

$V$  and  $I$

Does  $V$  have an inverse?

V and I

### Definition

Let  $C \subset k^n$  and let  $R = k[X_1, \dots, X_n]$ .

## V and I

### Definition

Let  $C \subset k^n$  and let  $R = k[X_1, \dots, X_n]$ .

Define  $I(C) := \{f \in R : f(x) = 0, \forall x \in C\}$ .

## V and I

### Definition

Let  $C \subset k^n$  and let  $R = k[X_1, \dots, X_n]$ .

Define  $I(C) := \{f \in R : f(x) = 0, \forall x \in C\}$ .

This is indeed an ideal.

## V and I

### Definition

Let  $C \subset k^n$  and let  $R = k[X_1, \dots, X_n]$ .

Define  $I(C) := \{f \in R : f(x) = 0, \forall x \in C\}$ .

This is indeed an ideal.

Let  $f_1, f_2 \in I(C)$ .

Then  $\forall x \in C, (f_1 + f_2)(x) = 0$ . So  $f_1 + f_2 \in I(C)$ .

## V and I

### Definition

Let  $C \subset k^n$  and let  $R = k[X_1, \dots, X_n]$ .

Define  $I(C) := \{f \in R : f(x) = 0, \forall x \in C\}$ .

This is indeed an ideal.


Let  $f_1, f_2 \in I(C)$ .

Then  $\forall x \in C, (f_1 + f_2)(x) = 0$ . So  $f_1 + f_2 \in I(C)$ .

Let  $f \in I(C), r \in R$ .

Then  $(fr)(x) := f(x)r(x) = 0, \forall x \in C$ . So  $fr \in I(C)$ .

V and I




$$I(C) =$$

Let  $C = \{x \in \mathbb{R}^2 : |x| = 1\}$ .



V and I




Let  $C = \{x \in \mathbb{R}^2 : |x| = 1\}$ .

$$I(C) = (x^2 + y^2 - 1).$$

$V$  and  $I$


Are  $V$  and  $I$  inverses of each other?

## A Computation




$$V(X^2 + Y^2 - 1), V(Y - 1)$$

## A Computation



$$V(X^2 + Y^2 - 1, Y - 1)$$

## A Computation



$$J = (X^2 + Y^2 - 1, Y - 1)$$
$$I(V(J)) = ?$$

## A Computation

$$I(\{(0, 1)\}) = ?$$

## A Computation

$$I(\{(0, 1)\}) = ?$$

Candidate ideal:  $(X, Y - 1)$ .

## A Computation

$$I(\{(0, 1)\}) = ?$$

Candidate ideal:  $(X, Y - 1)$ .

Let  $f = p_1X + p_2(Y - 1)$ .  $f(0, 1) =$



## A Computation

$$I(\{(0, 1)\}) = ?$$

Candidate ideal:  $(X, Y - 1)$ .

Let  $f = p_1X + p_2(Y - 1)$ .  $f(0, 1) = 0 + 0 = 0$ . So  $f \in I(\{(0, 1)\})$

## A Computation

$$I(\{(0, 1)\}) = ?$$

Candidate ideal:  $(X, Y - 1)$ .

Let  $f = p_1X + p_2(Y - 1)$ .  $f(0, 1) = 0 + 0 = 0$ . So  $f \in I(\{(0, 1)\})$

Let  $f \notin (X, Y - 1)$ . Then,  $f$  cannot be written  $p_1X + p_2(Y - 1)$ .

## A Computation

$$I(\{(0, 1)\}) = ?$$

Candidate ideal:  $(X, Y - 1)$ .

Let  $f = p_1X + p_2(Y - 1)$ .  $f(0, 1) = 0 + 0 = 0$ . So  $f \in I(\{(0, 1)\})$

Let  $f \notin (X, Y - 1)$ . Then,  $f$  cannot be written  $p_1X + p_2(Y - 1)$ .

Then,  $f = p_1X + p_2(Y - 1) + p_3$ , where  $p_3$  satisfies...

## A Computation

$$I(\{(0, 1)\}) = ?$$

Candidate ideal:  $(X, Y - 1)$ .

Let  $f = p_1X + p_2(Y - 1)$ .  $f(0, 1) = 0 + 0 = 0$ . So  $f \in I(\{(0, 1)\})$

Let  $f \notin (X, Y - 1)$ . Then,  $f$  cannot be written  $p_1X + p_2(Y - 1)$ .

Then,  $f = p_1X + p_2(Y - 1) + p_3$ , where  $p_3$  satisfies...

Neither  $X$  nor  $(Y - 1)$  divides  $p_3$ .

## A Computation

$$I(\{(0, 1)\}) = ?$$

Candidate ideal:  $(X, Y - 1)$ .

Let  $f = p_1X + p_2(Y - 1)$ .  $f(0, 1) = 0 + 0 = 0$ . So  $f \in I(\{(0, 1)\})$

Let  $f \notin (X, Y - 1)$ . Then,  $f$  cannot be written  $p_1X + p_2(Y - 1)$ .

Then,  $f = p_1X + p_2(Y - 1) + p_3$ , where  $p_3$  satisfies...

Neither  $X$  nor  $(Y - 1)$  divides  $p_3$ .

$p_3 = a_0 + a_1Y + \dots + a_nY^n$ , where  $(Y - 1)$  does not divide  $p_3$ .

## A Computation

$$I(\{(0, 1)\}) = ?$$

Candidate ideal:  $(X, Y - 1)$ .

Let  $f = p_1X + p_2(Y - 1)$ .  $f(0, 1) = 0 + 0 = 0$ . So  $f \in I(\{(0, 1)\})$

Let  $f \notin (X, Y - 1)$ . Then,  $f$  cannot be written  $p_1X + p_2(Y - 1)$ .

Then,  $f = p_1X + p_2(Y - 1) + p_3$ , where  $p_3$  satisfies...

Neither  $X$  nor  $(Y - 1)$  divides  $p_3$ .

$p_3 = a_0 + a_1Y + \dots + a_nY^n$ , where  $(Y - 1)$  does not divide  $p_3$ .

By a MATH 578 proof, we know  $p_3$  has a root at  $y = 1$  iff  $y - 1$  divides  $p_3$ .

## A Computation

$$I(\{(0, 1)\}) = ?$$

Candidate ideal:  $(X, Y - 1)$ .

Let  $f = p_1X + p_2(Y - 1)$ .  $f(0, 1) = 0 + 0 = 0$ . So  $f \in I(\{(0, 1)\})$

Let  $f \notin (X, Y - 1)$ . Then,  $f$  cannot be written  $p_1X + p_2(Y - 1)$ .

Then,  $f = p_1X + p_2(Y - 1) + p_3$ , where  $p_3$  satisfies...

Neither  $X$  nor  $(Y - 1)$  divides  $p_3$ .

$p_3 = a_0 + a_1Y + \dots + a_nY^n$ , where  $(Y - 1)$  does not divide  $p_3$ .

By a MATH 578 proof, we know  $p_3$  has a root at  $y = 1$  iff  $y - 1$  divides  $p_3$ .

So  $p_3(X, 1) \neq 0$ .

## A Computation

$$I(\{(0, 1)\}) = ?$$

Candidate ideal:  $(X, Y - 1)$ .

Let  $f = p_1X + p_2(Y - 1)$ .  $f(0, 1) = 0 + 0 = 0$ . So  $f \in I(\{(0, 1)\})$

Let  $f \notin (X, Y - 1)$ . Then,  $f$  cannot be written  $p_1X + p_2(Y - 1)$ .

Then,  $f = p_1X + p_2(Y - 1) + p_3$ , where  $p_3$  satisfies...

Neither  $X$  nor  $(Y - 1)$  divides  $p_3$ .

$p_3 = a_0 + a_1Y + \dots + a_nY^n$ , where  $(Y - 1)$  does not divide  $p_3$ .

By a MATH 578 proof, we know  $p_3$  has a root at  $y = 1$  iff  $y - 1$  divides  $p_3$ .

So  $p_3(X, 1) \neq 0$ .

Thus,  $f(0, 1) = 0 + 0 + p_3(0, 1) \neq 0$ . So  $f \notin I(\{(0, 1)\})$ .



## A Computation

So  $J = (X^2 + Y^2 - 1, Y - 1)$ .

And  $I(V(J)) = (X, Y - 1)$ .

## A Computation

So  $J = (X^2 + Y^2 - 1, Y - 1)$ .

And  $I(V(J)) = (X, Y - 1)$ .

$I(V(J)) \neq J$  by a similar proof.

## A Computation

So  $J = (X^2 + Y^2 - 1, Y - 1)$ .

And  $I(V(J)) = (X, Y - 1)$ .

$I(V(J)) \neq J$  by a similar proof.

$I$  and  $V$  are not strict inverses of each other.

## A Computation

Likewise, if  $J = (X^2)$ ,  
 $V(J) =$

## A Computation

Likewise, if  $J = (X^2)$ ,

$V(J)$  = the y-axis.

$I(V(J)) =$

## A Computation

Likewise, if  $J = (X^2)$ ,

$V(J)$  = the y-axis.

$I(V(J)) = (X)$ .

## A Computation

Likewise, if  $J = (X^2)$ ,

$V(J)$  = the y-axis.

$I(V(J)) = (X)$ .

$X \in (X)$  but  $X \notin (X^2)$ .

## Nullstellensatz

Let  $k$  be an algebraically closed field (such as  $\mathbb{C}$ ).



## Nullstellensatz

Let  $k$  be an algebraically closed field (such as  $\mathbb{C}$ ).

- ▶ a) Every maximal ideal of the polynomial ring  $A = k[X_1, \dots, X_n]$  is  $\dots I(\{(c_1, \dots, c_n)\})$  for some  $(c_1, \dots, c_n) \in k^n$ .

# Nullstellensatz

Let  $k$  be an algebraically closed field (such as  $\mathbb{C}$ ).

- ▶ a) Every maximal ideal of the polynomial ring  $A = k[X_1, \dots, X_n]$  is  $\dots I(\{(c_1, \dots, c_n)\})$  for some  $(c_1, \dots, c_n) \in k^n$ .
- ▶ b) Let  $J \subset A$  be an ideal,  $J \neq (1)$ ; then  $V \neq \emptyset$ .
  - ▶ Weak Nullstellensatz

# Nullstellensatz

Let  $k$  be an algebraically closed field (such as  $\mathbb{C}$ ).

- ▶ a) Every maximal ideal of the polynomial ring  $A = k[X_1, \dots, X_n]$  is  $\dots I(\{(c_1, \dots, c_n)\})$  for some  $(c_1, \dots, c_n) \in k^n$ .
- ▶ b) Let  $J \subset A$  be an ideal,  $J \neq (1)$ ; then  $V \neq \emptyset$ .
  - ▶ Weak Nullstellensatz
- ▶ c) For any  $f \in I(V(J))$ ,  $\exists n \in \mathbb{N}$  such that  $f^n \in J$ .
  - ▶ Strong Nullstellensatz

# Nullstellensatz

In an algebraically closed field,

- ▶ If a polynomial is **non-constant**, then it **has a zero** (FTA).
- ▶ If the ideal generated by a set of polynomials is **not the ideal of a constant**, then the **vanishing set of those polynomials is nonempty** (Weak NSS).

## Nullstellensatz

(a) Every maximal ideal of the polynomial ring  $A = k[X_1, \dots, X_n]$  is  $\dots I(\{(c_1, \dots, c_n)\})$  for some  $(c_1, \dots, c_n) \in k^n$ .

► Given  $M$ , use  $\phi : k \rightarrow k[X_1, \dots, X_n] \rightarrow k[X_1, \dots, X_n]/M$

## Nullstellensatz

(a) Every maximal ideal of the polynomial ring  $A = k[X_1, \dots, X_n]$  is  $\dots I(\{(c_1, \dots, c_n)\})$  for some  $(c_1, \dots, c_n) \in k^n$ .

- ▶ Given  $M$ , use  $\phi : k \rightarrow k[X_1, \dots, X_n] \rightarrow k[X_1, \dots, X_n]/M$
- ▶ Hard Fact:  $\phi$  is an isomorphism.

## Nullstellensatz

(a) Every maximal ideal of the polynomial ring  $A = k[X_1, \dots, X_n]$  is  $\dots I(\{(c_1, \dots, c_n)\})$  for some  $(c_1, \dots, c_n) \in k^n$ .

- ▶ Given  $M$ , use  $\phi : k \rightarrow k[X_1, \dots, X_n] \rightarrow k[X_1, \dots, X_n]/M$
- ▶ Hard Fact:  $\phi$  is an isomorphism.
- ▶  $X_i \mapsto b_i$ . Let  $a_i = \phi^{-1}(b_i)$ . Then  $X_i - a_i \in \ker f_2 = M$ .

## Nullstellensatz

(a) Every maximal ideal of the polynomial ring  $A = k[X_1, \dots, X_n]$  is  $\dots I(\{(c_1, \dots, c_n)\})$  for some  $(c_1, \dots, c_n) \in k^n$ .

- ▶ Given  $M$ , use  $\phi : k \rightarrow k[X_1, \dots, X_n] \rightarrow k[X_1, \dots, X_n]/M$
- ▶ Hard Fact:  $\phi$  is an isomorphism.
- ▶  $X_i \mapsto b_i$ . Let  $a_i = \phi^{-1}(b_i)$ . Then  $X_i - a_i \in \ker f_2 = M$ .
- ▶ Thus,  $(X_1 - a_1, \dots, X_n - a_n) \subset M$ , and thus  $= M$ .



## Nullstellensatz

(b) Let  $J \subset A$  be an ideal,  $J \neq (1)$ ; then  $V \neq \emptyset$ . (Weak Nullstellensatz)

►  $J \neq R \implies \exists M$  maximal ideal w/  $J \subset M$ .

## Nullstellensatz

(b) Let  $J \subset A$  be an ideal,  $J \neq (1)$ ; then  $V \neq \emptyset$ . (Weak Nullstellensatz)

- ▶  $J \neq R \implies \exists M$  maximal ideal w/  $J \subset M$ .
- ▶  $V(M) = \{P\}$ , and  $J \subset M$ , so  $V(J) \ni P$ .

## Nullstellensatz

(c) For any  $J \subset A$ ,  $I(V(J)) = \sqrt{J}$ . (Strong Nullstellensatz)

- ▶ Rabinowitsch Trick
- ▶ Look it up on Wikipedia!

# Thank You!

Acknowledgements:

- ▶ Hunter Dinkins, my DRP Mentor
- ▶ The DRP Committee

## Citation

Miles Reid. *Undergraduate Algebraic Geometry*. Cambridge University Press, Cambridge, 1989.