

CONSTRAINTS ON N-LATTICE SIMPLICES OF VOLUME $\frac{1}{n!}$ IN N-CUBE, UNIQUE UNDER SYMMETRY


DAVID SNIDER

ROSE-HULMAN INSTITUTE OF TECHNOLOGY
MATHEMATICS REU



JULY 20, 2022

MENTORED BY MCCABE OLSEN


Introduction



Introduction



Introduction




Definition

A **(convex) polytope** in \mathbb{R}^n is the convex hull of finitely many points in \mathbb{R}^n . That is,


$$\text{conv}\{v_1, \dots, v_d\} = \left\{ \sum_{i=1}^d \lambda_i v_i : \sum_{i=1}^d \lambda_i = 1, \lambda_i \geq 0 \right\}$$

What is a triangulation of a polytope \mathcal{P} ?

What is a triangulation of a polytope \mathcal{P} ?




What is a triangulation of a polytope \mathcal{P} ?




- use vertices of \mathcal{P}

What is a triangulation of a polytope \mathcal{P} ?




- use vertices of \mathcal{P}
- (UP) - all points in \mathcal{P} are contained in a triangle

What is a triangulation of a polytope \mathcal{P} ?




- use vertices of \mathcal{P}
- (UP) - all points in \mathcal{P} are contained in a triangle
- (IP) - two triangles intersect in a face of smaller dimension

What is a triangulation of a polytope \mathcal{P} ?



- use vertices of \mathcal{P}
- (UP) - all points in \mathcal{P} are contained in a triangle
- (IP) - two triangles intersect in a face of smaller dimension



Using triangles to triangulate higher dimensional polytopes?

Using triangles to triangulate higher dimensional polytopes?

This would violate condition (UP).

Using triangles to triangulate higher dimensional polytopes?

This would violate condition (UP).

Definition

A **d-simplex** in \mathbb{R}^n is the convex hull of $d + 1$ affinely independent points in \mathbb{R}^n .

Using triangles to triangulate higher dimensional polytopes?

This would violate condition (UP).

Definition

A **d-simplex** in \mathbb{R}^n is the convex hull of $d + 1$ affinely independent points in \mathbb{R}^n .

$\{v_1, \dots, v_{d+1}\}$ are **affinely independent** if $\{v_1 - v_{d+1}, \dots, v_d - v_{d+1}\}$ is linearly independent.


Definition

A **triangulation** \mathcal{T} of a full dimensional polytope \mathcal{P} in \mathbb{R}^n is a finite collection of n -simplices whose vertices are vertices of \mathcal{P} and that satisfies the following two properties:



$$\bigcup_{\Delta \in \mathcal{T}} \Delta = \mathcal{P} \quad (\text{UP})$$

Given $\Delta_1, \Delta_2 \in \mathcal{T}$, $\Delta_1 \cap \Delta_2$ is a common face (possibly empty)
(IP)


Introduction





Introduction






Introduction




Introduction




Introduction




Introduction



Introduction





Introduction



Volume = $1/6$

Introduction



Volume = $1/3$

Definition

A triangulation of an n -dimensional polytope in \mathbb{R}^n is **unimodular** if all its n -simplices are translations of linear transformations of the standard simplex $\text{conv}\{0, e_1, \dots, e_n\}$, where the linear transformation's components are integers, and its determinant is ± 1 .

How many n -simplices of volume $\frac{1}{n!}$ fit in $[0, 1]^n$, unique under symmetry?

How many n -simplices of volume $\frac{1}{n!}$ fit in $[0, 1]^n$, unique under symmetry?

A counting problem.

Question

How many n -simplices of volume $\frac{1}{n!}$ fit in $[0, 1]^n$, unique under symmetry?

A counting problem.

Why ask this question?

How many n -simplices of volume $\frac{1}{n!}$ fit in $[0, 1]^n$, unique under symmetry?

A counting problem.

Why ask this question?

- Applications in Linear Algebra [1]


How many n -simplices of volume $\frac{1}{n!}$ fit in $[0, 1]^n$, unique under symmetry?

A counting problem.


Why ask this question?

- Applications in Linear Algebra [1]
- Unimodular triangulations not well understood.

The Representation



$$\left\{ \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}$$



$$\left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} \right\}$$

The Representation

- Want a representation that respects equivalence under symmetry.

The Representation

- Want a representation that respects equivalence under symmetry.
- Symmetries preserve distance.

The Representation

- Want a representation that respects equivalence under symmetry.
- Symmetries preserve distance.
- \implies The important information is the distance between vertices.


Definition

Let x, y be elements of $[0, 1]^n \cap \mathbb{Z}^n$. Define the **taxi-cab distance** of x, y , denoted $\delta(x, y)$, as the minimum number of standard basis vectors that must be added to or subtracted from x to get y .

The Representation

Definition

Let x, y be elements of $[0, 1]^n \cap \mathbb{Z}^n$. Define the **taxi-cab distance** of x, y , denoted $\delta(x, y)$, as the minimum number of standard basis vectors that must be added to or subtracted from x to get y .




$$\delta((1, 0, 0), (0, 1, 1)) = 3$$

The Representation

Definition

Let x, y be elements of $[0, 1]^n \cap \mathbb{Z}^n$. Define the **taxi-cab distance** of x, y , denoted $\delta(x, y)$, as the minimum number of standard basis vectors that must be added to or subtracted from x to get y .




$$\delta((1, 0, 0), (0, 1, 1)) = 3$$

The Representation

Definition

Let x, y be elements of $[0, 1]^n \cap \mathbb{Z}^n$. Define the **taxi-cab distance** of x, y , denoted $\delta(x, y)$, as the minimum number of standard basis vectors that must be added to or subtracted from x to get y .




$$\delta((1, 0, 0), (0, 1, 1)) = 3$$

The Representation

Definition

Let x, y be elements of $[0, 1]^n \cap \mathbb{Z}^n$. Define the **taxi-cab distance** of x, y , denoted $\delta(x, y)$, as the minimum number of standard basis vectors that must be added to or subtracted from x to get y .



$$\delta((1, 0, 0), (0, 1, 1)) = 3$$

Definition

Let Δ be an d -simplex in \mathbb{Z}^n . Its **adjacency matrix** is

$$(a_{ij}) = \begin{cases} 0 & i = j \\ \delta(v_i, v_j) & i \neq j \end{cases}$$

Definition


Let Δ be an d -simplex in \mathbb{Z}^n . Its **adjacency matrix** is

$$(a_{ij}) = \begin{cases} 0 & i = j \\ \delta(v_i, v_j) & i \neq j \end{cases}$$


Definition

Define the multigraph $G = (V, E)$ of Δ as the graph of $d + 1$ vertices with a_{ij} edges between v_i, v_j .

The Representation




The Representation




$$\begin{pmatrix} 0 & 1 & 1 & 2 \\ 1 & 0 & 2 & 3 \\ 1 & 2 & 0 & 1 \\ 2 & 3 & 1 & 0 \end{pmatrix}$$

The Representation



$$\begin{pmatrix} 0 & 1 & 1 & 2 \\ 1 & 0 & 2 & 3 \\ 1 & 2 & 0 & 1 \\ 2 & 3 & 1 & 0 \end{pmatrix}$$



- Already, we've developed a constraint that Δ must follow.

- Already, we've developed a constraint that Δ must follow.
 - $n = 3$:

- Already, we've developed a constraint that Δ must follow.
 - $n = 3$:
 - $\binom{8}{4} = 70$ ways to choose 4 points

- Already, we've developed a constraint that Δ must follow.
 - $n = 3$:
 - $\binom{8}{4} = 70$ ways to choose 4 points
 - 6 multigraph representations

- Already, we've developed a constraint that Δ must follow.
 - $n = 3$:
 - $\binom{8}{4} = 70$ ways to choose 4 points
 - 6 multigraph representations
 - 3 “nice” simplices

- Already, we've developed a constraint that Δ must follow.
 - $n = 3$:
 - $\binom{8}{4} = 70$ ways to choose 4 points
 - 6 multigraph representations
 - 3 “nice” simplices
- In search of more constraints, we encountered interesting properties.

Theorem

Let Δ be an n -lattice simplex in $[0, 1]^n$ with adjacency matrix A and characteristic polynomial $(-1)^{n+1}\lambda^{n+1} + k_n\lambda^n + \dots + k_1\lambda + k_0$.

Theorem

Let Δ be an n -lattice simplex in $[0, 1]^n$ with adjacency matrix A and characteristic polynomial $(-1)^{n+1}\lambda^{n+1} + k_n\lambda^n + \dots + k_1\lambda + k_0$. Let $\mathcal{F} = \{f\}$ be the set of faces of Δ of dimension $\leq n - 1$, and $\mathcal{F} = \mathcal{F}_{-1} \cup \mathcal{F}_0 \cup \dots \cup \mathcal{F}_{n-1}$ partition the faces of \mathcal{F} by dimension, with \mathcal{F}_{-1} consisting only of the empty set. Then,

Theorem

Let Δ be an n -lattice simplex in $[0, 1]^n$ with adjacency matrix A and characteristic polynomial $(-1)^{n+1}\lambda^{n+1} + k_n\lambda^n + \dots + k_1\lambda + k_0$. Let $\mathcal{F} = \{f\}$ be the set of faces of Δ of dimension $\leq n-1$, and $\mathcal{F} = \mathcal{F}_{-1} \cup \mathcal{F}_0 \cup \dots \cup \mathcal{F}_{n-1}$ partition the faces of \mathcal{F} by dimension, with \mathcal{F}_{-1} consisting only of the empty set. Then,

$$k_0 = \det A$$

Theorem

Let Δ be an n -lattice simplex in $[0, 1]^n$ with adjacency matrix A and characteristic polynomial $(-1)^{n+1}\lambda^{n+1} + k_n\lambda^n + \dots + k_1\lambda + k_0$. Let $\mathcal{F} = \{f\}$ be the set of faces of Δ of dimension $\leq n-1$, and $\mathcal{F} = \mathcal{F}_{-1} \cup \mathcal{F}_0 \cup \dots \cup \mathcal{F}_{n-1}$ partition the faces of \mathcal{F} by dimension, with \mathcal{F}_{-1} consisting only of the empty set. Then,

$$\begin{aligned} k_0 &= \det A \\ -k_1 &= \sum_{f \in \mathcal{F}_{n-1}} \det A_f \end{aligned}$$

Theorem

Let Δ be an n -lattice simplex in $[0, 1]^n$ with adjacency matrix A and characteristic polynomial $(-1)^{n+1}\lambda^{n+1} + k_n\lambda^n + \dots + k_1\lambda + k_0$. Let $\mathcal{F} = \{f\}$ be the set of faces of Δ of dimension $\leq n-1$, and $\mathcal{F} = \mathcal{F}_{-1} \cup \mathcal{F}_0 \cup \dots \cup \mathcal{F}_{n-1}$ partition the faces of \mathcal{F} by dimension, with \mathcal{F}_{-1} consisting only of the empty set. Then,

$$\begin{aligned} k_0 &= \det A \\ -k_1 &= \sum_{f \in \mathcal{F}_{n-1}} \det A_f \\ &\dots \\ (-1)^q k_q &= \sum_{f \in \mathcal{F}_{n-q}} \det A_f \end{aligned}$$

Theorem

Let Δ be an n -lattice simplex in $[0, 1]^n$ with adjacency matrix A and characteristic polynomial $(-1)^{n+1}\lambda^{n+1} + k_n\lambda^n + \dots + k_1\lambda + k_0$. Let $\mathcal{F} = \{f\}$ be the set of faces of Δ of dimension $\leq n-1$, and $\mathcal{F} = \mathcal{F}_{-1} \cup \mathcal{F}_0 \cup \dots \cup \mathcal{F}_{n-1}$ partition the faces of \mathcal{F} by dimension, with \mathcal{F}_{-1} consisting only of the empty set. Then,

$$\begin{aligned}
 k_0 &= \det A \\
 -k_1 &= \sum_{f \in \mathcal{F}_{n-1}} \det A_f \\
 &\dots \\
 (-1)^q k_q &= \sum_{f \in \mathcal{F}_{n-q}} \det A_f \\
 &\dots \\
 k_n &= 0.
 \end{aligned} \tag{1}$$

Proof:

Let A be the adjacency matrix of G , and let $A - \lambda I = (a_{ij})$.

Proof:

Let A be the adjacency matrix of G , and let $A - \lambda I = (a_{ij})$. The characteristic polynomial of G is

$$\begin{aligned} p_G(\lambda) &= (-1)^{n+1} \lambda^{n+1} + k_n \lambda^n + \dots + k_1 \lambda + k_0 \\ &= \sum_{\sigma \in S_{n+1}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(n+1)\sigma(n+1)}, \end{aligned}$$

Proof:

Let A be the adjacency matrix of G , and let $A - \lambda I = (a_{ij})$. The characteristic polynomial of G is

$$\begin{aligned} p_G(\lambda) &= (-1)^{n+1} \lambda^{n+1} + k_n \lambda^n + \dots + k_1 \lambda + k_0 \\ &= \sum_{\sigma \in S_{n+1}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(n+1)\sigma(n+1)}, \end{aligned}$$

where S_{n+1} is the symmetric group on $n + 1$ elements, and

$$(-1)^\sigma := \begin{cases} 1 & \sigma \text{ is even} \\ -1 & \sigma \text{ is odd} \end{cases}$$

Proof:

Let A be the adjacency matrix of G , and let $A - \lambda I = (a_{ij})$. The characteristic polynomial of G is

$$\begin{aligned} p_G(\lambda) &= (-1)^{n+1} \lambda^{n+1} + k_n \lambda^n + \dots + k_1 \lambda + k_0 \\ &= \sum_{\sigma \in S_{n+1}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(n+1)\sigma(n+1)}, \end{aligned}$$

where S_{n+1} is the symmetric group on $n + 1$ elements, and

$$(-1)^\sigma := \begin{cases} 1 & \sigma \text{ is even} \\ -1 & \sigma \text{ is odd} \end{cases}$$

The coefficient k_n is the sum of terms that satisfy the condition that σ maps all elements except one to itself (Why?).

Proof:

Let A be the adjacency matrix of G , and let $A - \lambda I = (a_{ij})$. The characteristic polynomial of G is

$$\begin{aligned} p_G(\lambda) &= (-1)^{n+1} \lambda^{n+1} + k_n \lambda^n + \dots + k_1 \lambda + k_0 \\ &= \sum_{\sigma \in S_{n+1}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(n+1)\sigma(n+1)}, \end{aligned}$$

where S_{n+1} is the symmetric group on $n + 1$ elements, and

$$(-1)^\sigma := \begin{cases} 1 & \sigma \text{ is even} \\ -1 & \sigma \text{ is odd} \end{cases}$$

The coefficient k_n is the sum of terms that satisfy the condition that σ maps all elements except one to itself (Why?). But there are no such terms, since such maps do not exist in S_{n+1} .

$$\begin{aligned} p_G(\lambda) &= (-1)^{n+1} \lambda^{n+1} + k_n \lambda^n + \dots + k_1 \lambda + k_0 \\ &= \sum_{\sigma \in S_{n+1}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(n+1)\sigma(n+1)}. \end{aligned}$$

The coefficient k_1 is the sum of terms such that σ maps one and only one element to itself.

$$\begin{aligned} p_G(\lambda) &= (-1)^{n+1} \lambda^{n+1} + k_n \lambda^n + \dots + k_1 \lambda + k_0 \\ &= \sum_{\sigma \in S_{n+1}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(n+1)\sigma(n+1)}. \end{aligned}$$

The coefficient k_1 is the sum of terms such that σ maps one and only one element to itself. Call the set of such permutations, where k is that element, $S_{n+1,k}$, and let $\{A_{n-1,k}\}_{k=1}^{n+1}$ be the adjacency matrices of the $(n-1)$ -dimensional faces.

$$\begin{aligned} p_G(\lambda) &= (-1)^{n+1} \lambda^{n+1} + k_n \lambda^n + \dots + k_1 \lambda + k_0 \\ &= \sum_{\sigma \in S_{n+1}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(n+1)\sigma(n+1)}. \end{aligned}$$

The coefficient k_1 is the sum of terms such that σ maps one and only one element to itself. Call the set of such permutations, where k is that element, $S_{n+1,k}$, and let $\{A_{n-1,k}\}_{k=1}^{n+1}$ be the adjacency matrices of the $(n-1)$ -dimensional faces. Then,

$$\begin{aligned} k_1 \lambda &= \sum_{k=1}^{n+1} \sum_{\sigma \in S_{n+1,k}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(n+1)\sigma(n+1)} \\ &= -\lambda \sum_{k=1}^{n+1} \sum_{\sigma \in S_{n+1,k}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(k-1)\sigma(k-1)} a_{(k+1)\sigma(k+1)} \dots a_{(n+1)\sigma(n+1)} \\ &= -\lambda \sum_{k=1}^{n+1} \det A_{n-1,k}. \end{aligned}$$

$$\begin{aligned} p_G(\lambda) &= (-1)^{n+1} \lambda^{n+1} + k_n \lambda^n + \dots + k_1 \lambda + k_0 \\ &= \sum_{\sigma \in S_{n+1}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(n+1)\sigma(n+1)}. \end{aligned}$$

The coefficient k_1 is the sum of terms such that σ maps one and only one element to itself. Call the set of such permutations, where k is that element, $S_{n+1,k}$, and let $\{A_{n-1,k}\}_{k=1}^{n+1}$ be the adjacency matrices of the $(n-1)$ -dimensional faces. Then,

$$\begin{aligned} k_1 \lambda &= \sum_{k=1}^{n+1} \sum_{\sigma \in S_{n+1,k}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(n+1)\sigma(n+1)} \\ &= -\lambda \sum_{k=1}^{n+1} \sum_{\sigma \in S_{n+1,k}} (-1)^\sigma a_{1\sigma(1)} \dots a_{(k-1)\sigma(k-1)} a_{(k+1)\sigma(k+1)} \dots a_{(n+1)\sigma(n+1)} \\ &= -\lambda \sum_{k=1}^{n+1} \det A_{n-1,k}. \end{aligned}$$

Theorem

Let Δ be an n -lattice simplex in $[0, 1]^n$, and let $\mathcal{F} = \{f\}$ be the set of faces of Δ of dimension $\leq n - 1$, including the empty set, which we assign the trivial dimension -1 and polynomial $p_\emptyset = 1$. Let A_Δ be the adjacency matrix of Δ . Then,

$$p_\Delta(\lambda) = \det A_\Delta - \sum_{f \in \mathcal{F}} \lambda^{n - \dim f} p_f(\lambda) \quad (2)$$

Proof is similar.

Conjecture

Let Δ be an n -lattice simplex in $[0, 1]^n$ with normalized volume 1 and adjacency matrix A .

Conjecture

Let Δ be an n -lattice simplex in $[0, 1]^n$ with normalized volume 1 and adjacency matrix A . Then

$$\det A = (-1)^n 2^{n-1} n. \quad (3)$$

Conjecture

Let Δ be an n -lattice simplex in $[0, 1]^n$ with normalized volume 1 and adjacency matrix A . Then

$$\det A = (-1)^n 2^{n-1} n. \quad (3)$$

- Tested on non-regular triangulation given by de Loera [2].

Conjecture

Let Δ be an n -lattice simplex in $[0, 1]^n$ with normalized volume 1 and adjacency matrix A . Then

$$\det A = (-1)^n 2^{n-1} n. \quad (3)$$

- Tested on non-regular triangulation given by de Loera [2].
- Interpretation

Conjecture

Let Δ be an n -lattice simplex in $[0, 1]^n$ with normalized volume 1 and adjacency matrix A . Then

$$\det A = (-1)^n 2^{n-1} n. \quad (3)$$

- Tested on non-regular triangulation given by de Loera [2].
- Interpretation
- Tighter bound on simplices unique under symmetry.

Conjecture

Let Δ be an n -lattice simplex in $[0, 1]^n$ with normalized volume 1 and adjacency matrix A . Then

$$\det A = (-1)^n 2^{n-1} n. \quad (3)$$

- Tested on non-regular triangulation given by de Loera [2].
- Interpretation
- Tighter bound on simplices unique under symmetry.

Proposition

To prove the conjecture, it suffices to show that if $A^{-1} = (b_{ij})$,

$$-\left(\sum_{i,j=1,\dots,n+1} b_{ij}\right) = 2/n. \quad (4)$$

Conjecture

Let Δ be an n -lattice simplex in $[0, 1]^n$ with normalized volume 1 and adjacency matrix A . Then

$$\det A = (-1)^n 2^{n-1} n. \quad (3)$$

- Tested on non-regular triangulation given by de Loera [2].
- Interpretation
- Tighter bound on simplices unique under symmetry.

Proposition

To prove the conjecture, it suffices to show that if $A^{-1} = (b_{ij})$,

$$-\left(\sum_{i,j=1,\dots,n+1} b_{ij}\right) = 2/n. \quad (4)$$

Proof: Uses Cayley-Menger determinant and a block matrix identity from J.R. Sylvester [3].

Acknowledgements

Thanks to

- McCabe Olsen
- Wayne Tarrant
- Fellow students at RHIT's REU

- [1] Huggins et al., *The hyperdeterminant and triangulations of the 4-cube*, Mathematics of Computation **77(263)** (2008), 1653–1679.
- [2] J.A. de Loera, *Nonregular triangulations of products of simplices*, Discrete Comput Geom **15** (1996), 262–263.
- [3] J.R. Sylvester, *Determinants of block matrices*, Math Gaz **84(501)** (2000), 460–467.