

# class09\_\_david\_ma

David Ma

## Section 1: Intro to RPDB

Q1: What percentage of structures in the PDB are solved by X-ray and electron microscopy?

```
db <- read.csv("Data_Export_Summary.csv", row.names = 1)

numerics <- function(x){

  #Takes the sum after removing commas and converting strings into integers
  sum(as.numeric(gsub(",", "", x)))
}
```

For Xray:

```
numerics(db$X.ray) / numerics(db$Total)
```

```
[1] 0.8553721
```

For EM:

```
numerics(db$EM) / numerics(db$Total)
```

```
[1] 0.07455763
```

Q2: What proportion of structures in the PDB are protein?

```
numerics(db[1, "Total"]) / numerics(db$Total)
```

```
[1] 0.8681246
```

## Section 2: Visualizing HIV-1

Q3: Type HIV in the PDB website search box on the home page and determine how many HIV-1 protease structures are in the current PDB?



Figure 1: HIV-PR structure from MERK with a bound drug

Q4: Water molecules normally have 3 atoms. Why do we see just one atom per water molecule in this structure?

The structure is too low a resolution to see H atoms. You need a sub 1 Angstrom resolution to see Hydrogen.

Q5: There is a critical “conserved” water molecule in the binding site. Can you identify this water molecule? What residue number does this water molecule have?

HOH 308

## Section 3: Working with Structures in R

We can use the `bio3d` package to read and perform bioinformatics calculations on PDB structures.

```
library(bio3d)
```

Warning: package 'bio3d' was built under R version 4.2.3

```
pdb <- read.pdb("1hsg")
```

Note: Accessing on-line PDB file

```
pdb
```

```
Call: read.pdb(file = "1hsg")
```

```
Total Models#: 1
```

```
Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)
```

```
Protein Atoms#: 1514 (residues/Calpha atoms#: 198)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 172 (residues: 128)
```

```
Non-protein/nucleic resid values: [ HOH (127), MK1 (1) ]
```

```
Protein sequence:
```

```
PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD  
QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE  
ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP  
VNIIGRNLLTQIGCTLNF
```

```
+ attr: atom, xyz, seqres, helix, sheet,  
      calpha, remark, call
```

Q7: How many amino acid residues are in this PDB object?

198

Q8: Name one of the two-non protein residues?

MK1

Q9: How many protein chains are in this structure?

2

```
attributes(pdb)
```

```
$names
```

```
[1] "atom" "xyz" "seqres" "helix" "sheet" "calpha" "remark" "call"
```

```
$class
```

```
[1] "pdb" "sse"
```

```
head(pdb$atom)
```

	type	eleno	elety	alt	resid	chain	resno	insert	x	y	z	o	b
1	ATOM	1	N	<NA>	PRO	A	1	<NA>	29.361	39.686	5.862	1	38.10
2	ATOM	2	CA	<NA>	PRO	A	1	<NA>	30.307	38.663	5.319	1	40.62
3	ATOM	3	C	<NA>	PRO	A	1	<NA>	29.760	38.071	4.022	1	42.64
4	ATOM	4	O	<NA>	PRO	A	1	<NA>	28.600	38.302	3.676	1	43.40
5	ATOM	5	CB	<NA>	PRO	A	1	<NA>	30.508	37.541	6.342	1	37.87
6	ATOM	6	CG	<NA>	PRO	A	1	<NA>	29.296	37.591	7.162	1	38.40

	segid	elesy	charge
1	<NA>	N	<NA>
2	<NA>	C	<NA>
3	<NA>	C	<NA>
4	<NA>	O	<NA>
5	<NA>	C	<NA>
6	<NA>	C	<NA>

Read an ADK structure

```
adk <- read.pdb("6s36")
```

Note: Accessing on-line PDB file

PDB has ALT records, taking A only, rm.alt=TRUE

```
adk
```

```
Call: read.pdb(file = "6s36")
```

```
Total Models#: 1
```

```
Total Atoms#: 1898, XYZs#: 5694 Chains#: 1 (values: A)
```

```
Protein Atoms#: 1654 (residues/Calpha atoms#: 214)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 244 (residues: 244)
```

```
Non-protein/nucleic resid values: [ CL (3), HOH (238), MG (2), NA (1) ]
```

```
Protein sequence:
```

```
MRIILLGAPGAGKGTQAQFIMEKYGIPQISTGDMRLRAAVKSGSELGKQAKDIMDAGKLV  
DELVIALVKERIAQEDCRNGFLLDGFPRTPQADAMKEAGINVDYVLEFDVPDELIVDKI  
VGRRVHAPSGRVYHVKFNPVKVEGKDDVTGEELTTRKDDQEETVRKRLVEYHQM  
TAPLIG  
YYSKEAEAGNTKYAKVDGTPVAEVRADLEKILG
```

```
+ attr: atom, xyz, seqres, helix, sheet,  
      calpha, remark, call
```


Perform a prediction of flexibility with a technique called NMA (normal mode analysis)

```
# Perform flexibility prediction  
m <- nma(adk)
```

```
Building Hessian... Done in 0.03 seconds.
```

```
Diagonalizing Hessian... Done in 0.32 seconds.
```

```
plot(m)
```



Write out a “movie” (a.k.a trajectory) of the motion for viewing in M0lstar.

```
mktrj(m, file="adk_m7.pdb")
```