

City General Hospital

30-Day Readmission Risk Assessment

Patient ID:	20251102-Typ-b15f68	Generated:	November 02, 2025 11:53
Patient Name:	JOSHUA CO	Age:	50 years
Sex:	Male	Disease:	Type 2 Diabetes

30-Day Readmission Risk: **9.0%** | Classification: **LOW RISK** (Threshold: 45%)

Index Admission Summary

Length of Stay:	N/A	Discharge Destination:	N/A
Prior Admissions (90d):	N/A	Comorbidities Count:	N/A
Follow-up Scheduled:	N/A	Admission Type:	Emergency

Top Contributing Risk Factors (SHAP Analysis)

Feature	Value	Contribution	Impact	Interpretation
HbA1c	0.00	-3.301	↓ Lower Risk	HbA1c influences readmission risk. (0.00% — below ref 4.0-6.5%). This feature has a major effect and reduces the readmission risk. Optimize long-te...
comorbidities_count	0.00	-1.039	↓ Lower Risk	comorbidities_count influences readmission risk.. This feature has a major effect and reduces the readmission risk.
Fasting_glucose	0.00	-0.635	↓ Lower Risk	Fasting_glucose influences readmission risk.. This feature has a moderate effect and reduces the readmission risk.
age	50.00	-0.255	↓ Lower Risk	age influences readmission risk.. This feature has a minor effect and reduces the readmission risk.
BMI	0.00	+0.232	↑ Higher Risk	BMI influences readmission risk. (0.00kg/m² — below ref 18.5-25kg/m²). This feature has a minor effect and increases the readmission risk. Reinforc...
Creatinine	0.00	-0.149	↓ Lower Risk	Creatinine influences readmission risk. (0.00mg/dL — below ref 0.6-1.3mg/dL). This feature has a minimal effect and reduces the readmission risk. A...
prior_admissions_90d	0.00	-0.073	↓ Lower Risk	prior_admissions_90d influences readmission risk.. This feature has a minimal effect and reduces the readmission risk.
sex	nan	-0.051	↓ Lower Risk	sex influences readmission risk.. This feature has a minimal effect and reduces the readmission risk.

Clinical Summary

This 50 years male patient presents with a **low risk 30-day readmission risk (9.0%)** following discharge for Type 2 Diabetes management. The risk assessment threshold for this condition is 45%, placing this patient below the high-risk threshold.

Primary Risk Drivers: HbA1c, comorbidities_count, Fasting_glucose are identified as major contributing factors.

Recommendation: Close follow-up and aggressive management of identified risk factors is recommended to prevent readmission.

Clinical Management Recommendations

- **Primary Disease Management:** Review and optimize current treatment plan for Type 2 Diabetes. Consider consultation with appropriate specialists.
- **Medication Reconciliation:** Complete medication review at discharge. Ensure patient understands all medications, dosing, and timing.
- **Care Coordination:** Schedule follow-up appointment within 7 days of discharge. Consider home health services for high-risk patients.
- **Patient Education:** Reinforce medication adherence, warning signs requiring immediate attention, and lifestyle modifications.
- **Laboratory Monitoring:** Schedule appropriate lab work based on disease-specific guidelines and medication monitoring requirements.

City General Hospital

Medication Recommendations

Patient ID:	20251102-Typ-b15f68	Patient Name:	JOSHUA CO
Age / Sex:	50 years / Male	Disease:	Type 2 Diabetes

Important: The following medication recommendations are based on current clinical guidelines and the patient's risk profile. All medications must be reviewed, prescribed, and adjusted by the attending physician based on individual patient factors, allergies, drug interactions, and institutional protocols.

Recommended Medication Protocol:

First-Line Therapy:

- Metformin 500-1000mg twice daily (adjust for renal function)
- SGLT2 inhibitors (e.g., Empagliflozin 10-25mg daily) if GFR >45
- GLP-1 agonists (e.g., Semaglutide) for cardiovascular benefit

Glucose Control:

- Insulin therapy if HbA1c >9% or symptomatic hyperglycemia
- DPP-4 inhibitors (e.g., Sitagliptin 100mg daily) as add-on
- Basal insulin (e.g., Glargine) starting 10 units at bedtime

Complications:

- ACE inhibitors/ARBs for nephroprotection (e.g., Lisinopril 10-40mg)
- Statin therapy (e.g., Atorvastatin 20-40mg) for cardiovascular risk
- Aspirin 81mg daily if cardiovascular disease present

Monitoring:

- Check HbA1c every 3 months, annual kidney function, regular foot exams

Note: These are general guidelines. All medications should be prescribed and adjusted by the attending physician based on individual patient factors.

City General Hospital

Potential Disease Progression & Related Conditions

Patient ID:	20251102-Typ-b15f68	Patient Name:	JOSHUA CO
Age / Sex:	50 years / Male	Disease:	Type 2 Diabetes

Potential Disease Progression & Related Conditions:

[HIGH RISK] High-Risk Conditions (Requires Active Prevention):

1. Diabetic Nephropathy

- *Risk Factors:* elevated creatinine, proteinuria, poor HbA1c control
- *Typical Time Frame:* 5-10 years
- *Prevention Strategy:* Strict BP control, ACE-I/ARB therapy, HbA1c <7%

2. Diabetic Retinopathy

- *Risk Factors:* HbA1c >8%, hypertension, diabetes duration >10 years
- *Typical Time Frame:* 10-15 years
- *Prevention Strategy:* Annual eye exams, optimal glucose control

3. Cardiovascular Disease

- *Risk Factors:* high LDL, obesity, smoking, hypertension
- *Typical Time Frame:* 10-20 years
- *Prevention Strategy:* Statin therapy, aspirin, lifestyle modification

4. Peripheral Neuropathy

- *Risk Factors:* poor glycemic control, long diabetes duration
- *Typical Time Frame:* 5-10 years
- *Prevention Strategy:* Glucose control, B12 supplementation, foot care

[MODERATE RISK] Moderate-Risk Conditions (Monitor Closely):

1. Diabetic Foot Ulcers

- *Risk Factors:* neuropathy, poor circulation, foot deformity
- *Prevention:* Daily foot inspections, proper footwear

2. Gastroparesis

- *Risk Factors:* autonomic neuropathy, poor glucose control
- *Prevention:* Controlled glucose, smaller frequent meals

Personalized Risk Assessment:

- Current markers within acceptable ranges. Continue monitoring.

Disclaimer: This report is generated by a machine learning model for clinical decision support. All recommendations should be reviewed and approved by qualified healthcare professionals. Medication dosages and treatment plans must be individualized based on patient-specific factors, comorbidities, and current clinical guidelines.