


# COMBINATORIAL TOOLS IN EQUIVARIANT ALGEBRAIC TOPOLOGY

slides available at [www.davidmehrlle.com/csu.pdf](http://www.davidmehrlle.com/csu.pdf)

PART I  
THE CONJECTURES

## KERVAIRE PROBLEM (1960)

Can every smooth  $n$ -manifold become a sphere via surgery?



## SURGERY

Replace a **submanifold** by **another** with the same boundary

## KERVAIRE PROBLEM (1960)

Can every smooth  $n$ -manifold become a sphere via surgery?

(1963–1984) counterexamples with  $n = 2, 6, 14, 30, 62$

(1969) Browder: counterexamples must have  $n = 2^k - 2$

## KERVAIRE PROBLEM (1960)

Can every smooth  $n$ -manifold become a sphere via surgery?

(1963–1984) counterexamples with  $n = 2, 6, 14, 30, 62$

(1969) Browder: counterexamples must have  $n = 2^k - 2$

THEOREM (Hill–Hopkins–Ravenel, 2016)

The only counterexamples have  $n = 2, 6, 14, 30, 62$ , or maybe 126

Key ingredient: **equivariant homotopy theory**

## KERVAIRE PROBLEM (1960)

Can every smooth  $n$ -manifold become a sphere via surgery?

(1963–1984) counterexamples with  $n = 2, 6, 14, 30, 62$

(1969) Browder: counterexamples must have  $n = 2^k - 2$

THEOREM (Hill–Hopkins–Ravenel, 2016)

The only counterexamples have  $n = 2, 6, 14, 30, 62$ , or maybe 126

Key ingredient: **equivariant homotopy theory**

THEOREM (Lin–Wang–Xu, December 2024)

There exist counterexample 126-manifolds (nonconstructive)

TELESCOPE CONJECTURE (Ravenel, 1984)

Two ways of computing  $\pi_k S^n$  are the same

THEOREM (Burklund–Hahn–Levy–Schlank, 2023)

The Telescope Conjecture is false!

A key ingredient: equivariant homotopy theory

TELESCOPE CONJECTURE (Ravenel, 1984)

Two ways of computing  $\pi_k S^n$  are the same

THEOREM (Burklund–Hahn–Levy–Schlank, 2023)

The Telescope Conjecture is false!

A key ingredient: **equivariant homotopy theory**

Will equivariant homotopy theory solve all our problems?



## TELESCOPE CONJECTURE (Ravenel, 1984)

Two ways of computing  $\pi_k S^n$  are the same

THEOREM (Burklund–Hahn–Levy–Schlank, 2023)

The Telescope Conjecture is false!

A key ingredient: **equivariant homotopy theory**

Will equivariant homotopy theory solve all our problems?

Not yet.

- Group actions are rigid; homotopy theory is floppy
- Computations are tough, even for small finite groups
- Needs new algebraic tools: Mackey/Tambara functors

## TELESCOPE CONJECTURE (Ravenel, 1984)

Two ways of computing  $\pi_k S^n$  are the same

THEOREM (Burklund–Hahn–Levy–Schlank, 2023)

The Telescope Conjecture is false!

A key ingredient: **equivariant homotopy theory**

Will equivariant homotopy theory solve all our problems?

Not yet.

- Group actions are rigid; homotopy theory is floppy
- Computations are tough, even for small finite groups
- Needs new algebraic tools: Mackey/Tambara functors

The better we understand these tools, the more we can do!

PART II  
EQUIVARIANT  
ALGEBRAIC TOPOLOGY

Let  $G$  be a finite group.

DEFINITION

A  **$G$ -space** is a topological space  $X$  with a  $G$ -action  $G \times X \rightarrow X$ :

$$1 \cdot x = x$$

$$g \cdot (h \cdot x) = (gh) \cdot x$$

such that  $x \mapsto g \cdot x$  is continuous for all  $g \in G$ .

Let  $G$  be a finite group.

#### DEFINITION

A  **$G$ -space** is a topological space  $X$  with a  $G$ -action  $G \times X \rightarrow X$ :

$$1 \cdot x = x$$

$$g \cdot (h \cdot x) = (gh) \cdot x$$

such that  $x \mapsto g \cdot x$  is continuous for all  $g \in G$ .


#### DEFINITION

A  **$G$ -equivariant map**  $f: X \rightarrow Y$  is a continuous map such that


$$f(g \cdot x) = g \cdot f(x)$$

for all  $g \in G$  and all  $x \in X$ .


# EXAMPLES



Rotation  
 $\mathbb{Z}/n$



Covering Spaces  
 $\mathbb{Z}/3$



$\text{PbCr}_3\text{S}_4$  crystals  
Dihedral Group  $D_6$

In algebraic topology, we study spaces using **invariants**

# DEFINITION

An **invariant** is data  $F(X)$  built from a space  $X$  such that

$$X \simeq Y \implies F(X) \cong F(Y)$$

In algebraic topology, we study spaces using **invariants**

# DEFINITION

An **invariant** is data  $F(X)$  built from a space  $X$  such that

$$X \simeq Y \implies F(X) \cong F(Y)$$

We mostly use the contrapositive:  $F(X) \not\cong F(Y) \implies X \not\simeq Y$



In algebraic topology, we study spaces using **invariants**

## DEFINITION


An **invariant** is data  $F(X)$  built from a space  $X$  such that

$$X \simeq Y \implies F(X) \cong F(Y)$$

We mostly use the contrapositive:  $F(X) \not\cong F(Y) \implies X \not\simeq Y$

## EXAMPLE

The **fundamental group**  $\pi_1(X)$  is the group of homotopy classes of based loops in  $X$ .



$$\pi_1(S^1) \cong \mathbb{Z}$$

In algebraic topology, we study spaces using **invariants**

## DEFINITION


An **invariant** is data  $F(X)$  built from a space  $X$  such that

$$X \simeq Y \implies F(X) \cong F(Y)$$


We mostly use the contrapositive:  $F(X) \not\cong F(Y) \implies X \not\simeq Y$

## EXAMPLE

The **fundamental group**  $\pi_1(X)$  is the group of homotopy classes of based loops in  $X$ .



$$\pi_1(S^1) \cong \mathbb{Z}$$



$$\pi_1(T) \cong \mathbb{Z} \times \mathbb{Z}$$

In algebraic topology, we study spaces using **invariants**

## DEFINITION


An **invariant** is data  $F(X)$  built from a space  $X$  such that

$$X \simeq Y \implies F(X) \cong F(Y)$$


We mostly use the contrapositive:  $F(X) \not\cong F(Y) \implies X \not\simeq Y$

## EXAMPLE


The **fundamental group**  $\pi_1(X)$  is the group of homotopy classes of based loops in  $X$ .



$$\pi_1(S^1) \cong \mathbb{Z}$$




$$\pi_1(T) \cong \mathbb{Z} \times \mathbb{Z}$$




$$\pi_1(S^2) = 0$$

## QUESTION

How do you see group actions on spaces using invariants?




$\mathbb{Z}/2$  acts by reflection




$\mathbb{Z}/2$  acts trivially

## QUESTION

How do you see group actions on spaces using invariants?



$\mathbb{Z}/2$  acts by reflection  
fixed points  $S^0$



$\mathbb{Z}/2$  acts trivially  
fixed points  $S^1$

## ANSWER

Consider the fixed point subspaces  $X^G$  as well!



$$X^G := \left\{ x \in X \mid g \cdot x = x \text{ for all } g \in G \right\} \subseteq X$$

# THE DATA PIPELINE

$G \curvearrowright X$

G-space

# THE DATA PIPELINE






THEOREM (Elmendorf, Piacenza, 1991)

For the purposes of homotopy theory, we may replace a  $G$ -space  $X$  by its collection of fixed point subspaces  $\{X^H\}_{H \subseteq G}$

# THE DATA PIPELINE

$G \curvearrowright X$





THEOREM (Elmendorf, Piacenza, 1991)

For the purposes of homotopy theory, we may replace a  $G$ -space  $X$  by its collection of fixed point subspaces  $\{X^H\}_{H \subseteq G}$



# THE DATA PIPELINE




THEOREM (Elmendorf, Piacenza, 1991)


For the purposes of homotopy theory, we may replace a  $G$ -space  $X$  by its collection of fixed point subspaces  $\{X^H\}_{H \subseteq G}$

# EXAMPLE

reflection  
action




trivial  
action



# EXAMPLE


reflection  
action



$$(S^1)^{\mathbb{Z}/2}$$

$$\downarrow$$
  
 $S^1$

trivial  
action




$$(S^1)^{\mathbb{Z}/2}$$

$$\downarrow$$
  
 $S^1$

# EXAMPLE

reflection  
action




$S^0$




$S^1$

trivial  
action




$S^1$



$S^1$

# EXAMPLE

reflection  
action




$S^0$



$S^1$

$\pi_1^{\text{st}}$




$\mathbb{Z}/2$



$\mathbb{Z}$

trivial  
action




$S^1$



$S^1$

$\pi_1^{\text{st}}$



$\mathbb{Z}$



$\mathbb{Z}$

DEFINITION (Dress, 1971)

A Mackey Functor for  $G$  is the data:

DEFINITION (Dress, 1971)

A **Mackey Functor for  $G$**  is the data:

- an abelian group  $M(H)$  for each subgroup  $H \subseteq G$

$$M(G)$$

$$M(H)$$

$$M(K)$$

$$M(gKg^{-1})$$


$$M(1)$$

## DEFINITION (Dress, 1971)

A **Mackey Functor** for  $G$  is the data:

- an abelian group  $M(H)$  for each subgroup  $H \subseteq G$
- **restriction homomorphisms**

$$\text{res}_K^H : M(H) \rightarrow M(K)$$





## DEFINITION (Dress, 1971)


A **Mackey Functor** for  $G$  is the data:

- an abelian group  $M(H)$  for each subgroup  $H \subseteq G$
- **restriction homomorphisms**

$$\text{res}_K^H: M(H) \rightarrow M(K)$$

- **transfer homomorphisms**

$$\text{tr}_K^H: M(K) \rightarrow M(H)$$



# DEFINITION (Dress, 1971)

A **Mackey Functor** for  $G$  is the data:

- an abelian group  $M(H)$  for each subgroup  $H \subseteq G$

- **restriction homomorphisms**

$$\text{res}_K^H: M(H) \rightarrow M(K)$$


- **transfer homomorphisms**

$$\text{tr}_K^H: M(K) \rightarrow M(H)$$

- **conjugation isomorphisms**

$$c_g: M(K) \cong M(gKg^{-1})$$

with a “double coset formula” for  $\text{res} \circ \text{tr}$ , and other conditions



# SMALL GROUPS

A Mackey functor for  $G = \mathbb{1}$  is just an abelian group

# SMALL GROUPS

A Mackey functor for  $G = \mathbb{1}$  is just an abelian group

A Mackey functor  $M$  for  $G = C_p$  (cyclic, order  $p$ ) is the data:

$$\begin{array}{c} M(C_p) \\ \begin{array}{c} \text{tr}_{\mathbb{1}}^{C_p} \curvearrowright \\ \text{res}_{\mathbb{1}}^{C_p} \downarrow \end{array} \\ M(\mathbb{1}) \\ \begin{array}{c} \hookrightarrow \\ C_p \end{array} \end{array}$$

# SMALL GROUPS

A Mackey functor for  $G = \mathbb{1}$  is just an abelian group

A Mackey functor  $M$  for  $G = C_p$  (cyclic, order  $p$ ) is the data:

$$\begin{array}{c} M(C_p) \\ \begin{array}{c} \text{blue curved arrow } \text{tr}_{\mathbb{1}}^{C_p} \uparrow \\ \text{orange straight arrow } \text{res}_{\mathbb{1}}^{C_p} \downarrow \end{array} \\ M(\mathbb{1}) \\ \begin{array}{c} \text{curved arrow} \\ C_p \end{array} \end{array}$$

$$\text{res}_{\mathbb{1}}^{C_p} \circ \text{tr}_{\mathbb{1}}^{C_p}(m) = \sum_{g \in C_p} g \cdot m$$

## QUESTION

Where do the transfers come from?

## QUESTION

Where do the transfers come from?

Let  $E/F$  be a Galois extension with (finite) Galois group  $\text{Gal}(E/F)$

There is a  $\text{Gal}(E/F)$ -Mackey functor  $M$  with  $M(H) = (E^H, +)$

## QUESTION

Where do the transfers come from?

Let  $E/F$  be a Galois extension with (finite) Galois group  $\text{Gal}(E/F)$

There is a  $\text{Gal}(E/F)$ -Mackey functor  $M$  with  $M(H) = (E^H, +)$

**Restrictions** are inclusions:

$$\text{res}_K^H: E^H \hookrightarrow E^K$$



## QUESTION

Where do the transfers come from?

Let  $E/F$  be a Galois extension with (finite) Galois group  $\text{Gal}(E/F)$

There is a  $\text{Gal}(E/F)$ -Mackey functor  $M$  with  $M(H) = (E^H, +)$

**Restrictions** are inclusions:

$$\text{res}_K^H: E^H \hookrightarrow E^K$$

**Transfers** are sums over orbits (field-theoretic traces):

$$\begin{aligned} \text{tr}_K^H: E^K &\longrightarrow E^H \\ a &\longmapsto \sum_{gK \in H/K} g \cdot a \end{aligned}$$

## QUESTION

Where do the transfers come from?

Let  $E/F$  be a Galois extension with (finite) Galois group  $\text{Gal}(E/F)$

There is a  $\text{Gal}(E/F)$ -Mackey functor  $M$  with  $M(H) = (E^H, +)$

**Restrictions** are inclusions:

$$\text{res}_K^H: E^H \hookrightarrow E^K$$

**Transfers** are sums over orbits (field-theoretic traces):

$$\begin{aligned} \text{tr}_K^H: E^K &\longrightarrow E^H \\ a &\longmapsto \sum_{gK \in H/K} g \cdot a \end{aligned}$$

## SLOGAN

Mackey Functors  $\pi_1^{\text{st}}(X^H)$  encode a “Galois theory” of  $G$ -spaces

PART III

EQUIVARIANT ALGEBRA

The category of  $G$ -Mackey functors is:

- abelian
- symmetric monoidal
- generated by finitely many projectives

The category of  $G$ -Mackey functors is:

- abelian
- symmetric monoidal
- generated by finitely many projectives

$\implies$  algebra!

The category of  $G$ -Mackey functors is:

- abelian  $\implies$  algebra!
- symmetric monoidal  $\implies$  commutative algebra!
- generated by finitely many projectives

The category of  $G$ -Mackey functors is:

- abelian  $\implies$  algebra!
- symmetric monoidal  $\implies$  commutative algebra!
- generated by finitely many projectives  $\implies$  homological algebra!

The category of  $G$ -Mackey functors is:

- abelian  $\implies$  algebra!
- symmetric monoidal  $\implies$  commutative algebra!
- generated by finitely many projectives  $\implies$  homological algebra!

#### EXAMPLE

When  $G = \mathbb{1}$ , this is just abelian groups and commutative rings



## QUESTION

What is a commutative ring for  $G$ -Mackey functors?

## QUESTION

What is a commutative ring for  $G$ -Mackey functors?

Let  $E/F$  be a Galois extension with (finite) Galois group  $\text{Gal}(E/F)$

There is a  $\text{Gal}(E/F)$ -Mackey functor  $M$  with  $M(H) = E^H$

## QUESTION

What is a commutative ring for  $G$ -Mackey functors?

Let  $E/F$  be a Galois extension with (finite) Galois group  $\text{Gal}(E/F)$

There is a  $\text{Gal}(E/F)$ -Mackey functor  $M$  with  $M(H) = E^H$

**Restrictions** are inclusions:

$$\text{res}_K^H: E^H \hookrightarrow E^K$$

**Transfers** are sums over orbits (field-theoretic traces):

$$\text{tr}_K^H: E^K \longrightarrow E^H$$

$$a \longmapsto \sum_{\sigma \in \text{Gal}(E^K/E^H)} \sigma(a)$$

## QUESTION

What is a commutative ring for  $G$ -Mackey functors?

Let  $E/F$  be a Galois extension with (finite) Galois group  $\text{Gal}(E/F)$

There is a  $\text{Gal}(E/F)$ -Mackey functor  $M$  with  $M(H) = E^H$

**Restrictions** are inclusions:

$$\text{res}_K^H: E^H \hookrightarrow E^K$$

**Transfers** are sums over orbits (field-theoretic traces):

$$\text{tr}_K^H: E^K \longrightarrow E^H$$

$$a \longmapsto \sum_{\sigma \in \text{Gal}(E^K/E^H)} \sigma(a)$$

Also have field-theoretic **norms**:

$$a \longmapsto \prod_{\sigma \in \text{Gal}(E^K/E^H)} \sigma(a)$$

DEFINITION (Tambara, 1993)

A **Tambara Functor for  $G$**  is the data:

DEFINITION (Tambara, 1993)

A **Tambara Functor for  $G$**  is the data:

- a commutative ring  $T(H)$  for each subgroup  $H \subseteq G$

$$T(G)$$

$$T(H)$$

$$T(K)$$


$$T(\mathbb{1})$$

# DEFINITION (Tambara, 1993)

A **Tambara Functor** for  $G$  is the data:

- a commutative ring  $T(H)$  for each subgroup  $H \subseteq G$
- restriction (ring) homomorphisms

$$\text{res}_K^H: T(H) \rightarrow T(K)$$



# DEFINITION (Tambara, 1993)


A **Tambara Functor** for  $G$  is the data:

- a commutative ring  $T(H)$  for each subgroup  $H \subseteq G$
- **restriction (ring) homomorphisms**

$$\text{res}_K^H: T(H) \rightarrow T(K)$$

- **transfer homomorphisms** (for  $+$ )

$$\text{tr}_K^H: T(K) \rightarrow T(H)$$





## DEFINITION (Tambara, 1993)

A **Tambara Functor** for  $G$  is the data:

- a commutative ring  $T(H)$  for each subgroup  $H \subseteq G$
- **restriction (ring) homomorphisms**


$$\text{res}_K^H: T(H) \rightarrow T(K)$$

- **transfer homomorphisms** (for  $+$ )

$$\text{tr}_K^H: T(K) \rightarrow T(H)$$

- **norm homomorphisms** (for  $\times$ )

$$\text{nm}_K^H: T(K) \rightarrow T(H)$$



# DEFINITION (Tambara, 1993)

A **Tambara Functor** for  $G$  is the data:

- a commutative ring  $T(H)$  for each subgroup  $H \subseteq G$
- **restriction (ring) homomorphisms**

$$\text{res}_K^H: T(H) \rightarrow T(K)$$

- **transfer homomorphisms** (for  $+$ )


$$\text{tr}_K^H: T(K) \rightarrow T(H)$$

- **norm homomorphisms** (for  $\times$ )

$$\text{nm}_K^H: T(K) \rightarrow T(H)$$

- conjugation isomorphisms

with “double coset formulas” for  $\text{res} \circ \text{tr}$  and  $\text{res} \circ \text{nm}$ , and ...




# TAMBARA FUNCTORS FOR SMALL GROUPS

A Tambara functor for  $G = \mathbb{1}$  is a commutative ring

# TAMBARA FUNCTORS FOR SMALL GROUPS

A Tambara functor for  $G = \mathbb{1}$  is a commutative ring


A Tambara functor  $T$  for  $G = C_p$  is the data:



# TAMBARA FUNCTORS FOR SMALL GROUPS

A Tambara functor for  $G = \mathbb{1}$  is a commutative ring

A Tambara functor  $T$  for  $G = C_p$  is the data:



$$\text{nm}(0) = 0$$

$$\text{res} \circ \text{tr}(x) = \sum_{g \in C_p} g \cdot x$$

$$\text{res} \circ \text{nm}(x) = \prod_{g \in C_p} g \cdot x$$

(and other conditions)


## EXAMPLE: BURNSIDE FUNCTOR $\mathbb{A}$

For each  $G$ , there is a **Burnside functor**  $\mathbb{A}$  with origin in topology. Each  $\mathbb{A}(H)$  is built from finite  $G$ -sets.

# EXAMPLE: BURNSIDE FUNCTOR $\mathbb{A}$

For each  $G$ , there is a **Burnside functor**  $\mathbb{A}$  with origin in topology. Each  $\mathbb{A}(H)$  is built from finite  $G$ -sets.

$\mathbb{A}$  for  $C_p$ :



## EXAMPLE: BURNSIDE FUNCTOR $\mathbb{A}$

For each  $G$ , there is a **Burnside functor**  $\mathbb{A}$  with origin in topology. Each  $\mathbb{A}(H)$  is built from finite  $G$ -sets.

$\mathbb{A}$  for  $C_p$ :

$$\begin{array}{ccc} & \mathbb{Z}[t]/(t^2 - pt) & \\ \text{blue } \curvearrowleft x \mapsto tx & \downarrow \text{orange } t \mapsto p & \text{green } \curvearrowright x \mapsto x + \left(\frac{x^p - x}{p}\right)t \\ & \mathbb{Z} & \\ & \downarrow \text{trivial} & \end{array}$$

This Tambara functor  $\mathbb{A}$  plays the role of  $\mathbb{Z}$ .

- Every Mackey functor is an  $\mathbb{A}$ -module
- $\mathbb{A}$  is the initial Tambara functor
- $\mathbb{A}$  is the unit for the tensor product



# REPRESENTATION THEORY

## DEFINITION

Let  $X$  be a  $G$ -space. The space  $X^n$  has actions of both  $G^n$  and  $S_n$ , which combine to an action of the **wreath product**  $G \wr S_n$ .

# REPRESENTATION THEORY

## DEFINITION

Let  $X$  be a  $G$ -space. The space  $X^n$  has actions of both  $G^n$  and  $S_n$ , which combine to an action of the **wreath product**  $G \wr S_n$ .

THEOREM (Cornelius–Dominguez–Modi–Mehrle–Rose–Stapleton, 2024)

Connect  $\mathbb{A}(G \wr S_n)$  to the representation theory of  $S_n$ ; use this to find computationally effective formulas for **nm** in  $\mathbb{A}$ .

# REPRESENTATION THEORY

## DEFINITION

Let  $X$  be a  $G$ -space. The space  $X^n$  has actions of both  $G^n$  and  $S_n$ , which combine to an action of the **wreath product**  $G \wr S_n$ .

THEOREM (Cornelius–Dominguez–Modi–Mehrle–Rose–Stapleton, 2024)

Connect  $\mathbb{A}(G \wr S_n)$  to the representation theory of  $S_n$ ; use this to find computationally effective formulas for **nm** in  $\mathbb{A}$ .

WORK-IN-PROGRESS (Calle–Chan–Mehrle–Quigley–Spitz–Van Niel)

Every Tambara functor  $T$  has a character theory: a Tambara functor  $\Gamma(T)$  with  $T \hookrightarrow \Gamma(T)$ , where  $\Gamma(T)$  is easier to study.

# COMMUTATIVE ALGEBRA

THEOREM (Nakaoka, 2014)

There is a robust theory of prime ideals for Tambara functors.

# COMMUTATIVE ALGEBRA

THEOREM (Nakaoka, 2014)

There is a robust theory of prime ideals for Tambara functors.

THEOREM (Chan–Mehrlé–Quigley–Spitz–Van Niel, 2024)

Use  $\Gamma(\mathbb{A})$  for  $G = C_p$  to describe the Tambara affine line, and relate affine Tambara algebraic geometry to invariant theory.


LONG-TERM GOAL

Algebraic geometry and invariant theory of Tambara functors

PART IV  
HOMOTOPICAL  
COMBINATORICS


# A NON-EXAMPLE

$\mathbb{Z}/2$  acts on  $S^2$  by  
reflection in equator



# A NON-EXAMPLE


$\mathbb{Z}/2$  acts on  $S^2$  by  
reflection in equator






# A NON-EXAMPLE

$\mathbb{Z}/2$  acts on  $S^2$  by  
reflection in equator




# A NON-EXAMPLE

$\mathbb{Z}/2$  acts on  $S^2$  by  
reflection in equator




# A NON-EXAMPLE


$\mathbb{Z}/2$  acts on  $S^2$  by  
reflection in equator



$S^1$




$\mathbb{Z}$




# A NON-EXAMPLE

$\mathbb{Z}/2$  acts on  $S^2$  by  
reflection in equator



$S^1$




$\mathbb{Z}$






$0 = 1$

in the ring  $0$



# A NON-EXAMPLE

$\mathbb{Z}/2$  acts on  $S^2$  by  
reflection in equator






$$\implies \begin{matrix} 0 = 1 \\ nm(0) = nm(1) \end{matrix}$$

in the ring 0

# A NON-EXAMPLE

$\mathbb{Z}/2$  acts on  $S^2$  by  
reflection in equator





$$\begin{aligned} & 0 = 1 \quad \text{in the ring } 0 \\ \implies & \text{nm}(0) = \text{nm}(1) \\ \implies & 0 = 1 \quad \text{in } \mathbb{Z} \end{aligned}$$

**Contradiction!** No map  $\text{nm}$  can exist

# A NON-EXAMPLE

$\mathbb{Z}/2$  acts on  $S^2$  by reflection in equator



$$\begin{aligned} & 0 = 1 && \text{in the ring } 0 \\ \implies & \text{nm}(0) = \text{nm}(1) \\ \implies & 0 = 1 && \text{in } \mathbb{Z} \end{aligned}$$

**Contradiction!** No map  $\text{nm}$  can exist

Want to allow Tambara functors with a subset of the norms

## QUESTION

Which combinations of **norms** are allowable?

## DEFINITION (Rubin, 2020)

A **G-transfer system** is a partial order  $\rightarrow$  on subgroups of  $G$ :

- (refinement) if  $K \rightarrow H$ , then  $K \subseteq H$
- (conjugation) if  $K \rightarrow H$ , then  $gKg^{-1} \rightarrow gHg^{-1}$  for all  $g \in G$
- (restriction) if  $K \rightarrow H$  and  $L \subseteq H$ , then  $K \cap L \rightarrow L$



## QUESTION

Which combinations of **norms** are allowable?

## DEFINITION (Rubin, 2020)

A  **$G$ -transfer system** is a partial order  $\rightarrow$  on subgroups of  $G$ :

- (refinement) if  $K \rightarrow H$ , then  $K \subseteq H$
- (conjugation) if  $K \rightarrow H$ , then  $gKg^{-1} \rightarrow gHg^{-1}$  for all  $g \in G$
- (restriction) if  $K \rightarrow H$  and  $L \subseteq H$ , then  $K \cap L \rightarrow L$

Transfer systems for  $G = C_p$ :

$C_p$

$\mathbb{1}$

$C_p$

$\uparrow$   
 $\mathbb{1}$

## QUESTION


Which combinations of **norms** are allowable?

## DEFINITION (Rubin, 2020)


A  **$G$ -transfer system** is a partial order  $\rightarrow$  on subgroups of  $G$ :

- (refinement) if  $K \rightarrow H$ , then  $K \subseteq H$
- (conjugation) if  $K \rightarrow H$ , then  $gKg^{-1} \rightarrow gHg^{-1}$  for all  $g \in G$
- (restriction) if  $K \rightarrow H$  and  $L \subseteq H$ , then  $K \cap L \rightarrow L$


Transfer systems for  $G = C_p$ :




Transfer systems for  $G = C_{p^2}$ :



Transfer systems for  $G = C_{p^2}$ :



Transfer systems for  $G = C_{p^2}$ :



THEOREM (Balchin–Barnes–Roitzheim, 2021)

There are  $\text{Cat}(n + 1) = \frac{1}{n+2} \binom{2n+2}{n+1}$  transfer systems for  $C_{p^n}$ .

# INCOMPLETE TAMBARA FUNCTORS



## DEFINITION

Let  $\tau$  be a transfer system for  $G$ . A  $\tau$ -Tambara functor is a Tambara functor with only those norms parameterized by  $\tau$ .

# INCOMPLETE TAMBARA FUNCTORS

## DEFINITION

Let  $\tau$  be a transfer system for  $G$ . A  $\tau$ -Tambara functor is a Tambara functor with only those **norms** parameterized by  $\tau$ .



# FREE ALGEBRAS

$\mathbb{Z}[x]$  is a free algebra with one generator



# FREE ALGEBRAS

$\mathbb{Z}[x]$  is a free algebra with one generator

The free  $\tau$ -Tambara functor  $\mathbb{A}[\tau, H]$  is determined by:

- a transfer system  $\tau$
- a subgroup  $H \subseteq G$


# FREE ALGEBRAS

$\mathbb{Z}[x]$  is a free algebra with one generator

The free  $\tau$ -Tambara functor  $\mathbb{A}[\tau, H]$  is determined by:

- a transfer system  $\tau$
- a subgroup  $H \subseteq G$

$\mathbb{A}[\tau, C_p] :$




# FREE ALGEBRAS

$\mathbb{Z}[x]$  is a free algebra with one generator

The free  $\tau$ -Tambara functor  $\mathbb{A}[\tau, H]$  is determined by:

- a transfer system  $\tau$
- a subgroup  $H \subseteq G$

$\mathbb{A}[\tau, C_p] :$




# FREE ALGEBRAS

$\mathbb{Z}[x]$  is a free algebra with one generator

The free  $\tau$ -Tambara functor  $\mathbb{A}[\tau, H]$  is determined by:

- a transfer system  $\tau$
- a subgroup  $H \subseteq G$

$\mathbb{A}[\tau, C_p] :$




# FREE ALGEBRAS

$\mathbb{Z}[x]$  is a free algebra with one generator

The free  $\tau$ -Tambara functor  $\mathbb{A}[\tau, H]$  is determined by:

- a transfer system  $\tau$
- a subgroup  $H \subseteq G$

$\mathbb{A}[\tau, C_p] :$




# FREE ALGEBRAS

$\mathbb{Z}[x]$  is a free algebra with one generator

The free  $\tau$ -Tambara functor  $\mathbb{A}[\tau, H]$  is determined by:

- a transfer system  $\tau$
- a subgroup  $H \subseteq G$

$\mathbb{A}[\tau, C_p] :$



$$\mathbb{A}(C_{p^2})[\text{tr}(x_i), \text{nm}(x_i)] / \sim$$

$$\begin{array}{ccc} \curvearrowright \text{tr} & \downarrow \text{res} & \curvearrowleft \text{nm} \\ \mathbb{A}(C_p)[x_1, \dots, x_{[G:H]}] \\ \curvearrowright \text{tr} & \downarrow \text{res} & \\ \mathbb{A}(1)[\text{res}(x_i)] \end{array}$$

$\mathbb{Z}[x]$  is a free  $\mathbb{Z}$ -algebra with one generator

$\mathbb{Z}[x]$  is a free  $\mathbb{Z}$ -module with basis  $\{1, x, x^2, x^3, \dots\}$

$\mathbb{Z}[x]$  is a free  $\mathbb{Z}$ -algebra with one generator

$\mathbb{Z}[x]$  is a free  $\mathbb{Z}$ -module with basis  $\{1, x, x^2, x^3, \dots\}$

## QUESTION

Is the free  $\tau$ -Tambara functor  $\mathbb{A}[\tau, H]$  free as a Mackey functor?



$\mathbb{Z}[x]$  is a free  $\mathbb{Z}$ -algebra with one generator

$\mathbb{Z}[x]$  is a free  $\mathbb{Z}$ -module with basis  $\{1, x, x^2, x^3, \dots\}$

## QUESTION

Is the free  $\tau$ -Tambara functor  $\mathbb{A}[\tau, H]$  free as a Mackey functor?

## THEOREM (Hill–Mehrlé–Quigley, 2023)

Let  $G$  be solvable. Then  $\mathbb{A}[\tau, H]$  is free as a Mackey functor iff:

- (a)  $H \rightarrow G$  in  $\tau$ ,
- (b)  $\tau$  has no arrows below  $H$ .


# THEOREM (Hill–Mehrle–Quigley, 2023)

Let  $G$  be solvable. Then  $\mathbb{A}[\tau, H]$  is free as a Mackey functor iff:


- (a)  $H \rightarrow G$  in  $\tau$ ,
- (b)  $\tau$  has no arrows below  $H$ .




not  
free



free



not  
free



free

## COMBINATORIAL QUESTION

How many  $\mathbb{A}[\tau, H]$  are free?

Group	# subgroups $H$	# transfer systems $\tau$	# pairs $(\tau, H)$	# free	% free
$C_p$	2	2	4	2	50
$C_{p^2}$	3	5	15	4	$\approx 27$
$C_{p^3}$	4	14	56	9	$\approx 16$
$C_{p^4}$	5	42	210	23	$\approx 11$
	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$

# COMBINATORIAL QUESTION

How many  $\mathbb{A}[\tau, H]$  are free?

Group	# subgroups $H$	# transfer systems $\tau$	# pairs $(\tau, H)$	# free	% free
$C_p$	2	2	4	2	50
$C_{p^2}$	3	5	15	4	$\approx 27$
$C_{p^3}$	4	14	56	9	$\approx 16$
$C_{p^4}$	5	42	210	23	$\approx 11$
	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$C_{p^n}$	$n + 1$	$\text{Cat}(n + 1)$	$P(n)$	$F(n)$	$\frac{F(n)}{P(n)}$

$$P(n) = (n + 1) \text{Cat}(n + 1)$$

$$F(n) = \sum_{i=0}^n \text{Cat}(i)$$

# COMBINATORIAL QUESTION

How many  $\mathbb{A}[\tau, H]$  are free?

Group	# subgroups $H$	# transfer systems $\tau$	# pairs $(\tau, H)$	# free	% free
$C_p$	2	2	4	2	50
$C_{p^2}$	3	5	15	4	$\approx 27$
$C_{p^3}$	4	14	56	9	$\approx 16$
$C_{p^4}$	5	42	210	23	$\approx 11$
	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$C_{p^n}$	$n + 1$	$\text{Cat}(n + 1)$	$P(n)$	$F(n)$	$\frac{F(n)}{P(n)}$

$$P(n) = (n + 1) \text{Cat}(n + 1)$$

$$F(n) = \sum_{i=0}^n \text{Cat}(i)$$

$$\lim_{n \rightarrow \infty} \frac{\sum_{i=0}^n F(i)}{\sum_{i=0}^n P(i)} = 0$$

THEOREM (Hill–Mehrlé–Quigley, 2023)

Fix a bijection  $\mathbb{G}: \mathbb{N} \rightarrow \{\text{isomorphism classes of finite groups}\}$ .

Let  $F(G)$  be the number of pairs  $(\tau, H)$  such that  $\mathbb{A}[\tau, H]$  is free.

Let  $P(G)$  be the total number of pairs  $(\tau, H)$  for  $G$ .

$$\lim_{n \rightarrow \infty} \frac{\sum_{i=1}^n F(\mathbb{G}(i))}{\sum_{i=1}^n P(\mathbb{G}(i))} = 0$$

## THEOREM (Hill–Mehrlé–Quigley, 2023)

Fix a bijection  $\mathbb{G}: \mathbb{N} \rightarrow \{\text{isomorphism classes of finite groups}\}$ .

Let  $F(G)$  be the number of pairs  $(\tau, H)$  such that  $\mathbb{A}[\tau, H]$  is free.

Let  $P(G)$  be the total number of pairs  $(\tau, H)$  for  $G$ .

$$\lim_{n \rightarrow \infty} \frac{\sum_{i=1}^n F(\mathbb{G}(i))}{\sum_{i=1}^n P(\mathbb{G}(i))} = 0$$

## SLOGAN

“Free incomplete Tambara functors are almost never free.”

## THEOREM (Hill–Mehrle–Quigley, 2023)

Fix a bijection  $\mathbb{G}: \mathbb{N} \rightarrow \{\text{isomorphism classes of finite groups}\}$ .

Let  $F(G)$  be the number of pairs  $(\tau, H)$  such that  $\mathbb{A}[\tau, H]$  is free.

Let  $P(G)$  be the total number of pairs  $(\tau, H)$  for  $G$ .

$$\lim_{n \rightarrow \infty} \frac{\sum_{i=1}^n F(\mathbb{G}(i))}{\sum_{i=1}^n P(\mathbb{G}(i))} = 0$$

## SLOGAN

“Free incomplete Tambara functors are almost never free.”

## WORK-IN-PROGRESS (Bingham-Franchere-Jones-Mehrle-Shoults-Yousef)

Computer code and recursive formulas to enumerate free transfer systems for  $C_{p^n}$ ,  $C_{pq^n}$ ,  $C_{p^2q^n}$ ,  $\dots$



THEOREM (Mehrle–Quigley–Stahlhauer, 2024)

Let  $G$  be a cyclic  $p$ -group for an odd prime  $p$ . If  $\mathbb{A}[\tau, H]$  is free, we construct well-behaved Koszul resolutions.

THEOREM (Mehrle–Quigley–Stahlhauer, 2024)

Let  $G$  be a cyclic  $p$ -group, any prime  $p$ . If  $\mathbb{A}[\tau, H]$  is *not* free, then it is infinite dimensional: there is a module with no finite resolution.

GOAL

A theory of minimal resolutions for Tambara functors

# TAKEAWAYS

Will equivariant algebraic topology solve all our problems?

Not yet.

# TAKEAWAYS

Will equivariant algebraic topology solve all our problems?

Not yet. But sooner rather than later!

- Renewed interest in the field
- We understand the tools much better than 10 years ago
- New computational aids, e.g. homotopical combinatorics

# TAKEAWAYS

Will equivariant algebraic topology solve all our problems?

Not yet. But sooner rather than later!

- Renewed interest in the field
- We understand the tools much better than 10 years ago
- New computational aids, e.g. homotopical combinatorics

WORK-IN-PROGRESS (Guillou–Keyes–Mehrlé)


Apply what we've learned about Tambara functors to make new calculations in equivariant homotopy theory.

Thank you!

## BONUS: MACKEY FUNCTORS

$\mathcal{F}\text{in}^G$  = category of finite  $G$ -sets and  $G$ -equivariant functions

$\text{Span}(\mathcal{F}\text{in}^G)$  = category of finite  $G$ -sets and **spans** of finite  $G$ -sets



### DEFINITION

A **Mackey functor** is a product-preserving functor

$$M: \text{Span}(\mathcal{F}\text{in}^G) \rightarrow \mathcal{A}\text{b}$$

$$M(H) := M(G/H)$$

$$\text{res}_K^H := M \left( G/H \begin{array}{c} \text{orange} \\ \llcorner \end{array} G/K \xrightarrow{\text{id}} G/K \right)$$

$$\text{tr}_K^H := M \left( G/K \xleftarrow{\text{id}} G/K \begin{array}{c} \text{blue} \\ \lrcorner \end{array} G/H \right)$$

# BONUS: TAMBARA FUNCTORS

$\text{Bispan}(\text{Fin}^G)$  = category of finite  $G$ -sets & **bispans** of finite  $G$ -sets



## DEFINITION

A **Tambara functor** is a product-preserving functor

$$T: \text{Bispan}(\text{Fin}^G) \rightarrow \text{Set}$$

such that each  $T(U)$  is a commutative ring

$$\text{res}_K^H := T \left( G/H \begin{array}{c} \text{orange} \\ \llcorner \end{array} G/K \xrightarrow{\text{id}} G/K \xrightarrow{\text{id}} G/K \right)$$

$$\text{tr}_K^H := T \left( G/K \xleftarrow{\text{id}} G/K \xrightarrow{\text{id}} G/K \begin{array}{c} \text{blue} \\ \gg \end{array} G/H \right)$$

$$\text{nm}_K^H := T \left( G/K \xleftarrow{\text{id}} G/K \begin{array}{c} \text{green} \\ \gg \end{array} G/H \xrightarrow{\text{id}} G/H \right)$$