

Due at the beginning of class on 4 March 2025

- Your answers should be neatly written and logically organized.
- You may collaborate on solving the problems, but the solutions you turn in should be your own.
- You may use any resource you find online (or elsewhere), but you must cite any resource you use.

Reading: [Mal23, Sections 2.1, 2.2, and 2.3].

- (1) Let $X_0 \xrightarrow{f_0} X_1 \xrightarrow{f_1} X_2 \xrightarrow{f_2} \dots$ be a sequence of spectra. Show that the homotopy colimit [Mal23, Definition 2.3.17] of this sequence commutes with stable homotopy groups, in the sense that

$$\pi_k(\operatorname{hocolim}_n X_n) = \operatorname{colim}_n \pi_k(X_n).$$

SOLUTION: We'll start with the right hand side and produce a chain of isomorphisms leading to the left hand side. Let $X_{n,m}$ be the m -th space in the spectrum X_n .

$$\begin{aligned} \operatorname{colim}_n \pi_k(X_n) &= \operatorname{colim}_n \operatorname{colim}_m \pi_{k+m}(X_{n,m}) && \text{definition of } \pi_k \text{ of a spectrum} \\ &\cong \operatorname{colim}_m \operatorname{colim}_n \pi_{k+m}(X_{n,m}) && \text{colimits commute with colimits} \\ &\cong \operatorname{colim}_m \pi_{k+m}(\operatorname{hocolim}_n X_{n,m}) && \text{the same fact for spaces (PSet 3, Q4)} \\ &\cong \pi_k(\operatorname{hocolim}_n X_n) && \text{definition of } \pi_k(\operatorname{hocolim}_n X_n) \end{aligned}$$

- (2) Eilenberg-MacLane spectra are characterized by their homotopy groups: if any other spectrum X satisfies $\pi_i X = 0$ for $i \neq 0$, then $X \simeq H(\pi_0 X)$.

- (a) Prove that $H(\mathbb{Z}/p)$ is stably equivalent to the homotopy cofiber of the map obtained by applying the functor H to $p: \mathbb{Z} \rightarrow \mathbb{Z}$.

SOLUTION: Consider the long exact sequence in homotopy of the cofiber sequence

$$H\mathbb{Z} \xrightarrow{Hp} H\mathbb{Z} \rightarrow \operatorname{cof}(Hp)$$

Because the homotopy of $H\mathbb{Z}$ is concentrated in degree zero, this boils down to an exact sequence

$$0 \rightarrow \pi_1 \operatorname{cof}(Hp) \rightarrow \mathbb{Z} \xrightarrow{p} \mathbb{Z} \rightarrow \pi_0 \operatorname{cof}(Hp) \rightarrow 0$$

But $p: \mathbb{Z} \rightarrow \mathbb{Z}$ is injective, so $\pi_1 \operatorname{cof}(Hp) = 0$, and we have a short exact sequence with third term isomorphic to \mathbb{Z}/p . Hence, $\operatorname{cof}(Hp)$ has one nonzero homotopy group $H\mathbb{Z}/p$, so it is stably equivalent to $H(\mathbb{Z}/p)$.

- (b) The rationalization $S_{\mathbb{Q}}$ of the sphere spectrum S is the homotopy colimit of the diagram

$$S \xrightarrow{1} S \xrightarrow{2} S \xrightarrow{3} S \xrightarrow{4} S \xrightarrow{5} S \rightarrow \dots$$

Prove that $S_{\mathbb{Q}}$ is stably equivalent to $H\mathbb{Q}$.

SOLUTION: The rationalization of S is the spectrum

$$S_{\mathbb{Q}} = \operatorname{hocolim}(S \xrightarrow{1} S \xrightarrow{2} S \xrightarrow{3} S \xrightarrow{4} \dots).$$

Recall that stable homotopy groups commute with homotopy colimits (turning the homotopy colimits to colimits in $\mathcal{A}b$). So,

$$\pi_n(S_{\mathbb{Q}}) = \operatorname{colim}(\pi_n S \xrightarrow{1} \pi_n S \xrightarrow{2} \pi_n S \xrightarrow{3} \dots)$$

This is isomorphic to $\pi_n S \otimes \mathbb{Q}$. Recall from the last problem set that $\pi_n S$ is a finite abelian group for $n > 0$ and $\pi_n S = \mathbb{Z}$ for $n = 0$. Therefore, $\pi_n(S_{\mathbb{Q}}) = \pi_n S \otimes \mathbb{Q} = 0$ for $n > 0$ and $\pi_0(S_{\mathbb{Q}}) = \mathbb{Z} \otimes \mathbb{Q} = \mathbb{Q}$. So this has one nonzero homotopy group, so $S_{\mathbb{Q}} \simeq H\mathbb{Q}$.

See also [Mal23, Chapter 1, Exercise 39] and [Mal23, Example 2.5.30].

(3) Let $\operatorname{ev}_0 : \mathcal{S}p \rightarrow \mathcal{T}op_*$ be the functor that evaluates a spectrum at its zeroth space: $\operatorname{ev}_0 X = X_0$.

(a) Prove that Σ^∞ is left adjoint to ev_0 .

SOLUTION: We must show there is a natural isomorphism

$$\operatorname{Hom}_{\mathcal{S}p}(\Sigma^\infty X, Y) \cong \operatorname{Hom}_{\mathcal{T}op_*}(X, Y_0)$$

Recall that a map of spectra $\Sigma^\infty \rightarrow Y$ is a sequence of pointed maps $f_n : \Sigma^n X \rightarrow Y_n$ such that the following squares commute for all n :

$$\begin{array}{ccc} \Sigma(\Sigma^n X) & \xrightarrow{\Sigma f_n} & \Sigma Y_n \\ \downarrow = & & \downarrow \\ \Sigma^{n+1} X & \xrightarrow{f_{n+1}} & Y_{n+1} \end{array}$$

This is naturally determined by the map $f_0 : X \rightarrow Y_0$. Denote the composite structure maps of Y by $y_n : \Sigma^n Y_0 \rightarrow Y_n$. Note that the diagram above forces $f_1 = y_1 \circ \Sigma f_0$. Suppose for induction that $f_n = y_n \circ \Sigma f_0$, then the above square commuting forces $f_{n+1} = y_{n+1} \circ \Sigma f_0$. So the map $\Sigma^\infty X \rightarrow Y$ uniquely determines the map $f_0 : X \rightarrow Y_0$ by the structure maps of Y . Going the other way, any map $g : X \rightarrow Y_0$ can naturally ascend to a map of spectra $\Sigma^\infty X \rightarrow Y$ by setting $g_0 = g$ and $g_n = y_n \circ \Sigma g_0$. These constructions are clearly natural in X and Y , so this completes the proof of the adjunction.

(b) For any spectrum X , let $\Omega^\infty X := \operatorname{ev}_0 R X$, where R is the fibrant replacement functor [Mal23, Proposition 2.2.9]. Use [Rie14, Exercise 2.2.15] to prove that there is an adjunction

$$\Sigma^\infty : \operatorname{ho}(\mathcal{T}op_*) \rightleftarrows \operatorname{ho}(\mathcal{S}p) : \Omega^\infty.$$

SOLUTION: First, observe that Σ^∞ is homotopical. For any space X , the stable homotopy groups of $\Sigma^\infty X$ are the homotopy groups of X , and if $X \xrightarrow{f} Y$ is a weak equivalence of spaces, the maps induced by $\Sigma^\infty f$ between the stable homotopy groups are exactly the isomorphisms induced by f between the homotopy groups of X and Y . Hence, Σ^∞ is a stable equivalence.

Note that the above is true so long as our spaces are well-based, so that $S^1 \wedge X$ agrees with the homotopy pushout of $* \leftarrow X \rightarrow *$. If the spaces are not well based, then these need not agree. See this mathOverflow answer: [Law13].

A derived functor for ev_0 can be computed by pre-composing with a right deformation of $\mathcal{S}p$. In particular, the functor R in the exercise is such a deformation. ev_0 is homotopical on the full subcategory of $\mathcal{S}p$ generated by the image of R : if $f : X \rightarrow Y$ is a stable equivalence between two Ω -spectra, then $f_0 : X_0 \rightarrow Y_0$ is a weak equivalence because $\pi_m \cong \pi_m(X_0)$, $\pi_m(Y) \cong \pi_m(Y_0)$ and since $\{f_m\}_{m \in \mathbb{N}}$ commute with the bonding maps, the isomorphisms $\pi_m(X) \rightarrow \pi_m(Y)$ induced by f is exactly the map induced by f_0 from $\pi_m X_0$ to $\pi_m(Y_0)$.

So $\Omega^\infty = \operatorname{ev}_0 \circ R$ is a right derived functor for ev_0 . By proposition 2.2.13 in Riehl, a derived functor computed thanks to a deformation is an absolute Kan extension. Therefore, by [Rie14, Exercise 2.2.15], Σ^∞ and Ω^∞ are an adjoint pair.

REFERENCES

- [Law13] Tyler Lawson. Must a weak homotopy equivalence induce an isomorphism between stable homotopy groups? "<https://mathoverflow.net/questions/148963/must-a-weak-homotopy-equivalence-induce-an-isomorphism-between-stable-homotopy-g>", 2013.
- [Mal23] Cary Malkiewich. Spectra and stable homotopy theory. http://people.math.binghamton.edu/malkiewich/spectra_book_draft.pdf, October 2023.
- [Rie14] Emily Riehl. *Categorical homotopy theory*, volume 24 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2014.