

MATH 472: INTRODUCTION TO TOPOLOGY

David Mehrle
david.mehrle@colostate.edu

Colorado State University
Fall 2026

Last updated September 9, 2026.

Contents

1	Topological Spaces and Continuous Functions	5
1.1	Metric Spaces	5
1.2	Bases for a Topology	6
1.3	Topologies	8
1.4	Closed Sets	10
1.4.1	Infinitely Many Prime Numbers	11
1.5	Continuous Functions	11
1.5.1	Building Continuous Functions	12
1.5.2	Homeomorphisms	13

Contents by Lecture

Lecture 00 on 24 August 2026	3
Lecture 01 on 26 August 2026	5
Lecture 02 on 28 August 2026	6
Lecture 03 on 31 August 2026	8
Lecture 04 on 02 September 2026	9
Lecture 05 on 04 September 2026	11
Lecture 06 on 9 September 2026	12

What is topology?

At its core, topology is the study of continuity, not only continuity of functions, but also continuous deformations of spaces, shapes, and surfaces. The popular conception of topology (Möbius strips, coffee cups = donuts, knots, etc.), however, fails to see what Kyle Ormsby calls the "...stunning breadth and extreme utility of topology in contemporary mathematics." [Orm, Week 1 Lecture Notes]. Here is a woefully incomplete list of places in mathematics and other fields of science where topology arises:

- (1) Manifold geometry: the study of manifolds and higher-dimensional surfaces requires passage from local regions to global geometry. Questions about gluing and connection.
- (2) Algebraic geometry: the Zariski topology, intersection theory, moduli spaces, complex varieties as topological spaces, etc.
- (3) Number Theory: there's a whole set of number-theoretic metrics out there called p-adic metrics, whereby two numbers are close to each other if they share the same divisors. Number theorists do analysis using this measure of distance.
- (4) Dynamical Systems and Flows: long-term behavior of differential equations, attractors, and qualitative features that persist under perturbation.
- (5) Combinatorics of Geometric Objects: graphs, posets, simplicial complexes; topological invariants constrain which objects exist and how they behave.
- (6) Topological Data Analysis: studying high-dimensional point clouds via their topology and using this to create summary statistics about data sets with homology.
- (7) Physics: Phase spaces, gauge symmetry, and homotopy theory in high energy physics.
- (8) Paleontology (and biology): I just read a book about modern paleontology and lots of the ideas therein seemed decidedly topological (although I doubt the author thought about them like that!). For example, distinguishing species of dinosaur by the number of holes in their skulls, or grouping species based on the number of shared characteristics (this is a notion of distance – hence continuity!)

The breadth of applications here implies that topology is a very abstract subject, and that is true in many ways. That's how topology shows up, but here's some of the ideas that characterize modern topology.

- (1) **Properties over construction:** when we study objects that are the same under continuous deformation, it often doesn't matter exactly how they are built. What we want to know is that no matter how you build them, they have the same topological properties. This leads naturally to the idea of category theory and universal properties, where we will *define* objects by the properties they have rather than by their construction.
- (2) **Topological equivalence:** there are many different ways to define a circle. Is it the image of $t \mapsto e^{2\pi i t}$ in $\mathbb{C}$? Or is it the image of $\theta \mapsto (\cos \theta, \sin \theta)$ in $\mathbb{R}^2$? Or is it what results from taking the unit interval

$[0, 1]$ and gluing 0 to 1? These are all really the same thing, and we say that they are topologically equivalent. The object that we really want to study is independent of the model chosen. There are multiple ways to measure topological equivalence, some more rigid than others.

- (3) **Invariants:** Given many different topologically equivalent avatars for the same object, we need some way to make differentiate between objects even when they're presented differently. Enter invariants: an invariant is a piece of data that stays the same (is "invariant") when replacing an object by a topologically equivalent one. We can use invariants to tell objects apart: if object X has invariant $I(X) = 1$ and object Y has invariant $I(Y) = 0$, then they cannot be topologically equivalent.

Chapter 1

Topological Spaces and Continuous Functions

1.1 Metric Spaces

The first principle above is properties over construction. With that in mind, our first task is to extract the properties of continuity that we learned from analysis and determine which properties determine them. We did this a little bit with the property of distance on the worksheet.

Definition 1.1.1. Let X be a set. A *metric* on X is a function $d: X \times X \rightarrow [0, \infty)$ with the following properties:

- For any $x \in X$, $d(x, x) = 0$.
- (positivity) For any $x, y \in X$, if $x \neq y$, then $d(x, y) > 0$
- (symmetry) For any $x, y \in X$, $d(x, y) = d(y, x)$
- (triangle inequality) For any $x, y, z \in X$, $d(x, z) \leq d(x, y) + d(y, z)$

Definition 1.1.2. A *metric space* is a pair (X, d) of a set X and a metric d on X .

Example 1.1.3. Here are some examples of metric spaces.

- (a) The metric space $(\mathbb{R}^n, d)$ where d is the standard Euclidean metric. For vectors $\vec{u}$ and $\vec{v}$, the distance between them is the length of their difference:

$$d(\vec{u}, \vec{v}) = \|\vec{u} - \vec{v}\|.$$

In $\mathbb{R}^2$ with vectors $\vec{u} = (u_0, u_1)$ and $\vec{v} = (v_0, v_1)$, we get

$$d(\vec{u}, \vec{v}) = \sqrt{(u_0 - v_0)^2 + (u_1 - v_1)^2}.$$

- (b) The taxicab metric on $\mathbb{R}^n$.
- (c) The L^p metric on $\mathbb{R}^n$.
- (d) The L^∞ metric on $\mathbb{R}^n$.
- (e) The set of binary strings of length n and Hamming distance: the number of positions at which the strings differ.

(f) The discrete metric on any space.

Given a notion of distance, we can talk about open balls in a metric space.

Definition 1.1.4. Let (X, d) be a metric space. Let $x \in X$ and let $r > 0$ be a positive real number. The *open ball of radius r around x* is the set

$$B_r(x) := \{y \in X \mid d(x, y) < r\}.$$

We saw some examples of this on the worksheet.

Example 1.1.5. In the hamming distance example, the open ball of radius 2 around any string is the set of all strings that differ from it by at most one character. The open ball of radius 1 around a given string is just the string itself (remember that it's a strict $<$).

Definition 1.1.6. Let (X, d) be a metric space. An *open set* in X is a set U such that for all $x \in U$, there exists some $r > 0$ such that $B_r(x) \subseteq U$.

Lemma 1.1.7. The open ball $B_r(x)$ is an open set.

Proof. Let $y \in B_r(x)$. Then $d(x, y) < r$ by the definition of an open ball. Let $\varepsilon = r - d(x, y)$. We claim that $B_\varepsilon(y) \subseteq B_r(x)$. Let $z \in B_\varepsilon(y)$. Then

$$\begin{aligned} d(x, z) &\leq d(x, y) + d(y, z) \\ &< d(x, y) + (r - d(x, y)) \\ &= r. \end{aligned}$$

Therefore, $d(x, z) < r$, so $z \in B_r(x)$. Hence, $B_\varepsilon(y) \subseteq B_r(x)$. □

Example 1.1.8. Draw some pictures of open sets in $\mathbb{R}^2$ with various metrics.

Question 1.1.9. Are there sets in $\mathbb{R}^2$ that are open in the Euclidean metric that are not open in the taxicab metric, or vice versa?

Example 1.1.10. Every subset of $\mathbb{R}^2$ is open for the discrete metric, because the open balls of radius 1 are just points!

1.2 Bases for a Topology

A topology on a set X is the collection of open sets on X (these must follow some rules, which we'll get to soon). For metric spaces, we can define the open sets by those sets U such that around each point $x \in U$, there is an open ball $B_\varepsilon(x)$ completely contained in U . This suggests that the important thing in the topology is not just the open sets, but the open balls in particular. The notion of a basis generalizes the collection of open balls, without any reference to distances.

Recall that if X is a set, then $\mathcal{P}(X)$ is the powerset of X : the set of all subsets of X .

Definition 1.2.1. Let X be a set. A (topological) basis for X is a collection $\beta \subseteq \mathcal{P}(X)$ of subsets of X satisfying the following properties:

- (covering) For each $x \in X$, there exists some $B \in \beta$ such that $x \in B$.
- (filtered) For all $B_1, B_2 \in \beta$ and all $x \in B_1 \cap B_2$, there exists $B_3 \in \beta$ such that $x \in B_3$ and $B_3 \subseteq B_1 \cap B_2$.

Another way to say the covering property is that the union of all sets in β is X . In symbols:

$$\bigcup_{B \in \beta} B = X.$$

Example 1.2.2. The set of open balls in a metric space forms a basis for that topology. You will prove this on your homework. The proof uses similar ideas to the ones that we used in 1.1.7.

Example 1.2.3. For $a, b \in \mathbb{Z}$ with $a \neq 0$, let $S(a, b) = \{an + b \mid n \in \mathbb{Z}\}$. This is called an *arithmetic progression*. Then a topological basis on $\mathbb{Z}$ is given by the set of all arithmetic progressions

$$\beta = \{S(a, b) \mid a, b \in \mathbb{Z}\}.$$

Let's check that this is a basis:

- (covering) Let $n \in \mathbb{Z}$. We must find some arithmetic progression $S(a, b) \in \beta$ such that $n \in S(a, b)$. One option is $S(1, n)$, which contains n because $n = 0(1) + n$.
- (filtered) Suppose that we have basis sets $S(a, b)$ and $S(c, d)$. Let $x \in S(a, b) \cap S(c, d)$. We want to find a basic open set that contains x and is contained in both $S(a, b)$ and $S(c, d)$. The arithmetic progression $S(\text{lcm}(a, c), x)$ is such a basic open set, and $S(\text{lcm}(a, c), x) \subseteq S(a, b) \cap S(c, d)$.

The topology generated by this basis (see below) is called the *Furstenberg topology*. You can use this topology to give a proof of the infinitude of primes different from Euler's proof.

Given a basis, we can define open sets just as we did with the open balls, but with no reference whatsoever to a metric.

Definition 1.2.4. Let X be a set with a basis β . A subset $U \subseteq X$ is *open in X* if for each $x \in U$, there exists some $B \in \beta$ such that $x \in B$ and $B \subseteq U$.

Example 1.2.5. Let X be a set with a topological basis β . Then $\emptyset$ is open (vacuously) and X is open (by the covering axiom).

With the notion of a topological basis, we have axiomatized the idea of open balls to something more general than metric spaces. But we can go even further, and axiomatize the open sets themselves. Let's first think about some of the properties of open sets.

Lemma 1.2.6. Let X be a set with topological basis β . Let I be a set, and for each $i \in I$, let $U_i \subseteq X$ be an open set in X . Then the union U of all of the U_i ,

$$U := \bigcup_{i \in I} U_i$$

is open as well.

Proof. Let $U = \bigcup_{i \in I} U_i$. We must show that for each $x \in U$, there is some $B \in \beta$ such that $x \in B$ and $B \subseteq U$. To that end, note that if $x \in U$, then $x \in U_j$ for some j . Since U_j is open, there is some $B \in \beta$ such that $x \in B$ and $B \subseteq U_j$. But note that $U_j \subseteq U$ (U_j is a subset of the union of all of the U_i s), and therefore this B is the one we search for. In particular, $x \in B$ and $B \subseteq U_j \subseteq U$. $\square$

Lemma 1.2.7. Let X be a set with topological basis β . Let U and V be open sets in X . Then $U \cap V$ is open.

Proof. To show that $U \cap V$ is open, we must pick an arbitrary $x \in U \cap V$ and then find some $B \in \beta$ such that $x \in B$ and $B \subseteq U \cap V$.

To that end, let $x \in U \cap V$. Then $x \in U$, and U is open, so there exists some $B_1 \in \beta$ such that $x \in B_1$ and $B_1 \subseteq U$.

Similarly, $x \in V$, and V is open, so there exists some $B_2 \in \beta$ such that $x \in B_2$ and $B_2 \subseteq V$.

Now we have $x \in B_1$ and $x \in B_2$, which means that $x \in B_1 \cap B_2$. By the filtering axiom of a basis, this means that there exists some $B_3 \in \beta$ such that $x \in B_3$ and $B_3 \subseteq B_1 \cap B_2$.

This is the B that we were searching for: since $B_1 \subseteq U$ and $B_2 \subseteq V$, we have

$$B_1 \cap B_2 \subseteq U \cap V.$$

Then $B_3 \subseteq B_1 \cap B_2$ implies that $B_3 \subseteq U \cap V$. We also have $x \in B_3$, so we're done! $\square$

Lemma 1.2.8. Let X be a set with topological basis β . Let $U_1, \dots, U_n$ be open sets in X . Then the intersection

$$\bigcap_{i=1}^n U_i$$

is open.

Proof. Proof by induction on n . For $n = 0$ the empty intersection is all of X , which we already showed was open in 1.2.5. For $n = 1$, the intersection of one open set is open again. For $n = 2$, this is the previous lemma 1.2.7. This handles the base cases.

For the inductive step, assume that the intersection of any $n - 1$ open sets is open. Let $V = \bigcap_{i=1}^{n-1} U_i$. Then by the previous lemma 1.2.7, $U_n \cap V$ is the intersection of two opens, and is therefore open. But

$$U_n \cap V = U_n \cap \bigcap_{i=1}^{n-1} U_i = \bigcap_{i=1}^n U_i,$$

which is what we were trying to prove is open. $\square$

Example 1.2.9. We can't get away with arbitrary intersections of open sets being open. For example, consider the intersection of all open balls of radius $1/n$ in the Euclidean topology on $\mathbb{R}^2$:

$$\bigcap_{n=1}^{\infty} B_{\frac{1}{n}}((0,0)).$$

This set contains just one point: the origin. And a single point isn't open.

1.3 Topologies

Knowing what we know about the open sets in X with respect to a basis β , we can axiomatize the open sets.

Definition 1.3.1. Let X be a set. A *topology* τ on X is a collection $\tau \subseteq \mathcal{P}(X)$ of subsets of X such that

- Arbitrary unions of sets in τ are again in τ . In symbols: if I is a set and $U_i \in \tau$ for all $i \in I$, then

$$\bigcup_{i \in I} U_i \in \tau.$$

- Finite intersections of sets in τ are again in τ . In symbols: if $U_1, \dots, U_n \in \tau$ for some nonnegative integer n , then

$$\bigcap_{i=1}^n U_i \in \tau.$$

We call the elements of τ the *open sets* of the topology.

Definition 1.3.2. A *topological space* is a pair (X, τ) of a set X and a topology τ on X .

Remark 1.3.3. Some authors also demand the axiom that $\emptyset \in \tau$ and $X \in \tau$. But these follow from the axioms given above! Can you see why?*

This is even more abstract than the abstract bases we talked about earlier! But we already have some examples given by sets with bases.

*The answer is that the union of zero sets is the empty set, so $\emptyset \in \tau$ follows from the unions axiom. Similarly, the intersection of zero sets is all of X , so $X \in \tau$ follows from the finite intersection axiom.

Theorem 1.3.4. Let X be a set with a topological basis β . Let τ be the set

$$\tau := \left\{ U \subseteq X \mid U \text{ is open with respect to the basis } \beta \right\}.$$

We call this the topology generated by the basis β . Then τ is a topology on X .

Proof. This is what we proved in the last section. See 1.2.6 and 1.2.8 □

But there are many, many more examples.

Example 1.3.5. On any set X , the *discrete topology* is the topology τ_{disc} where every set is open. Then τ_{disc} is the whole powerset of X : $\tau_{\text{disc}} := \mathcal{P}(X)$

Example 1.3.6. On any set X , the *indiscrete topology* or the *concrete topology* is the topology $\tau_{\text{conc}} := \{\emptyset, X\}$. That is, the only open sets in this topology are empty or the whole set.

Example 1.3.7. The set $\{0, 1\}$ has a topology given by

$$\tau = \{\emptyset, \{1\}, \{0, 1\}\}.$$

This is not the discrete topology (because $\{0\}$ isn't open), and it's not the concrete topology (because $\{1\}$ is open). So this is a genuinely new topology on $\{0, 1\}$. Are there any others?

Example 1.3.8. Given a topological space (X, τ) and a subset $A \subseteq X$, we can build a new topology on A by saying that the open sets in A are the sets of the form $U \cap A$ for some open set U of X . In symbols, the topology on A is

$$\tau' = \left\{ U \cap A \mid U \in \tau \right\}.$$

This is called the *subspace topology*.

Example 1.3.9. Cofinite topology on $\mathbb{N}$. The topology τ_{cof} is

$$\tau_{\text{cof}} := \left\{ U \subseteq \mathbb{N} \mid \mathbb{N} \setminus U \text{ is finite} \right\} \cup \{\emptyset\}.$$

In words, the nonempty open sets in this topology are the subsets of $\mathbb{N}$ whose complement is finite. Note that this is different than infinite sets being open! The odd numbers is an infinite set whose complement (the even numbers) is still infinite.

Example 1.3.10. The poset topology. Let $(P, \leq)$ be a poset. A subset $U \subseteq P$ is called *upwards closed* if for all $x \in U$ and $y \in P$ such that $x \leq y$, then $y \in U$ as well. In other words, U is upwards closed if $x \in U$ implies that $y \in U$ for all $y \geq x$.

Let τ be the set of all upwards closed subsets of P . Then τ is a topology on P .

$$\tau = \left\{ U \subseteq P \mid U \text{ is upwards closed} \right\}.$$

Frequently in the previous few sections, we made use of open balls around a point. But when we're talking about general topological spaces, we no longer have open balls. So we use the idea of a neighborhood instead. This is very important vocabulary that we will use often!

Definition 1.3.11. Let (X, τ) be a topological space and let $x \in X$. An *(open) neighborhood* of x is an open set $U \in \tau$ such that $x \in U$.

1.4 Closed Sets

Another important idea in topology is the notion of a closed set.

Definition 1.4.1. Let (X, τ) be a topological space. A subset $K \subseteq X$ is *closed* if $X \setminus K$ is open.

Example 1.4.2. Since $\emptyset$ and X are always open, then $X = X \setminus \emptyset$ and $\emptyset = X \setminus X$ are always closed as well. This illustrates that closed and open are not mutually exclusive!

It is false to say that if a set is not open, then it is closed, as the following example shows.

Example 1.4.3. In the Euclidean topology on $\mathbb{R}$, open intervals (a, b) are open sets, and closed intervals $[a, b]$ are closed sets. But there are intervals of the form $[a, b)$ and $(a, b]$ that are neither open nor closed.

Example 1.4.4. In the cofinite topology on $\mathbb{N}$, the closed sets are either finite sets or all of $\mathbb{N}$.

You can also define a topology by specifying the closed sets. In fact, this is probably the better way to define some topologies, like the cofinite topology.

Theorem 1.4.5. Let X be a set. Let $\kappa \subseteq \mathcal{P}(X)$ be a collection of subsets of X such that

- (a) κ is closed under finite unions: if $K_1, \dots, K_n \in \kappa$, then $\bigcup_{i=1}^n K_i \in \kappa$.
- (b) κ is closed under arbitrary intersections: if I is a set and $K_i \in \kappa$ for each $i \in I$, then $\bigcap_{i \in I} K_i \in \kappa$.

Then

$$\tau = \left\{ X \setminus K \mid K \in \kappa \right\}.$$

is a topology on X .

Proof. To prove that τ is a topology on X , we must show that τ is closed under finite intersections and arbitrary unions.

- (τ is closed under finite intersection) Let $U_1, \dots, U_n \in \tau$. Because every set in τ is the complement of a set in κ , then for each $i = 1, \dots, n$ there is some $K_i \in \kappa$ such that $U_i = X \setminus K_i$. Then

$$\bigcap_{i=1}^n U_i = \bigcap_{i=1}^n (X \setminus K_i) = X \setminus \left(\bigcup_{i=1}^n K_i \right).$$

Since κ is closed under finite union, this shows that the intersection of all the U_i s is again a complement of a set in κ . Hence, $\bigcap_{i=1}^n U_i \in \tau$.

- (τ is closed under arbitrary union) Let I be a set and let $U_i \in \tau$ for each $i \in I$. Because every set in τ is the complement of a set in κ , then for each $i \in I$, there is some $K_i \in \kappa$ such that $U_i = X \setminus K_i$. Then

$$\bigcup_{i \in I} U_i = \bigcup_{i \in I} (X \setminus K_i) = X \setminus \left(\bigcap_{i \in I} K_i \right).$$

Since κ is closed under arbitrary intersection, this shows that the union of all of the U_i s is again a complement of a set in κ . Hence, $\bigcup_{i \in I} U_i \in \tau$.

This shows that τ is both closed under finite intersection and arbitrary union, and hence τ is a topology. $\square$

1.4.1 Infinitely Many Prime Numbers

We can return to Example 1.2.3 and use it to prove that there are infinitely many prime numbers. Recall that we can define a topology on $\mathbb{Z}$ by stating that the basic open sets are the arithmetic progressions $S(a, b)$ with $a \neq 0$, where

$$S(a, b) = \{an + b \mid n \in \mathbb{Z}\}$$

That is, a set $U \subseteq \mathbb{Z}$ is open in this topology if for all $x \in U$, there is some $S(a, b)$ such that $x \in S(a, b)$ and $S(a, b) \subseteq U$.

Lemma 1.4.6. $S(a, b)$ is a closed set.

Proof. We must write $S(a, b)$ as the complement of an open set to show that it is closed.

$$S(a, b) = \mathbb{Z} \setminus \left(\bigcup_{i=1}^{a-1} S(a, b+i) \right)$$

□

This is an example of a topology in which there are many sets that are both open and closed!

Lemma 1.4.7. Every nonempty open set in this topology is infinite.

Proof. If U is open and nonempty, then there is some $x \in U$. Since U is open, then there is some $S(a, b)$ such that $x \in S(a, b)$ and $S(a, b) \subseteq U$. Since $S(a, b)$ is an infinite set, then U must be infinite as well. □

Theorem 1.4.8. There are infinitely many prime numbers.

Proof. Proof by contradiction. Assume that there are finitely many prime numbers.

By Lemma 1.4.7, the set $\{-1, +1\}$ cannot be open. Therefore, $\mathbb{Z} \setminus \{-1, +1\}$ cannot be a closed set. However, we can write

$$\mathbb{Z} \setminus \{-1, +1\} = \bigcup_{p \text{ prime}} S(p, 0),$$

which by Lemma 1.4.6 is a finite union of closed sets, and therefore closed. (Remember we are operating under the assumption of finitely many primes.) This contradicts the fact that $\mathbb{Z} \setminus \{-1, +1\}$ cannot be closed, so our assumption must be false and there are infinitely many primes. □

1.5 Continuous Functions

Topology appears whenever we have a notion of closeness, whether we define it via a metric or a topological basis or a topology.

Definition 1.5.1. Let X and Y be sets, and let $f: X \rightarrow Y$ be a function.

(a) Let $B \subseteq Y$. The *preimage* of B under f is the set

$$f^{-1}(B) = \{x \in X \mid f(x) \in B\}.$$

(b) Let $A \subseteq X$. The *image* of A under f is the set

$$f(A) = \{f(a) \mid a \in A\}.$$

Definition 1.5.2. Let X and Y be topological spaces, and let $f: X \rightarrow Y$ be a function. The function f is *continuous* if for each open subset V of Y , $f^{-1}(V)$ is open.

Let's see how this definition of continuity matches up with the definition of continuity you learned in analysis.

Proposition 1.5.3. *Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function, meaning that for all $x_0 \in \mathbb{R}$ and all $\varepsilon > 0$, there is some $\delta > 0$ such that if $|x - x_0| < \delta$, then $|f(x) - f(x_0)| < \varepsilon$.*

Then f is continuous in the sense of topology (1.5.2).

Proof. Assume that $f: \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function. Let $V \subseteq \mathbb{R}$ be an open set. We want to show that $f^{-1}(V)$ is open.

To show that $f^{-1}(V)$ is open, we must pick some point $x_0 \in f^{-1}(V)$ and show that we can fit an open ball around x_0 entirely inside $f^{-1}(V)$.

Since V is open, we can fit an open ball around any point inside of V . In particular, $f(x_0) \in V$ by the definition of $x_0 \in f^{-1}(V)$. Therefore, there is an open ball $B_\varepsilon(f(x_0)) \subseteq V$.

The analysis definition of continuity shows us that if $x \in B_\delta(x_0)$, then $f(x) \in B_\varepsilon(f(x_0))$. In other words,

$$B_\delta(x_0) \subseteq f^{-1}(B_\varepsilon(f(x_0))).$$

Since $B_\varepsilon(f(x_0)) \subseteq V$, we see then that $f^{-1}(B_\varepsilon(f(x_0))) \subseteq f^{-1}(V)$. Therefore,

$$B_\delta(x_0) \subseteq f^{-1}(B_\varepsilon(f(x_0))) \subseteq f^{-1}(V).$$

This finds an open set around x_0 that is contained in $f^{-1}(V)$. Since our choice of x_0 was arbitrary, $f^{-1}(V)$ is open. Therefore, f is continuous in the sense of topology. $\square$

Exercise 1.5.4. Prove that if $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous in the sense of topology (1.5.2), then $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous in the sense of analysis.

Example 1.5.5. • identity functions are continuous

- every function whose domain is discrete is continuous
- from X one topology to X with another topology
- taxicab geometry and regular geometry

1.5.1 Building Continuous Functions

How can you go about building continuous functions from one space to another? There are a couple of standard techniques.

Proposition 1.5.6. *Let X and Y be topological spaces and let $y \in Y$. Then the constant function $f: X \rightarrow Y$ defined by $f(x) = y$ for all $x \in X$ is continuous.*

Proof. Let $V \subseteq Y$ be open. If $y \in V$, then $f^{-1}(V) = X$, which is open. If $y \notin V$, then $f^{-1}(V) = \emptyset$, which is also open. Therefore, the preimage of any open in Y is open, which means that f is continuous. $\square$

Proposition 1.5.7. *Let X, Y , and Z be topological spaces with continuous functions $f: X \rightarrow Y$ and $g: Y \rightarrow Z$. Then $g \circ f: X \rightarrow Z$ is continuous.*

Proof. To show that $g \circ f: X \rightarrow Z$ is continuous, we need to show that for each open set $W \subseteq Z$, $(g \circ f)^{-1}(W)$ is open in X . But notice that

$$(g \circ f)^{-1}(W) = f^{-1}(g^{-1}(W)).$$

Since g is continuous and W is open, then $g^{-1}(W)$ is open. Since f is continuous and $g^{-1}(W)$ is open, then $f^{-1}(g^{-1}(W))$ is open. Therefore, $(g \circ f)^{-1}(W)$ is open, so $g \circ f$ is continuous. $\square$

Lemma 1.5.8. Let X be a topological space with a subspace A . Let $i: A \hookrightarrow X$ be the inclusion function. Then $i: A \rightarrow X$ is continuous.

Proof. Let $U \subseteq X$ be open. Then $i^{-1}(U) = U \cap A$, which is open in A by the definition of the subspace topology. $\square$

Proposition 1.5.9. Let $f: X \rightarrow Y$ be a continuous function between topological spaces. Let $A \subseteq X$ be a subspace with the subspace topology. Then if we restrict the domain of f to A , it is still continuous.

Proof. Let $i: A \hookrightarrow X$ be the inclusion function. Then restricting the domain of f to A is the same as composing $f \circ i$. Since f is continuous (by assumption) and i is continuous (by previous lemma), then $f \circ i$ is continuous by a previous proposition. $\square$

Lemma 1.5.10. Let X be a topological space with subspace A . Give A the subspace topology.

- (a) If A is open in X and $U \subseteq A$ is open in A , then U is open in X .
- (b) If A is closed in X and $K \subseteq A$ is closed in A , then K is closed in X .

Theorem 1.5.11 (Pasting). Let X be a topological space such that $X = A \cup B$, where A and B are closed in X . Then let $f: A \rightarrow Y$ and $g: B \rightarrow Y$ be continuous functions, where A and B are given the subspace topology. If $f(x) = g(x)$ for all $x \in A \cap B$, then f and g can be combined to give a continuous function $h: X \rightarrow Y$ defined by

$$h(x) = \begin{cases} f(x) & \text{if } x \in A \\ g(x) & \text{if } x \in B. \end{cases}$$

Proof. We will use the formulation of continuity where $f: Y \rightarrow Z$ is continuous if and only if for each closed subset $K \subseteq Z$, $f^{-1}(K)$ is closed in Y .

Let C be a closed set in Y . Then $h^{-1}(C) = f^{-1}(C) \cup g^{-1}(C)$. Since f is continuous, $f^{-1}(C)$ is closed in A , and therefore closed in X by the previous lemma. Since g is continuous, $g^{-1}(C)$ is closed in B , and therefore closed in X by the previous lemma. Hence, $h^{-1}(C)$ is the union of two closed sets, and therefore closed. So h sends closed sets to closed sets and is therefore continuous. $\square$

Example 1.5.12. The function

$$h(x) = \begin{cases} 2x & \text{if } x \leq 0 \\ x/2 & \text{if } x \geq 0 \end{cases}$$

is defined by two functions $f: (-\infty, 0] \rightarrow \mathbb{R}$, $f(x) = 2x$ and $g: [0, \infty) \rightarrow \mathbb{R}$, $g(x) = x/2$ on two closed sets $A = (-\infty, 0]$ and $B = [0, \infty)$. The functions f and g agree on $A \cap B = \{0\}$. The pasting theorem guarantees that h is continuous.

1.5.2 Homeomorphisms

Definition 1.5.13. A function $f: X \rightarrow Y$ between topological spaces is a *homeomorphism* if

- f is continuous
- f is a bijection with inverse function g
- the inverse function $g: Y \rightarrow X$ is continuous

Definition 1.5.14. If $f: X \rightarrow Y$ is a homeomorphism, then we say that X and Y are *homeomorphic* and write $X \cong Y$.

It's very important that the inverse function is also continuous! It's not automatic that the inverse is continuous, as the following example shows.

Example 1.5.15. $[0, 1) \rightarrow S^1$ given by $t \mapsto e^{2\pi i t}$ is continuous and a bijection, but no continuous inverse. In particular, $[0, \frac{1}{2})$ is open in the interval $[0, 1)$ under the subspace topology but its preimage in the circle S^1 is not open in the subspace topology on $\mathbb{C}$; it includes $z = 1$ along the real axis.

There's another useful characterization of homeomorphisms.

Proposition 1.5.16. *Let $f: X \rightarrow Y$ be a function between topological spaces. Then the following are equivalent:*

- (a) $f: X \rightarrow Y$ is a homeomorphism
- (b) $f: X \rightarrow Y$ is a bijection and V is open in Y if and only if $f^{-1}(V)$ is open in X

Proof. ($\implies$). Let $f: X \rightarrow Y$ be a homeomorphism, and let U be an open set in Y . Since f is continuous, then $f^{-1}(U)$ is open in X .

Since f is a homeomorphism, it has an inverse $g: Y \rightarrow X$, and g is continuous. So if $f^{-1}(V)$ is open in X , then $g^{-1}(f^{-1}(V))$ is open in Y . But $g^{-1}(f^{-1}(V)) = (f \circ g)^{-1}(V) = V$, so V is open in Y .

($\impliedby$). Since $f: X \rightarrow Y$ is a bijection, we just need to show that f and its inverse $g: Y \rightarrow X$ are both continuous.

We assume that V is open in Y if and only if $f^{-1}(V)$ is open in X . In particular, this shows that f is continuous.

Let $U \subseteq X$ be open. Then $g^{-1}(U)$ is the same thing as $f(U)$, because f and g are inverses. Then $f^{-1}(f(U)) = U$ is open in X , so $f(U)$ must be open in Y . But $f(U) = g^{-1}(U)$, so $g^{-1}(U)$ is open in Y , and therefore g is continuous.

Hence, f is a homeomorphism. □

This shows that a homeomorphism $f: X \rightarrow Y$ not only sends points in X bijectively to points in Y , but also open sets in X bijectively to open sets in Y . For that reason, whenever there is any property of X that can be expressed entirely in terms of the topology of X (that is, entirely in terms of the open sets of X), then that property applies to Y as well. Such a property of X is called a *topological property*.

Definition 1.5.17. A *topological property* P is a property of topological spaces such that a space X has property P and $X \cong Y$ implies that Y has property P as well.

In other words, a topological property is a property that is *homeomorphism invariant*.

Bibliography

- [Arm83] M. A. Armstrong. *Basic Topology*. Undergraduate Texts in Mathematics. Springer Science+Business Media, New York, NY, 1 edition, July 1983. <https://math.mit.edu/~hrm/palestine/armstrong-basic-topology.pdf>.
- [AT00] C. L. Adler and James Tanton. π is the minimum value for pi. *The College Mathematics Journal*, 31(2):102–106, 2000. <https://doi.org/10.1080/07468342.2000.11974122>.
- [BBRS21] Clark Bray, Adrian Butscher, and Simon Rubinstein-Salzedo. *Algebraic Topology*. Springer, Cham, 1 edition, 2021.
- [ES99] Russell Euler and Jawad Sadek. The π s go full circle. *Mathematics Magazine*, 72(1):59–63, 1999. <https://doi.org/10.1080/0025570X.1999.11996701>.
- [Goo05] Sue E. Goodman. *Beginning Topology*, volume 10 of *Pure and Applied Undergraduate Texts*. American Mathematical Society, Providence, RI, 2005.
- [Hat02] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002. <https://pi.math.cornell.edu/~hatcher/AT/ATpage>.
- [Mat] Showing that the triangle inequality holds for the l-infinity norm as a metric. <https://math.stackexchange.com/questions/1151794/showing-that-the-triangle-inequality-holds-for-the-l-infty-norm-as-a-metric>. Accessed 2026-08-26.
- [Orm] Kyle Ormsby. MATH 342 Spring 2026. <https://e-infinity.space/math342spring26/course/>. Course webpage, accessed 2026-08-23.
- [Proa] ProofWiki. Standard Discrete Metric is Metric. https://proofwiki.org/wiki/Standard_Discrete_Metric_is_Metric. Accessed 2026-08-26.
- [Prob] ProofWiki. Taxicab Metric is Metric. https://proofwiki.org/wiki/Taxicab_Metric_is_Metric. Accessed 2026-08-26.