Why is i^i a real number?

David Meyer

dmm613@gmail.com

Last update: November 17, 2022

1 Introduction

There are a few ways to think about this question. In Section 3 we use the fact that complex conjugation is an automorphism of $\mathbb{C}$. In Section 4 we use the power series expansion of i^i to show that $i^i \in \mathbb{R}$, and in Section 5 we rely on Euler's formula to show that $i^i \in \mathbb{R}$. Section 6 does a bit of arithmetic to show what i^i equals numerically. Finally Section 7 offers a few conclusions.

Before launching into all of this, note that we will make heavy use of the exponential function $\exp(z)$ (and therefore as you might expect, this means heavy use of $\log(z)$, where by $\log(z)$ we mean $\log_e(z)$ aka $\ln(z)$). The important point for this discussion is that $\exp(z)$ is injective over $\mathbb{R}$ but not over $\mathbb{C}$. More specifically, for a complex number z the the complex logarithm $\log(z)$ is defined as the inverse function to the exponential function. That is, it satisfies $e^{\log z} \equiv z$. Now since $re^{i\theta} = re^{i\theta+2\pi ik}$ for all $k \in \mathbb{Z}$ we have that for any choice of k, $\log(re^{i\theta+2\pi ki})$ is a valid inverse for $re^{i\theta}$. The logarithm is therefore a multivalued function and each value of k defines what is called a different *branch* of the logarithm. The principal branch [8] usually corresponds to the choice $k = 0$.

2 First: Why does a complex number z equal $r\mathrm{e}^{i\theta}$?

Our first question regards the nature of a complex number z, where the length of the line from the origin to the point z, $|z|$, equals r. In particular, why does a complex number $z = a + bi = r\mathrm{e}^{i\theta}$?

Well, consider the complex plane, shown in Figure 1:



Figure 1: The Complex Plane



Figure 2: Euler's Formula, the Unit Circle, and the Complex Plane

Figures 1 and 2 make it easy to see why $z = re^{i\theta}$. Specifically

$$\begin{aligned}
z &= a + ib && \text{\# definition of a point } z \text{ in the complex plane} \\
&= r\cos\theta + ir\sin\theta && \text{\# switch to polar coordinates: } a = r\cos\theta \text{ and } b = r\sin\theta \\
&= r\left(\cos\theta + i\sin\theta\right) && \text{\# factor out } r \\
&= re^{i\theta} && \text{\# } e^{i\theta} = \cos\theta + i\sin\theta \text{ (Euler's formula [7])}
\end{aligned}$$

So we see that $z = re^{i\theta}$. ■

3 Automorphism of $\mathbb{C}$

In this section we look at solving this puzzle using the fact that complex conjugation is an automorphism of $\mathbb{C}$ [3]. For this approach we need the following facts:

1. If a complex number z equals its complex conjugate $\overline{z}$ then $z \in \mathbb{R}$
2. $-i = i^{-1}$
3. $\overline{z} = re^{-i\theta}$

To see 1., consider the following argument: Let z be a complex number so that $z = a + bi$. Then $\overline{z} = a - bi$. So $z = \overline{z} \Rightarrow a + bi = a - bi$. Subtracting a from both sides of the right hand side of this implication gives us $bi = -bi$ or $2bi = 0$. Since we know that $2 \neq 0$ and $i \neq 0$ it must be the case that $b = 0$. Said another way: $\text{Im}(z) = 0$ and $\text{Re}(z) = z$. That is, $z \in \mathbb{R}$. So if $z = \overline{z}$ we know that $z \in \mathbb{R}$.

To see 2., notice that $-(i \cdot i) = -i^2 = -(-1) = 1$ and $-i \cdot i = 1 \Rightarrow -i = i^{-1}$.

To see 3., consider

$$\begin{aligned}
\overline{z} &= \overline{r \cdot (\cos\theta + i\sin\theta)} && \text{\# since } z = r(\cos\theta + i\sin\theta) \text{ (Figure 1)} \\
&= \overline{r} \cdot \overline{(\cos\theta + i\sin\theta)} && \text{\# since } \overline{z_1 \cdot z_2} = \overline{z_1} \cdot \overline{z_2} \text{ (product rule for complex conjugates [4])} \\
&= r \cdot \overline{(\cos\theta + i\sin\theta)} && \text{\# since } \overline{r} = \overline{r + 0i} = r - 0i = r \ (\overline{r} = r \text{ for } r \in \mathbb{R}) \\
&= r \cdot (\cos\theta - i\sin\theta) && \text{\# since } \overline{a + bi} = a - bi \\
&= re^{-i\theta} && \text{\# since } e^{-i\theta} = \cos\theta - i\sin\theta \text{ (corollary to Euler's formula [2])}
\end{aligned}$$

Now, we want to show that $\overline{i^i} = i^i$, which would show that $i^i \in \mathbb{R}$ by the above argument. So consider:

$$\begin{aligned}
\overline{i^i} &= \overline{i}^{\overline{i}} && \text{\# see proof below} \\
&= (-i)^{-i} && \text{\# since } i = 0 + 1i \text{ and } \overline{0 + 1i} = -1i = -i \\
&= (i^{-1})^{-i} && \text{\# since } -i = i^{-1} \text{ (see 2. above)} \\
&= i^{(-1 \cdot -i)} && \text{\# since } (x^m)^n = x^{mn} \\
&= i^i && \text{\# since } -1 \cdot -i = i
\end{aligned}$$

So we see that $\overline{i^i} = i^i$ and so $i^i \in \mathbb{R}$. ■

All ok, but why does $\overline{i^i} = \overline{i}^{\overline{i}}$? To see why first notice that

$$\begin{aligned}
\log(\overline{z}) &= \log(\overline{re^{i\theta}}) && \text{\# since } z = re^{i\theta} \text{ (Figure 1)} \\
&= \log(re^{-i\theta}) && \text{\# by item 3. above} \\
&= \log\left(\frac{r}{e^{i\theta}}\right) && \text{\# since } a^{-b} = \frac{1}{a^b} \\
&= \log(r) - \log(e^{i\theta}) && \text{\# by the quotient rule for logarithms [6]}
\end{aligned}$$

So

$$\log(\overline{z}) = \log(r) - \log(e^{i\theta}) \tag{1}$$

Next, notice that

$$\begin{aligned}
\overline{\log(z)} &= \overline{\log(re^{i\theta})} && \text{\# since } z = re^{i\theta} \\
&= \overline{\log(r) + \log(e^{i\theta})} && \text{\# by the product rule for logarithms} \\
&= \overline{\log(r)} + \overline{\log(e^{i\theta})} && \text{\# by sum rule for conjugates [5]} \\
&= \log(r) + \overline{\log(e^{i\theta})} && \text{\# since } \log(r) \in \mathbb{R} \\
&= \log(r) + \overline{i\theta} && \text{\# since } \log(e^{i\theta}) = i\theta \\
&= \log(r) - i\theta && \text{\# since } \overline{i\theta} = \overline{0 + i\theta} = -i\theta \\
&= \log(r) - \log(e^{i\theta}) && \text{\# since } i\theta = \log(e^{i\theta}) \\
&= \log(\overline{z}) && \text{\# by Equation (1)}
\end{aligned}$$

Now we know that

$$\log(\overline{z}) = \overline{\log(z)} \tag{2}$$

and so we can see that

$$\begin{aligned}
\log(\overline{z^w}) &= \overline{\log(z^w)} && \text{\# Equation (2)} \\
&= \overline{w \log(z)} && \text{\# by the power rule for logarithms} \\
&= \overline{w} \cdot \overline{\log(z)} && \text{\# by the product rule for conjugation} \\
&= \overline{w} \cdot \log(\overline{z}) && \text{\# by Equation (2)} \\
&= \log(\overline{z}^{\overline{w}}) && \text{\# by the power rule for logarithms}
\end{aligned}$$

So now we know that

$$\log(\overline{z^w}) = \log(\overline{z}^{\overline{w}}) \tag{3}$$

Exponentiating both sides of Equation (3) gives us

$$\overline{z^w} = \overline{z}^{\overline{w}} \tag{4}$$

So now if we set $z = w = i$ in Equation (4) we see that $\overline{i^i} = \overline{i}^{\overline{i}}$. ■

Note that this argument relies on the fact that complex conjugation is an automorphism of $\mathbb{C}$. So one "problem" with this approach is that you would first need to know not only that conjugation is an automorphism of $\mathbb{C}$ but also that the automorphism is equipped with exponentiation. So one could argue that this is proof is incomplete since it may not be true for branches (of the logarithm) other than the principal branch [8].

4 Power Series

We know that the Taylor series [9] for $e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}$. We also know from Euler's formula [1] that $i = e^{i\frac{\pi}{2}}$ (since $e^{i\frac{\pi}{2}} = \cos\frac{\pi}{2} + i\sin\frac{\pi}{2} = i$). Note that the same result can be obtained with $x \in \left\{\frac{\pi}{2}, \frac{5\pi}{2}, \frac{9\pi}{2}, \ldots\right\}$, so in general $x = \frac{\pi}{2} \cdot (1 + 4n)$ where $n \in \mathbb{N} \cup \{0\}$.

So we see that

$$\begin{aligned}
i^t &= \left(e^{\frac{i\pi}{2}}\right)^t && \text{\# Euler's formula for } i\text{, raised to the power } t \\
&= e^{\frac{it\pi}{2}} && \text{\# since } (x^m)^n = x^{mn} \\
&= \sum_{k=0}^{\infty} \frac{\left(\frac{it\pi}{2}\right)^k}{k!} && \text{\# since } e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!} \text{ and } x = \frac{it\pi}{2}
\end{aligned}$$

If we let $t = i$ we get $i^i = e^{i^2\frac{\pi}{2}} = e^{-\frac{\pi}{2}}$. So $i^i = e^{-\frac{\pi}{2}}$ and since $e^{-\frac{\pi}{2}} \in \mathbb{R}$ we know that $i^i \in \mathbb{R}$. Fortunately this agrees with the result we find in Section 5. ■

5 Euler's Formula

Another approach to answering this question is to consider Euler's formula evaluated at $x = \frac{\pi}{2}$:

$$\begin{aligned}
e^{ix} &= \cos x + i\sin x && \text{\# Euler's formula} \\
&\Rightarrow e^{i\frac{\pi}{2}} = \cos\frac{\pi}{2} + i\sin\frac{\pi}{2} && \text{\# set } x = \frac{\pi}{2} \\
&\Rightarrow e^{i\frac{\pi}{2}} = 0 + i \cdot 1 && \text{\# } \cos\frac{\pi}{2} = 0 \text{ and} \sin\frac{\pi}{2} = 1 \\
&\Rightarrow e^{i\frac{\pi}{2}} = i && \text{\# simplify} \\
&\Rightarrow \left(e^{i\frac{\pi}{2}}\right)^i = i^i && \text{\# raise both sides to } i \\
&\Rightarrow e^{\frac{i^2\pi}{2}} = i^i && \text{\# } (x^m)^n = x^{mn} \\
&\Rightarrow e^{-\frac{\pi}{2}} = i^i && \text{\# } i^2 = -1 \\
&\Rightarrow e^{-\frac{\pi}{2}} \in \mathbb{R} \Rightarrow i^i \in \mathbb{R} && \text{\# } i^i \text{ is a real number}
\end{aligned}$$

It is important to notice that this expression is multivalued, and when we evaluate it at $\frac{\pi}{2}$ we are on the principle branch. However, the same result can be obtained with $x \in \left\{\frac{\pi}{2}, \frac{5\pi}{2}, \frac{9\pi}{2}, \ldots\right\}$. In general $x = \frac{\pi}{2} \cdot (1 + 4n)$ for $n \in \mathbb{N} \cup \{0\}$. ■

6 Ok, $i^i \in \mathbb{R}$, but what does it equal

We saw above that $i^i = e^{-\frac{\pi}{2}}$. So $i^i = e^{-\frac{\pi}{2}} \approx 0.20788$.



7 Conclusions

Acknowledgements

Thanks to Dima Pasechnik (@dimpase@mathstodon.xyz), Andrés E. Caicedo (@AndresCaicedo@mathstodon.xyz), Ben Reiniger (@bmreiniger@mathstodon.xyz) and Dave Neary (@dneary@mastodon.ie) for their many helpful comments.

LaTeX Source

`https://www.overleaf.com/read/cfjttjzdbfnc`

References

[1] David Meyer. A Few Notes on Euler's Formula and Euler's Identity. `https://davidmeyer.github.io/qc/euler.pdf`, 2021. [See `https://davidmeyer.github.io/qc/`].

[2] Proof Wiki. Corollary to Euler's Formula. `https://proofwiki.org/wiki/Euler%27s_Formula/Corollary`, September 2018. [Online; accessed 17-November-2022].

[3] Proof Wiki. Complex Conjugation is Automorphism. `https://proofwiki.org/wiki/Complex_Conjugation_is_Automorphism`, September 2020. [Online; accessed 15-November-2022].

[4] Proof Wiki. Product of Complex Conjugates. `https://proofwiki.org/wiki/Product_of_Complex_Conjugates`, September 2020. [Online; accessed 17-November-2022].

[5] Proof Wiki. Sum of Complex Conjugates. `https://proofwiki.org/wiki/Sum_of_Complex_Conjugates`, September 2020. [Online; accessed 17-November-2022].

[6] shsu.edu. Properties of Logarithms. `https://www.shsu.edu/kws006/Precalculus/3.3_Logarithms_files/S&Z%206.2%20&%206.3.pdf`. [Online; accessed 17-November-2022].

[7] Wikipedia Contributors. Euler's Formula — Wikipedia, The Free Encyclopedia. `https://en.wikipedia.org/w/index.php?title=Euler%27s_formula&oldid=866429907`, 2018. [Online; accessed 11-November-2018].

[8] Wikipedia Contributors. Principal Branch — Wikipedia, The Free Encyclopedia. `https://en.wikipedia.org/w/index.php?title=Principal_branch&oldid=1009085215`, 2021. [Online; accessed 15-November-2022].

[9] Wikipedia Contributors. Taylor Series — Wikipedia, The Free Encyclopedia. `https://en.wikipedia.org/w/index.php?title=Taylor_series&oldid=1033135703`, 2021. [Online; accessed 6-August-2021].