A Few Notes on the Brachistochrone Problem

David Meyer
dmm@1-4-5.net

Last update: September 27, 2021

1 Introduction

Johann Bernoulli posed the "problem of the brachistochrone" to the readers of *Acta Eruditorum* in June, 1696 [1], which asks the following question:

> Given two points A and B in a vertical plane, what is the curve traced out by a point acted on only by gravity, which starts at A and reaches B in the shortest time?

The scenario Bernoulli envisioned is depicted in Figure 1.



Figure 1: The Brachistochrone Problem Setup

2 A Bit of History

The problem posed by Johann Bernoulli was the called the Brachistochrone[1] problem, or the path of fastest descent. Bernoulli issued his challenge in *Acta Eruditorum* [1], writing

> I, Johann Bernoulli, address the most brilliant mathematicians in the world. Nothing is more attractive to intelligent people than an honest, challenging problem, whose possible solution will bestow fame and remain as a lasting monument. Following the example set by Pascal, Fermat, etc., I hope to gain the gratitude of the whole scientific community by placing before the finest mathematicians of our time a problem which will test their methods and the strength of their intellect. If someone communicates to me the solution of the proposed problem, I shall publicly declare him worthy of praise.

[1] The word brachistochrone comes from the Ancient Greek: *brachis* (shortest) + *chronos* (time) [7].

Bernoulli wrote the problem statement as:

> Given two points A and B in a vertical plane, what is the curve traced out by a point acted on only by gravity, which starts at A and reaches B in the shortest time.

Figure 1 attempts to capture Bernoulli's problem statement.

Galileo had attempted to solve this problem in his *Two New Sciences* [6] and had concluded, based on geometric arguments, that the solution was a circular path. Galileo's approach is shown in Figure 2. Apparently Galileo hedged a bit on his solution. In fact, it is said that he had reservations about his conclusion and suggested that a "higher mathematics" would possibly find a better solution.



Figure 2: Galileo considered a mass falling along different chords of a circle starting at A. He proved that the path along ABG was quicker than along AG, and ABCG was quicker than ABG, and ABCDG was quicker than ABCG, etc. In this way he showed that the path along the circular arc was quicker than any set of chords. From this he (incorrectly) inferred that the circle was the path of quickest descent. Galileo held out reservations about his solution, and rightfully so.

Then in 1659, while studying the physics of pendulums and time keeping, Christiaan Huygens [3] recognized that a perfect harmonic oscillator, one whose restoring force was linear in the displacement of the oscillator, would produce the perfect time piece. Unfortunately, while the pendulum (as proposed by Galileo) was the simplest oscillator to construct, Huygens knew that it was not a perfect harmonic oscillator. In particular, he knew that the period of oscillation became smaller when the amplitude of the oscillation became larger. In order to "fix" the pendulum, he searched for a curve of equal time, called the *tautochrone* [5], that would allow all amplitudes of the pendulum to have the same period. He found the solution and recognized it to be a cycloid curve. A generic cycloid curve is shown in Figure 3.



Figure 3: The Cycloid Curve

Almost thirty years later, during the infancy of the infinitesimal calculus, the Tautochrone problem was held up as a master example of an "optimal" solution whose derivation should yield to the much more powerful and elegant methods of the calculus. Jakob Bernoulli, Johann's brother, succeeded in deriving the Tautochrone problem in 1690 using the calculus, using the term "integral" for the first time in print, but it was not at first clear what other problems could yield in a similar way.

Then, in 1696, Johann Bernoulli posed the Brachistrochrone problem in the pages of *Acta Eruditorum*.

2.1 A (Very) Brief Timeline of the Brachistrochrone Problem

- 1638 – Galileo proposes arc of a circle as the least time solution (Figure 2)
- 1658 – Pascal puts forth a challenge involving finding the area under a segment of a cycloid
- 1659 – Huygens shows experimentally that the cycloid is the solution to the Tautochrone problem
- 1662 – Fermat's Principle [4] (Principle of Least Time)
- 1696 – Johann Bernoulli poses the Brachistrochrone problem in *Acta Eruditorum* (edited by Leibniz)
- 1697 – Solutions provided by the Bernoulli brothers, L'Hôpital, Leibniz, and Newton (anonymously)
- 1707 – Euler is born
- 1726 – Newton dies
- 1726 – Euler completes his Ph.D under Johann Bernoulli
- 1755 – Lagrange, at age 19, provides an analytic solution to the Tautochrone problem
- 1755 – Euler drops his own (partly geometric) approach in favor of Lagrange's purely analytic approach
- 1756 – Euler renames the subject the "Calculus of Variations" in his lecture *Elementa Calculi Variationum* [2]

3 The Calculus of Variations and the Brachistrochrone Problem

4 Conclusions

Acknowledgements

LaTeX Source

`https://www.overleaf.com/read/vpwbqqpnjywj`

References

[1] Gottfried Wilhelm Leibniz, editor. *Acta eruditorum*. Acta eruditorum. Christoph Günther, 1696.

[2] The Open University. Introduction to the Calculus of Variations. `https://www.open.edu/openlearn/ocw/pluginfile.php/1118521/mod_resource/content/3/Introduction%20to%20the%20calculus%20of%20variations_ms327.pdf`, 2021. [Online; accessed 8-September-2021].

[3] Wikipedia contributors. Christiaan Huygens — Wikipedia, The Free Encyclopedia. `https://en.wikipedia.org/w/index.php?title=Christiaan_Huygens&oldid=1046113855`, 2021. [Online; accessed 27-September-2021].

[4] Wikipedia contributors. Fermat's Principle — Wikipedia, The Free Encyclopedia. `https://en.wikipedia.org/w/index.php?title=Fermat%27s_principle&oldid=1045474088`, 2021. [Online; accessed 27-September-2021].

[5] Wikipedia contributors. Tautochrone Curve — Wikipedia, The Free Encyclopedia. `https://en.wikipedia.org/w/index.php?title=Tautochrone_curve&oldid=1032330850`, 2021. [Online; accessed 27-September-2021].

[6] Wikipedia contributors. Two New Sciences — Wikipedia, The Free Encyclopedia. `https://en.wikipedia.org/w/index.php?title=Two_New_Sciences&oldid=1040947163`, 2021. [Online; accessed 27-September-2021].

[7] Wikisource. 1911 Encyclopædia Britannica/Brachistochrone — Wikisource, 2016. [Online; accessed 27-September-2021].