

# ECE 345 / ME 380: Introduction to Control Systems

## Collaborative Quiz #1 Grading Sheet

Dr. Oishi

September 10, 2020

This quiz is open-note and open-book. Calculators and Matlab are allowed. No partial credit will be awarded. For each of the questions, clearly write the correct answer.

### In-Class Questions

1. (d)  $A_P = \begin{bmatrix} 0 & 1 \\ -\frac{mgl}{I} & 0 \end{bmatrix}$ ,  $B_P = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ ,  $C_P = \begin{bmatrix} -\frac{1}{I} & 0 \end{bmatrix}$ ,  $D_P = 0$   
\_\_\_\_\_
2. (a)  $s^2 + mgl/I$   
\_\_\_\_\_
3. (b)  $G(s) = H(s)$ , because a transfer function can be represented by many different state-space models. please see next page for equations  
\_\_\_\_\_
4. (d) The input indirectly affects the generated torque  $\tau(t)$  and the flywheel angular velocity  $\dot{\theta}(t)$ .  
\_\_\_\_\_
5. We would require a lower inertia to allow the flywheel to spin up at higher acceleration,  $z(t_{\text{hop}})$  is inversely proportional to the flywheel inertia.  
\_\_\_\_\_  
\_\_\_\_\_

### Statement of Effort

By providing my name below, I pledge that I have written this quiz as per the indicated instructions, and fully participated in the group.

Nathan Burt	burtn@unm.edu
Name	Email @unm.edu
Noah Jackson	njackson18@unm.edu
Name	Email @unm.edu
David Kirby	davidkirby@unm.edu
Name	Email @unm.edu

$$I s^2 \theta(s) = -mgl \theta(s) - \tau(s)$$

$$I s^2 \theta(s) + mgl \theta(s) = -\tau(s)$$

$$\theta(s) [I s^2 + mgl] = -\tau(s)$$

$$G(s) = \frac{\theta(s)}{\tau(s)} = -\frac{1}{I s^2 + mgl}$$

$$= -\frac{\frac{1}{I}}{s^2 + \frac{mgl}{I}}$$

$$H(s) = C(sI - A)^{-1}B + D$$

$$= \begin{bmatrix} -\frac{1}{I} & 0 \end{bmatrix} \begin{bmatrix} s & -1 \\ \frac{mgl}{I} & s \end{bmatrix}^{-1} \begin{bmatrix} 0 \\ 1 \end{bmatrix} + \underset{0}{0}$$

$$= \begin{bmatrix} -\frac{1}{I} & 0 \end{bmatrix} \frac{1}{s^2 + \frac{mgl}{I}} \begin{bmatrix} s & 1 \\ -\frac{mgl}{I} & s \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} -\frac{1}{I} & 0 \end{bmatrix} \frac{1}{s^2 + \frac{mgl}{I}} \begin{bmatrix} (s)(0) + (1)(1) \\ (-\frac{mgl}{I})(0) + s \end{bmatrix}$$

$$= \begin{bmatrix} -\frac{1}{I} & 0 \end{bmatrix} \frac{1}{s^2 + \frac{mgl}{I}} \begin{bmatrix} 1 \\ s \end{bmatrix} = -\frac{1}{I} \left( \frac{1}{s^2 + \frac{mgl}{I}} \right)$$

$$G(s) = H(s)$$