



Course Home Module 3 - Support vector machines for classification

Take Test: Quiz 3. Support vector classification



Take Test: Quiz 3. Support vector classification

Test Information

Description

Instructions

Multiple Attempts This test allows multiple attempts.

Force Completion This test can be saved and resumed later.

QUESTION 1

0.014 points

Saved

The SVM criterion takes into account the structural risk and the empirical risk. The optimization is intended to minimize both at the same time.

☒ True

☐ False

QUESTION 2

0.014 points

Saved

The SVM optimization is performed through a Lagrange analysis. It leads to

☒ a dual functional that depends on the corresponding Lagrange multipliers. The solution is always guaranteed but it is not unique.

☐ The solution of the dual is always guaranteed because the matrix of the functional is non negative definite.

☐ The solution is guaranteed because the functional is the sum of a convex and a concave functions.

☐ The solution is never guaranteed.

▼ Question Completion Status:

In a SVM classifier, the weight vector is a linear combination of the training data.

- ☒ True
- ☐ False

QUESTION 4

0.014 points

Saved

In a Support Vector Machine, the solution depends on a subset of the training data.

- ☒ True
- ☐ False

QUESTION 5

0.015 points

Saved

The KKT conditions

- ☒ are obtained from nulling the gradient of the lagrange functional with respect to the primal variables (except the complementary ones).
- ☐ The KKT conditions are obtained by nulling the gradient of the Lagrange functional wrt the primal and dual variables (except the complementary ones).
- ☒ The complementary KKT conditions say that either the constraints or the corresponding Lagrange multiplier is zero.
- ☒ The KKT assure that the constraints vanish from the Lagrangian at the optimal point.

QUESTION 6

0.015 points

Saved

The dual variables or Lagrange multipliers

- ☒ of the samples that are properly classified and outside the margin are zero.
- ☐ of the samples that are properly classified and inside the margin have a value strictly less than C.
- ☒ of the samples that are inside the margin have a value equal to C.
- ☒ of the samples that are misclassified and outside the margin are C

margin?

QUESTION 7**0.014 points****Saved**

All the dual variables are either 0 or C



True



False

Click Save and Submit to save and submit. Click Save All Answers to save all answers.

Save All Answers

Save and Submit