

Causal Statistical Decision Theory|What are interventions?

David Johnston

October 20, 2020


Contents

1	Theories of causal inference	2
2	Causal Questions	5
2.1	Formalising Causal Statistical Decision Problems: See-Do Maps .	6
2.2	D-causation	8
2.3	D-causation vs Heckerman and Shachter	9
2.4	Properties of D-causation	12
3	Notation	13
4	What is the difference between Causal Bayesian Networks and See-Do models?	14
4.1	Influence Diagrams vs See-Do models	14
4.2	Influence diagrams vs Causal Bayesian Networks	17
4.3	What is meant by “variables”?	18
4.4	Necessary relationships	18
4.5	Recursive Structural Causal Models	19
4.6	Recursive Structural Causal Models with Necessary Relationships	20
4.7	Cyclic Structural Causal Models	25
4.8	Not all variables have well-defined interventions	27
4.8.1	Necessary relationships in cyclic SCMs	27
5	Definitions and key notation	31
5.1	Standard Symbols	31
5.2	Probability Theory	32
5.3	Product Notation	33
5.4	String Diagrams	33
5.4.1	Comparison of notations	37
5.4.2	Working With String Diagrams	38
5.5	Random Variables	40
5.5.1	Conditional Independence	49

6 Decision makers and decision models 52

6.1 Expected utility maximisers 53

Note on terminology: I am trying the name “See-Do model” to describe the following:



I was calling it a “causal theory” before. Reasons for the change: I think “See-Do” helps to understand what the model does, and the name doesn’t make premature claims to explain causality. Also, it’s only two syllables which I like.

1 Theories of causal inference

Feedback start here

Beginning in the 1930s, a number of associations between cigarette smoking and lung cancer were established: on a population level, lung cancer rates rose rapidly alongside the prevalence of cigarette smoking. Lung cancer patients were far more likely to have a smoking history than demographically similar individuals without cancer and smokers were around 40 times as likely as demographically similar non-smokers to go on to develop lung cancer. In laboratory experiments, cells which were introduced to tobacco smoke developed *ciliastasis*, and mice exposed to cigarette smoke tars developed tumors(Proctor, 2012). Nevertheless, until the late 1950s, substantial controversy persisted over the question of whether the available data was sufficient to establish that smoking cigarettes *caused* lung cancer. Cigarette manufacturers famously argued against any possible connection (Oreskes and Conway, 2011) and Roland Fisher in particular argued that the available data was not enough to establish that smoking actually caused lung cancer (Fisher, 1958). Today, it is widely accepted that cigarettes do cause lung cancer, along with other serious conditions such as vascular disease and chronic respiratory disease (World Health Organisation, 2018; Wiblin, 2016).

The question of a causal link between smoking and cancer is a very important one. Individuals who enjoy smoking (or think they might) may wish to avoid smoking if cigarettes pose a severe health risk, so they are interested in knowing whether or not it is so. Potential investors in cigarette manufacturers want to know if the product they are backing is likely to see limited adoption due to health concerns. People holding investments in cigarette manufacturing firms want the world to be such that cigarettes do not pose a substantial health risk, as this increases the value of their investment. Governments and organisations with a responsibility for public health may see themselves as having responsibility to discourage smoking as much as possible if smoking is severely detrimental to health. The costs and benefits of poor decisions about smoking are large: 8 million annual deaths are attributed to cigarette-caused cancer and vascular disease in 2018(World Health Organisation, 2018) while global cigarette

sales were estimated at US\$711 million in 2020, while (Statista, 2020) (a figure which might be substantially larger if cigarettes were not widely believed to be harmful).

The question of whether or not cigarette smoking causes cancer illustrates two key facts about causal questions: First, having the right answers to some causal questions is of tremendous importance to huge numbers of people. Second, even when large amounts of data show unambiguous associations between phenomena of interest, it is still difficult to know when a causal conclusion is justified.

Causal conclusions are often justified on the basis of ad-hoc reasoning. For example Krittanawong et al. (2020) states:

[...] the potential benefit of increased chocolate consumption, reducing coronary artery disease (CAD) risk is not known. We aimed to explore the association between chocolate consumption and CAD.

It is not clear whether Krittanawong et. al. mean that a negative association between chocolate consumption and CAD implies that increased chocolate consumption is likely to reduce coronary artery disease, or that an association may be relevant to the question and the reader should draw their own conclusions. Whether the implication is being suggested by Krittanawong et. al. or merely imputed by naïve readers, it is being drawn on an ad-hoc basis – no argument for the implication can be found in this paper. As Pearl (2009) has forcefully argued, additional assumptions are always required to answer causal questions from associational facts, and stating these assumptions explicitly allows those assumptions to be productively scrutinised.

Theories of causal inference exist to enable formal rather than ad-hoc reasoning about causal questions. Instead of posing informal causal question and answering them based on ad-hoc reasoning, within a theory of causal inference we ask about properties of “causal models” (which are simply mathematical types defined by the theory) subject to certain assumptions we are willing to make. A successful theory of causal inference should enable causal models that “adequately represent” the original informal question, and the assumptions we invoke should be more accessible to scrutiny than ad-hoc assertions made in the course of answering the informal question.

As well as defining causal models, which represent *claims about causation*, theories of causal inference also formalise the problem of *inferring the correct causal model* - this is the problem of taking some observational data and concluding which causal models are “possible” or “appropriate to use for the given purpose”.

Defining causal models is difficult. While philosophical theories of causation are not heavily discussed in the applied literature on causal inference, the principles used to motivate the definitions of causal models are widely discussed in the philosophical literature. Applied theories of causal inference:

1. “ X_i causes X_j ” means that there exist different *ideal interventions* that result in different values of X_i , hold other “causally sufficient” variables

constant, do not directly affect X_j but nonetheless entail different values of X_j

2. “ X_i causes X_j ” means that the *counterfactual value* of X_j would be different “if X_i had taken a different value”

In practice, most theories of causal inference seem to be based on the notion of *ideal interventions*. Even “counterfactual” theories of causal inference (such as the theory based on “potential outcomes” notation) tend to define counterfactual values as “values that a variable would have taken were it exposed to an ideal intervention”, if they are defined at all (Morgan and Winship, 2014; Rubin, 2005; Richardson and Robins, 2013). There are, however, alternative definitions of counterfactual values such as Lewis’ closest world semantics (Lewis, 1986), though these definitions have serious difficulties.

“Ideal interventions” are difficult to define. The structural model approach of Pearl (2009) defines ideal interventions in terms of “causally sufficient models”. However, this definition ends up circular:

- An $[X_i, X_j]$ -ideal intervention is an operation whose result is determined by applying the do-calculus to a causally sufficient triple $((\Omega, \mathcal{F}, \mathbb{Q}), \mathcal{G}, \mathbf{U})$
- A triple $((\Omega, \mathcal{F}, \mathbb{Q}), \mathcal{G}, \mathbf{U})$ is $[X_i, X_j]$ -causally sufficient if \mathbf{U} contains X_i , X_j and “all intervenable variables” that *cause* (definition (1)) both X_i and X_j

Circularity is a recognised problem with interventional definitions of causation (Woodward, 2016). An alternative approach is to designate certain real-world events – such as flipping coins, querying random number generators and so forth – as prototypical “ideal interventions”. The main problem with this definition is that it fails to answer any causal query that hasn’t been subject to a randomised trial, which seems to leave most causal questions unanswerable, despite the fact that many of them have apparent answers (Pearl, 2018a), and the causal questions represented under this definition are often considered to be inadequate representations of the original questions to which answers were sought (Deaton and Cartwright, 2018; Heckman, 1991). Additional challenges posed by “ideal interventions” are discussed in Section 4.

An account of ideal interventions has not yet been offered that stands up to scrutiny. This difficulty is not without precedent, and is certainly no reason to abandon theories built upon ideal interventions. It is also difficult to provide an account of what it means for data to be “distributed according to probability distribution \mathbb{P} ” (Hájek, 2019) that stands up under scrutiny, but the usefulness of probability distributions in representing data regularities is widely accepted.

However, in light of the difficulties with theories of causation, it is notable that many causal questions can be formalised without appealing to a theory of causation at all, an observation also made by Dawid (2020). Causal statistical decision theory (CSDT), introduced here, shows how the type definition

¹Weaker conditions for causal sufficiency are possible, but they are still premised on causal relationships, so the charge of circularity stands (Shpitser and Pearl, 2008).

of “causal models” is almost completely determined by the given *causal decision problem*. Causal decision problems are problems such as: Should I smoke? Should cigarettes be taxed? Such problems are ubiquitous, and causal inferences are instrumental in answering them. As with existing theories of causal inference, determining the relation between observational probabilities and causal models is the key problem in CSDT, and existing theories of causal inference have developed many useful approaches to this. CSDT additionally offers the opportunity to understand what remains true of causal decision problems regardless of the theory of causation in use and the opportunity to consider theories of causation which may be useful in solving particular causal decision problems but need not be universally valid.

2 Causal Questions

Pearl and Mackenzie (2018) has proposed three types of causal question:

1. Association: How are X and Y related? How would observing X change my beliefs about Y ?
2. Intervention: What would happen if I do ... ? How can I make E happen?
3. Counterfactual: Was X responsible for Y ? What if I had done ... instead of what I actually did?

I focus on a particular type of question here: “How can I make E happen?”. Pearl calls this type of question an “interventional” question but we observe that this query is in fact a *causal decision problem* (I also wish to avoid the term “intervention” as it has a technical meaning which that I touched on above). A causal decision problem is a problem of the following form:

Given my available options D and the way I would like the world to turn out, which options are likely to have a desirable result?

When one asks “How can I make E happen?”, I suppose that the answer will be one or more elements of a set of available options D . Observing that E corresponds to “the way I would like the world to turn out”, we can see that this question is a causal decision problem.

I choose to focus on this type of problem because the problem of choosing a decision given desired results is ubiquitous, and Definition 2 is a broadly applicable prototype for this kind of problem. It’s possible that a similar approach can succeed for counterfactual queries, but obviously a different prototype problem would be required. Enough questions are raised by focusing on causal decision problems to restrict our attention to this type of problem for now.

Causal *statistical* causal decision problems extend causal decision problems by introducing data and assumptions about how the world works:

Given my available options D , data \mathbf{X} and preferences u , which options are likely to have a desirable result?

2.1 Formalising Causal Statistical Decision Problems: See-Do Maps

I introduce *see-do maps* which formalise causal statistical decision problems introduced in Definition 2. I don't derive see-do maps from first principles - rather, I make use of probability theory and expected utility theory because they're the standard tools used to formalise models of noisy observations and choice under uncertainty.

Section 6 (currently unfinished) considers more fundamental assumptions that might lead to see-do maps

First, we formally define *causal statistical decision problems*.

Definition 2.1 (Causal Statistical Decision Problem). A *causal statistical decision problem* (CSDP) is a tuple (D, x, X, Y, u) where D is a set of *decisions* that may be taken, $x \in X$ is a *data sequence* of observations, Y is the space of possible *consequences* of decisions and $u : Y \rightarrow \mathbb{R}$ is a *utility function* where $u(y) \geq u(y')$ implies that y is at least as desirable as y' .

We place a number of additional requirements on causal statistical decision problems:

- D is a denumerable set
- The principle of *expected utility maximisation* is acceptable, so the desirability of a probability distribution $\mathbb{P}_Y \in \Delta(\mathcal{Y})$ can be found by evaluating $\mathbb{E}_{\mathbb{P}_Y}[u]$
- X and Y are *standard measurable* (D is automatically standard measurable due to it being denumerable)
- A decision maker may select any probability distribution in $\Delta(\mathcal{D})$ as its stochastic decision. A deterministic distribution δ_d is equivalent to directly choosing $d \in D$, while deciding on a non-deterministic distribution involves consulting a source of randomness to select a particular decision in accordance with the probabilities chosen

Definition 2.1 are the elements of a CSDP that we regard as fixed.

A *decision maker* brings prior knowledge to the problem in the form of a *see-do map*, which captures the relation between the *observed probability distribution* and the *consequence map*.

Definition 2.2 (Observed probability distribution). Given a CSDP (D, x, X, Y, u) , under hypothesis θ the data x are “distributed according to” some *observed probability* $\mathbb{O}_\theta \in \Delta(\mathcal{X})$.

Definition 2.3 (Consequence map). Given a decision set D and a consequence set Y , under hypothesis θ a *consequence map* $\mathbb{C}_\theta : D \rightarrow \Delta(\mathcal{Y})$ is a Markov kernel mapping decisions to probability distributions on the consequence space. For a stochastic decision $\gamma \in \Delta(\mathcal{D})$, $\gamma\mathbb{C}$ is the *consequence* of γ with respect to \mathbb{C}_θ .

“Distributed according to” is a weasel phrase with no particularly satisfactory interpretation, but I'd like to avoid worrying about it at this point (Hájek, 2019).

⓪.. However, in general we don't require that ⓪ and ℂ are related in this manner. This assumption will be revisited in

A section I haven't written yet

2.2 D-causation

While we take D to be a primitive element of causal decision problems, and therefore a primitive of see-do maps. Causes are not primitive, but we can offer a secondary notion of causation. We call this D -causation to stress the fact that it arises in a theory of causal inference in which the set D of available decisions is primitive. A similar idea is discussed extensively in Heckerman and Shachter (1995). The main differences are that what we call “consequence maps” map decisions to probability distributions over possible consequences while Heckerman and Shachter work with “states” that map decisions deterministically to consequences. In addition, while we define D -causation relative to a particular consequence map \mathbb{C}_θ , Heckerman and Shachter define it with respect to a *set* of states.

Section 4 explores the difficulty of defining “objective causation” without reference to a set of basic decisions, acts or operations. D need not be interpreted as the set of decisions an agent may make, but whatever interpretation it is assigned, all existing examples of causal models seem to require a “domain set”.

See Section 5.5 for the definition of random variables.

Add definition of conditional independence, revise wire label definitions

One way to motivate the notion of D -causation is to observe that for many decision problems, the full set D may be extremely large. Suppose I aim to have my light switched on, and there is a switch that controls the light. Often, the relevant choice of acts for such a problem would appear to be $D_0 = \{\text{flip the switch, don't flip the switch}\}$. However, in principle I have a much larger range of options to choose from. For simplicity's sake, suppose I have instead the following set of options:

$D_1 := \{\text{“walk to the switch and press it with my thumb”},$
 $\text{“trip over the lego on the floor, hop to the light switch and stab my finger at it”},$
 $\text{“stay in bed”}\}$

If having the light turned on is all that matters, I could consider any acts in D_1 to be equivalent if they have the same ultimate impact on the position of the light switch. D_0 is a quotient over D_1 under this equivalence relation.

If I hypothesize that, relative to D_1 , the ultimate state of the light switch is all that matters to determine the ultimate state of the light, I can say that the light switch D_1 -causes the state of the light. Given this D_1 -causation, the D_1 decision problem can (subject to my hypothesis) be reduced to a D_0 decision between states of the light switch.

If I consider an even larger set of possible acts D_2 , I might not accept the hypothesis of D_2 -causation. Let D_2 be the following acts:

$D_2 := \{$ “walk to the switch and press it with my thumb”,
 “trip over the lego on the floor, hop to the light switch and stab my finger at it”,
 “stay in bed”,

“toggle the mains p

In this case, it would be unreasonable to hypothesize that all acts that left the light switch in the “on” position would also result in the light being “on”. Thus the switch does not D_2 -cause the light to be on.

Formally, D -causation is defined in terms of conditional independence:

Definition 2.7 (D -causation). Given a consequence map $\mathbb{C}_\theta : D \rightarrow \Delta(\mathcal{Y})$, random variables $Y_1 : Y \times D \rightarrow Y_1$, $Y_2 : Y \times D \rightarrow Y_2$ and domain variable $D : Y \times D \rightarrow D$ (Definition 5.8), Y_1 D -causes Y_2 iff $Y_2 \perp\!\!\!\perp_{\mathbb{C}_\theta} D | Y_1$.

2.3 D-causation vs Heckerman and Shachter

Heckerman and Shachter study deterministic “consequence maps”. Furthermore, what we call hypotheses $\theta \in \Theta$, Heckerman and Schachter call states $s \in S$. One could consider a state to be a hypothesis that is specific enough to yield a deterministic map from decisions to outcomes. Heckerman and Shachter’s notion of causation is defined by *limited unresponsiveness* rather than *conditional independence*, which depends on a partition of states rather than a particular hypothesis.

Definition 2.8 (Limited unresponsiveness). Given states S , deterministic consequence maps $\mathbb{C}_s : D \rightarrow \Delta(F)$ for each $s \in A$ and a random variables $X : F \rightarrow X$, $Y : F \rightarrow Y$, Y is unresponsive to D in states limited by X if $\mathbb{C}_{(s,d)}^{X|D} = \mathbb{C}_{(s,d')}^{X|DS} \implies \mathbb{C}_{(s,d)}^{Y|DS} = \mathbb{C}_{(s,d')}^{Y|DS}$ for all $d, d' \in D$, $s \in S$. Write $Y \not\prec_X D$

Lemma 2.9 (Limited unresponsiveness implies D -causation). *For deterministic consequence maps, $Y \not\prec_X D$ implies X D -causes Y in every state $s \in S$.*

Proof. By the assumption of determinism, for each $s \in S$ and $d \in D$ there exists $x(s, d)$ and $y(s, d)$ such that $\mathbb{C}_{d,s}^{XY|DS} = \delta_{x(s,d)} \otimes \delta_{y(s,d)}$.

By the assumption of limited unresponsiveness, for all d, d' such that $x(s, d) = x(s, d')$, $y(s, d) = y(s, d')$ also. Define $f : X \times S \rightarrow Y$ by $(s, x) \mapsto y(s, [x(s, \cdot)]^{-1}(x(s, d)))$ where $[x(s, \cdot)]^{-1}(a)$ is an arbitrary element of $\{d | x(s, d) = a\}$. For all s, d , $f(x(s, d), s) = y(s, d)$. Define $\mathbb{M} : X \times D \times S \rightarrow \Delta(\mathcal{Y})$ by $(x, d, s) \mapsto \delta_{f(x,s)}$. \mathbb{M} is a version of $\mathbb{C}^{Y|X,D,S}$ because, for all $A \in \mathcal{X}$, $B \in \mathcal{Y}$, $s \in S$, $d \in D$:

$$\mathbb{C}_{(d,s)}^{\mathbf{X}|\mathbf{DS}} \Upsilon(\mathbb{M} \otimes \text{Id}) = \int_A \mathbb{M}(x', d, s; B) d\delta_{x(s,d)}(x') \quad (2)$$

$$= \int_A \delta_{f(x',s)}(B) d\delta_{x(s,d)}(x') \quad (3)$$

$$= \delta_{f(x(s,d),s)}(B) \delta_{x(s,d)}(A) \quad (4)$$

$$= \delta_{y(s,d)}(B) \delta_{x(s,d)}(A) \quad (5)$$

$$= \delta_{x(s,d)} \otimes \delta_{y(s,d)}(A \times B) \quad (6)$$

\mathbb{M} is also independent of \mathbf{D} , given the obvious labeling of inputs. Therefore $\mathbf{Y} \perp\!\!\!\perp_{\mathbb{C}_s} \mathbf{D} | \mathbf{X}$. \square

However, despite limited unresponsiveness implying D -causation within every state, it does not imply D -causation in mixtures of states. Suppose $D = \{0, 1\}$ where 1 stands for “toggle light switch” and 0 stands for “do nothing”. Suppose $S = \{[0, 0], [0, 1], [1, 0], [1, 1]\}$ where $[0, 0]$ represents “switch initially off, mains off” the other states generalise this in the obvious way. Finally, $\mathbf{F} \in \{0, 1\}$ is the final position of the switch and $\mathbf{L} \in \{0, 1\}$ is the final state of the light. We have

$$\mathbb{C}_{d,[i,m]}^{\mathbf{LF}|\mathbf{DS}} = \delta_{(d \text{ XOR } i) \text{ AND } m} \otimes \delta_{(d \text{ XOR } i) \text{ AND } m} \quad (7)$$

Within states $[0, 0]$ and $[1, 0]$, the light is always off, so $\mathbf{F} = a \implies \mathbf{L} = 0$ for any a . In states $[0, 1]$ and $[1, 1]$, $\mathbf{F} = 1 \implies \mathbf{L} = 1$ and $\mathbf{F} = 0 \implies \mathbf{L} = 0$. Thus $\mathbf{L} \not\prec_{\mathbf{F}} \mathbf{D}$. However, suppose we take a mixture of consequence maps:

$$\mathbb{C}_\gamma = \frac{1}{4}\mathbb{C}_{\cdot,[0,0]} + \frac{1}{4}\mathbb{C}_{\cdot,[0,1]} + \frac{1}{2}\mathbb{C}_{\cdot,[1,1]} \quad (8)$$

$$\mathbb{C}_\gamma^{\mathbf{FL}|\mathbf{D}} = \frac{1}{4} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \otimes \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} + \frac{1}{4} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \otimes \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \otimes \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad (9)$$

Then

$$[1, 0]\mathbb{C}_\gamma^{\mathbf{FL}|\mathbf{D}} = \frac{1}{4}[0, 1] \otimes [1, 0] + \frac{1}{4}[0, 1] \otimes [0, 1] + \frac{1}{2}[1, 0] \otimes [1, 0] \quad (10)$$

$$[1, 0]\Upsilon(\mathbb{C}_\gamma^{\mathbf{F}|\mathbf{D}} \otimes \mathbb{C}_\gamma^{\mathbf{L}|\mathbf{D}}) = (\frac{1}{2}[0, 1] + \frac{1}{2}[1, 0]) \otimes (\frac{1}{4}[0, 1] + \frac{3}{4}[1, 0]) \quad (11)$$

$$\implies [1, 0]\mathbb{C}_\gamma^{\mathbf{FL}|\mathbf{D}} \neq [1, 0]\Upsilon(\mathbb{C}_\gamma^{\mathbf{F}|\mathbf{D}} \otimes \mathbb{C}_\gamma^{\mathbf{L}|\mathbf{D}}) \quad (12)$$

Thus under hypothesis mixture γ , \mathbf{F} does not D -cause \mathbf{L} even though \mathbf{F} D -causes \mathbf{L} in all states S . The definition of D -causation was motivated by the idea that we could reduce a difficult decision problem with a large set D to a simpler problem with a smaller “effective” set of decisions by exploiting conditional independence. Even if \mathbf{X} D -causes \mathbf{Y} in every $\theta \in S$, \mathbf{X} does not

necessarily D -cause Y in mixtures of states in S . For this reason, we do not say that X D -causes Y in S if X D -causes Y in every $\theta \in S$, and in this way we differ substantially from Heckerman and Shachter (1995).

Instead, we simply extend the definition of D -causation to mixtures of hypotheses: if $\gamma \in \Delta(\Theta)$ is a mixture of hypotheses, define $\mathbb{C}_\gamma := (\gamma \otimes \mathbf{Id})\mathbb{C}$. Then X D -causes Y relative to γ iff $Y \perp\!\!\!\perp_{\mathbb{C}_\gamma} D|X$.

Theorem 2.10 shows that under some conditions, D -causation can hold for arbitrary mixtures over subsets of the hypothesis class Θ .

Theorem 2.10 (Universal D -causation). *If $\mathbb{C}_\theta^{X|D} = \mathbb{C}_{\theta'}^{X|D}$ for all $\theta, \theta' \in S \subset \Theta$ and X D -causes Y in all $\theta \in S$, then X D -causes Y with respect to all mixed consequence maps \mathbb{C}_γ for all $\gamma \in \Delta(\Theta)$ with $\gamma(S) = 1$.*

Proof. For $\gamma \in \Delta(\Theta)$, define the mixture

$$\mathbb{C}_\gamma := \begin{array}{c} \triangle \gamma \\ \swarrow \\ D \end{array} \begin{array}{c} \boxed{\mathbb{C}} \\ \longrightarrow F \end{array} \quad (13)$$

Because $\mathbb{C}_\theta^{X|D} = \mathbb{C}_{\theta'}^{X|D}$ for all $\theta, \theta' \in \Theta$, we have

$$\begin{array}{c} \triangle \gamma \\ \swarrow \\ D \end{array} \begin{array}{c} \boxed{\mathbb{C}^{X|D\Theta}} \\ \longrightarrow X \end{array} \begin{array}{c} \longrightarrow \Theta \end{array} = \begin{array}{c} \triangle \gamma \\ \swarrow \\ D \end{array} \begin{array}{c} \boxed{\mathbb{C}^{X|D\Theta}} \\ \longrightarrow X \end{array} \begin{array}{c} \longrightarrow \Theta \end{array} \quad (14)$$

Also

$$\mathbb{C}_\gamma^{XY|D} = \begin{array}{c} \triangleleft \gamma \\ D \text{---} \boxed{C} \text{---} \boxed{F^{X \otimes Y}} \text{---} \begin{array}{c} X \\ Y \end{array} \end{array} \quad (15)$$

$$= \begin{array}{c} \triangleleft \gamma \\ D \text{---} \boxed{C^{XY|D\theta}} \text{---} \begin{array}{c} X \\ Y \end{array} \end{array} \quad (16)$$

$$= \begin{array}{c} \triangleleft \gamma \\ D \text{---} \boxed{C^{X|D\theta}} \text{---} \boxed{C^{Y|XD\theta}} \text{---} \begin{array}{c} Y \\ X \end{array} \end{array} \quad (17)$$

$$\stackrel{Y \perp\!\!\!\perp_{\mathbb{C}_\gamma} D | X \theta}{=} \begin{array}{c} \triangleleft \gamma \\ D \text{---} \boxed{C^{X|D\theta}} \text{---} \boxed{C^{Y|X\theta}} \text{---} \begin{array}{c} Y \\ X \end{array} \end{array} \quad (18)$$

$$\stackrel{14}{=} \begin{array}{c} \triangleleft \gamma \\ D \text{---} \boxed{C^{X|D\theta}} \text{---} \triangleleft \gamma \text{---} \boxed{C^{Y|X\theta}} \text{---} \begin{array}{c} Y \\ X \end{array} \end{array} \quad (19)$$

$$\stackrel{14}{=} \begin{array}{c} \triangleleft \gamma \\ D \text{---} \boxed{C_\gamma^{X|D\theta}} \text{---} \triangleleft \gamma^* \text{---} \boxed{C^{Y|X\theta}} \text{---} \begin{array}{c} Y \\ X \end{array} \end{array} \quad (20)$$

Equation 20 establishes that $(\gamma \otimes \text{Id}_X \otimes *_D)C^{Y|X\theta}$ is a version of $\mathbb{C}_\gamma^{Y|XD}$, and thus $Y \perp\!\!\!\perp_{\mathbb{C}_\gamma} D | X$.

This can also be derived from the semi-graphoid rules:

$$\theta \perp\!\!\!\perp D \wedge \theta \perp\!\!\!\perp X | D \implies \theta \perp\!\!\!\perp XD \quad (21)$$

$$\implies \theta \perp\!\!\!\perp D | X \quad (22)$$

$$D \perp\!\!\!\perp \theta | X \wedge D \perp\!\!\!\perp Y | X \theta \implies D \perp\!\!\!\perp Y | X \quad (23)$$

$$\implies Y \perp\!\!\!\perp D | X \quad (24)$$

□

2.4 Properties of D-causation

If X D-causes Y relative to \mathbb{C}_θ , then the following holds:

$$\mathbb{C}_\theta^{X|D} = D \text{---} \boxed{C^{X|D}} \text{---} \boxed{C^{Y|X}} \text{---} Y \quad (25)$$

This follows from version (2) of Definition 5.28:

$$\mathbb{C}_\theta^{\mathbf{X}|\mathbf{D}} = \mathbf{D} - \boxed{\mathbb{C}^{\mathbf{X}|\mathbf{D}}} - \boxed{\mathbb{C}^{\mathbf{Y}|\mathbf{XD}}} - \mathbf{Y} \quad (26)$$

$$= \mathbf{D} - \boxed{\mathbb{C}^{\mathbf{X}|\mathbf{D}}} - \boxed{\mathbb{C}^{\mathbf{Y}|\mathbf{X}}} - \mathbf{Y} \quad (27)$$

$$= \mathbf{D} - \boxed{\mathbb{C}^{\mathbf{X}|\mathbf{D}}} - \boxed{\mathbb{C}^{\mathbf{Y}|\mathbf{X}}} - \mathbf{Y} \quad (28)$$

D-causation is not transitive: if \mathbf{X} D-causes \mathbf{Y} and \mathbf{Y} D-causes \mathbf{Z} then \mathbf{X} doesn't necessarily D-cause \mathbf{Z} .

3 Notation

- \mathbf{X} is a random variable, X is its codomain and \mathcal{X} is the σ -algebra on X
- Bold letters \mathbf{X} may be used for product spaces, \mathbf{x} for elements of product spaces, \mathbf{f} for vector valued functions and \mathbf{X} for random variables taking values in product spaces. The absence of bold font does not imply the absence of product space structure
- Nodes in a graph are italic sans serif X
- Given an indexed product space $\prod_{i \in \mathcal{I}} X_i$, $\pi_i : \prod_{i \in \mathcal{I}} X_i \rightarrow X_i$ is the projection map $(x_1, \dots, x_i, \dots, x_n) \mapsto x_i$
- Given a product space $X \times Y$, $\pi_X : X \times Y \rightarrow X$ is the projection map $(x, y) \mapsto x$
- Given an index set \mathcal{I} , $\mathbf{X}_{\mathcal{I}}$ is the indexed product space $\prod_{i \in \mathcal{I}} X_i$
- $[n]$ is the index set $\{1, \dots, n\}$ for $n \in \mathbb{N}$
- Given an indexed product space $\mathbf{X}_{[n]}$, $\mathbf{X}_{< j}$ is the set $\prod_{i=1}^j X_i$
- $\underline{\otimes}$ is the coupled tensor product, see Definition 5.9
- Given a random variable \mathbf{X} , $F_{\mathbf{X}}$ is the associated Markov kernel, see Definition 5.1
- See section 5.4.2 for rules of string diagram manipulation
- $*_X : X \rightarrow \Delta(\{*\})$ is the discard map defined in Equation 122

4 What is the difference between Causal Bayesian Networks and See-Do models?

See-Do models and Causal Bayesian Networks (and related models such as SCMs) are quite different in their appearance and in the interpretation of various elements. In defining See-Do models, we assume that there is a decision problem that fixes in advance the observation space E , the space of consequences F and the set of available decisions D . By including D , See-Do models have an agent “baked in” to the definition. In contrast, Causal Bayesian Networks assume that a set of observed variables \mathbf{X} and a set of unobserved variables \mathbf{U} is fixed by nature. A subtlety here is that \mathbf{U} is generally *not* known, but it may be assumed that whatever variables actually comprise \mathbf{U} , they may be generically representable by some known set of variables \mathbf{U}' without loss. Because a set of decisions D is absent, Causal Bayesian Networks appear to model agent-independent “causal relationships”.

Despite the fact that Causal Bayesian Networks don’t seem to be built to model the consequences of an agent’s decision making, they are nonetheless considered to be appropriate for this purpose. This is because the “do-operations” that Causal Bayesian Networks support are considered to have some relationship to any set D decisions some agent might want to consider.

The question of how this relationship might be determined in general is one that I have not seen addressed anywhere. Typically, the approach taken is “I know it when I see it”. For example, if I were a doctor and I could either A) hand a patient a prescription or B) not hand the patient the prescription, and if I had a set of observational data of past patients including a variable S representing whether or not they had received a prescription for the drug, I could consider option A to correspond to $do(S = 1)$ and option B to correspond to $do(S = 0)$ in some causal model (perhaps in the “true” causal model). While this might appear to be reasonable, we should be cautious: this is a completely *ad-hoc* assumption.

In fact, given a sufficiently rich set of variables $\mathbf{U} \cup \mathbf{X}$, I argue that $do(S)$ will almost never even approximate the consequences of an action known in advance to fix the value of some variable S . For this reason, it is valuable to have a theory like CSDT that is concerned only with the consequences of actions.

4.1 Influence Diagrams vs See-Do models

Influence diagrams are used to represent causal models in a manner similar to, but not quite the same as, Causal Bayesian Networks. Using the version found in Dawid (2002), an influence diagram is a directed acyclic graph (DAG) with two node types: “chance” nodes and “decision” nodes. For example:



(29)

Is an influence diagram with the decision node D and the chance node F . When nodes are associated with sets representing possible values, influence diagrams represent sets of Markov kernels. For example, if we associate the measurable set (D, \mathcal{D}) with D and (F, \mathcal{F}) with F , then we could take Diagram 29 to represent the set of all Markov kernels $D \rightarrow \Delta(\mathcal{F})$, or if we add some additional assumptions it might represent a particular subset $S \subset \Delta(\mathcal{F})^D$ of these kernels that share certain properties. Compare this to a string diagram:

$$D - \boxed{\mathbb{K}} - F \quad (30)$$

This diagram represents a *particular* Markov kernel $\mathbb{K} : D \rightarrow \Delta(\mathcal{F})$. A set such as that represented by Diagram 29 could be constructed by creating a set Θ and a function $f : \Theta \rightarrow S$ that indexes each element of S with $\theta \in \Theta$. Then $S = \{f(\theta) | \theta \in \Theta\}$. If f is measurable then there is a Markov kernel $\mathbb{T} : \Theta \times D \rightarrow \Delta(\mathcal{F})$ such that $f(\theta) = \mathbb{T}_{\theta, \cdot}$. In this sense, for any additional assumptions that are combined with Diagram 29 to yield the set S of Markov kernels, there exists a single Markov kernel $\mathbb{T} : \Theta \times D \rightarrow \Delta(\mathcal{F})$ that generates S . \mathbb{T} can be drawn as

notation

$$\begin{array}{c} \Theta \\ \diagdown \\ D \end{array} - \boxed{\mathbb{T}} - F \quad (31)$$

Diagram 31 has a few more elements than Diagram 29 - Θ and \mathbb{T} in particular. If I were to define exactly what D , Θ , \mathbb{T} and F were, Diagram 31 would represent a unique Markov kernel. On the other hand, Diagram 29 along with a precise definition of D and F could represent many different sets of Markov kernels, depending on whatever additional properties I want elements of S to share. Loosely, we can say that for any kernel \mathbb{T} , there is a diagram 29 along with additional assumptions that represents “the same thing”.

Influence diagrams *as typically used in causal modelling* usually additional assumptions. For example, Dawid (2002) proposes that D contains a special “do nothing” element $o \in D$ such that the observations in state θ are given by $\mathbb{T}_{\theta, o}$ and consequences in state θ are given by $\mathbb{T}_{\theta, \cdot}$. This corresponds to the See-Do model

Under the condition that the choice of index “doesn’t matter” somehow

notation

$$\begin{array}{c} \Theta \\ \diagdown \\ D \end{array} - \begin{array}{c} \boxed{\mathbb{T}_{\cdot, o}} \\ \boxed{\mathbb{T}} \end{array} - F \quad (32)$$

While this is a feature of influence diagrams in Dawid (2002), *in general* a diagram like 29 can represent an arbitrary set of typed Markov kernels. See-Do models generate Θ -indexed sets of Markov kernels. We can therefore represent generic See-Do models with influence diagrams.

There is some kind of measurability condition needed to make the two representations coincide

Concretely, given the influence diagram

$$\mathcal{J} = \begin{array}{c} \textcircled{E} \\ \textcircled{D} \rightarrow \textcircled{F} \end{array} \quad (33)$$

and any See-Do model $\mathbb{T} : \Theta \times D \rightarrow \Delta(\mathcal{E} \otimes \mathcal{F})$ there exists a set of auxiliary conditions A such that the model (\mathcal{J}, A) is equivalent to \mathbb{T} . To illustrate how influence diagrams and string diagrams compare, \mathbb{T} can be drawn:

subject to irrelevance of index Θ

$$\mathbb{T} := \begin{array}{c} \Theta \text{ --- } \boxed{\mathbb{H}} \text{ --- } F \\ \quad \quad \quad \diagup \quad \diagdown \\ D \text{ --- } \boxed{\mathbb{C}} \text{ --- } E \end{array} \quad (34)$$

Maybe move the following to an appendix?

Definition 4.1 (Markov kernel/influence diagram compatibility). Given a Markov kernel $\mathbb{K} : E \rightarrow \Delta(\mathcal{F})$, an influence diagram $\mathcal{J} = (S, A, E)$ and an injective $f : X \cup A \rightarrow \mathcal{F} \otimes \mathcal{E}$ which assigns each node to exactly one random variable in $\mathcal{F} \otimes \mathcal{E}$, if for all $X_1, X_2 \in \mathbf{X}$ we have $X_1 \perp_{\mathcal{J}} X_2 \implies f(X_1) \perp_{\mathbb{K}} f(X_2)$

still need to *define* conditional independence WRT kernels, though I think I can do so

Consider an arbitrary See-Do model $\mathbb{T} : D \times \Theta \rightarrow \Delta(\mathcal{E}_1 \otimes \mathcal{E}_2)$ and random variables $D := \pi_D, E := \pi_E, F := \pi_F$ on $\mathcal{D} \otimes \mathcal{E} \otimes \mathcal{F}$. For any $\theta \in \Theta$, \mathbb{T}_θ is compatible with the influence diagram $\mathcal{J} = (\{E, F\}, \{D\}, \{D \rightarrow F\})$ with respect to the injective function

$$f : \begin{cases} A \mapsto D \\ E \mapsto E \\ F \mapsto F \end{cases} \quad (35)$$

There is always some $\mathbb{H} : \Theta \rightarrow \Delta(\mathcal{E})$ and $\mathbb{C} : \Theta \times D \rightarrow \Delta(\mathcal{E} \otimes \mathcal{F})$ such that: \mathbb{T}_θ is equal to

$$\mathbb{T}_\theta = \begin{array}{c} \boxed{\mathbb{H}_\theta} \text{ --- } E \\ D \text{ --- } \boxed{\mathbb{C}_\theta} \text{ --- } F \end{array} \quad (36)$$

Which implies $E \perp_{\mathbb{T}_\theta} F$ and $E \perp_{\mathbb{T}_\theta} D$.

The influence diagram



Features the d-separations $E \perp_{\mathcal{J}} F$ and $E \perp_{\mathcal{J}} D$ (Peters et al., 2017; Woodward, 2016; Dawid, 2002). Thus \mathbb{T}_{θ} is compatible with \mathcal{I} for all $\theta \in \Theta$.

4.2 Influence diagrams vs Causal Bayesian Networks

See-Do models can always be represented by influence diagrams with auxiliary assumptions. We can then learn something about how Causal Bayesian Networks compare to See-Do models by asking how they compare to influence diagrams. The key difference between Causal Bayesian Networks and influence diagrams is that the diagrams do not contain decision nodes. Instead of Diagram 29, a Causal Bayesian Network for the same system might be



The difference between Diagram 29 and Diagram 38 is that the former demands a set D to be bound to the decision node D and a set F to be bound to F , while Diagram 38 demands only F . Instead of explicitly representing decisions that can be chosen, Causal Bayesian Networks *by default* feature a set of *do-interventions* on the chance nodes which seem to have a role similar to decisions in influence diagrams and See-Do models (in fact, Pearl (2009) pg 108 suggests that do-interventions and decisions are the same thing). This default set of do-interventions is what allows CBNs to avoid explicitly requiring a set D . If a set of decisions is required that is not equivalent to the set of do-interventions, this can be specified via auxiliary assumptions, although in practice influence diagrams are usually adopted such as in (Yang et al., 2018).

Dealing with a set of decisions D can be troublesome. It can easily be the case that, for example, I might be tasked with inferring a consequence map that someone else might use and I am not privy to the decisions that they might be able to make. In this case, I'd need to pick a set of decisions D which I am pretty sure covers all the possibilities.

Alternatively, the set D might be unworkably large. The set of *all* the decisions you could in principle make at this moment in as much detail as you can - this is clearly something that's far too big to write down and work with in solving an inference problem.

Speculatively, the fact that Causal Bayesian Networks default to do-interventions might help with the problems of unknown or unworkably large decision sets. If any possible decision must be resolvable to do-interventions of some type, and the effects of do-interventions are well defined, then could provide a basis for

some more examples

partially solving decision problems while remaining ignorant of the particular set of decisions that will ultimately be selected from.

as in, doing the inference but not picking the best option

Beyond this, causal effects as they are informally understood *seem* to refer to universal things, not things that depend on the set of decisions one has available. While not an especially strong reason to avoid specifying D in causal models, it is a reason nonetheless.

Do-interventions, however, cannot solve these problems. If there are no restrictions on the variables that may be included in a CBN model, then as we show do-operations and equation surgery frequently produce invalid results. To ensure that any do-interventions at all are well defined, Causal Bayesian Networks require the specification of “intervenable variables” or “basic interventions”, a requirement that is analogous to the requirement of a set of decisions D in See-Do models and influence diagrams. Specifying D may be difficult, but the do-intervention paradigm provides no solutions to this difficulty; it merely sweeps it under the rug.

4.3 What is meant by “variables”?

Not sure where to put it, but Pearl pp 162-163 puts his models on the hook for including arbitrary variables

4.4 Necessary relationships

The relationship between a person’s body mass index, their weight and their height defines what body mass index is. A fundamental claim of ours is that any causal model that defines “the causal effect of body mass index” should do so without reference to any submodel that violates this definitional relationship violation of the definition. This is an important assumption, and it rests on a judgement of what causal models ought to do. I think it is quite clear that when anyone asks for a causal effect, they expect that any operations required to define the causal effect *do not change the definitions of the variables they are employing*. While theories of causality have a role in sharpening our understanding of the term *causal effect*, the thing called a “causal effect” in an SCM should still respect some of our pre-theoretic intuitions about what causal effects are or else it should be called something else. “Causal effects” that depend on redefining variables do not respect pre-theoretic intuitions about what causal effects are:

- If I ask for the “causal effect of a person’s BMI”, I do not imagine that I am asking what would happen if someone’s BMI were defined to be something other than their weight divided by their height
- If I ask for the “causal effect of a person’s weight”, I do not imagine that I am asking what would happen if someone’s weight were not equal to their volume multiplied by their density
- If I ask for the “causal effect of a person’s weight”, I also do not imagine that I am asking what would happen if their weight were not equal to the

weight of fat in their body plus the weight of all non-fat parts of their body

- If I ask for the “causal effect of taking a medicine”, I do not imagine that I am asking what would happen if a person were declared to have taken a medicine independently of whatever substances have actually entered their body and how they entered

We will call relationships that have to hold *necessary relationships*. We provide the example of relationships that have to hold by definition as examples, but definitions may not be the only variety of necessary relationships. For example, one might also wish to stipulate that certain laws of physics are required to hold in all submodels.

If a causal model contains variables that are necessarily related, then an intervention on one of them must always change another variable in the relationship. If I change a person’s weight, their height or BMI must change (or both). If I change their height, their weight or BMI must change and if I change their BMI then their weight or height must change. This conflicts with the usual acyclic definition of causal models, where the proposition that **A** causes **B** rules out the possibility that **B** or any of its descendents are a cause of **A**. Thus in an acyclic model it isn’t possible for for an intervention on BMI to change weight or height and interventions on weight and height to also change BMI. Theroem 4.11 formalises this conflict for recursive structural causal models: for any set of variables that are necessarily related by a cyclic relationship, at least one of them has no hard interventions defined.

4.5 Recursive Structural Causal Models

We begin by showing that necessary relationships are incompatible with structural causal models.

Definition 4.2 (Recursive Structural Causal Model). A recursive structural causal model (SCM) is a tuple

$$\mathcal{M} := \langle N, M, \mathbf{X}_{[N]}, \mathbf{E}_{[M]}, \{f_i | i \in [N]\}, \mathbb{P}_{\mathcal{E}} \rangle \quad (39)$$

where

- $N \in \mathbb{N}$ is the number of *endogenous variables* in the model
- $M \in \mathbb{N}$ is the number of *exogenous variables* in the model
- $\mathbf{X}_{[N]} := \{X_i | i \in [N]\}$ where, for each $i \in [N]$, (X_i, \mathcal{X}_i) is a standard measurable space taking and the codomain of the i -th endogenous variable
- $\mathbf{E}_{[M]} := \{E_j | j \in [M]\}$ where, for $j \in [M]$, E_j is a standard measurable space and the codomain of the j -th exogenous variable
- $f_i : \mathbf{X}_{<i} \times \mathbf{E}_{\mathcal{J}} \rightarrow X_i$ is a measurable function which we call *the causal mechanism controlling the i -th endogenous variable*

- $\mathbb{P}_{\mathcal{E}} \in \Delta(\mathbf{E}_{\mathcal{I}})$ is a probability measure on the space of exogenous variables

Definition 4.3 (Observable kernel). Given an SCM \mathcal{M} with causal mechanisms $\{f_i | i \in [N]\}$, define the *observable kernel* $G_i : E \rightarrow \Delta(\mathbf{X}_{[i]})$ recursively:

$$G_1 = \mathbf{E}_{[M]} \text{---} \boxed{F_{f_1}} \text{---} X_1 \quad f_1 \quad (40)$$

$$G_{n+1} = \mathbf{E}_{[M]} \text{---} \boxed{G_n} \text{---} \mathbf{X}_{<n+1} \quad \boxed{F_{f_{n+1}}} \text{---} X_{n+1} \quad (41)$$

Definition 4.4 (Joint distribution on endogenous variables). The *joint distribution on endogenous variables* defined by \mathcal{M} is $\mathbb{P}_{\mathcal{M}} := \mathbb{P}_{\mathcal{E}} G_N$ (which is the regular kernel product, see Definition 5.3). For each $i \in [N]$ define the random variable $\mathbf{X}_i : \mathbf{X}_{[N]} \rightarrow X_i$ as the projection map $\pi_i : (x_1, \dots, x_i, \dots, x_N) \mapsto x_i$. By Lemma 4.5, $\underline{\otimes}_{i \in [N]} \mathbf{X}_i = \text{Id}_{\mathbf{X}_{[N]}}$, and so $\mathbb{P}_{\mathcal{M}}$ is the joint distribution of the variables $\{\mathbf{X}_i | i \in [N]\}$.

I use the notation $\mathbb{P}_{\mathcal{M}}$ rather than $\mathbb{P}_{\mathbf{X}_{[N]}}$ to emphasize the dependence on the model \mathcal{M} .

Lemma 4.5 (Coupled product of all random variables is the identity). $\underline{\otimes}_{i \in [N]} \mathbf{X}_i = \text{Id}_{\mathbf{X}_{[N]}}$

Proof. for any $\mathbf{X} \in \mathbf{X}_{[N]}$,

$$\underline{\otimes}_{i \in [N]} \mathbf{X}_i(\mathbf{X}) = (\pi_1(\mathbf{X}), \dots, \pi_N(\mathbf{X})) \quad (42)$$

$$= (x_1, \dots, x_n) \quad (43)$$

$$= \mathbf{X} \quad (44)$$

□

Definition 4.6 (Hard Interventions). Let \mathcal{M} be the set of all *SCMs* sharing the indices, spaces and measure $\langle N, M, \mathbf{X}_{\mathcal{I}}, \mathbf{E}_{[M]}, \mathbb{P}_{\mathcal{E}} \rangle$. Note that the causal mechanisms are not fixed.

Given an SCM $\mathcal{M} = \langle N, M, \mathbf{X}_{\mathcal{I}}, \mathbf{E}_{[M]}, \{f_i | i \in [N]\}, \mathbb{P}_{\mathcal{E}} \rangle$ and $\mathcal{S} \subset [N]$, a *hard intervention* on $\mathbf{X}_{\mathcal{S}}$ is a map $Do_{\mathcal{S}} : \mathbf{X}_{\mathcal{S}} \times \mathcal{M} \rightarrow \mathcal{M}$ such that for $\mathbf{a} \in \mathbf{X}_{\mathcal{S}}$, $Do_{\mathcal{S}}(\mathbf{a}, \mathcal{M}) = \langle N, M, \mathbf{X}_{\mathcal{I}}, \mathbf{E}_{[M]}, \{f'_i | i \in [N]\}, \mathbb{P}_{\mathcal{E}} \rangle$ where

$$f'_i = f_i \quad i \notin \mathcal{S} \quad (45)$$

$$f'_i = \pi_i(\mathbf{a}) \quad i \in \mathcal{S} \quad (46)$$

To match standard notation, we will write $\mathcal{M}^{do(\mathbf{X}_{\mathcal{S}}=\mathbf{a})} := Do_{\mathcal{S}}(\mathbf{a}, \mathcal{M})$

4.6 Recursive Structural Causal Models with Necessary Relationships

Necessary relationships are extra constraints on the joint distribution on endogenous variables defined by an SCM. For example, given an SCM \mathcal{M} if the

variable X_1 represents weight, X_2 represents height and X_3 represents BMI then we want to impose the constraint that

$$X_3 = \frac{X_1}{X_2} \quad (47)$$

$\mathbb{P}_{\mathcal{M}}$ -almost surely.

Definition 4.7 (Constrained Recursive Structural Causal Model (CSCM)). A CSCM $\mathcal{M} := \langle N, M, \mathbf{X}_{[N]}, \mathbf{E}_{[M]}, \{f_i | i \in [N]\}, \{r_i | i \in [N]\}, \mathbb{P}_{\mathcal{E}} \rangle$ is an SCM along with a set of *constraints* $r_i : \mathbf{X}_{[N]} \rightarrow X_i$.

If $X_i = r_i(\mathbf{X}_{[N]})$ $\mathbb{P}_{\mathcal{M}}$ -almost surely then \mathbf{M} is *valid*, otherwise it is *invalid*.

We can recover regular SCMs by imposing only trivial constraints:

Lemma 4.8 (CSCM with trivial constraints is always valid). *Let \mathcal{M} be a CSCM with the trivial constraints $r_i = \pi_i$ for all $i \in [N]$. Then \mathcal{M} is valid.*

Proof. By definition 4.7, we require $X_i = \pi_i$, $\mathbb{P}_{\mathcal{M}}$ -almost surely. $X_i(\mathbf{X}) = \pi_i(\mathbf{X})$ for all $\mathbf{X} \in \mathbf{X}_{[N]}$ and $P_{\mathcal{M}}(\mathbf{X}_{[N]}) = 1$, therefore \mathcal{M} is valid. \square

Call a constraint r_i *cyclic* if $X_i = r_i(\mathbf{X}_{[N]})$ implies there exists an index set $O \subset [N]$, $O \ni i$, such that for each $j \in O$, $\mathbf{b} \in \mathbf{X}_{O \setminus \{j\}}$ there exists $a \in X_j$ such that

$$X_{O \setminus \{j\}} = \mathbf{b} \quad (48)$$

$$\implies X_j = a \quad (49)$$

BMI is an example of a cyclic constraint if we insist that weight and height are always greater than 0. If $X_3 = \frac{X_1}{X_2}$ then we have:

$$[X_1, X_2] = [b_1, b_2] \quad (50)$$

$$\implies X_3 = \frac{b_1}{b_2} \quad (51)$$

$$[X_2, X_3] = [b_2, b_3] \quad (52)$$

$$\implies X_1 = b_2 b_3 \quad (53)$$

$$[X_1, X_3] = [b_1, b_3] \quad (54)$$

$$\implies X_2 = \frac{b_1}{b_3} \quad (55)$$

The following is a generally useful lemma that should probably be in basic definitions of Markov kernel spaces

Lemma 4.9 (Projection and selectors). *Given an indexed product space $\mathbf{X} := \prod_{i \in \mathcal{I}} X_i$ with ordered finite index set $\mathcal{I} \ni i$, let $\pi_i : \mathbf{X} \rightarrow X_i$ be the projection of the i -indexed element of $\mathbf{X} \in \mathbf{X}$.*

Let $F_{\pi_i} : \mathbf{X} \rightarrow \Delta(\mathcal{X}_i)$ be the Markov kernel associated with the function π_i , $F_{\pi_i} : \mathbf{X} \mapsto \delta_{\pi_i(\mathbf{X})}$. Given $O \subset \mathcal{I}$, define the selector S_i^O :

$$S_i^O = \begin{cases} \text{Id}_{X_i} & i \in O \\ *_{X_i} & i \notin O \end{cases} \quad (56)$$

Then $\underline{\otimes}_{i \in O} F_{\pi_i} = \underline{\otimes}_{i \in \mathcal{I}} S_i^O$.

Proof. Suppose O is the empty set. Then the empty tensor product $\underline{\otimes}_{i \in \emptyset} S_i$ and the empty coupled tensor product $\underline{\otimes}_{i \in \emptyset} F_{\pi_i}$ are both equal to $*_{\mathbf{X}}$.

By definition of F_{π_i} , $F_{\pi_i} = \underline{\otimes}_{i \in \mathcal{I}} S_i^{\{i\}}$.

Suppose for $P \subsetneq O$ with greatest element k we have $\underline{\otimes}_{i \in P} F_{\pi_i} = \underline{\otimes}_{i \in \mathcal{I}} S_i^P$, and suppose that j is the next element of O not in P .

$$(\underline{\otimes}_{i \in P} F_{\pi_i}) \underline{\otimes} F_{\pi_j} = \begin{array}{c} \mathbf{X} \begin{array}{l} \swarrow \boxed{\underline{\otimes}_{i \in P} F_{\pi_i}} - \mathbf{X}_P \\ \searrow \boxed{F_{\pi_j}} - X_j \end{array} \end{array} \quad (57)$$

$$= \begin{array}{c} \mathbf{X}_{<j} \quad X_j \quad \mathbf{X}_{>j} \begin{array}{l} \swarrow \boxed{\underline{\otimes}_{i \in P} F_{\pi_i}} - \mathbf{X}_P \\ \searrow \boxed{F_{\pi_j}} - X_j \end{array} \end{array} \quad (58)$$

$$= \begin{array}{c} \mathbf{X}_{<j} \quad X_j \quad \mathbf{X}_{>j} \begin{array}{l} \swarrow \boxed{\underline{\otimes}_{i \in P} F_{\pi_i}} - \mathbf{X}_P \\ \searrow * - X_j \end{array} \end{array} \quad (59)$$

$$= \begin{array}{c} \mathbf{X}_{<j} \quad X_j \quad \mathbf{X}_{>j} \begin{array}{l} \swarrow \boxed{\underline{\otimes}_{i \in P} F_{\pi_i}} - \mathbf{X}_P \\ \searrow - X_j \end{array} \end{array} \quad (60)$$

$$= \begin{array}{c} \mathbf{X}_{<j} \quad X_j \quad \mathbf{X}_{>j} \begin{array}{l} \swarrow \boxed{\underline{\otimes}_{i \in \mathcal{I}} S_i^P} - \mathbf{X}_P \\ \searrow - X_j \end{array} \end{array} \quad (61)$$

Because all elements of P are less than j , the selector S_k^P resolves to the discard

map for $k > j$:

$$= \begin{array}{c} \mathbf{X}_{<j} \\ X_j \\ \mathbf{X}_{>j} \end{array} \begin{array}{l} \text{---} \text{ } \otimes_{i<j} S_i^P \text{ ---} \mathbf{X}_P \\ \text{---} \text{ } * \text{ ---} \\ \text{---} \text{ } \times \text{ ---} \\ \text{---} \text{ } \cdot \text{ ---} \end{array} \begin{array}{l} \text{---} \mathbf{X}_P \\ \text{---} \\ \text{---} \\ \text{---} X_j \end{array} \quad (62)$$

$$\begin{array}{c}
\mathbf{X}_{<j} \text{ --- } \boxed{\otimes_{i<j} S_i^P} \text{ --- } \mathbf{X}_P \\
X_j \text{ --- } X_j \\
= \mathbf{X}_{>j} \text{ --- } *
\end{array} \quad (63)$$

$$\begin{array}{c}
\mathbf{X}_{<j} \\
X_j \\
\mathbf{X}_{>j}
\end{array}
= \left[\bigotimes_{i \in \mathcal{I}} S_i^{P \cup \{j\}} \right] \begin{array}{c} \\ \\ \end{array} \begin{array}{c} \mathbf{X}_P \\ X_j \end{array} \quad (64)$$

Where 64 follows from the definition of the selector $S_i^{P \cup \{j\}}$. The proof follows by induction on the elements of O .

1

Lemma 4.10 (Hard interventions do not affect the joint distributions of earlier variables). *Given a CSCM $\mathcal{M} = \langle N, M, \mathbf{X}_{[N]}, \mathbf{E}_{[M]}, \{f_i | i \in [N]\}, \{r_i | i \in [N]\}, \mathbb{P}_{\mathcal{E}} \rangle$, any $k \in [N]$ and any $O \subset [k-1]$, $P_{\mathcal{M}}(\mathbf{X}_O) = P_{\mathcal{M}}^{do(\mathbf{X}_k)=a}(\mathbf{X}_O)$ for all $a \in X_k$.*

Proof. Let $G_i^{\mathcal{Q}}$, $i \in [N]$ be the i -th iteration of the kernel defined in Equations 40 and 41 with respect to model \mathcal{Q} . Note that from Equation 41

$$\begin{array}{c} \textbf{E}_{[M]} - \boxed{G_i^Q} - * \\ \quad \quad \quad \downarrow \\ \quad \quad \quad \textbf{X}_{<i} \end{array} = \begin{array}{ccc} & \boxed{G_{i-1}^Q} & \textbf{X}_{<i} \\ \textbf{E}_{[M]} - & \swarrow & \downarrow \\ & & \boxed{F_{f_i}} - * \end{array} \quad (65)$$

$$= G_{i-1}^Q \quad (66)$$

It follows that

$$\mathbf{E}_{[M]} \text{---} \boxed{G_N} \text{---} \begin{array}{c} \mathbf{X}_{<i} \\ * \end{array} = G_{i-1} \quad (67)$$

Because $f_i = f_i^{do(X_k=a)}$ for $i < k$, we have

$$G_i^{\mathcal{M}} = G_i^{\mathcal{M}^{do(X_k=a)}} \quad (68)$$

for all $i < k$. By lemma 4.9, for any $O \subset [k-1]$ we have $F_{X_O} = \otimes_{i \in [N]} S_i^O$. As there are no elements of O greater than or equal to k , the selector S_i^O resolves to the discard for all $i \geq k$. Thus $F_{X_O} = (\otimes_{i \in [k-1]} S_i^O) \otimes *_{\mathbf{x}_{[N] \setminus [k-1]}}$. Defining $S_{[k-1]}^O := \otimes_{i \in [k-1]} S_i^O$, we have:

$$F_{\mathbf{X}_O} = \begin{array}{c} \mathbf{X}_{[k-1]} \text{ --- } \boxed{S_{[k-1]}^O} \text{ --- } \mathbf{X}_O \\ \mathbf{X}_{[N] \setminus [k-1]} \text{ --- } * \end{array} \quad (69)$$

Thus

$$\mathbb{P}_{\mathcal{M}}(\mathbf{X}_O) = \mathbb{P}_{\mathcal{E}} G_N^{\mathcal{M}} F_{\mathbf{X}_O} \quad (70)$$

$$\stackrel{69}{=} \begin{array}{c} \triangleleft \mathbb{P}_{\mathcal{E}} \text{ --- } \boxed{G_N^{\mathcal{M}}} \text{ --- } \boxed{S_{[k-1]}^O} \text{ --- } \mathbf{X}_O \\ * (\mathbf{X}_{[N] \setminus [k-1]}) \end{array} \quad (71)$$

$$\stackrel{67}{=} \begin{array}{c} \triangleleft \mathbb{P}_{\mathcal{E}} \text{ --- } \boxed{G_{k-1}^{\mathcal{M}}} \text{ --- } \boxed{S_{[k-1]}^O} \text{ --- } \mathbf{X}_O \end{array} \quad (72)$$

$$\stackrel{68}{=} \begin{array}{c} \triangleleft \mathbb{P}_{\mathcal{E}} \text{ --- } \boxed{G_{k-1}^{\mathcal{M}^{do(\mathbf{X}_k=a)}}} \text{ --- } \boxed{S_{[k-1]}^O} \text{ --- } \mathbf{X}_O \end{array} \quad (73)$$

$$\stackrel{67}{=} \begin{array}{c} \triangleleft \mathbb{P}_{\mathcal{E}} \text{ --- } \boxed{G_N^{\mathcal{M}^{do(\mathbf{X}_k=a)}}} \text{ --- } \boxed{S_{[k-1]}^O} \text{ --- } \mathbf{X}_O \\ * (\mathbf{X}_{[N] \setminus [k-1]}) \end{array} \quad (74)$$

$$= P_{\mathcal{M}^{do(\mathbf{X}_k=a)}}(\mathbf{X}_O) \quad (75)$$

□

Theorem 4.11 (Undefined hard interventions with cyclic constraints). *Consider a CSCM $\mathcal{M} = \langle N, M, \mathbf{X}_{[N]}, \mathbf{E}_{[M]}, \{f_i | i \in [N]\}, \{r_i | i \in [N]\}, \mathbb{P}_{\mathcal{E}} \rangle$ with r_i a cyclic constraint with respect to $O \subset [N]$ and the rest of the constraints trivial: $r_j = \pi_j$, $j \neq i$, and suppose \mathcal{M} is valid.*

If for each $k \in O$, $\exists A \in \mathcal{X}_i$ such that $0 < \mathbb{P}_{\mathcal{M}}(\mathbf{X}_i \in A) < 1$ then for at least one $k \in O$ all models given by a hard intervention on \mathbf{X}_k are invalid.

Proof. Choose k to be the maximum element of O . By the assumption \mathcal{M} is valid, we have $\mathbf{X}_i = r_i(\mathbf{X})$, $\mathbb{P}_{\mathcal{M}}$ -almost surely. Let $B^A = \{\mathbf{b} \in \mathbf{X}_{O \setminus k} | \mathbf{X}_{O \setminus k} = \mathbf{b} \implies \mathbf{X}_k \in A\}$ and $B^{A^C} = \{\mathbf{b} \in \mathbf{X}_{O \setminus k} | \mathbf{X}_{O \setminus k} = \mathbf{b} \implies \mathbf{X}_k \notin A\}$.

r_i holds on a set of measure 1, and wherever it holds $\mathbf{X}_{O \setminus \{k\}}$ is either in B^A or B^{A^C} . Thus $\mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in B^A \cup B^{A^C}) = 1$.

B^A and B^{A^C} are disjoint.

By construction, $\mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in B^A \ \& \ \mathbf{X}_k \in A) = \mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in B^A)$ and $\mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in B^{A^C} \ \& \ \mathbf{X}_k \in A^C) = \mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in B^{A^C})$.

By additivity, $\mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in B^A \ \& \ \mathbf{X}_k \in A) + \mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \notin B^A \ \& \ \mathbf{X}_k \in A) = P_{\mathcal{M}}(\mathbf{X}_k \in A)$.

By additivity again

$$\mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \notin B^A \ \& \ \mathbf{X}_k \in A) = \mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in B^{A^C} \ \& \ \mathbf{X}_k \in A) \quad (76)$$

$$+ \mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in (B^{A^C} \cup B^A)^C \ \& \ \mathbf{X}_k \in A) \quad (77)$$

$$<= 0 + P_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in (B^{A^C} \cup B^A)^C) \quad (78)$$

$$= 0 \quad (79)$$

Thus $\mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in B^A \ \& \ \mathbf{X}_k \in A) = P_{\mathcal{M}}(\mathbf{X}_k \in A) = P_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in B^A)$ and by an analogous argument $\mathbb{P}_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in B^{A^C}) = P_{\mathcal{M}}(\mathbf{X}_k \in A^C)$.

Choose some $a \in A$, and consider the hard intervention $\mathcal{M}^{do(\mathbf{X}_k=a)}$. Suppose $\mathcal{M}^{do(\mathbf{X}_k=a)}$ is also valid. Then, as before, $\mathbb{P}_{\mathcal{M}^{do(\mathbf{X}_k=a)}}(\mathbf{X}_{O \setminus \{k\}} \in B^{A^C}) = \mathbb{P}_{\mathcal{M}^{do(\mathbf{X}_k=a)}}(\mathbf{X}_k \in A^C)$.

By definition of hard interventions, $f_k^{do(\mathbf{X}_k=a)} = a$. Thus $G_N^{\mathcal{M}^{do(\mathbf{X}_k=a)}} F_{\mathbf{X}_k}$ is the kernel $\mathbf{X} \mapsto \delta_a$ and it follows that $\mathbb{P}_{\mathcal{M}^{do(\mathbf{X}_k=a)}}(\mathbf{X}_k) = \delta_a$.

By lemma 4.10, $\mathbb{P}_{\mathcal{M}^{do(\mathbf{X}_k=a)}}(\mathbf{X}_{O \setminus \{k\}} \in B^{A^C}) = P_{\mathcal{M}}(\mathbf{X}_{O \setminus \{k\}} \in B^{A^C}) = P_{\mathcal{M}}(\mathbf{X}_k \in A^C) > 0$. But $P_{\mathcal{M}^{do(\mathbf{X}_k=a)}}(\mathbf{X}_k \in A^C) = \delta_z(\mathbf{X}_k \in A^C) = 0$, contradicting the assumption of validity of $\mathcal{M}^{do(\mathbf{X}_k=a)}$.

An analogous argument shows that all hard interventions $a' \in A^C$ are also invalid. \square

4.7 Cyclic Structural Causal Models

It is not very surprising that acyclic causal models cannot accommodate cyclic constraints. Can cyclic causal models do so? While Bongers et al. (2016) has developed a theory of cyclic causal models, cyclic are generally far less well understood than acyclic models. I show that the theory of cyclic models that Bongers has developed also fails to define hard interventions on variables subject to cyclic constraints. This does not rule out the possibility that there is some other way to define cyclic causal models that do handle these constraints, but I have not taken it upon myself to develop such a theory.

Haven't done any work from here on

We adopt the framework of cyclic structural causal models to make our arguments, adapted from Bongers et al. (2016). This is somewhat non-standard, but allows us to make a stronger argument for the impossibility of modelling arbitrary sets of variables using structural interventional models.

Definition 4.12 (Structural Causal Model). A structural causal model (SCM) is a tuple

$$\mathcal{M} := \langle \mathcal{I}, \mathcal{J}, \mathbf{X}_{\mathcal{I}}, \mathbf{E}_{\mathcal{J}}, \mathbf{f}_{\mathcal{I}}, \mathbb{P}_{\mathcal{E}}, \mathbf{E}_{\mathcal{J}} \rangle \quad (80)$$

where

- \mathcal{I} is a finite index set of *endogenous variables*
- \mathcal{J} is a finite index set of *exogenous variables*
- $\mathbf{X}_{\mathcal{I}} := \{X_i\}_{\mathcal{I}}$ where, for each $i \in \mathcal{I}$, (X_i, \mathcal{X}_i) is a standard measurable space taking and the codomain of the i -th endogenous variable
- $\mathbf{E}_{\mathcal{J}} := \{E_j\}_{\mathcal{J}}$ where, for $j \in \mathcal{J}$, E_j is a standard measurable space and the codomain of the j -th endogenous variable
- $\mathbf{f}_{\mathcal{I}} = \bigotimes_{i \in \mathcal{I}} f_i$ is a measurable function, and $f_i : \mathbf{X}_{\mathcal{I}} \times \mathbf{E}_{\mathcal{J}} \rightarrow X_i$ is the causal mechanism controlling X_i
- $\mathbb{P}_{\mathcal{E}} \in \Delta(\mathbf{E}_{\mathcal{J}})$ is a probability measure on the space of exogenous variables
- $\mathbf{E}_{\mathcal{J}} = \bigotimes_{j \in \mathcal{J}} E_j$ is the set of exogenous variables, with $\mathbb{P}_{\mathcal{E}} = \mathbf{E}_{\mathcal{J}\#} P_{\mathcal{E}}$ and E_j is the j -th exogenous variable with marginal distribution given by $\mathbf{E}_{j\#} \mathbb{P}_{\mathcal{E}}$

If for $\mathbb{P}_{\mathcal{E}}$ -almost every $\mathbf{e} \in \mathbf{E}_{\mathcal{J}}$ there exists $\mathbf{X} \in \mathbf{X}_{\mathcal{I}}$ such that

$$\mathbf{X} = \mathbf{f}_{\mathcal{I}}(\mathbf{X}, \mathbf{e}) \quad (81)$$

Then an SCM \mathcal{M} induces a unique probability space $(\mathbf{X}_{\mathcal{I}} \times \mathbf{E}_{\mathcal{J}}, \mathcal{X}_{\mathcal{I}} \otimes \mathcal{E}_{\mathcal{J}}, \mathbb{P}_{\mathcal{M}})$ (Bongers et al., 2016). If no such solution exists then we will say an SCM is invalid, as it imposes mutually incompatible constraints on the endogenous variables. It may be also the case that multiple solutions exist.

If an SCM induces a unique probability space then there exist random variables $\{X_i\}_{i \in \mathcal{I}}$ such that, $P_{\mathcal{M}}$ almost surely Bongers et al. (2016):

$$X_i = f_i(\mathbf{X}_{\mathcal{I}}, \mathbf{E}_{\mathcal{J}}) \quad (82)$$

Where $\mathbf{X}_{\mathcal{I}} = \bigotimes_{i \in \mathcal{I}} X_i$.

A structural causal model can be transformed by *mechanism surgery*. Given $\mathcal{S} \subset \mathcal{I}$ and a set of new functions $\mathbf{f}_{\mathcal{S}}^I : \mathbf{X}_{\mathcal{S}} \times \mathbf{E}_{\mathcal{J}} \rightarrow \mathbf{X}_{\mathcal{S}}$, mechanism surgery “replaces” the corresponding parts of $\mathbf{f}_{\mathcal{I}}$ with $\mathbf{f}_{\mathcal{S}}^I$.

Definition 4.13 (Mechanism surgery). Let \mathcal{M} be the set of SCMs with elements $\langle \mathcal{I}, \mathcal{J}, \mathbf{X}_{\mathcal{I}}, \mathbf{E}_{\mathcal{J}}, _, \mathbb{P}_{\mathcal{E}}, \mathbf{f}_{\mathcal{I}} \rangle$ (note that the causal mechanisms are unspecified). Mechanism surgery is an operation $I : \mathbf{X}_{\mathcal{I}}^{\mathbf{X}_{\mathcal{I}} \times \mathbf{E}_{\mathcal{J}}} \times \mathcal{M} \rightarrow \mathcal{M}$ that takes a causal model \mathcal{M} with arbitrary causal mechanisms and a set of causal mechanisms $\mathbf{f}_{\mathcal{I}}'$ and maps it to a model $\mathcal{M}' = \langle \mathcal{I}, \mathcal{J}, \mathbf{X}_{\mathcal{I}}, \mathbf{E}_{\mathcal{J}}, \mathbf{f}_{\mathcal{I}}', \mathbb{P}_{\mathcal{E}}, \mathbf{f}_{\mathcal{I}}' \rangle$.

If \mathcal{M} has causal mechanisms $\mathbf{f}_{\mathcal{I}}$ and $\mathcal{O} \subset \mathcal{I}$ is the largest set such that $\pi_{\mathcal{O}} \circ \mathbf{f}_{\mathcal{I}} = \pi_{\mathcal{O}} \circ \mathbf{f}_{\mathcal{I}}'$ then we say I is an *intervention* on $\mathcal{L} := \mathcal{I} \setminus \mathcal{O}$. We will use the special notation $\mathcal{M}^{I(\mathcal{L}), \mathbf{f}_{\mathcal{L}}'} := I(\mathcal{M}, \mathbf{f}_{\mathcal{L}}')$ to denote an SCM related to \mathcal{M} by intervention on a subset of \mathcal{I} .

If furthermore $\pi_{\mathcal{L}} \mathbf{f}_{\mathcal{L}}'$ is a constant function equal to \mathbf{a} , then we say I is a *hard intervention* on \mathcal{L} . We use the special notation $\mathcal{M}^{Do(\mathcal{L})=\mathbf{a}} := I(\mathcal{M}, \mathbf{f}_{\mathcal{L}}')$ to denote SCMs related to \mathcal{M} by hard interventions. We also say that the *causal effect* of \mathcal{L} is the set of SCNMs $\{\mathcal{M}^{Do(\mathcal{L})=\mathbf{a}} | \mathbf{a} \in \mathbf{X}_{\mathcal{L}}\}$.

We say a *causal model* is any kind of model that defines causal effects. An SCM \mathcal{M} in combination with hard interventions defines causal effects, so an SCM is a causal model. Call each interventional model $\mathcal{M}^{do(X_i=x)}$ a *submodel* of \mathcal{M} .

Strictly, the random variables X_i depend on the probability space induced by a particular model \mathcal{M} , they are intended to refer to “the same variable” across different models that are related by mechanism surgery. We will abuse notation and use X_i to refer to the *family* of random variables induced by a set of models related by mechanism surgery, and rely on explicitly noting the measure $\mathbb{P}_{\dots}(\dots)$ to specify exactly which random variables we are talking about.

In practice, we typically specify a “small” SCM containing a few endogenous variables \mathcal{I} (called a “marginal SCM” by Bongers et al. (2016)) which is understood to summarise the relevant characteristics of a “large” SCM containing many variables \mathcal{I}^* . We will argue that without restrictions on the large set of variables \mathcal{I}^* , surgically transformed SCMs will usually be invalid.

Incidentally, this messiness with random variables can be solved if we use See-Do models.

4.8 Not all variables have well-defined interventions

A long-running controversy about causal inference concerns the question of “the causal effect of body mass index on mortality”. On the one hand, Hernán and Taubman (2008) and others claim that there is no well-defined causal effect of a person’s body mass index (BMI), defined as their weight divided by their height, and their risk of death. Pearl claims, in defence of Causal Bayesian Networks, that the causal effect of *obesity* is well-defined, though it is not clear whether he defends the proposition that BMI itself has a causal effect:

That BMI is merely a coarse proxy of obesity is well taken; obesity should ideally be described by a vector of many factors, some are easy to measure and others are not. But accessibility to measurement has no bearing on whether the effect of that vector of factors on morbidity is “well defined” or whether the condition of consistency is violated when we fail to specify the interventions used to regulate those factors. (Pearl, 2018b)

We argue that BMI does *not* have a well-defined causal effect, and without further assumptions neither does any variable.

4.8.1 Necessary relationships in cyclic SCMs

If an SCM contains variables that are necessarily related, we wish to impose the additional restriction that these necessary relationships hold for every submodel. This can be done by extending the previous definition:

Definition 4.14 (SCM with necessary relationships). An SCM with necessary relationships (SCNM) is a tuple $\mathcal{M} := \langle \mathcal{I}, \mathcal{J}, \mathbf{X}_{\mathcal{I}}, \mathbf{E}_{\mathcal{J}}, \mathbf{f}_{\mathcal{I}}, \mathbf{g}_{\mathcal{I}}, \mathbb{P}_{\mathcal{E}}, \mathbf{E}_{\mathcal{J}} \rangle$, which is an SCM with the addition of a vector function of *necessary relationships* $\mathbf{g}_{\mathcal{I}} := \bigotimes_{i \in \mathcal{I}} g_i$ where each $g_i : \mathbf{X}_{\mathcal{I}} \rightarrow X_i$ is a necessary relationship involving X_i .

An SCM with necessary induces a unique probability space if for $\mathbb{P}_{\mathcal{E}}$ -almost every $e \in \mathcal{E}$ there exists a unique $\mathbf{X} \in \mathbf{X}_{\mathcal{I}}$ such that simultaneously

$$\mathbf{X} = \mathbf{f}_{\mathcal{I}}(\mathbf{X}, \mathbf{e}) \quad (83)$$

$$\mathbf{X} = \mathbf{g}_{\mathcal{I}}(\mathbf{X}) \quad (84)$$

If no such \mathbf{X} exists then an SCNM is invalid.

Mechanism surgery for SCNMs involves modification of $\mathbf{f}_{\mathcal{I}}$ only, just like SCMs.

If we wish to stipulate that a particular variable X_i has no causal relationships or necessary relationships we can specify this with the trivial mechanisms $f_i : (\mathbf{X}, \mathbf{e}) \mapsto x_i$ and $g_i : \mathbf{X} \mapsto x_i$ respectively. An SCNM \mathcal{M} with the trivial necessary relationship $\mathbf{g}_{\mathcal{I}} : \mathbf{X} \mapsto \mathbf{X}$ induces the equivalent probability spaces as the SCM obtained by removing $\mathbf{g}_{\mathcal{I}}$ from \mathcal{M} .

Because BMI is always equal height/weight, given some SCNM \mathcal{M} containing endogenous variables X_h , X_w and X_b representing height, weight and BMI respectively, it should be possible to construct a more “primitive” SCNM \mathcal{M}^p containing every variable \mathcal{M} does except X_b that agrees with \mathcal{M} on all interventions except those on X_b .

Definition 4.15 (Marginal model). Given an SCNM

$$\mathcal{M} = \langle \mathcal{I}, \mathcal{J}, \mathbf{X}_{\mathcal{I}}, \mathbf{E}_{\mathcal{J}}, \mathbf{f}_{\mathcal{I}}, \mathbf{g}_{\mathcal{I}}, \mathbb{P}_{\mathcal{E}}, \mathbf{E}_{\mathcal{J}} \rangle$$

a marginal model over $\mathcal{L} \subset \mathcal{I}$ is an SCNM

$$\mathcal{M}^{\ast \mathcal{L}} = \langle \mathcal{O}, \mathcal{J}, \mathbf{X}_{\mathcal{O}}, \mathbf{E}_{\mathcal{J}}, \mathbf{f}_{\mathcal{O}}^{\mathcal{L}}, \mathbf{g}_{\mathcal{O}}^{\mathcal{L}}, \mathbb{P}_{\mathcal{E}}, \mathbf{E}_{\mathcal{J}} \rangle$$

such that $(\mathbb{P}_{\mathcal{M}})^{\ast \mathcal{L}} = \mathbb{P}_{(\mathcal{M}^{\ast \mathcal{L}})}$ and for all interventions $\mathbf{f}'_{\mathcal{O}}$ on $\mathcal{O} := \mathcal{I} \setminus \mathcal{L}$ that do not depend on \mathcal{L}

$$(\mathbb{P}_{\mathcal{M}^{I(\mathcal{O}), \mathbf{f}'_{\mathcal{O}}}})^{\ast \mathcal{L}} = \mathbb{P}_{(\mathcal{M}^{\ast \mathcal{L}}, I(\mathcal{O}), \mathbf{f}'_{\mathcal{O}} \circ \pi_{\mathcal{O}})}$$

A *primitive model* is a special case of a marginal model where any intervention that depended only on endogenous variables in the original model can be replicated with some intervention that depends only on endogenous variables in the marginal model. If the endogenous variables represent *observed* variables, then the plausible intervention operations may only be allowed to depend on these variables. In general, there may be interventions that are possible in the original model that are not possible in the marginal model.

Definition 4.16 (Primitive model). A *primitive model* \mathcal{M}^p is a marginal model of \mathcal{M} with respect to some \mathcal{L} such that for all interventions $\mathbf{f}'_{\mathcal{O}}$ that do not depend on \mathcal{J} there exists some $\mathbf{g}'_{\mathcal{O}} : \mathbf{X}_{\mathcal{O}} \times \mathbf{E}_{\mathcal{J}} \rightarrow \mathbf{X}_{\mathcal{O}}$ that does not depend on \mathcal{J} such that

$$(\mathbb{P}_{\mathcal{M}^{I(\mathcal{O}), \mathbf{f}'_{\mathcal{O}}}})^{\ast \mathcal{L}} = \mathbb{P}_{(\mathcal{M}^{\ast \mathcal{L}}, I(\mathcal{O}), \mathbf{g}'_{\mathcal{O}})}$$

We claim that given any SCNM \mathcal{M} containing endogenous variables X_h, X_w and X_b representing height, weight and BMI there should be a primitive model \mathcal{M}^p of \mathcal{M} with respect to $\{p\}$.

Lemma 4.17 (Primitive models). *\mathcal{M}^p is a primitive model of \mathcal{M} with respect to $\mathcal{L} \subset \mathcal{I}$ iff $S(\pi_{\mathcal{O}} \mathbf{f}_{\mathcal{I}}) \stackrel{a.s.}{=} S(\mathbf{f}_{\mathcal{O}}^p)$ for $\mathcal{O} := \mathcal{I} \setminus \mathcal{L}$ and for all $\mathbf{X} \in \mathbf{X}_{\mathcal{I}}, \mathbf{g}$*

However, as Theroem 4.19 shows, if an SCNM with height, weight and BMI can be derived from an SCNM containing just height and weight then there are no valid hard interventions on BMI.

Definition 4.18 (Derived model). Given a SCNM $\mathcal{M} := \langle \mathcal{I}, \mathcal{J}, \mathbf{X}_{\mathcal{I}}, \mathbf{E}_{\mathcal{J}}, \mathbf{f}_{\mathcal{I}}, \mathbf{g}_{\mathcal{I}}, \mathbb{P}_{\mathcal{E}}, \mathbf{E}_{\mathcal{J}} \rangle$, say $\mathcal{M}' = \langle \mathcal{I}', \mathcal{J}, \mathbf{X}_{\mathcal{I}'}, \mathbf{E}_{\mathcal{J}}, \mathbf{f}_{\mathcal{I}'}, \mathbf{g}_{\mathcal{I}'}, \mathbb{P}_{\mathcal{E}}, \mathbf{E}_{\mathcal{J}} \rangle$ is *derived* from \mathcal{M} if there exists some additional index/variable/relationships $i' \notin \mathcal{I}, X_{i'}$ such that

$$\mathcal{I}' = \mathcal{I} \cup \{i'\} \quad (85)$$

$$\mathbf{X}_{\mathcal{I}'} = \mathbf{X}_{\mathcal{I}} \cup X_{i'} \quad (86)$$

and, defining $\pi_{\mathcal{I}' \setminus i'} : \mathbf{X}_{\mathcal{I}'} \rightarrow \mathbf{X}_{\mathcal{I}}$ as the projection map that “forgets” $X_{i'}$, for any $\mathbf{e} \in \mathbf{E}_{\mathcal{J}}$ we have

$$\mathbf{X}' = \mathbf{f}_{\mathcal{I}'}(\mathbf{X}', \mathbf{e}) \quad (87)$$

$$\text{and } \mathbf{X}' = \mathbf{g}_{\mathcal{I}'}(\mathbf{X}') \implies \pi_{\mathcal{I}' \setminus i'}(\mathbf{X}') = \mathbf{f}_{\mathcal{I}}(\pi_{\mathcal{I}' \setminus i'}(\mathbf{X}'), \mathbf{e}) \quad (88)$$

$$\text{and } \pi_{\mathcal{I}' \setminus i'}(\mathbf{X}') = \mathbf{g}_{\mathcal{I}'}(\pi_{\mathcal{I}' \setminus i'}(\mathbf{X}')) \quad (89)$$

Theorem 4.19 (Interventions and necessary relationships don’t mix). *If \mathcal{M}' is derived from \mathcal{M} with the additional elements $i', X_{i'}, f_{i'}, g_{i'}$ and both \mathcal{M} and \mathcal{M}' are uniquely solvable and $\mathbb{P}_{\mathcal{X}' \otimes \mathcal{E}}(X_{i'})$ is not single valued then no hard interventions on $X_{i'}$ are possible.*

Proof. Because \mathcal{M} is uniquely solvable, for $\mathbb{P}_{\mathcal{E}}$ almost every \mathbf{e} there is a unique \mathbf{X}^e such that

$$\mathbf{X}^e = \mathbf{f}_{\mathcal{I}}(\mathbf{X}^e, \mathbf{e}) \quad (90)$$

$$\mathbf{X}^e = \mathbf{g}_{\mathcal{I}}(\mathbf{X}^e) \quad (91)$$

Because \mathcal{M}' is also uniquely solvable, for $\mathbb{P}_{\mathcal{E}}$ almost every \mathbf{e} we have $\mathbf{X}'^e \in \mathbf{X}_{\mathcal{I}'}$ such that $\pi_{\mathcal{I}' \setminus i'}(\mathbf{X}'^e) = \mathbf{X}^e$ and

$$x_{i'}'^e = \mathbf{g}_{i'}(\mathbf{X}'^e) \quad (92)$$

Because $\mathbb{P}_{\mathcal{X}' \otimes \mathcal{E}}(X_{i'})$ is not single valued there are non-null sets $A, B \in \mathcal{E}$ such that $e_a \in A, e_b \in B$ implies

$$\mathbf{g}_{i'}(\mathbf{X}'^{e_a}) \neq \mathbf{g}_{i'}(\mathbf{X}'^{e_b}) \quad (93)$$

Therefore there exists no $a \in X_{i'}$ that can simultaneously satisfy 92 for almost every \mathbf{e} . However, any hard intervention $\mathcal{M}', do(X_{i'}=a)$ requires such an a in order to be solvable. \square

Corollary 4.20. *Either there are no hard interventions defined on BMI or there is no SCNM containing height and weight with a unique solution from which an SCNM containing height, weight and BMI can be derived.*

I can formalise the following, but I'm just writing it out so I can get to the end for now

The problem posed by Theorem 4.19 can be circumvented to some extent by joint interventions. Consider the variables X_1 and X_2 where $X_1 = -X_2$ necessarily. While Theorem 4.19 disallows interventions on X_2 individually (supposing we can obtain a unique model featuring only X_1), it does not disallow interventions that jointly set X_1 and X_2 to permissible values. In this case, this is unproblematic as the only joint intervention that sets X_1 to 1 must also set X_2 to -1 .

If we have non-invertible necessary relationships such as $X_1 = X_2 + X_3$, however, there are now *multiple* joint interventions on X_1 that can be performed. I regard this as the most plausible solution to the difficulties raised so far: for variables that are in non-invertible necessary relationships, there is a set of operations associated with the “intervention” that sets $X_1 = 1$.

However, we still need to make sure the interventions that we have supposed comprise the operations associated with setting $X_1 = 1$ exist themselves. It is sufficient that the SCNM with X_1 is derived from a higher order *uniquely solvable SCM* with X_2 and X_3 only.

And necessary? There might be “degenerate” necessary relationships that don't harm the possibility of defining interventions, and I'd need to show an equivalence to an SCM in this case

If any variables are included in a causal model that are necessarily related to other variables (and honestly, is there any variable that isn't?), it is not enough to suppose that the model being used is a marginalisation of some larger causal model. Rather, it must be obtained by derivation and marginalisation from some model that represents the basic interventions that are possible, which we call the *atomic model*.

Definition 4.21 (Atomic model). Given an SCNM \mathcal{M} , the *atomic model* $\mathcal{M}_{\text{atom}}$ is a uniquely solvable SCM such that there exists a model \mathcal{M} is derived from of $\mathcal{M}_{\text{atom}}$.

Typically, in order to get an actually usable model you'll also need to marginalize, but I think this complication can be avoided

Definition 4.22 (Causal universality hypothesis). There exists a uniquely solvable SCM $\mathcal{M}_{\text{atom}}$ which is the atomic model that correctly represents all decision problems

I don't know how to define “correctly represents” or “causal problem”, but it seems like something like the universality hypothesis is necessary if you want to define “the causal effect of X ” independent of any atomic model

because interventions are defined in uniquely solvable SCMs and derivation preserves interventions on the old variables

what does that mean?

or causal problems?

Relate decisions to interventions on atomic model. Decisions->atomic model is straightforward, but the reverse direction is not so obvious

Causal effects are uniquely defined via atoms iff they are defined via decisions

Are there any plausible ways to construct atomic models?

5 Definitions and key notation

We use three notations for working with probability theory. The “elementary” notation makes use of regular symbolic conventions (functions, products, sums, integrals, unions etc.) along with the expectation operator \mathbb{E} . This is the most flexible notation which comes at the cost of being verbose and difficult to read. Secondly, we use a semi-formal string diagram notation extending the formal diagram notation for symmetric monoidal categories Selinger (2010). Objects in this diagram refer to stochastic maps, and by interpreting diagrams as symbols we can, in theory, be just as flexible as the purely symbolic approach. However, we avoid complex mixtures of symbols and diagrams elements, and fall back to symbolic representations if it is called for. Finally, we use a matrix-vector product convention that isn’t particularly expressive but can compactly express some common operations.

5.1 Standard Symbols

Symbol	Meaning
$[n]$	The natural numbers $\{1, \dots, n\}$
$f : a \mapsto b$	Function definition, equivalent to $f(a) := b$
Dots appearing in function arguments: $f(\cdot, \cdot, z)$	The “curried” function $(x, y) \mapsto f(x, y, z)$
Capital letters: A, B, X	sets
Script letters: $\mathcal{A}, \mathcal{B}, \mathcal{X}$	σ -algebras on the sets A, B, X respectively
Script \mathcal{G}	A directed acyclic graph made up of nodes V and edges
Blackboard \mathbb{P}, μ, ν :	Probability measures
δ_x	The Dirac delta measure: $\delta_x(A) = 1$ if $x \in A$ and 0 otherwise
Capital delta: $\Delta(\mathcal{E})$	The set of all probability measures on \mathcal{E}
Blackboard capitals: \mathbb{A}	Markov kernel $\mathbb{A} : X \times \mathcal{Y} \rightarrow [0, 1]$ (stochastic maps)
Subscripted Markov kernels: \mathbb{A}_x	The probability measure given by the curried Markov kernel \mathbb{A}_x
$A \rightarrow \Delta(\mathcal{B})$	Markov kernel signature, treated as equivalent to $A \times \mathcal{B}$
$\mathbb{A} : x \mapsto \nu$	Markov kernel definition, equivalent to $\mathbb{A}(x, B) = \nu(B)$ for all $B \in \mathcal{B}$
Sans serif capitals: A, X	Measurable functions; we will also call them random variables
\mathbb{F}^X	The Markov kernel associated with the function X : $\mathbb{F}^X \equiv \mathbb{A}_x$ for all x
$\mathbb{N}_{A B}$	The conditional probability (disintegration) of A given B
$\nu \mathbb{F}^X$	The marginal distribution of X under ν

5.2 Probability Theory

Given a set A , a σ -algebra \mathcal{A} is a collection of subsets of A where

- $A \in \mathcal{A}$ and $\emptyset \in \mathcal{A}$
- $B \in \mathcal{A} \implies B^C \in \mathcal{A}$
- \mathcal{A} is closed under countable unions: For any countable collection $\{B_i | i \in Z \subset \mathbb{N}\}$ of elements of \mathcal{A} , $\cup_{i \in Z} B_i \in \mathcal{A}$

A measurable space (A, \mathcal{A}) is a set A along with a σ -algebra \mathcal{A} . Sometimes the sigma algebra will be left implicit, in which case A will just be introduced as a measurable space.

Common σ algebras For any A , $\{\emptyset, A\}$ is a σ -algebra. In particular, it is the only sigma algebra for any one element set $\{*\}$.

For countable A , the power set $\mathcal{P}(A)$ is known as the discrete σ -algebra.

Given A and a collection of subsets of $B \subset \mathcal{P}(A)$, $\sigma(B)$ is the smallest σ -algebra containing all the elements of B .

Let T be all the open subsets of \mathbb{R} . Then $\mathcal{B}(\mathbb{R}) := \sigma(T)$ is the *Borel σ -algebra* on the reals. This definition extends to an arbitrary topological space A with topology T .

A *standard measurable set* is a measurable set A that is isomorphic either to a discrete measurable space A or $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$. For any A that is a complete separable metric space, $(A, \mathcal{B}(A))$ is standard measurable.

Given a measurable space (E, \mathcal{E}) , a map $\mu : \mathcal{E} \rightarrow [0, 1]$ is a *probability measure* if

- $\mu(E) = 1$, $\mu(\emptyset) = 0$
- Given countable collection $\{A_i\} \subset \mathcal{E}$, $\mu(\cup_i A_i) = \sum_i \mu(A_i)$

Write by $\Delta(\mathcal{E})$ the set of all probability measures on \mathcal{E} .

Given a second measurable space (F, \mathcal{F}) , a *stochastic map* or *Markov kernel* is a map $\mathbb{M} : E \times \mathcal{F} \rightarrow [0, 1]$ such that

- The map $\mathbb{M}(\cdot; A) : x \mapsto \mathbb{M}(x; A)$ is \mathcal{E} -measurable for all $A \in \mathcal{F}$
- The map $\mathbb{M}_x : A \mapsto \mathbb{M}(x; A)$ is a probability measure on F for all $x \in E$

Extending the subscript notation above, for $\mathbb{C} : X \times Y \rightarrow \Delta(\mathcal{Z})$ and $x \in X$ we will write \mathbb{C}_x for the “curried” map $y \mapsto \mathbb{C}_{x,y}$.

The map $x \mapsto \mathbb{M}_x$ is of type $E \rightarrow \Delta(\mathcal{F})$. We will abuse notation somewhat to write $\mathbb{M} : E \rightarrow \Delta(\mathcal{F})$, which captures the intuition that a Markov kernel maps from elements of E to probability measures on \mathcal{F} . Note that we “reverse” this idea and consider Markov kernels to map from elements of \mathcal{F} to measurable functions $E \rightarrow [0, 1]$, an interpretation found in Clerc et al. (2017), but (at this stage) we don’t make use of this interpretation here.

Given an indiscrete measurable space $(\{*\}, \{\{*\}, \emptyset\})$, we identify Markov kernels $\mathbb{N} : \{*\} \rightarrow \Delta(\mathcal{E})$ with the probability measure \mathbb{N}_* . In addition, there is a unique Markov kernel $* : E \rightarrow \Delta(\{\{*\}, \emptyset\})$ given by $x \mapsto \delta_*$ for all $x \in E$ which we will call the “discard” map.

5.3 Product Notation

We can use a notation similar to the standard notation for matrix-vector products to represent operations with Markov kernels. Probability measures $\mu \in \Delta(\mathcal{X})$ can be read as row vectors, Markov kernels as matrices and measurable functions $\mathbb{T} : Y \rightarrow T$ as column vectors. Defining $\mathbb{M} : X \rightarrow \Delta(\mathcal{Y})$ and $\mathbb{N} : Y \rightarrow \Delta(\mathcal{Z})$, the measure-kernel product $\mu \mathbb{A}(G) := \int \mathbb{A}_x(G) d\mu(x)$ yields a probability measure $\mu \mathbb{A}$ on \mathcal{Z} , the kernel-kernel product $\mathbb{M} \mathbb{N}(x; H) = \int_Y \mathbb{B}(y; H) d\mathbb{A}_x$ yields a kernel $\mathbb{M} \mathbb{N} : X \rightarrow \Delta(\mathcal{Z})$ and the kernel-function product $\mathbb{A} \mathbb{T}(x) := \int_Y \mathbb{T}(y) d\mathbb{A}_x$ yields a measurable function $X \rightarrow T$. Kernel products are associative (Çinlar, 2011).

The tensor product $(\mathbb{M} \otimes \mathbb{N})(x, y; G, H) := \mathbb{M}(x; G) \mathbb{N}(y; H)$ yields a kernel $(\mathbb{M} \otimes \mathbb{N}) : X \times Y \rightarrow \Delta(\mathcal{Y} \otimes \mathcal{Z})$.

5.4 String Diagrams

Some constructions are unwieldy in product notation; for example, given $\mu \in \Delta(\mathcal{E})$ and $\mathbb{M} : E \rightarrow (\mathcal{F})$, it is not straightforward to construct a measure $\nu \in \Delta(\mathcal{E} \otimes \mathcal{F})$ that captures the “joint distribution” given by $A \times B \mapsto \int_A \mathbb{M}(x; B) d\mu$.

Such constructions can, however, be straightforwardly captured with string diagrams, a notation developed for category theoretic probability. Cho and Jacobs (2019) also provides an extensive introduction to the notation discussed here.

Some key ideas of string diagrams:

- Basic string diagrams can always be interpreted as a mixture of kernel-kernel products and tensor products of Markov kernels
 - Extended string diagrams can be interpreted as a mixture of kernel-kernel products, kernel-function products, tensor products of kernels and functions and scalar products
- String diagrams are the subject of a coherence theorem: taking a string diagram and applying a planar deformation yields a string diagram that represents the same kernel (Selinger, 2010). This also holds for a number of additional transformations detailed below

A kernel $\mathbb{M} : X \rightarrow \Delta(\mathcal{Y})$ is written as a box with input and output wires, probability measures $\mu \in \Delta(\mathcal{X})$ are written as triangles “closed on the left” and measurable functions (which are only elements of the “extended” notation) $\mathbb{T} : Y \rightarrow T$ as triangles “closed on the right”. For this introduction we will label wires with the names of their corresponding spaces, but in practice we will

usually name them with corresponding *random variables*, though additional care is required when using random variables as labels (see paragraph ??).

For $\mathbb{M} : X \rightarrow \Delta(\mathcal{Y})$, $\mu \in \Delta(\mathcal{X})$ and $f : X \rightarrow W$:

$$X \text{ --- } \boxed{\mathbb{M}} \text{ --- } Y \quad \triangleleft_{\mu} \text{ --- } X \quad X \text{ --- } \triangleright_f \quad (94)$$

Elementary operations We can compose Markov kernels with appropriate spaces - the equivalent operation of the “matrix products” of product notation. Given $\mathbb{M} : X \rightarrow \Delta(\mathcal{Y})$ and $\mathbb{N} : Y \rightarrow \Delta(\mathcal{Z})$, we have

$$\mathbb{M}\mathbb{N} := X \text{ --- } \boxed{\mathbb{M}} \text{ --- } \boxed{\mathbb{N}} \text{ --- } Z \quad (95)$$

Probability measures are distinguished in that that they only admit “right composition” while functions only admit “left composition”. For $\mu \in \Delta(\mathcal{E})$, $h : F \rightarrow X$:

$$\mu\mathbb{M} := \triangleleft_{\mu} \text{ --- } \boxed{\mathbb{M}} \text{ --- } Z \quad (96)$$

$$\mathbb{M}f := X \text{ --- } \boxed{\mathbb{M}} \text{ --- } \triangleright_f \quad (97)$$

A diagram that is closed on the right and the left is an expectation:

$$\mathbb{E}_{\mu\mathbb{M}}(f) = \mu\mathbb{M}f \quad (98)$$

$$:= \triangleleft_{\mu} \text{ --- } \boxed{\mathbb{M}} \text{ --- } \triangleright_f \quad (99)$$

We can also combine Markov kernels using tensor products, which we represent with vertical juxtaposition. For $\mathbb{O} : Z \rightarrow \Delta(\mathcal{W})$:

$$\mathbb{M} \otimes \mathbb{N} := \begin{array}{c} X \text{ --- } \boxed{\mathbb{M}} \text{ --- } Y \\ Z \text{ --- } \boxed{\mathbb{O}} \text{ --- } W \end{array} \quad (100)$$

Product spaces can be represented either by two parallel wires or a single wire:

$$X \times Y \cong \text{Id}_X \otimes \text{Id}_Y := \begin{array}{c} X \text{ --- } X \\ Y \text{ --- } Y \end{array} \quad (101)$$

$$= X \times Y \text{ --- } X \times Y \quad (102)$$

Because a product space can be represented by parallel wires, a kernel $\mathbb{L} : X \rightarrow \Delta(\mathcal{Y} \otimes \mathcal{Z})$ can be written using either two parallel output wires or a single output wire:

$$X \text{ --- } \boxed{\mathbb{L}} \text{ --- } \begin{array}{c} Y \\ Z \end{array} \quad (103)$$

$$\equiv \quad (104)$$

$$X \text{ --- } \boxed{\mathbb{L}} \text{ --- } Y \times Z \quad (105)$$

Probability measures, Markov kernels and functions One has to exercise special care when including functions in diagrammatic notation. While any diagram that includes only probability measures (triangles pointing to the left) and Markov kernels (rectangles) is automatically a Markov kernel itself, while diagrams that include functions (triangles pointing to the right) only represent Markov kernels if they are correctly normalised, which is not a property that can be checked just by looking at the shape of the diagram.

Markov kernels with special notation A number of Markov kernels are given special notation distinct from the generic “box” representation above. These special representations facilitate intuitive graphical interpretations.

The identity kernel $\mathbf{Id} : X \rightarrow \Delta(X)$ maps a point x to the measure δ_x that places all mass on the same point:

$$\mathbf{Id} : x \mapsto \delta_x \equiv X \text{ --- } X \quad (106)$$

The identity kernel acts as the identity under left or right products:

$$(\mathbb{K}\mathbf{Id})_w(A) = \int_X \mathbf{Id}_x(A) d\mathbb{K}_w(x) \quad (107)$$

$$= \int_X \delta_x(A) d\mathbb{K}_w(x) \quad (108)$$

$$= \int_A d\mathbb{K}_w(x) \quad (109)$$

$$= \mathbb{K}_w(A) \quad (110)$$

$$(\mathbf{Id}\mathbb{K})_w(A) = \int_X \mathbb{K}_x(A) d\mathbf{Id}_w(x) \quad (111)$$

$$= \int_X \mathbb{K}_x(A) d\delta_w(x) \quad (112)$$

$$= \mathbb{K}_w(A) \quad (113)$$

The copy map $\Upsilon : X \rightarrow \Delta(\mathcal{X} \times \mathcal{X})$ maps a point x to two identical copies of x :

$$\Upsilon : x \mapsto \delta_{(x,x)} \equiv \begin{array}{c} X \\ \diagdown \quad \diagup \\ X \end{array} \quad (114)$$

The copy map “copies” its arguments under an integral:

$$\int_{(\cdot)} X \times X f(x, x', x'') d\Upsilon_x(x', x'') = \int_{(\cdot)} X \times X f(x, x', x'') d\delta_{(x,x)}(x', x'') \quad (115)$$

$$= f(x, x, x) \quad (116)$$

$$\int_W \int_{(\cdot)} X \times X f(x', x'') d\Upsilon_w(x', x'') d\mu(w) \quad (117)$$

$$= \int_W f(w, w) d\mu(w) \quad (118)$$

The swap map $\sigma : X \times Y \rightarrow \Delta(\mathcal{Y} \otimes \mathcal{X})$ swaps its inputs:

$$\sigma := (x, y) \mapsto \delta_{(y,x)} \equiv \begin{array}{c} Y \\ \diagdown \quad \diagup \\ X \end{array} \quad (119)$$

The swap map swaps its arguments under an integral:

$$\int_{(\cdot)} X \times X f(x, x') d\sigma_{(x_0, x_1)}(x, x') = \int_{(\cdot)} X \times X f(x, x') d\delta_{(x_1, x_0)}(x, x') \quad (120)$$

$$= f(x_1, x_0) \quad (121)$$

The discard map $*$: $X \rightarrow \Delta(\{*\})$ maps every input to δ_* . Note that the only non-empty event in $\{\emptyset, \{*\}\}$ must have probability 1.

$$* : x \mapsto \delta_* \equiv X \longrightarrow * \quad (122)$$

Any measurable function $F \rightarrow X$ has an associated Markov kernel $F \rightarrow \Delta(\mathcal{X})$. The Markov kernel associated with a function is different to the function itself - while the product of a probability measure μ with a function f is an expectation μf (see Definition 99), the product of a probability measure with the associated Markov kernel is the pushforward measure $f_{\#}\mu$.

Definition 5.1 (Function induced kernel). Given a measurable function $g : F \rightarrow X$, define the function induced kernel $\mathbb{F}^g : F \rightarrow \Delta(\mathcal{X})$ to be the the Markov kernel $a \mapsto \delta_{g(a)}$ for all $a \in X$.

Definition 5.2 (Pushforward kernel). Given a kernel $\mathbb{M} : E \rightarrow \Delta(\mathcal{F})$ and a measurable function $g : F \rightarrow X$, the *pushforward kernel* $g_{\#}\mathbb{M} : E \rightarrow \Delta(\mathcal{X})$ is the kernel such that $g_{\#}\mathbb{M}(a; B) = \mathbb{M}(a; g^{-1}(B))$.

If E is the one element space $\{*\}$, then $\mathbb{M} : \{*\} \rightarrow \Delta(\mathcal{F})$ can be identified with the probability measure \mathbb{M}_* and the pushforward kernel $g_{\#}\mathbb{M}$ identified with the pushforward measure $g_{\#}\mathbb{M}_*$, so pushforward kernels reduce to pushforward measures.

Lemma 5.3 (Pushforward kernels are functional kernel products). *Given a kernel $\mathbb{M} : E \rightarrow \Delta(\mathcal{F})$ and a measurable function $g : F \rightarrow X$, the pushforward $g_{\#}\mathbb{M} = \mathbb{M}\mathbb{F}^g$.*

Proof.

$$\mathbb{M}\mathbb{F}^g(a; B) = \int_F \delta_{g(y)}(B) d\mathbb{M}_a(y) \quad (123)$$

$$= \int_F \delta_y(g^{-1}(B)) d\mathbb{M}_a(y) \quad (124)$$

$$= \int_{g^{-1}(B)} d\mathbb{M}_a(y) \quad (125)$$

$$= g_{\#}\mathbb{M}(a; B) \quad (126)$$

□

5.4.1 Comparison of notations

We are in a position to compare the three introduced notations using a few examples. Given $\mu \in \Delta(X)$, $\mathbb{A} : X \rightarrow \Delta(Y)$ and $A \in \mathcal{X}$, $B \in \mathcal{Y}$, the following correspondences hold, where we express the same object in elementary notation, product notation and string notation respectively:

$$\nu := A \times B \mapsto \int_A A(x; B) d\mu(x) \equiv \mu \vee (\mathbf{Id}_X \otimes \mathbb{A}) \equiv \begin{array}{c} \text{---} X \\ \swarrow \quad \searrow \\ \triangleleft \mu \quad \boxed{\mathbb{A}} \text{---} Y \end{array} \quad (127)$$

Where the resulting object is a probability measure $\nu \in \Delta(\mathcal{X} \otimes \mathcal{Y})$. Note that the elementary notation requires a function definition here, while the product and string notations can represent the measure without explicitly addressing its action on various inputs and outputs. Cho and Jacobs (2019) calls this construction “integrating \mathbb{A} with respect to μ ”.

Define the marginal $\nu_Y \in \Delta(\mathcal{Y}) : B \mapsto \nu(X \times B)$ for $B \in \mathcal{Y}$ and similarly for ν_X . We can then express the result of marginalising 127 over X in our three

separate notations as follows:

$$\nu_Y(B) = \nu(X \times B) = \int_X A(x; B) d\mu(x) \quad (128)$$

$$\nu_Y = \mu \mathbb{A} = \mu \vee (\mathbf{Id}_X \otimes \mathbb{A})(\ast \otimes \mathbf{Id}_Y) \quad (129)$$

$$\nu_Y = \begin{array}{c} \triangleleft \mu \\ \hline \boxed{\mathbb{A}} - Y \end{array} = \begin{array}{c} \triangleleft \mu \\ \hline \boxed{\mathbb{A}} - Y \end{array} \begin{array}{c} \ast \end{array} \quad (130)$$

The elementary notation 128 makes the relationship between ν_Y and ν explicit and, again, requires the action on each event to be defined. The product notation 129 is, in my view, the least transparent but also the most compact in the form $\mu \mathbb{A}$, and does not demand the explicit definition of how ν_Y treats every event. The graphical notation is the least compact in terms of space taken up on the page, but unlike the product notation it shows a clear relationship to the graphical construction in 127, and displays a clear graphical logic whereby marginalisation corresponds to “cutting off branches”. Like product notation, it also allows for the definition of derived measures such as ν_Y without explicit definition of the handling of all events. It also features a much smaller collection of symbols than does elementary notation.

String diagrams often achieve a good balance between being ease of understanding at a glance and expressive power. On the downside, they can be time consuming to typeset, and formal reasoning with them takes some practice.

5.4.2 Working With String Diagrams

todo:

- Functional generalisation
- Conditioning
- Infinite copy map
- De Finetti’s representation theorem

There are a relatively small number of manipulation rules that are useful for string diagrams. In addition, we will define graphically analogues of the standard notions of *conditional probability*, *conditioning*, and infinite sequences of exchangeable random variables.

Axioms of Symmetric Monoidal Categories For the following, we either omit labels or label diagrams with their domain and codomain spaces, as we are discussing identities of kernels rather than identities of components of a conditional probability space. Recalling the unique Markov kernels defined above, the following equivalences, known as the *commutative comonoid axioms*, hold among string diagrams:

$$\begin{array}{c} \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} := \begin{array}{c} \text{---} \end{array} \begin{array}{c} \text{---} \end{array} \quad (131)$$

$$\begin{array}{c} \text{---} \end{array} \begin{array}{c} * \\ \text{---} \end{array} = \begin{array}{c} \text{---} \end{array} \begin{array}{c} \text{---} \\ * \end{array} = \text{---} \quad (132)$$

$$\begin{array}{c} \text{X} \end{array} \begin{array}{c} \text{---} \end{array} \begin{array}{c} \text{X} \\ \text{X} \end{array} = \begin{array}{c} \text{---} \end{array} \begin{array}{c} \text{---} \end{array} \quad (133)$$

The discard map $*$ can “fall through” any Markov kernel:

$$\text{---} \boxed{\mathbb{A}} \text{---} * = \text{---} * \quad (134)$$

Combining 132 and 134 we can derive the following: integrating $\mathbb{A} : X \rightarrow \Delta(\mathcal{Y})$ with respect to $\mu \in \Delta(\mathcal{X})$ and then discarding the output of \mathbb{A} leaves us with μ :

$$\begin{array}{c} \triangleleft \mu \end{array} \begin{array}{c} \text{---} \end{array} \begin{array}{c} \text{---} \end{array} \boxed{\mathbb{A}} \text{---} * = \begin{array}{c} \triangleleft \mu \end{array} \begin{array}{c} \text{---} \end{array} \begin{array}{c} \text{---} \end{array} * = \begin{array}{c} \triangleleft \mu \end{array} \text{---} \quad (135)$$

In elementary notation, this is equivalent to the fact that, for all $B \in \mathcal{X}$, $\int_B \mathbb{A}(x; B) d\mu(x) = \mu(B)$.

The following additional properties hold for $*$ and \curlyvee :

$$X \times Y \text{---} * = \begin{array}{c} X \\ Y \end{array} \text{---} * \quad (136)$$

$$X \times Y \text{---} \begin{array}{c} X \times Y \\ X \times Y \end{array} = \begin{array}{c} X \\ Y \end{array} \text{---} \begin{array}{c} X \\ Y \end{array} \quad (137)$$

A key fact that *does not* hold in general is

(138)

In fact, it holds only when \mathbb{A} is a *deterministic* kernel.

Definition 5.4 (Deterministic Markov kernel). A *deterministic* Markov kernel $\mathbb{A} : E \rightarrow \Delta(\mathcal{F})$ is a kernel such that $\mathbb{A}_x(B) \in \{0, 1\}$ for all $x \in E$, $B \in \mathcal{F}$.

Theorem 5.5 (Copy map commutes for deterministic kernels (Fong, 2013)). Equation 138 holds iff \mathbb{A} is *deterministic*.

5.5 Random Variables

The summary of this section is:

- Random variables are usually defined as measurable functions on a *probability space*
- It's possible to define them as measurable functions on a *Markov kernel space* instead
- It is useful to label wires with random variable names instead of names of spaces

Probability theory is primarily concerned with the behaviour of *random variables*. This behaviour can be analysed via a collection of probability measures and Markov kernels representing joint, marginal and conditional distributions of random variables of interest. In the framework developed by Kolmogorov, this collection of joint, marginal and conditional distributions is modeled by a single underlying *probability space*, and random variables by measurable functions on the probability space.

We use the same approach here, with a couple of additions. We are interested in variables whose outcomes depend both on random processes and decisions. Suppose that given a particular distribution over decision variables, a probability distribution over the decision variables and random variables is obtained. Such a model is described by a Markov kernel rather than a probability distribution. We therefore investigate *Markov kernel spaces*.

In the graphical notation that we are using, random variables can be thought of as a means of assigning unambiguous names to each wire in a set of diagrams. In order to do this, it is necessary to suppose that all diagrams in the set describe properties of an *ambient Markov kernel* or *ambient probability measure*. Consider the following example with the ambient probability measure $\mu \in \Delta(\mathcal{X} \otimes \mathcal{X})$. Suppose we have a Markov kernel $\mathbb{K} : X \rightarrow \Delta(\mathcal{X})$ such that the following holds:

(139)

Suppose that we also assign the names X_1 to the upper output wire and X_2 to the lower output wire in the diagram of μ :

$$\begin{array}{c} \triangleleft \mu \\ \hline X_1 \\ X_2 \end{array} \quad (140)$$

Then it seems sensible to call \mathbb{K} “the probability of X_2 given X_1 ”. We will make this precise so that it matches the usual notion of the probability of one variable given another (see Çinlar (2011) for a definition of this usual notion).

Definition 5.6 (Probability space, Markov kernel space). A *Markov kernel space* $(\mathbb{K}, \Omega, \mathcal{F}, D, \mathcal{D})$ is a Markov kernel $\mathbb{K} : D \rightarrow \Delta(\mathcal{D} \otimes \mathcal{F})$, called the *ambient kernel*, along with the sample space (Ω, \mathcal{F}) and the domain (D, \mathcal{D}) . We suppose that \mathbb{K} is such that there exists a *fundamental kernel* \mathbb{K}_0 satisfying

$$\mathbb{K} := \begin{array}{c} \boxed{\mathbb{K}_0} \\ \swarrow \quad \searrow \end{array} \quad (141)$$

For brevity, we will omit the σ -algebras in further definitions of Markov kernel spaces: (\mathbb{K}, Ω, D) .

A *probability space* $(\mathbb{P}, \Omega, \mathcal{F})$ is a probability measure $\mathbb{P} : \Delta(\Omega)$, which we call the *ambient measure*, along with the *sample space* Ω and the *events* \mathcal{F} . A probability space is equivalent to a Markov kernel space with domain $D = \{*\}$ - note that $\Omega \times \{*\} \cong \Omega$.

Definition 5.7 (Random variable). Given a Markov kernel space (\mathbb{K}, Ω, D) , a random variable X is a measurable function $\Omega \times D \rightarrow E$ for arbitrary measurable E .

Definition 5.8 (Domain variable). Given a Markov kernel space (\mathbb{K}, Ω, D) , the *domain variable* $D : \Omega \times D \rightarrow D$ is the distinguished random variable $D : (x, d) \mapsto d$.

Unlike random variables on probability spaces, random variables on Markov kernel spaces do not generally have unique marginal distributions. An analogous operation of *marginalisation* can be defined, but the result is generally a Markov kernel. We will define marginalisation via coupled tensor products.

Definition 5.9 (Coupled tensor product \otimes). Given two Markov kernels \mathbb{M} and \mathbb{N} or functions f and g with shared domain E , let $\mathbb{M} \otimes \mathbb{N} := \vee(\mathbb{M} \otimes \mathbb{N})$ and $f \otimes g := \vee(f \otimes g)$ where these expressions are interpreted using standard product notation. Graphically:

$$\underline{\mathbb{M}} \otimes \underline{\mathbb{N}} := \begin{array}{c} \begin{array}{c} \boxed{\mathbb{M}} - \mathbf{X} \\ \boxed{\mathbb{N}} - \mathbf{Y} \end{array} \\ E \text{ --- } \left. \begin{array}{c} \boxed{\mathbb{M}} \\ \boxed{\mathbb{N}} \end{array} \right\} \end{array} \quad (142)$$

$$f \otimes g := \begin{array}{c} \begin{array}{c} \triangle f \\ \triangle g \end{array} \\ E \text{ --- } \left. \begin{array}{c} \triangle f \\ \triangle g \end{array} \right\} \end{array} \quad (143)$$

The operation denoted by \otimes is associative (Lemma 5.10), so we can without ambiguity write $f \otimes g \otimes h = (f \otimes g) \otimes h = f \otimes (g \otimes h)$ for finite groups of functions or Markov kernels sharing a domain.

The notation $\otimes_{i \in [N]} f_i$ is taken to mean $f_1 \otimes f_2 \otimes \dots \otimes f_N$.

Lemma 5.10 (\otimes is associative). *For Markov kernels $\mathbb{L} : E \rightarrow \delta(\mathcal{F})$, $\mathbb{M} : E \rightarrow \delta(\mathcal{G})$ and $\mathbb{N} : E \rightarrow \delta(\mathcal{H})$, $(\underline{\mathbb{L}} \otimes \underline{\mathbb{M}}) \otimes \underline{\mathbb{N}} = \underline{\mathbb{L}} \otimes (\underline{\mathbb{M}} \otimes \underline{\mathbb{N}})$.*

Proof.

$$\underline{\mathbb{L}} \otimes (\underline{\mathbb{M}} \otimes \underline{\mathbb{N}}) = \begin{array}{c} \begin{array}{c} \boxed{\mathbb{L}} - F \\ \boxed{\mathbb{M}} - G \\ \boxed{\mathbb{N}} - H \end{array} \\ E \text{ --- } \left. \begin{array}{c} \boxed{\mathbb{L}} \\ \boxed{\mathbb{M}} \\ \boxed{\mathbb{N}} \end{array} \right\} \end{array} \quad (144)$$

$$= \begin{array}{c} \begin{array}{c} \boxed{\mathbb{L}} - F \\ \boxed{\mathbb{M}} - G \\ \boxed{\mathbb{N}} - H \end{array} \\ E \text{ --- } \left. \begin{array}{c} \boxed{\mathbb{L}} \\ \boxed{\mathbb{M}} \\ \boxed{\mathbb{N}} \end{array} \right\} \end{array} \quad (145)$$

$$= (\underline{\mathbb{L}} \otimes \underline{\mathbb{M}}) \otimes \underline{\mathbb{N}} \quad (146)$$

This follows directly from Equation 131. \square

Definition 5.11 (Marginal distribution, marginal kernel). Given a probability space $(\mathbb{P}, \Omega, \mathcal{F})$ and the random variable $\mathbf{X} : \Omega \rightarrow G$ the *marginal distribution* of \mathbf{X} is the probability measure $\mathbb{P}^{\mathbf{X}} := \mathbb{P}\mathbb{F}^{\mathbf{X}}$.

See Lemma 5.3 for the proof that this matches the usual definition of marginal distribution.

Given a Markov kernel space $(\mathbb{K}, \Omega, \mathcal{F}, D, \mathcal{D})$ and the random variable $\mathbf{X} : \Omega \rightarrow G$, the *marginal kernel* is $\mathbb{K}^{\mathbf{X}|D} := \mathbb{K}\mathbb{F}^{\mathbf{X}}$.

Definition 5.12 (Joint distribution, joint kernel). Given a probability space $(\mathbb{P}, \Omega, \mathcal{F})$ and the random variables $\mathbf{X} : \Omega \rightarrow G$ and $\mathbf{Y} : \Omega \rightarrow H$, the *joint distribution* of \mathbf{X} and \mathbf{Y} , $\mathbb{P}^{\mathbf{X}\mathbf{Y}} \in \Delta(\mathcal{G} \otimes \mathcal{H})$, is the marginal distribution of $\mathbf{X} \otimes \mathbf{Y}$. That is, $\mathbb{P}^{\mathbf{X}\mathbf{Y}} := \mathbb{P}\mathbb{F}^{\mathbf{X} \otimes \mathbf{Y}}$.

This is identical to the definition in Çinlar (2011) if we note that the random variable $(\mathbf{X}, \mathbf{Y}) : \omega \mapsto (\mathbf{X}(\omega), \mathbf{Y}(\omega))$ (Çinlar's definition) is precisely the same thing as $\mathbf{X} \otimes \mathbf{Y}$.

Analogously, the joint kernel $\mathbb{K}^{\mathbf{X}\mathbf{Y}|D}$ is the product $\mathbb{K}\mathbb{F}^{\mathbf{X} \otimes \mathbf{Y}}$.

Joint distributions and kernels have a nice visual representation, as a result of Lemma 5.13 which follows.

Lemma 5.13 (Product marginalisation interchange). *Given two functions, the kernel associated with their coupled product is equal to the coupled product of the kernels associated with each function.*

Given $\mathsf{X} : \Omega \rightarrow G$ and $\mathsf{Y} : \Omega \rightarrow H$, $\mathbb{F}^{\mathsf{X} \otimes \mathsf{Y}} = \mathbb{F}^{\mathsf{X}} \otimes \mathbb{F}^{\mathsf{Y}}$

Proof. For $a \in \Omega$, $B \in \mathcal{G}$, $C \in \mathcal{H}$,

$$\mathbb{F}^{\mathsf{X} \otimes \mathsf{Y}}(a; B \times C) = \delta_{\mathsf{X}(a), \mathsf{Y}(a)}(B \times C) \quad (147)$$

$$= \delta_{\mathsf{X}(a)}(B) \delta_{\mathsf{Y}(a)}(C) \quad (148)$$

$$= (\delta_{\mathsf{X}(a)} \otimes \delta_{\mathsf{Y}(a)})(B \times C) \quad (149)$$

$$= \mathbb{F}^{\mathsf{X}} \otimes \mathbb{F}^{\mathsf{Y}} \quad (150)$$

Equality follows from the monotone class theorem. \square

Corollary 5.14. *Given a Markov kernel space (\mathbb{K}, Ω, D) and random variables $\mathsf{X} : \Omega \times D \rightarrow X$, $\mathsf{Y} : \Omega \times D \rightarrow Y$, the following holds:*

$$D \text{ --- } \boxed{\mathbb{K}^{\mathsf{X}\mathsf{Y}|D}} \text{ --- } \begin{matrix} X \\ Y \end{matrix} = D \text{ --- } \boxed{\mathbb{K}} \text{ --- } \left(\begin{matrix} \boxed{\mathbb{F}^{\mathsf{X}}} \text{ --- } X \\ \boxed{\mathbb{F}^{\mathsf{Y}}} \text{ --- } Y \end{matrix} \right) \quad (151)$$

We will now define wire labels for “output” wires.

Definition 5.15 (Wire labels - joint kernels). Suppose we have a Markov kernel space (\mathbb{K}, Ω, D) , random variables $\mathsf{X} : \Omega \times D \rightarrow X$, $\mathsf{Y} : \Omega \times D \rightarrow Y$ and a Markov kernel $\mathbb{L} : D \rightarrow \Delta(\mathcal{X} \times \mathcal{Y})$. The following *output labelling* of \mathbb{L} :

$$D \text{ --- } \boxed{\mathbb{L}} \text{ --- } \begin{matrix} \textcolor{blue}{X} \\ \textcolor{blue}{Y} \end{matrix} \quad (152)$$

is *valid* iff

$$\mathbb{L} = \mathbb{K}_{\mathsf{X}\mathsf{Y}|D} \quad (153)$$

and

$$D \text{ --- } \boxed{\mathbb{L}} \text{ --- } \begin{matrix} \textcolor{blue}{X} \\ * \end{matrix} = \mathbb{K}^{\mathsf{X}|D} \quad (154)$$

and

$$D \text{ --- } \boxed{\mathbb{L}} \text{ --- } \begin{matrix} * \\ \textcolor{blue}{Y} \end{matrix} = \mathbb{K}^{\mathsf{Y}|D} \quad (155)$$

The second and third conditions are nontrivial: suppose \mathbf{X} takes values in some product space $\text{Range}(\mathbf{X}) = W \times Z$, and \mathbf{Y} takes values in Y . Then we could have $\mathbb{L} = \mathbb{K}^{\mathbf{X}\mathbf{Y}|\mathbf{D}}$ and draw the diagram

$$D - \boxed{\mathbb{L}} - \begin{array}{c} W \\ Z \times Y \end{array} \quad (156)$$

For *this* diagram, properties 154 and 155 do not hold, even though 153 does.

Lemma 5.16 (Output label assignments exist). *Given Markov kernel space (\mathbb{K}, Ω, D) , random variables $\mathbf{X} : \Omega \times D \rightarrow X$ and $\mathbf{Y} : \Omega \times D \rightarrow Y$ then there exists a diagram of $\mathbb{L} := \mathbb{K}^{\mathbf{X}\mathbf{Y}|\mathbf{D}}$ with a valid output labelling assigning \mathbf{X} and \mathbf{Y} to the output wires.*

Proof. By definition, \mathbb{L} has signature $D \rightarrow \Delta(\mathcal{X} \otimes \mathcal{Y})$. Thus, by the rule that tensor product spaces can be represented by parallel wires, we can draw

$$D - \boxed{\mathbb{L}} - \begin{array}{c} X \\ Y \end{array} \quad (157)$$

By Corollary 5.14, we have

$$D - \boxed{\mathbb{L}} - \begin{array}{c} X \\ Y \end{array} = D - \boxed{\mathbb{K}} - \left(\begin{array}{c} \boxed{\mathbb{F}^{\mathbf{X}}} - X \\ \boxed{\mathbb{F}^{\mathbf{Y}}} - Y \end{array} \right) \quad (158)$$

Therefore

$$D - \boxed{\mathbb{K}} - \left(\begin{array}{c} \boxed{\mathbb{F}^{\mathbf{X}}} - X \\ \boxed{\mathbb{F}^{\mathbf{Y}}} - * \end{array} \right) = \mathbb{K}\mathbb{F}^{\mathbf{X}} \quad (159)$$

$$= \mathbb{K}^{\mathbf{X}|\mathbf{D}} \quad (160)$$

$$D - \boxed{\mathbb{K}} - \left(\begin{array}{c} \boxed{\mathbb{F}^{\mathbf{X}}} - * \\ \boxed{\mathbb{F}^{\mathbf{Y}}} - Y \end{array} \right) = \mathbb{K}\mathbb{F}^{\mathbf{Y}} \quad (161)$$

$$= \mathbb{K}^{\mathbf{Y}|\mathbf{D}} \quad (162)$$

□

In all further work, wire labels will be used without special colouring.

Definition 5.17 (Disintegration). Given a probability space $(\mathbb{P}, \Omega, \mathcal{F})$, and random variables X and Y , we say that $\mathbb{M} : E \rightarrow \Delta(\mathcal{F})$ is a *Y on X disintegration* of \mathbb{P} iff

$$\triangleleft^{\mathbb{P}^{XY}} = \triangleleft^{\mathbb{P}^X} * \boxed{\mathbb{M}} \quad (163)$$

\mathbb{M} is a version of $\mathbb{P}^{Y|X}$, “the probability of Y given X ”. Let $\mathbb{P}^{\{Y|X\}}$ be the set of all kernels that satisfy 163 and $\mathbb{P}^{Y|X}$ an arbitrary member of $\mathbb{P}^{\{Y|X\}}$.

Given a Markov kernel space (\mathbb{K}, Ω, D) and random variables $X : \Omega \times D \rightarrow X$, $Y : \Omega \times D \rightarrow Y$, $\mathbb{M} : D \times E \rightarrow \Delta(\mathcal{F})$ is a *Y on DX disintegration* of $\mathbb{K}^{YX|D}$ iff

$$\boxed{\mathbb{K}^{YX|D}} = \triangleleft^{\mathbb{K}^{YX|D}} * \boxed{\mathbb{M}} \quad (164)$$

Write $\mathbb{K}^{\{Y|XD\}}$ for the set of kernels satisfying 164 and $\mathbb{K}^{Y|XD}$ for an arbitrary member of $\mathbb{K}^{\{Y|XD\}}$.

Definition 5.18 (Wire labels – input). An input wire is *connected* to an output wire if it is possible to trace a path from the start of the input wire to the end of the output wire without passing through any boxes, erase maps or right facing triangles.

If an input wire is connected to an output wire and that output wire has a valid label X , then it is valid to label the input wire with X .

For example, if the following are valid output labels with respect to (\mathbb{P}, Ω) :

$$\triangleleft^{\boxed{\mathbb{L}}} \quad (165)$$

i.e. if $\mathbb{L} \in \mathbb{P}^{\{XY|Y\}}$, then the following is a valid input label:

$$Y \triangleleft^{\boxed{\mathbb{L}}} \quad (166)$$

An input wire in a diagram for \mathbb{M} may be labeled X *if and only if* copy and identity maps can be inserted to yield a diagram in which the input wire labeled X is connected to an output wire with valid label X .

So, if $\mathbb{M} \in \mathbb{P}^{\{X|Y\}}$, then it is straightforward to show that

$$\triangleleft^{\boxed{\mathbb{M}}} \in \mathbb{P}^{\{XY|Y\}} \quad (167)$$

and hence the output labels are valid. Diagram 167 is constructed by taking the product of the copy map with $\mathbb{M} \otimes \mathbf{Id}$. Thus it is valid to label \mathbb{M} with

$$\mathbf{Y} - \boxed{\mathbb{M}} - \mathbf{X} \quad (168)$$

Lemma 5.19 (Labeling of disintegrations). *Given a kernel space (\mathbb{K}, Ω, D) , random variables \mathbf{X} and \mathbf{Y} , domain variable D and disintegration $\mathbb{L} \in \mathbb{K}^{\{\mathbf{Y}|\mathbf{X}D\}}$, there is a diagram of \mathbb{L} with valid input labels \mathbf{X} and \mathbf{D} and valid output label \mathbf{Y} .*

Proof. Note that for any variable $W : \Omega \times D \rightarrow W$ and the domain variable $D : \Omega \times D \rightarrow D$ we have by definition of \mathbb{K} :

$$\boxed{\mathbb{K}^{WD|D}} \begin{array}{c} W \\ D \end{array} = \begin{array}{c} \text{---} \boxed{\mathbb{K}_0} \text{---} \begin{array}{c} \boxed{F^W} \text{---} W \\ \boxed{F^D} \text{---} D \end{array} \end{array} \quad (169)$$

$$= \begin{array}{c} \text{---} \boxed{\mathbb{K}_0} \text{---} \boxed{F^W} \text{---} W \\ \text{---} D \end{array} \quad (170)$$

$$= \begin{array}{c} \text{---} \boxed{\mathbb{K}_0} \text{---} \boxed{F^W} \text{---} W \\ \text{---} D \end{array} \quad (171)$$

$$= \begin{array}{c} \text{---} \boxed{\mathbb{K}} \text{---} \boxed{F^W} \text{---} W \\ \text{---} D \end{array} \quad (172)$$

$$= \begin{array}{c} \text{---} \boxed{\mathbb{K}^{W|D}} \text{---} W \\ \text{---} D \end{array} \quad (173)$$

□

We also use the informal convention of labelling wires in quote marks “ \mathbf{X} ” if that wire is “supposed to” carry the label \mathbf{X} but the label may not be valid.

Theorem 5.20 (Iterated disintegration). *Given a kernel space (\mathbb{K}, Ω, D) , random variables \mathbf{X} , \mathbf{Y} and \mathbf{Z} and domain variable D ,*

$$\begin{array}{c} \text{“}D\text{”} \\ \text{“}X\text{”} \end{array} \begin{array}{c} \text{---} \boxed{\mathbb{K}^{Y|XD}} \text{---} \boxed{\mathbb{K}^{Z|XYD}} \text{---} \text{“}Z\text{”} \\ \text{---} \text{“}Y\text{”} \end{array} \in \mathbb{K}^{\{\mathbf{Z}|\mathbf{Y}|\mathbf{X}D\}} \quad (174)$$

Equivalently, for $d \in D$ and $x \in X$, $A \in \mathcal{Y}$, $B \in \mathcal{Z}$,

$$(d, x; A, B) \mapsto \int_A \mathbb{K}_{(x,y,d)}^{Z|XYD}(B) d\mathbb{K}_{(x,d)}^{Y|XD}(y) \in \mathbb{K}^{\{\mathbf{Z}|\mathbf{Y}|\mathbf{X}D\}} \quad (175)$$

Proof.

write this up

□

The existence of disintegrations of standard measurable probability spaces is well known.

Theorem 5.21 (Disintegration existence - probability space). *Given a probability measure $\mu \in \Delta(\mathcal{X} \otimes \mathcal{Y})$, if (F, \mathcal{F}) is standard then a disintegration $\mathbb{K} : X \rightarrow \Delta(\mathcal{Y})$ exists (Çinlar, 2011).*

In particular, if for all $x \in X$, $\mathbb{P}^X(X \in \{x\}) > 0$, then $\mathbb{P}_x^{Y|X}(Y \in A) = \frac{\mathbb{P}^{XY}(Y \in A \ \& \ X \in \{x\})}{\mathbb{P}^X(X \in \{x\})}$.

For Markov kernel spaces, we make the simplifying assumption that the domain space D is a discrete space. Given this assumption, there exists a positive definite probability $\mu \in \Delta(\mathcal{D})$. That is, for every $d \in D$, $\mu(\{d\}) > 0$. Given this assumption, for every Markov kernel space (\mathbb{K}, Ω, D) there is a probability space $(\mathbb{P}, \Omega \times D)$ such that \mathbb{K} can be uniquely defined as a disintegration of \mathbb{P} . For uncountable D , even if it is standard measurable, this is not possible (Hájek, 2003).

Definition 5.22 (Relative probability space).

better name

Given a Markov kernel space (\mathbb{K}, Ω, D) and a positive definite measure $\mu \in \Delta(\mathcal{D})$, $(\mu\mathbb{K}, \Omega \times D)$ is a *relative probability space*.

For any random variable $X : \Omega \times D \rightarrow X$ on (\mathbb{K}, Ω, D) , its relative on $(\mu\mathbb{K}, \Omega \times D)$ is given by the same measurable function, and we give it the same name X .

Lemma 5.23 (Agreement of disintegrations). *Given a Markov kernel space (\mathbb{K}, Ω, D) , any relative probability space $(\mu\mathbb{K}, \Omega \times D)$ and any random variables $X : \Omega \times D \rightarrow X$, $Y : \Omega \times D \rightarrow Y$, $\mathbb{K}^{\{Y|XD\}} = (\mu\mathbb{K})^{\{Y|XD\}}$ (note that this set equality).*

Proof. Define $\mathbb{P} := \mu\mathbb{K}$ and let \mathbb{M} be an arbitrary version of $\mathbb{K}^{\{Y|XD\}}$. Then

Diagram (176) shows a triangle with \mathbb{P}^{XYD} on the left. Its three output wires are labeled X , Y , and D from top to bottom. This triangle is followed by an equals sign and then a diagram consisting of a small triangle with μ inside, followed by a box labeled $\mathbb{K}^{XY|D}$. The μ triangle has two input wires from the left and one output wire that goes to the $\mathbb{K}^{XY|D}$ box. The $\mathbb{K}^{XY|D}$ box has two input wires (the top one from the μ triangle and the bottom one from the D wire of the first triangle) and two output wires labeled X and Y .

(176)

Diagram (177) shows a small triangle with μ inside, followed by a box labeled $\mathbb{K}^{X|D}$. The μ triangle has two input wires from the left and one output wire that goes to the $\mathbb{K}^{X|D}$ box. The $\mathbb{K}^{X|D}$ box has one input wire (from the μ triangle) and one output wire labeled X . This is followed by an equals sign and then a box labeled \mathbb{M} . The \mathbb{M} box has two input wires (the Y wire from the first triangle and the D wire from the first triangle) and two output wires labeled X and Y .

(177)

Diagram (178) shows a triangle with \mathbb{P}^{XD} on the left. Its three output wires are labeled X , Y , and D from top to bottom. This triangle is followed by an equals sign and then a box labeled \mathbb{M} . The \mathbb{M} box has two input wires (the Y wire and the D wire from the first triangle) and two output wires labeled X and Y .

(178)

Thus $\mathbb{M} \in \mathbb{P}^{\{Y|XD\}}$.

Let \mathbb{N} be an arbitrary version of $\mathbb{P}^{\{Y|XD\}}$. To show that $\mathbb{N} \in \mathbb{K}^{\{Y|XD\}}$, we will show for all $d \in D$

Diagram (179) shows a triangle with δ_d on the left. Its output wire goes to a box labeled $\mathbb{K}^{X|D}$. The $\mathbb{K}^{X|D}$ box has one input wire (from the δ_d triangle) and one output wire labeled X . This is followed by a box labeled \mathbb{N} . The \mathbb{N} box has two input wires (the X wire from the $\mathbb{K}^{X|D}$ box and the D wire from the first triangle) and two output wires labeled Y and X .

(179)

$\mathbb{Q} :=$

$= \mathbb{K}_d^{XYD|D}$

(180)

For $A \in \mathcal{X}, B \in \mathcal{Y}, d \in D$, we have $\mathbb{Q}(A \times B \times \emptyset) = 0 = \mathbb{K}_d^{XYD|D}(A \times B \times \emptyset)$, and for $\{d\} \in \mathcal{D}$ we have $\mu(\{d\}) > 0$ so:

$$\mathbb{Q}(A \times B \times \{d\}) = \int_{X^2} \int_X \int_{D^3} \mathbb{N}_{d'',x'}(A) \mathbf{Id}_{x''}(B) \mathbf{Id}_{d'''}(\{d\}) d\gamma_d(d', d'', d''') d\mathbb{K}_{d'}^{\mathbf{X}|\mathbf{D}}(x) d\gamma_x(x', x'') \quad (181)$$

$$= \delta_d(\{d\}) \int_X \mathbb{N}_{d,x}(A) \delta_x(B) d\mathbb{K}_d^{\mathbf{X}|\mathbf{D}}(x) \quad (182)$$

$$= \frac{1}{\mu(\{d\})} \int_{\{d\}} d\mu(d') \int_X \mathbb{N}_{d,x}(A) \delta_x(B) d\mathbb{K}_d^{\mathbf{X}|\mathbf{D}}(x) \quad (183)$$

$$= \frac{1}{\mu(\{d\})} \int_D \int_X \mathbb{N}_{d,x}(A) \delta_{d'}(\{d\}) \delta_x(B) d\mathbb{K}_d^{\mathbf{X}|\mathbf{D}}(a) d\mu(d') \quad (184)$$

$$= \frac{1}{\mu(\{d\})} \int_D \int_X \mathbb{N}_{d,x}(A) \delta_{d'}(\{d\}) \delta_x(B) d\mathbb{K}_{d'}^{\mathbf{X}|\mathbf{D}}(a) d\mu(d') \quad (185)$$

$$= \frac{1}{\mu(\{d\})} \mathbb{P}^{\mathbf{XYD}}(A \times B \times \{d\}) \quad (186)$$

$$= \frac{1}{\mu(\{d\})} \int_D \mathbb{K}_{d'}^{\mathbf{XYD}|\mathbf{D}}(A \times B \times \{d\}) d\mu(d') \quad (187)$$

$$= \frac{1}{\mu(\{d\})} \int_D \mathbb{K}_{d'}^{\mathbf{XY}|\mathbf{D}}(A \times B) \delta_{d'}(\{d\}) d\mu(d') \quad (188)$$

$$= \mathbb{K}_d^{\mathbf{XY}|\mathbf{D}}(A \times B) \quad (189)$$

$$= \mathbb{K}_d^{\mathbf{XY}|\mathbf{D}}(A \times B) \delta_d(\{d\}) \quad (190)$$

$$= \int_D \mathbb{K}_{d'}^{\mathbf{XY}}(A \times B) \delta_{d''}(\{d\}) d\gamma_d(d', d'') \quad (191)$$

$$= \mathbb{K}_d^{\mathbf{XYD}|\mathbf{D}}(A \times B \times \{d\}) \quad (192)$$

Equality follows from the monotone class theorem. Thus $\mathbb{N} \in \mathbb{K}^{\{\mathbf{Y}|\mathbf{XD}\}}$. \square

Thus any kernel conditional probability $\mathbb{K}^{\mathbf{Y}|\mathbf{XD}}$ can equally well be considered a regular conditional probability $\mathbb{P}^{\mathbf{Y}|\mathbf{XD}}$ for a related probability space $(\mathbb{P}, \Omega \times D)$ under the obvious identification of random variables, provided D is countable. Note that any conditional probability $\mathbb{P}^{\mathbf{Y}|\mathbf{X}}$ that is *not* conditioned on \mathbf{D} is undefined in the kernel space (\mathbb{K}, Ω, D) .

5.5.1 Conditional Independence

Definition 5.24 (Kernels constant in an argument). Given a kernel (\mathbb{K}, Ω, D) and random variables \mathbf{Y} and \mathbf{X} , we say a version of the disintegration $\mathbb{K}^{\mathbf{Y}|\mathbf{XD}}$ is constant in \mathbf{D} if for all $x \in X$, $d, d' \in D$, $\mathbb{K}_{(x,d)}^{\mathbf{Y}|\mathbf{XD}} = \mathbb{K}_{(x,d')}^{\mathbf{Y}|\mathbf{XD}}$.

Definition 5.25 (Domain Conditional Independence). Given a kernel space (\mathbb{K}, Ω, D) , relative probability space $(\mathbb{P}, \Omega \times D)$, variables \mathbf{X}, \mathbf{Y} and domain variable \mathbf{D} , \mathbf{X} is *conditionally independent* of \mathbf{D} given \mathbf{Y} , written $\mathbf{X} \perp_{\mathbb{K}} \mathbf{D} | \mathbf{Y}$ if any of the following equivalent conditions hold:

Almost sure equality

- $\mathbb{P}^{XD|Y} \sim \mathbb{P}^{X|Y} \underline{\otimes} \mathbb{P}^{D|Y}$
- For any version of $\mathbb{P}^{X|Y}$, $\mathbb{P}^{X|Y} \otimes *_D$ is a version of $\mathbb{K}^{X|YD}$
- There exists a version of $\mathbb{K}^{X|YD}$ constant in D

Theorem 5.26 (Definitions are equivalent). *(1) \implies (2): By Lemma 5.23, $\mathbb{P}^{Y|XD} = \mathbb{K}^{Y|XD}$. Thus it is sufficient to show that $\mathbb{P}^{X|Y} \otimes *$ is a version of $\mathbb{P}^{X|YD}$.*

$$(193)$$

$$(194)$$

$$(195)$$

$$(196)$$

(2) \implies (3)
 $\mathbb{P}^{X|Y} \otimes *_D$ is a version of $\mathbb{K}^{X|YD}$ by assumption, and is clearly constant in D .

(3) \implies (1)
 By lemma 5.23, there also exists a version of $\mathbb{P}^{X|YD}$ constant in D . Let $\mathbb{M} : Y \times D \rightarrow \Delta(\mathcal{X})$ be such a version. For arbitrary $d_0 \in D$, let $\mathbb{N} := \mathbb{M}_{(\cdot, d_0)} : Y \rightarrow \Delta(\mathcal{X})$ be the map $x \mapsto \mathbb{M}_{(x, d_0)}$. By constancy in D , $\mathbb{M} = * \otimes \mathbb{N}$. We wish to show $\mathbb{P}^{X|Y} \underline{\otimes} \mathbb{P}^{D|Y} \in \mathbb{P}^{XD|Y}$. By Theorem 5.20, we have

$$(197)$$

Definition 5.27 (Conditional probability existence). Given a kernel space (\mathbb{K}, Ω, D) and random variables X, Y , we say $\mathbb{K}^{\{Y|X\}}$ exists if $Y \perp\!\!\!\perp_{\mathbb{K}} D|X$. If $\mathbb{K}^{\{Y|X\}}$ exists then it is by definition equal to $\mathbb{P}^{\{Y|X\}}$ for any related probability space $(\mathbb{P}, \Omega \times D)$.

Note that $\mathbb{K}^{\{Y|XD\}}$ always exists.

Definition 5.28 (Conditional Independence). Given a kernel space (\mathbb{K}, Ω, D) , relative probability space $(\mathbb{P}, \Omega \times D)$, variables X, Y and Z , X is *conditionally independent* of Z given Y , written $X \perp\!\!\!\perp_{\mathbb{K}} Z|Y$ if $\mathbb{K}^{\{XY|Z\}}$ exists and any of the following equivalent conditions hold:

Almost sure equality

- $\mathbb{P}^{XZ|Y} \sim \mathbb{P}^{X|Y} \otimes \mathbb{P}^{Z|Y}$
- For any version of $\mathbb{P}^{\{X|Y\}}$, $\mathbb{P}^{X|Y} \otimes *_Z$ is a version of $\mathbb{K}^{\{X|YZ\}}$
- There exists a version of $\mathbb{K}^{\{X|YZ\}}$ constant in Z

Lemma 5.29 (Diagrammatic consequences of labels). *In general, diagram labels are “well behaved” with regard to the application of any of the special Markov kernels: identities 106, swaps 119, discards 122 and copies 114 as well as with respect to the coherence theorem of the CD category. They are not “well behaved” with respect to composition.*

Fix some Markov kernel space (\mathbb{K}, Ω, D) and random variables X, Y, Z taking values in X, Y, Z respectively. $\text{Sat} :$ indicates that a labeled diagram satisfies definitions 5.15 and 5.18 with respect to (\mathbb{K}, Ω, D) and X, Y, Z . The following always holds:

$$\text{Sat} : X - X \quad (198)$$

and the following implications hold:

$$\text{Sat} : \begin{array}{c} Z - \boxed{\mathbb{K}} - Y \\ \diagup \quad \diagdown \\ X \end{array} \implies \text{Sat} : \begin{array}{c} Z - \boxed{\mathbb{K}} - X \\ \diagup \quad \diagdown \\ * \end{array} \quad (199)$$

$$\text{Sat} : \begin{array}{c} Z - \boxed{\mathbb{K}} - Y \\ \diagup \quad \diagdown \\ X \end{array} \implies \text{Sat} : \begin{array}{c} Z - \boxed{\mathbb{K}} - X \\ \diagup \quad \diagdown \\ Y \end{array} \quad (200)$$

$$\text{Sat} : Z - \boxed{\mathbb{L}} - X \implies \text{Sat} : \begin{array}{c} Z - \boxed{\mathbb{L}} - \text{ } \\ \diagup \quad \diagdown \\ X \quad X \end{array} \quad (201)$$

$$\text{Sat} : Z - \boxed{\mathbb{K}} - Y \implies \text{Sat} : \begin{array}{c} \text{ } \\ \diagup \quad \diagdown \\ Z \quad \boxed{\mathbb{K}} - Y \end{array} \quad (202)$$

Proof. • Id_X is a version of $\mathbb{P}_{X|X}$ for all \mathbb{P} ; $\mathbb{P}_X \text{Id}_X = \mathbb{P}_X$

$$\bullet \mathbb{K} \text{Id} \otimes *) (w; A) = \int_{X \times Y} \delta_x(A) \mathbb{1}_Y(y) d\mathbb{K}_w(x, y) = \mathbb{K}_w(A \times Y) = \mathbb{P}_{X|Z}(w; A)$$

- $\int_{X \times Y} \delta_{\text{swap}(x,y)}(A \times B) d\mathbb{K}_w(x, y) = \mathbb{P}_{Y|Z}(w; A \times B)$
 - $\mathbb{K} \curlyvee(w; A \times B) = \int_X \delta_{x,x}(A \times B) d\mathbb{K}_w(x) = \mathbb{P}_{X|Z}(w; A \times B)$
- 202: Suppose \mathbb{K} is a version of $\mathbb{P}_{Y|Z}$. Then

$$\mathbb{P}_{ZY} = \text{Diagram: A triangle labeled } \mathbb{P}_Z \text{ with a wire from its top to a box labeled } \mathbb{K} \text{, which then has a wire to } Y \text{, and a curved wire from the top of } \mathbb{K} \text{ to } Z. \quad (203)$$

$$\mathbb{P}_{ZZY} = \text{Diagram: A triangle labeled } \mathbb{P}_Z \text{ with a wire from its top to a box labeled } \mathbb{K} \text{, which then has a wire to } Y \text{, and a curved wire from the top of } \mathbb{K} \text{ to } Z. \text{ There is an additional wire from } Z \text{ to } Y. \quad (204)$$

$$= \text{Diagram: A triangle labeled } \mathbb{P}_Z \text{ with a wire from its top to a box labeled } \mathbb{K} \text{, which then has a wire to } Y \text{, and a curved wire from the top of } \mathbb{K} \text{ to } Z. \text{ There is an additional wire from } Z \text{ to } Y. \quad (205)$$

Therefore $\curlyvee(\text{Id}_X \otimes \mathbb{K})$ is a version of $\mathbb{P}_{ZY|Z}$ by ?? □

The following property, on the other hand, does *not* generally hold:

$$\text{Sat} : Z - \boxed{\mathbb{K}} - Y, Y - \boxed{\mathbb{L}} - X \implies \text{Sat} : Z - \boxed{\mathbb{K}} - \boxed{\mathbb{L}} - X \quad (206)$$

Consider some ambient measure \mathbb{P} with $Z = X$ and $\mathbb{P}_{Y|X} = x \mapsto \text{Bernoulli}(0.5)$ for all $z \in Z$. Then $\mathbb{P}_{Z|Y} = y \mapsto \mathbb{P}_Z, \forall y \in Y$ and therefore $\mathbb{P}_{Y|Z} \mathbb{P}_{Z|Y} = x \mapsto \mathbb{P}_Z$ but $\mathbb{P}_{Z|X} = x \mapsto \delta_x \neq \mathbb{P}_{Y|Z} \mathbb{P}_{Z|Y}$.

This is an attempt to understand which assumptions lead to see-do models being represented by probability distributions.

6 Decision makers and decision models

We will consider a very general type of decision maker to motivate the construction of decision models. A decision maker takes some kind of evidence - for example, data, prior knowledge or a problem specification - and determines the consequences it believes each decision will have via some *inference rule*. It then examines the consequences of each decision and chooses a preferred decision via some *choice rule*.

More formally, let the set A be a set of possible pieces of evidence a decision maker could obtain, E be the set of consequences it may experience and D the set of decisions it may take. Call any function $C : D \rightarrow E$ a *consequence map*. A *inference rule* is a function $f : A \rightarrow E^D$ that takes a piece of evidence and returns a consequence map. A *choice rule* $\text{ch} : E^D \rightarrow D$ is a function that takes a consequence map and returns a preferred decision. A decision maker is specified by a particular choice of f and ch and implements the *decision function* $h := \text{ch} \circ f$.

Suppose $D = \{0, 1\}$ represents the decision to turn some switch to the on or off positions. Some examples of consequence maps follow:

- $E = \{0, 1\}$ is the state of a light connected to the switch and $\mathbf{C} = \text{Id}_{\{0,1\}}$ is the consequence function that implies the light is always in the same state as the switch
- $E = \{\{0\}, \{1\}, \{0, 1\}\}$ represents sets of possible outcomes and $\mathbf{C} : 0 \mapsto \{0\}, 1 \mapsto \{0, 1\}$ is the consequence function that implies the light is always off when the switch is off, but may take either state with the switch on (i.e. "the bulb may be out")

If $A = \{0, 1\}^{2N}$ is a sequence of joint states of the switch and lightbulb, then an inference function could be the map

$$f(s_1, l_1, \dots, s_N, l_N) = \begin{cases} 0 \mapsto \mathbb{I} \left[\frac{\sum_{i \in N} l_i(1-s_i)}{\sum_{i \in N} (1-s_i)} > 0.5 \right] \\ 1 \mapsto \mathbb{I} \left[\frac{\sum_{i \in N} l_i s_i}{\sum_{i \in N} s_i} > 0.5 \right] \end{cases} \quad (207)$$

That is, f returns the consequence function that maps each switch position to the lightbulb state more commonly observed in conjunction with that switch position in the given data.

Some examples of decision rules, where in each case $\mathbf{C} : D \rightarrow E$ is some consequence map:

- $E = \{\text{off}, \text{on}\}$ are states of the light bulb, $u : \text{off} \mapsto 0, \text{on} \mapsto 1$ is a utility function and $\text{ch}(\mathbf{C}) = \arg \min_{d \in D} u(\mathbf{C}(d))$
- $E = [0, 1]$ is a set of losses and $\text{ch}(\mathbf{C}) = \arg \min_{d \in D} (\mathbf{C}(d))$
- $E = \{\{0\}, \{1\}, \{0, 1\}\}$ are sets of possible outcomes and ch is the minimax operator: $\text{ch}(\mathbf{C}) = \arg \min_{d \in D} \max_{e \in \mathbf{C}(d)} (e)$

Another example of a decision maker is a learning algorithm that produces binary classifiers. A binary classifier takes some piece of data X and returns a class in $\{0, 1\}$. That is, the set of available decisions D is the set of functions from $X \rightarrow \{0, 1\}$. Suppose also that the given information is a set of N labeled examples $A = (X \times \{0, 1\})^N$. Then the inference function f might be an *empirical risk* functional that takes data $a \in A$ and a possible classifier $d \in D$ and returns the number of misclassified examples when d is applied to A . In such a case, ch might be the *argmin* functional, making our decision maker an *empirical risk minimiser*. This is not necessarily a good inference function - typically we are interested in the number of misclassified examples a classifier will produce on unseen data, not the number of misclassified examples it will produce on the data that has already been seen.

6.1 Expected utility maximisers

Expected utility maximisation (henceforth "EU") necessitates that the class of results models f have a particular signature. The EU choice rule compares a set of probability distributions over states of the world E and returns a decision associated with the preferred probability distribution. Thus, denoting by $\Delta(\mathcal{E})$

the set of probability distribution over E (now equipped with σ -algebra \mathcal{E}), f must have the signature $f : A \rightarrow \Delta(\mathcal{E})^D$. In particular f returns stochastic functions of the type $D \rightarrow \Delta(\mathcal{E})$.

So far we have not made any structural assumptions about the set D . One such assumption that we do want to make is that stochastic decisions are possible. That is, if we can choose $d_1 \in D$ and we can choose $d_2 \in D$, then it is also possible to choose d_1 with probability α and d_2 with probability $(1 - \alpha)$ for $0 \leq \alpha \leq 1$. We will make this assumption as follows: D represents an underlying set of “elementary” decisions, and the true set of decisions is the set $\Delta(\mathcal{D})$ of probability measures on D (equipped with some σ -algebra \mathcal{D}). Then, for example, if $D = \{d_1, d_2\}$ we can “choose” d_1 by selecting the measure δ_{d_1} , and we can choose a mixture of d_1 and d_2 by selecting the measure $\alpha\delta_{d_1} + (1 - \alpha)\delta_{d_2}$. Technically, all that we have done so far is assume that the set of decisions is isomorphic to $\Delta(\mathcal{D})$ - which, if D is standard Borel, ensures that the set of decisions is convex closed. In many cases, we also want the following to hold:

$$f_a(\alpha\delta_{d_1} + (1 - \alpha)\delta_{d_2}) = \alpha f_a(\delta_{d_1}) + (1 - \alpha)f_a(\delta_{d_2}) \quad (208)$$

That is, we assume that the results model f is *additive*. Informally, choosing to do d_1 with probability α and d_2 with probability $1 - \alpha$ will yield the result of d_1 with probability α and the result of d_2 with probability $1 - \alpha$. In addition, f_a must be continuous with respect to the total variation norm.

For arbitrary $\gamma \in \Delta(\mathcal{D})$, we can write for all $B \in \mathcal{D}$

$$\gamma(B) = \int_D \delta_d(B) d\gamma(d) \quad (209)$$

Can we also say

$$\gamma = \int_D \delta_d d\gamma(d) \quad (210)$$

For $G \in \mathcal{E}$, write $f_a(\delta_d; C)$ for the probability measure $f_a(\delta_d)$ evaluated at C , and $f_a(\cdot; G)$ for the map $d \mapsto f_a(d; C)$. If $f_a(\cdot; C)$ is measurable for all C and additive, then it is a property of the Lebesgue integral that for all $\gamma \in \Delta(\mathcal{D})$

$$\int f_a(\gamma; C) = \int f_a\left(\int_D \delta_{d'} d\gamma(d'); C\right) \quad (211)$$

$$= \int_D f_a(\delta_{d'}; C) d\gamma(d') \quad (212)$$

Define $\mathcal{C}_a : D \rightarrow \Delta(\mathcal{E})$ by $\mathcal{C}_a : d \mapsto f_a(\delta_d; \cdot)$. Then $\mathcal{C}_a(\cdot; B)$ is measurable and $\mathcal{C}_a(d; \cdot)$ is a probability measure - that is, \mathcal{C}_a is a Markov kernel. We can therefore adopt the product notation defined elsewhere to write the consequences of choosing a distribution γ as $\gamma\mathcal{C}_a$.

Have I made any assumptions here besides additivity, convex closedness of D ?

As we are chiefly interested in the set of elementary decisions D' , we will henceforth simply use (D, \mathcal{D}) for this set.

Assumption 208 still permits some unexpected behaviour with respect to randomised decisions. Consider the following variations of a problem:

Example 6.1 (Fighting children problem). A decision maker is mediating a dispute over a toy between two children. If he chooses to give the toy to the first child, the second child will cry, and if he chooses to give the toy to the second child, the first child will cry (the underlying decision set $D' = \{\text{give to first child, give to second child}\}$). If he chooses a random procedure:

Version 1 These children strongly object to randomness, and both will cry no matter the outcome

Version 2 These are children with a strong insistence on a version of procedural fairness which no doubt make a great deal of sense to them. In particular, if a procedure is used which gives the first child a chance p of receiving the toy, the first child will cry with probability p irrespective of the outcome. The second child, meanwhile, will cry iff the first child does not

Version 3 Whether or not the children cry depends only on whether or not they receive the toy

In version 1, assumption 208 is violated - the outcome when choosing a random procedure is not simply a mixture of the outcomes of each deterministic decision. In version 3, assumption 208 holds, and the results depend only on the outcome of randomisation, as is desired. In version 2, assumption 208 *also* holds - one child cries iff the other does not, and each child cries with the same probability as the probability that they receive the toy. However, the consequences of randomising and then giving the first child the toy are not the same as simply giving the first child the toy. To discount the possibility of this kind of behaviour, we need additional assumptions.

In the text of this example, we have spoken as if our decision had multiple consequences - whether or not each kid gets a toy, whether or not each kid cries - but the compliance of version 2 with 208 depends on considering only the children crying to be a consequence of our decision. To support the intuition that only version 3 is the kind of problem we want to deal with, we introduce the additional assumption that if we choose an extreme decision δ_d , then one consequence of this decision is that we will have chosen d with probability 1.

Formally, we assume the sample space can be written as $E \times D$ for some E , and, defining the random variable $D : E \times D \rightarrow D$ by the projection $D : (e, d) \mapsto d$, suppose

$$(\mathcal{C}_{(a,d)})_{\mathcal{D}} = \delta_d \quad (213)$$

We find in Joyce (2000) the assumption of a “supposition function” $\text{prob}(\cdot\|\cdot)$ that $\text{prob}(C\|C) = 1$ where C is “some distinguished condition”. Assumption 213 along with assumption 208 implies an analogous condition.

Theorem 6.2 ([Decision determinism implies decision certainty]).

That is, for any stochastic decision γ , the consequence conditional on \mathcal{D} is the consequence map f_a itself. This assumption rules out both versions 1 and 2 of the fighting children problem above - it assumes that if we randomise and the result of randomisation is some $d \in D$, this is the same as having chosen d to begin with.

Lemma 6.3 (Conditional agreement implies randomised decisions). *If we have*

$$(\gamma f_a)_{|\mathcal{D}} = f_a \quad (214)$$

Then for all $\gamma \in \Delta(\mathcal{D})$,

$$\gamma f_a(G) = \int_D f_a(d'; G) d\gamma(d') \quad (215)$$

Proof. By definition, for $\gamma \in \Delta(\mathcal{D})$ □

A consequence of assumption ?? is that for fixed $a \in A$, every stochastic decision $\gamma \in \Delta(\mathcal{D})$ leads to the same conditional probability $(\gamma f_a)_{|\mathcal{D}}$. In fact, as a result of the assumption that the codomain of f is a set of probability measures as well as assumptions 213 and ?? we find that the tuple $\langle E \times D, \mathcal{E} \otimes \mathcal{D}, \mathcal{D}, f_a \rangle$ for each $a \in A$ forms a *conditional probability space* (Rényi, 1955).

For our causal analysis we work with *subjunctive probability spaces*, a generalisation of the more familiar probability spaces. The subjunctive mood is used to describe hypothetical or supposed states of the world, and subjunctive probability spaces are models that ask for some hypothetical state and give us back a probability space. Subjunctive probability spaces are also different to *conditional probability spaces* Rényi (1956) as *hypothesising* or *supposing* (that is, those things described by the subjunctive mood) are different to *conditioning*, which is more closely analogous to *focussing your attention*.

We use subjunctive probability spaces because *supposition* is a core part of decision problems, one we cannot get away from. In contrast, modelling supposition with conditional probability (which would be necessary if we were to insist on using probability spaces) adds additional structure to our models which isn't clearly warranted and is potentially confusing.

Any decision problem must involve the comparison of different decisions. This comparison takes the form

- *Suppose* I choose the first decision - then the consequence would be X

Does this definition need to be here? I didn't know what this word meant before I looked it up

- Suppose I choose the second decision - then the consequence would be Y
- Etc.

If we prefer X to Y , then perhaps we should choose the first decision. We may be ambivalent as to what type of thing X and Y represent, how we ought to determine what values X and Y take or how we ought to determine what is preferable, but it is hard to do away with supposing that different decisions were taken and that these decisions come with their own consequences and still have what could reasonably be considered a decision problem. This process of supposition implicitly invokes a “subjunctive function” - we provide a hypothetical decision, and we are given a consequence of that hypothetical. Of particular interest are models that, for each hypothetical choice, return a probability distribution over space Ω . Such models are called *supposition functions* by Joyce (2000), but we will call them *consequence maps* in this work. We consider this particular type of subjunctive function - from possible decisions to probability distributions - to be attractive because we consider probability to be a sound choice for modelling uncertainty and stochasticity.

Formally, a consequence map \mathcal{C} is a stochastic function or *Markov kernel* from $D \rightarrow \Delta(\Omega)$, where $\Delta(\Omega)$ represents the set of all probability distributions on Ω . One might recall that, given two random variables $Y : \Omega \rightarrow Y$ and $X : \Omega \rightarrow X$ on some probability space $(\mathbb{P}, \Omega, \mathcal{F})$, the probability of Y conditional on X is a Markov kernel $X \rightarrow \Delta(\mathcal{Y})$. It is possible, in general, to define a probability distribution on the expanded space $\Omega \times D$ such that \mathcal{C} is a conditional probability. However, there are good reasons to keep the concepts of consequence maps and conditional probability separate. Firstly, there are technical issues such as the fact that it is not always possible to find a distribution on $\Omega \times D$ that yields \mathcal{C} as a *unique* conditional probability (Hájek, 2003) (though this requires an uncountable set of possible decisions, which is not a problem we consider in this work). Secondly, it is not clear that the way we ought to handle consequence maps is identical to the way we handle conditional probabilities. Consider an expanded set of options:

1. Suppose I choose the first decision d_1 - then the consequence would be P_1
2. Suppose I choose the second decision d_2 - then the consequence would be P_2
3. Suppose I choose either the first or second decision d_1 or d_2 - then the consequence would be ???
4. Suppose I choose d_1 with probability $0 \leq q \leq 1$ and d_2 otherwise - then the consequence would be ???

If we regard the consequence map as a conditional probability then it would be natural to consider the result of the third option to be a unique probability distribution obtained by conditioning on $\{d_1, d_2\}$, equal to $\alpha P_1 + (1 - \alpha)P_2$ for some fixed $0 \leq \alpha \leq 1$. However, there is no obvious reason that these should

be mixed in some fixed proportion α - it seems more appropriate to me to say that if d_1 or d_2 is chosen then the result will be P_1 or P_2 .

On the other hand, the result of the fourth option seems like it should be given by $qP_1 + (1 - q)P_2$, at least for most ordinary problems. If we were confronted with a mind reader who could tell the difference between us having chosen d_1 and us having randomised between d_1 and d_2 but come up with d_1 anyway then we might have reason to revise this assumption, but we will generally proceed under the assumption that such mind readers are absent. For any ambient probability measure, we can choose q not to be equal to α (as defined in the previous paragraph), and so the result will not be equal to the ambient measure conditioned on $\{d_1, d_2\}$.

In general, choosing the consequence map to be a conditional probability requires there to exist some joint distribution over the decision D and all other random variables in the problem. However, when we adopt hypotheses about what decisions we might make, we throw away key parts of this joint distribution (for example, we throw away the marginal distribution of D) and replace it with whatever we want to suppose instead. Instead of adopting a full joint distribution and then throwing away some parts to get hold of the consequence map, as we would if we were working with a probability space, a subjunctive probability space only supplies those parts which we intend to keep - i.e. in only supplies the consequence map.

References

- Stephan Bongers, Jonas Peters, Bernhard Schölkopf, and Joris M. Mooij. Theoretical Aspects of Cyclic Structural Causal Models. *arXiv:1611.06221 [cs, stat]*, November 2016. URL <http://arxiv.org/abs/1611.06221>. arXiv: 1611.06221.
- Erhan Çinlar. *Probability and Stochastics*. Springer, 2011.
- Kenta Cho and Bart Jacobs. Disintegration and Bayesian inversion via string diagrams. *Mathematical Structures in Computer Science*, 29(7):938–971, August 2019. ISSN 0960-1295, 1469-8072. doi: 10.1017/S0960129518000488.
- Florence Clerc, Fredrik Dahlqvist, Vincent Danos, and Ilias Garner. Pointless learning. *20th International Conference on Foundations of Software Science and Computation Structures (FoSSaCS 2017)*, March 2017. doi: 10.1007/978-3-662-54458-7_21. URL [https://www.research.ed.ac.uk/portal/en/publications/pointless-learning\(694fb610-69c5-469c-9793-825df4f8ddec\).html](https://www.research.ed.ac.uk/portal/en/publications/pointless-learning(694fb610-69c5-469c-9793-825df4f8ddec).html).
- A. P. Dawid. Influence Diagrams for Causal Modelling and Inference. *International Statistical Review*, 70(2):161–189, 2002. ISSN 1751-5823. doi: 10.1111/j.1751-5823.2002.tb00354.x. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1751-5823.2002.tb00354.x>.

- A. Philip Dawid. Decision-theoretic foundations for statistical causality. *arXiv:2004.12493 [math, stat]*, April 2020. URL <http://arxiv.org/abs/2004.12493>. arXiv: 2004.12493.
- Angus Deaton and Nancy Cartwright. Understanding and misunderstanding randomized controlled trials. *Social Science & Medicine*, 210:2–21, August 2018. ISSN 0277-9536. doi: 10.1016/j.socscimed.2017.12.005. URL <http://www.sciencedirect.com/science/article/pii/S0277953617307359>.
- Ronald A. Fisher. Cancer and Smoking. *Nature*, 182(4635):596–596, August 1958. ISSN 1476-4687. doi: 10.1038/182596a0. URL <https://www.nature.com/articles/182596a0>. Number: 4635 Publisher: Nature Publishing Group.
- Brendan Fong. Causal Theories: A Categorical Perspective on Bayesian Networks. *arXiv:1301.6201 [math]*, January 2013. URL <http://arxiv.org/abs/1301.6201>. arXiv: 1301.6201.
- D. Heckerman and R. Shachter. Decision-Theoretic Foundations for Causal Reasoning. *Journal of Artificial Intelligence Research*, 3:405–430, December 1995. ISSN 1076-9757. doi: 10.1613/jair.202. URL <https://www.jair.org/index.php/jair/article/view/10151>.
- James J. Heckman. Randomization and Social Policy Evaluation. SSRN Scholarly Paper ID 995151, Social Science Research Network, Rochester, NY, July 1991. URL <https://papers.ssrn.com/abstract=995151>.
- M. A. Hernán and S. L. Taubman. Does obesity shorten life? The importance of well-defined interventions to answer causal questions. *International Journal of Obesity*, 32(S3):S8–S14, August 2008. ISSN 1476-5497. doi: 10.1038/ijo.2008.82. URL <https://www.nature.com/articles/ijo200882>.
- Alan Hájek. What Conditional Probability Could Not Be. *Synthese*, 137(3): 273–323, December 2003. ISSN 1573-0964. doi: 10.1023/B:SYNT.0000004904.91112.16. URL <https://doi.org/10.1023/B:SYNT.0000004904.91112.16>.
- Alan Hájek. Interpretations of Probability. In Edward N. Zalta, editor, *The Stanford Encyclopedia of Philosophy*. Metaphysics Research Lab, Stanford University, fall 2019 edition, 2019. URL <https://plato.stanford.edu/archives/fall2019/entries/probability-interpret/>.
- James M. Joyce. Why We Still Need the Logic of Decision. *Philosophy of Science*, 67:S1–S13, 2000. ISSN 0031-8248. URL www.jstor.org/stable/188653.
- Chayakrit Krittanawong, Bharat Narasimhan, Zhen Wang, Joshua Hahn, Hafeez Ul Hassan Virk, Ann M. Farrell, HongJu Zhang, and WH Wilson Tang. Association between chocolate consumption and risk of coronary artery disease: a systematic review and meta-analysis. *European Journal of Preventive Cardiology*, July 2020. doi: 10.1177/2047487320936787. URL

- <http://journals.sagepub.com/doi/10.1177/2047487320936787>. Publisher: SAGE PublicationsSage UK: London, England.
- Finnian Lattimore and David Rohde. Replacing the do-calculus with Bayes rule. *arXiv:1906.07125 [cs, stat]*, December 2019. URL <http://arxiv.org/abs/1906.07125>. arXiv: 1906.07125.
- David K Lewis. Causation. *Journal of Philosophy*, 1986.
- Stephen L. Morgan and Christopher Winship. *Counterfactuals and Causal Inference: Methods and Principles for Social Research*. Cambridge University Press, New York, NY, 2 edition edition, November 2014. ISBN 978-1-107-69416-3.
- Naomi Oreskes and Erik M. Conway. *Merchants of Doubt: How a Handful of Scientists Obscured the Truth on Issues from Tobacco Smoke to Climate Change: How a Handful of Scientists ... Issues from Tobacco Smoke to Global Warming*. Bloomsbury Press, New York, NY, June 2011. ISBN 978-1-60819-394-3.
- Judea Pearl. *Causality: Models, Reasoning and Inference*. Cambridge University Press, 2 edition, 2009.
- Judea Pearl. Challenging the hegemony of randomized controlled trials: A commentary on Deaton and Cartwright. *Social Science & Medicine*, 2018a.
- Judea Pearl. Does Obesity Shorten Life? Or is it the Soda? On Non-manipulable Causes. *Journal of Causal Inference*, 6(2), 2018b. doi: 10.1515/jci-2018-2001. URL <https://www.degruyter.com/view/j/jci.2018.6.issue-2/jci-2018-2001/jci-2018-2001.xml>.
- Judea Pearl and Dana Mackenzie. *The Book of Why: The New Science of Cause and Effect*. Basic Books, New York, 1 edition edition, May 2018. ISBN 978-0-465-09760-9.
- Jonas Peters, Dominik Janzing, and Bernard Schölkopf. *Elements of Causal Inference*. MIT Press, 2017.
- Robert N. Proctor. The history of the discovery of the cigarettelung cancer link: evidentiary traditions, corporate denial, global toll. *Tobacco Control*, 21(2):87–91, March 2012. ISSN 0964-4563, 1468-3318. doi: 10.1136/tobaccocontrol-2011-050338. URL <https://tobaccocontrol.bmj.com/content/21/2/87>. Publisher: BMJ Publishing Group Ltd Section: The shameful past.
- Thomas S Richardson and James M Robins. Single world intervention graphs (swigs): A unification of the counterfactual and graphical approaches to causality. *Center for the Statistics and the Social Sciences, University of Washington Series. Working Paper*, 128(30):2013, 2013.

- Donald B. Rubin. Causal Inference Using Potential Outcomes. *Journal of the American Statistical Association*, 100(469):322–331, March 2005. ISSN 0162-1459. doi: 10.1198/016214504000001880. URL <https://doi.org/10.1198/016214504000001880>.
- A. Rényi. On Conditional Probability Spaces Generated by a Dimensionally Ordered Set of Measures. *Theory of Probability & Its Applications*, 1(1):55–64, January 1956. ISSN 0040-585X. doi: 10.1137/1101005. URL <https://epubs.siam.org/doi/abs/10.1137/1101005>.
- Alfréd Rényi. On a new axiomatic theory of probability. *Acta Mathematica Academiae Scientiarum Hungarica*, 6(3):285–335, September 1955. ISSN 1588-2632. doi: 10.1007/BF02024393. URL <https://doi.org/10.1007/BF02024393>.
- Peter Selinger. A survey of graphical languages for monoidal categories. *arXiv:0908.3347 [math]*, 813:289–355, 2010. doi: 10.1007/978-3-642-12821-9_4. URL <http://arxiv.org/abs/0908.3347>. arXiv: 0908.3347.
- Ilya Shpitser and Judea Pearl. Complete Identification Methods for the Causal Hierarchy. *Journal of Machine Learning Research*, 9(Sep):1941–1979, 2008. ISSN 1533-7928. URL <https://www.jmlr.org/papers/v9/shpitser08a.html>.
- Statista. Cigarettes - worldwide | Statista Market Forecast, 2020. URL <https://www.statista.com/outlook/50010000/100/cigarettes/worldwide>.
- Robert Wiblin. Why smoking in the developing world is an enormous problem and how you can help save lives, 2016. URL <https://80000hours.org/problem-profiles/tobacco/>.
- James Woodward. Causation and Manipulability. In Edward N. Zalta, editor, *The Stanford Encyclopedia of Philosophy*. Metaphysics Research Lab, Stanford University, winter 2016 edition, 2016. URL <https://plato.stanford.edu/archives/win2016/entries/causation-mani/>.
- World Health Organisation. Tobacco Fact sheet no 339, 2018. URL <https://www.webcitation.org/6gUXrCDKA>.
- Karren Yang, Abigail Katoff, and Caroline Uhler. Characterizing and Learning Equivalence Classes of Causal DAGs under Interventions. In *International Conference on Machine Learning*, pages 5537–5546, July 2018. URL <http://proceedings.mlr.press/v80/yang18a.html>.

Appendix: