

# iPiano: Inertial Proximal Algorithm for Non-Convex Optimization

David Stutz

April 13, 2016

# Table of Contents

# Problem

**Problem.** Minimize composite function

$$\min_{x \in \mathbb{R}^d} h(x) = \min_{n \in \mathbb{R}^d} (f(x) + g(x))$$

where

# Problem

**Problem.** Minimize composite function

$$\min_{x \in \mathbb{R}^d} h(x) = \min_{n \in \mathbb{R}^d} (f(x) + g(x)) \quad (1)$$

where

- $f : \mathbb{R}^d \rightarrow \mathbb{R} \in C^1$  with  $L$ -Lipschitz continuous gradient;

# Problem

**Problem.** Minimize composite function

$$\min_{x \in \mathbb{R}^d} h(x) = \min_{n \in \mathbb{R}^d} (f(x) + g(x)) \quad (1)$$

where

- $f : \mathbb{R}^d \rightarrow \mathbb{R} \in C^1$  with  $L$ -Lipschitz continuous gradient;
- $g : \text{dom}(g) \subset \mathbb{R}^d \rightarrow \mathbb{R}_\infty$  is proper closed convex and lower semicontinuous;

# Problem

**Problem.** Minimize composite function

$$\min_{x \in \mathbb{R}^d} h(x) = \min_{n \in \mathbb{R}^d} (f(x) + g(x)) \quad (1)$$

where

- $f : \mathbb{R}^d \rightarrow \mathbb{R} \in C^1$  with  $L$ -Lipschitz continuous gradient;
- $g : \text{dom}(g) \subset \mathbb{R}^d \rightarrow \mathbb{R}_\infty$  is proper closed convex and lower semicontinuous;
- and  $h$  coercive and bounded below by

$$-\infty < h_{\min} := \inf_{x \in \mathbb{R}^d} h(x).$$

# Problem

**Problem.** Minimize composite function

$$\min_{x \in \mathbb{R}^d} h(x) = \min_{n \in \mathbb{R}^d} (f(x) + g(x)) \quad (1)$$

where

- $f : \mathbb{R}^d \rightarrow \mathbb{R} \in C^1$  with  $L$ -Lipschitz continuous gradient;
- $g : \text{dom}(g) \subset \mathbb{R}^d \rightarrow \mathbb{R}_\infty$  is proper closed convex and lower semicontinuous;
- and  $h$  coercive and bounded below by

$$-\infty < h_{\min} := \inf_{x \in \mathbb{R}^d} h(x).$$

Ochs et al. [?] combine forward-backward splitting with an inertial force/momentum term to solve Equation (??) iteratively.

# Table of Contents



# Related Work

Gradient descent for  $h \in C^1$ :

$$x^{(n+1)} = x^{(n)} - \alpha_n \nabla h(x^{(n)}).$$

Gradient descent with inertial force/momentum term:

$$x^{(n+1)} = x^{(n)} - \alpha_n \nabla h(x^{(n)}) + \beta_n (x^{(n)} - x^{(n-1)}).$$

Proximal point for  $h$  being proper closed convex:

$$x^{(n+1)} = \text{prox}_{\alpha_n h}(x^{(n)}).$$

Forward-backward splitting for  $h = f + g$  with  $f \in C^1$  and  $f, g$  being proper closed convex:

$$x^{(n+1)} = \text{prox}_{\alpha_n g}(x^{(n)} - \alpha_n \nabla f(x^{(n)})).$$

# Related Work

Gradient descent for  $h \in C^1$ :

$$x^{(n+1)} = x^{(n)} - \alpha_n \nabla h(x^{(n)}).$$

Gradient descent with inertial force/momentum term:

$$x^{(n+1)} = x^{(n)} - \alpha_n \nabla h(x^{(n)}) + \beta_n (x^{(n)} - x^{(n-1)}).$$

Proximal point for  $h$  being proper closed convex:

$$x^{(n+1)} = \text{prox}_{\alpha_n h}(x^{(n)}).$$

Forward-backward splitting for  $h = f + g$  with  $f \in C^1$  and  $f, g$  being proper closed convex:

$$x^{(n+1)} = \text{prox}_{\alpha_n g}(x^{(n)} - \alpha_n \nabla f(x^{(n)})).$$

# Related Work

Gradient descent for  $h \in C^1$ :

$$x^{(n+1)} = x^{(n)} - \alpha_n \nabla h(x^{(n)}).$$

Gradient descent with inertial force/momentum term:

$$x^{(n+1)} = x^{(n)} - \alpha_n \nabla h(x^{(n)}) + \beta_n (x^{(n)} - x^{(n-1)}).$$

Proximal point for  $h$  being proper closed convex:

$$x^{(n+1)} = \text{prox}_{\alpha_n h}(x^{(n)}).$$

Forward-backward splitting for  $h = f + g$  with  $f \in C^1$  and  $f, g$  being proper closed convex:

$$x^{(n+1)} = \text{prox}_{\alpha_n g}(x^{(n)} - \alpha_n \nabla f(x^{(n)})).$$

# Related Work

Gradient descent for  $h \in C^1$ :

$$x^{(n+1)} = x^{(n)} - \alpha_n \nabla h(x^{(n)}).$$

Gradient descent with inertial force/momentum term:

$$x^{(n+1)} = x^{(n)} - \alpha_n \nabla h(x^{(n)}) + \beta_n (x^{(n)} - x^{(n-1)}).$$

Proximal point for  $h$  being proper closed convex:

$$x^{(n+1)} = \text{prox}_{\alpha_n h}(x^{(n)}).$$

Forward-backward splitting for  $h = f + g$  with  $f \in C^1$  and  $f, g$  being proper closed convex:

$$x^{(n+1)} = \text{prox}_{\alpha_n g}(x^{(n)} - \alpha_n \nabla f(x^{(n)})).$$

# Related Work

Gradient descent for  $h \in C^1$ :

$$x^{(n+1)} = x^{(n)} - \alpha_n \nabla h(x^{(n)}).$$

Gradient descent with inertial force/momentum term:

$$x^{(n+1)} = x^{(n)} - \alpha_n \nabla h(x^{(n)}) + \beta_n (x^{(n)} - x^{(n-1)}).$$

Proximal point for  $h$  being proper closed convex:

$$x^{(n+1)} = \text{prox}_{\alpha_n h}(x^{(n)}).$$

Forward-backward splitting for  $h = f + g$  with  $f \in C^1$  and  $f, g$  being proper closed convex:

$$x^{(n+1)} = \text{prox}_{\alpha_n g}(x^{(n)} - \alpha_n \nabla f(x^{(n)})).$$

# Table of Contents

# Algorithm – Iterates and Backtracking

Ochs et al. [?] combine forward-backward splitting with an inertial force/momentum term.

# Algorithm – Iterates and Backtracking

Ochs et al. [?] combine forward-backward splitting with an inertial force/momentum term:

$$x^{(n+1)} = \text{prox}_{\alpha_n g}(x^{(n)} - \alpha_n \nabla f(x^{(n)}) + \beta_n(x^{(n)} - x^{(n-1)}))$$

with step size parameters  $(\alpha_n)_{n \in \mathbb{N}}$  and momentum parameters  $(\beta_n)_{n \in \mathbb{N}}$ .



# Algorithm – Iterates and Backtracking

Ochs et al. [?] combine forward-backward splitting with an inertial force/momentum term

$$x^{(n+1)} = \text{prox}_{\alpha_n g}(x^{(n)} - \alpha_n \nabla f(x^{(n)}) + \beta_n(x^{(n)} - x^{(n-1)}))$$

with step size parameters  $(\alpha_n)_{n \in \mathbb{N}}$  and momentum parameters  $(\beta_n)_{n \in \mathbb{N}}$ .

Backtracking to estimate the local Lipschitz constant  $L_n$  such that

$$\begin{aligned} f(x^{(n+1)}) &\leq f(x^{(n)}) + \nabla f(x^{(n)})^T (x^{(n+1)} - x^{(n)}) \\ &\quad + \frac{L_n}{2} \|x^{(n+1)} - x^{(n)}\|_2^2 \end{aligned}$$

# Algorithm – Iterates and Backtracking

Ochs et al. [?] combine forward-backward splitting with an inertial force/momentum term

$$x^{(n+1)} = \text{prox}_{\alpha_n g}(x^{(n)} - \alpha_n \nabla f(x^{(n)}) + \beta_n(x^{(n)} - x^{(n-1)}))$$

with step size parameters  $(\alpha_n)_{n \in \mathbb{N}}$  and momentum parameters  $(\beta_n)_{n \in \mathbb{N}}$ .

Backtracking to estimate the local Lipschitz constant  $L_n$  such that

$$\begin{aligned} f(x^{(n+1)}) \leq & f(x^{(n)}) + \nabla f(x^{(n)})^T (x^{(n+1)} - x^{(n)}) \\ & + \frac{L_n}{2} \|x^{(n+1)} - x^{(n)}\|_2^2 \end{aligned}$$

# Algorithm – iPiano

---

## Algorithm iPiano.

---

```
1: choose  $c_1, c_2 > 0$  close to zero,  $L_{-1} > 0$ ,  $\eta > 1$ ,  $x^{(0)}$ 
2:  $x^{(-1)} := x^{(0)}$ 
3: for  $n = 1, \dots$  do
4:
5:
6:
7:
8:   choose  $\alpha_n \geq c_1$ ,  $\beta_n \geq 0$ 
9:
10:   $x^{(n+1)} = \text{prox}_{\alpha_n g} (x^{(n)} - \alpha_n \nabla f(x^{(n)}) + \beta_n (x^{(n)} - x^{(n-1)}))$ 
11:
12: end for
```

---

# Algorithm – iPiano

---

## Algorithm iPiano.

---

```
1: choose  $c_1, c_2 > 0$  close to zero,  $L_{-1} > 0$ ,  $\eta > 1$ ,  $x^{(0)}$ 
2:  $x^{(-1)} := x^{(0)}$ 
3: for  $n = 1, \dots$  do
4:
5:
6:
7:   repeat
8:     choose  $\alpha_n \geq c_1$ ,  $\beta_n \geq 0$ 
9:   until  $\delta_n := \frac{1}{\alpha_n} - \frac{L_n}{2} - \frac{\beta_n}{2\alpha_n} \geq \gamma_n := \frac{1}{\alpha_n} - \frac{L_n}{2} - \frac{\beta_n}{\alpha_n} \geq c_2$ 
10:   $x^{(n+1)} = \text{prox}_{\alpha_n g} \left( x^{(n)} - \alpha_n \nabla f(x^{(n)}) + \beta_n (x^{(n)} - x^{(n-1)}) \right)$ 
11:
12: end for
```

---

# Algorithm – iPiano

---

## Algorithm iPiano.

---

```
1: choose  $c_1, c_2 > 0$  close to zero,  $L_{-1} > 0$ ,  $\eta > 1$ ,  $x^{(0)}$ 
2:  $x^{(-1)} := x^{(0)}$ 
3: for  $n = 1, \dots$  do
4:    $L_n := \frac{1}{\eta} L_{n-1}$ 
5:   repeat
6:      $L_n := \eta L_n$ 
7:     repeat
8:       choose  $\alpha_n \geq c_1$ ,  $\beta_n \geq 0$ 
9:     until  $\delta_n := \frac{1}{\alpha_n} - \frac{L_n}{2} - \frac{\beta_n}{2\alpha_n} \geq \gamma_n := \frac{1}{\alpha_n} - \frac{L_n}{2} - \frac{\beta_n}{\alpha_n} \geq c_2$ 
10:     $x^{(n+1)} = \text{prox}_{\alpha_n g} (x^{(n)} - \alpha_n \nabla f(x^{(n)}) + \beta_n (x^{(n)} - x^{(n-1)}))$ 
11:  until (??) is satisfied for  $x^{(n+1)}$ 
12: end for
```

---

# Algorithm – Monotonically Decreasing $\delta_n$

## Lemma

*For each  $n \in \mathbb{N}$ , given  $L_n > 0$ , there exist  $\alpha_n < 2(1 - \beta_n)/L_n$  and  $0 \leq \beta_n < 1$  as in iPiano such that  $c_2 \leq \gamma_n \leq \delta_n$  and  $(\delta_n)_{n \in \mathbb{N}}$  is monotonically decreasing.*

# Algorithm – Monotonically Decreasing $\delta_n$

## Lemma

*For each  $n \in \mathbb{N}$ , given  $L_n > 0$ , there exist  $\alpha_n < 2(1 - \beta_n)/L_n$  and  $0 \leq \beta_n < 1$  as in iPiano such that  $c_2 \leq \gamma_n \leq \delta_n$  and  $(\delta_n)_{n \in \mathbb{N}}$  is monotonically decreasing.*

## Proof Sketch.

With  $b_n := (\delta_{n-1} + \frac{L_n}{2}) / (c_2 + \frac{L_n}{2})$ :

$$\gamma_n \geq c_2 \Leftrightarrow \alpha_n \leq \frac{1 - \beta_n}{c_2 + \frac{L_n}{2}} < \frac{2(1 - \beta_n)}{L_n}$$

$$\delta_{n-1} \geq \delta_n \Leftrightarrow \frac{1 - \beta_n}{c_2 + \frac{L_n}{2}} \geq \alpha_n \geq \frac{1 - \frac{\beta_n}{2}}{\delta_{n-1} + \frac{L_n}{2}} \Rightarrow \beta_n \leq \frac{b_n - 1}{b_n - \frac{1}{2}}$$



# Table of Contents



# Convergence – Overview

Convergence analysis is based on **three** requirements regarding

$$\begin{aligned} H_{\delta_{n+1}}(x^{(n+1)}, x^{(n)}) &:= h(x^{(n+1)}) + \delta_{n+1} \underbrace{\|x^{(n)} - x^{(n-1)}\|_2^2}_{\Delta_{n+1}^2} \\ &:= h(x^{(n+1)}) + \delta_{n+1} \end{aligned}$$

and the sequence

$$(z^{(n+1)})_{n \in \mathbb{N}} := (x^{(n+1)}, x^{(n)})_{n \in \mathbb{N}} \subset \mathbb{R}^{2d}$$

generated by iPiano.

# Convergence – Overview

Convergence analysis is based on **three** requirements regarding

$$\begin{aligned} H_{\delta_{n+1}}(x^{(n+1)}, x^{(n)}) &:= h(x^{(n+1)}) + \delta_{n+1} \underbrace{\|x^{(n)} - x^{(n-1)}\|_2^2}_{\Delta_{n+1}^2} \\ &:= h(x^{(n+1)}) + \delta_{n+1} \end{aligned}$$

and the sequence

$$(z^{(n+1)})_{n \in \mathbb{N}} := (x^{(n+1)}, x^{(n)})_{n \in \mathbb{N}} \subset \mathbb{R}^{2d}$$

generated by iPiano.

# Convergence – Overview

Convergence analysis is based on **three** requirements regarding

$$\begin{aligned} H_{\delta_{n+1}}(x^{(n+1)}, x^{(n)}) &:= h(x^{(n+1)}) + \delta_{n+1} \underbrace{\|x^{(n)} - x^{(n-1)}\|_2^2}_{\Delta_{n+1}^2} \\ &:= h(x^{(n+1)}) + \delta_{n+1} \end{aligned}$$

and the sequence

$$(z^{(n+1)})_{n \in \mathbb{N}} := (x^{(n+1)}, x^{(n)})_{n \in \mathbb{N}} \subset \mathbb{R}^{2d}$$

generated by iPiano.

# Convergence – Overview

Convergence analysis is based on **three** requirements regarding

$$\begin{aligned} H_{\delta_{n+1}}(x^{(n+1)}, x^{(n)}) &:= h(x^{(n+1)}) + \delta_{n+1} \underbrace{\|x^{(n)} - x^{(n-1)}\|_2^2}_{\Delta_{n+1}^2} \\ &:= h(x^{(n+1)}) + \delta_{n+1} \end{aligned}$$

and the sequence

$$(z^{(n+1)})_{n \in \mathbb{N}} := (x^{(n+1)}, x^{(n)})_{n \in \mathbb{N}} \subset \mathbb{R}^{2d}$$

generated by iPiano.

Furthermore,  $H_{\delta_n}$  is required to satisfy the Kurdyka-Lojasiewicz property [?, ?] at a critical point  $\tilde{z}$  of  $H_{\delta_n}$ .

# Convergence – Requirements

## Definition

Given  $a, b > 0$ .  $H : \mathbb{R}^{2d} \rightarrow \mathbb{R}_\infty$  and a sequence  $(z^{(n)})_{n \in \mathbb{N}} \subset \mathbb{R}^{2d}$  satisfy:

(H1) if for each  $n \in \mathbb{N}$ , it holds

$$H(z^{(n+1)}) + a\Delta_n^2 \leq H(z^{(n)});$$

# Convergence – Requirements

## Definition

Given  $a, b > 0$ .  $H : \mathbb{R}^{2d} \rightarrow \mathbb{R}_\infty$  and a sequence  $(z^{(n)})_{n \in \mathbb{N}} \subset \mathbb{R}^{2d}$  satisfy:

(H1) if for each  $n \in \mathbb{N}$ , it holds

$$H(z^{(n+1)}) + a\Delta_n^2 \leq H(z^{(n)});$$

(H2) if for each  $n \in \mathbb{N}$ , there exists  $w^{(n+1)} \in \partial H(z^{(n+1)})$  with

$$\|w^{(n+1)}\|_2 \leq \frac{b}{2}(\Delta_n + \Delta_{n+1});$$

# Convergence – Requirements

## Definition

Given  $a, b > 0$ .  $H : \mathbb{R}^{2d} \rightarrow \mathbb{R}_\infty$  and a sequence  $(z^{(n)})_{n \in \mathbb{N}} \subset \mathbb{R}^{2d}$  satisfy:

(H1) if for each  $n \in \mathbb{N}$ , it holds

$$H(z^{(n+1)}) + a\Delta_n^2 \leq H(z^{(n)});$$

(H2) if for each  $n \in \mathbb{N}$ , there exists  $w^{(n+1)} \in \partial H(z^{(n+1)})$  with

$$\|w^{(n+1)}\|_2 \leq \frac{b}{2}(\Delta_n + \Delta_{n+1});$$

(H3) if there exists a subsequence  $(z^{(n_j)})_{j \in \mathbb{N}}$  with  $z^{(n_j)} \rightarrow \tilde{z} = (\tilde{x}, \tilde{x})$  and  $H(z^{(n_j)}) \rightarrow H(\tilde{z})$  for  $j \rightarrow \infty$ .

# Convergence – Requirements, Condition ??

## Lemma

*$H_{\delta_n}$  and  $(z^{(n)})_{n \in \mathbb{N}}$  as generated by iPiano satisfy Condition ??, in particular for each  $n \in \mathbb{N}$  it holds*

$$H_{\delta_{n+1}}(z^{(n+1)}) + \gamma_n \Delta_n^2 \leq H_{\delta_n}(z^{(n)});$$



# Convergence – Requirements, Condition ??

## Lemma

*$H_{\delta_n}$  and  $(z^{(n)})_{n \in \mathbb{N}}$  as generated by iPiano satisfy Condition ??, in particular for each  $n \in \mathbb{N}$  it holds*

$$H_{\delta_{n+1}}(z^{(n+1)}) + \gamma_n \Delta_n^2 \leq H_{\delta_n}(z^{(n)});$$

## Proof Sketch.

Iteration (Equation (??))  $\Rightarrow$

$$w := \frac{x^{(n)} - x^{(n+1)}}{\alpha_n} - \nabla f(x^{(n)}) + \frac{\beta_n}{\alpha_n}(x^{(n)} - x^{(n-1)}) \in \partial g(x^{(n+1)})$$



# Convergence – Requirements, Condition ??

## Proof Sketch (cont'd).

With  $w \in \partial g(x^{(n+1)})$ , using the convexity of  $g$ ,

$$g(x^{(n+1)}) \leq g(x^{(n)}) - w^T(x^{(n)} - x^{(n-1)}),$$

and the  $L_n$ -Lipschitz continuity of  $\nabla f$ ,

$$f(x^{(n+1)}) \leq f(x^{(n)}) - \nabla f(x^{(n)})^T(x^{(n+1)} - x^{(n)}) + \frac{L_n}{2} \|x^{(n)} - x^{(n+1)}\|_2^2;$$

it can be shown

$$h(x^{(n+1)}) \leq h(x^{(n)}) - \delta_n \Delta_{n+1}^2 + \delta_n \Delta_n^2 - \gamma_n \Delta_n^2$$

which implies the claim as  $\delta_n$  is monotonically decreasing. □

# Convergence – Requirements, Condition ??

## Lemma

*$H_{\delta_n}$  and  $(z^{(n)})_{n \in \mathbb{N}}$  as generated by iPiano satisfy Condition ??, i.e. for each  $n \in \mathbb{N}$  there exists  $w^{(n+1)} \in \partial H_{\delta_{n+1}}(z^{(n+1)})$  such that  $\|w^{(n+1)}\|_2 \leq \frac{7}{c_1}(\Delta_n + \Delta_{n+1})$ .*

# Convergence – Requirements, Condition ??

## Lemma

*$H_{\delta_n}$  and  $(z^{(n)})_{n \in \mathbb{N}}$  as generated by iPiano satisfy Condition ??, i.e. for each  $n \in \mathbb{N}$  there exists  $w^{(n+1)} \in \partial H_{\delta_{n+1}}(z^{(n+1)})$  such that*

$$\|w^{(n+1)}\|_2 \leq \frac{7}{c_1}(\Delta_n + \Delta_{n+1}).$$

## Proof Sketch.

For  $w^{(n+1)} \in \partial H_{\delta_{n+1}}(z^{(n+1)})$  it is  $w^{(n+1)} = (w_1^{(n+1)}, w_2^{(n+1)})$  with

$$w_1^{(n+1)} \in \partial g(x^{(n+1)}) + \nabla f(x^{(n+1)}) + 2\delta_n(x^{(n+1)} - x^{(n)})$$

$$w_2^{(n+1)} = -2\delta_n(x^{(n+1)} - x^{(n)})$$

and

$$\|w^{(n+1)}\|_2 \leq \dots \leq \left(\frac{1}{\alpha_n} + 4\delta_n + L_n\right)\Delta_{n+1} + \frac{\beta_n}{\alpha_n}\Delta_n \leq \frac{7}{c_1}(\Delta_{n+1} + \Delta_n)$$

# Convergence – Requirements, Condition ??

## Lemma

*$H_{\delta_n}$  and  $(z^{(n)})_{n \in \mathbb{N}}$  as generated by iPiano satisfy Condition ??, i.e. there exists a subsequence  $(z^{(n_j)})_{j \in \mathbb{N}}$  with  $z^{(n_j)} \rightarrow \tilde{z} = (\tilde{x}, \tilde{x})$  and  $H_{\delta_{n_j}}(z^{(n_j)}) \rightarrow H_{\delta}(\tilde{z})$  for  $j \rightarrow \infty$ .*

# Convergence – Requirements, Condition ??

## Lemma

$H_{\delta_n}$  and  $(z^{(n)})_{n \in \mathbb{N}}$  as generated by iPiano satisfy Condition ??, i.e. there exists a subsequence  $(z^{(n_j)})_{j \in \mathbb{N}}$  with  $z^{(n_j)} \rightarrow \tilde{z} = (\tilde{x}, \tilde{x})$  and  $H_{\delta_{n_j}}(z^{(n_j)}) \rightarrow H_{\delta}(\tilde{z})$  for  $j \rightarrow \infty$ .

## Proof Sketch.

Claim 1: by summing Condition ?? and deducing  $\sum_{n=0}^{\infty} \Delta_n^2 < \infty$  it can be shown that  $\lim_{n \rightarrow \infty} \Delta_n = 0$ .

Claim 2: from the coercivity of  $h$  and the Bolzano-Weierstrass theorem it follows the existence of a subsequence  $(x^{(n_j)})_{j \in \mathbb{N}}$  with.

Then:

$$\lim_{j \rightarrow \infty} H_{\delta_{n_j+1}}(x^{(n_j+1)}, x^{(n_j)}) = H_{\delta}(\tilde{x}, \tilde{x}) = h(\tilde{x}).$$



# Convergence – Kurdyka-Lojasiewicz Property

The Kurdyka-Lojasiewicz property is intended to relate the behavior of the subdifferential  $\partial H$  to the function values.

# Convergence – Kurdyka-Lojasiewicz Property

The Kurdyka-Lojasiewicz property is intended to relate the behavior of the subdifferential  $\partial H$  to the function values.

## Definition (Informally)

For a point  $\tilde{z} \in \text{dom}(\partial H)$ ,  $H$  is said to satisfy the Kurdyka-Lojasiewicz property if there exists a concave  $\phi \in C^1$  with  $\phi(0) = 0$  and  $\phi' > 0$  such that

$$\phi'(H(z) - H(\tilde{z})) \inf_{\hat{z} \in \partial H(z)} \|\hat{z}\|_2 \geq 1$$

for all  $z$  in an appropriate neighborhood of  $\tilde{z}$ .

Intuitively, the inequality controls the difference in function values by the subdifferential.



# Convergence – Kurdyka-Lojasiewicz Property

The Kurdyka-Lojasiewicz property is intended to relate the behavior of the subdifferential  $\partial H$  to the function values.

## Definition (Informally)

For a point  $\tilde{z} \in \text{dom}(\partial H)$ ,  $H$  is said to satisfy the Kurdyka-Lojasiewicz property if there exists a concave  $\phi \in C^1$  with  $\phi(0) = 0$  and  $\phi' > 0$  such that

$$\phi'(H(z) - H(\tilde{z})) \inf_{\hat{z} \in \partial H(z)} \|\hat{z}\|_2 \geq 1$$

for all  $z$  in an appropriate neighborhood of  $\tilde{z}$ .

Intuitively, the inequality controls the difference in function values by the subdifferential.

# Convergence – Kurdyka-Lojasiewicz Property

The Kurdyka-Lojasiewicz property is intended to relate the behavior of the subdifferential  $\partial H$  to the function values.

## Definition (Informally)

For a point  $\tilde{z} \in \text{dom}(\partial H)$ ,  $H$  is said to satisfy the Kurdyka-Lojasiewicz property if there exists a concave  $\phi \in C^1$  with  $\phi(0) = 0$  and  $\phi' > 0$  such that

$$\phi'(H(z) - H(\tilde{z})) \inf_{\hat{z} \in \partial H(z)} \|\hat{z}\|_2 \geq 1$$

for all  $z$  in an **appropriate neighborhood** of  $\tilde{z}$ .

Intuitively, the inequality controls the difference in function values by the subdifferential.

# Convergence – Convergence Theorem

## Theorem

*Let  $H$  be proper lower semicontinuous, satisfying the Kurdyka-Lojasiewicz property at  $\tilde{z} = (\tilde{x}, \tilde{x})$  specified by Condition ??, and  $(z^{(n)})_{n \in \mathbb{N}}$ , satisfying Conditions ?? - ??. Then  $(x^{(n)})_{n \in \mathbb{N}}$  converges to  $\tilde{x}$  such that  $\tilde{z}$  is a critical point of  $H$ .*

# Convergence – Convergence Theorem

## Theorem

*Let  $H$  be proper lower semicontinuous, satisfying the Kurdyka-Lojasiewicz property at  $\tilde{z} = (\tilde{x}, \tilde{x})$  specified by Condition ??, and  $(z^{(n)})_{n \in \mathbb{N}}$ , satisfying Conditions ?? - ??. Then  $(x^{(n)})_{n \in \mathbb{N}}$  converges to  $\tilde{x}$  such that  $\tilde{z}$  is a critical point of  $H$ .*

It can further be shown that the convergence rate is  $\mathcal{O}(1/\sqrt{n})$  for the residual

$$r(x) := x - \text{prox}_g(x - \nabla f(x))$$

in  $L_2$  norm.

# Convergence – Convergence Theorem (cont'd)

## Proof Sketch.

The proof is based on the following claim:

$$\sum_{i=1}^n \Delta_i \leq \frac{1}{2}(\Delta_0 - \Delta_n) + \frac{b}{a} \left[ \phi(H(z^{(1)}) - H(\tilde{z})) - \phi(H(z^{(n+1)}) - H(\tilde{z})) \right]$$

which is shown by induction. Then, it follows  $\sum_{n=0}^{\infty} \Delta_n < \infty$  and  $x^{(n)} \rightarrow \tilde{x}$ . Using the Kurdyka-Lojasiewicz property it can be shown that  $H(z^{(n)}) \rightarrow H(\tilde{z})$ . With Condition ?? it also follows that  $\tilde{z}$  is a critical point of  $H$ . □

# Table of Contents

# Implementation – Initialization

Remember, derived bounds for  $\alpha_0$  and  $\beta_0$ :

$$\alpha_0 < \frac{2(1 - \beta_0)}{L_0};$$
$$\beta_0 \leq \frac{b_0 - 1}{b_0 - \frac{1}{2}} \quad \text{with} \quad b_0 := \frac{\delta_{-1} + \frac{L_n}{2}}{c_2 + \frac{L_n}{2}}.$$

Guessing an appropriate  $\beta_0$  is obviously easier than guessing  $\delta_{-1}$ , so fix  $\beta_0$  and estimate  $L_0$  using

$$\frac{\|\nabla f(x^{(0)}) - \nabla f(\hat{x})\|_2}{\|x^{(0)} - \hat{x}\|_2} \leq L_0$$

for  $\hat{x} = \text{prox}_g(x^{(0)} - \nabla f(x^{(0)}))$ .

# Implementation – Initialization

Remember, derived bounds for  $\alpha_0$  and  $\beta_0$ :

$$\boxed{\alpha_0 < \frac{2(1 - \beta_0)}{L_0};}$$
$$\beta_0 \leq \frac{b_0 - 1}{b_0 - \frac{1}{2}} \quad \text{with} \quad b_0 := \frac{\delta_{-1} + \frac{L_n}{2}}{c_2 + \frac{L_n}{2}}.$$

Guessing an appropriate  $\beta_0$  is obviously easier than guessing  $\delta_{-1}$ , so fix  $\beta_0$  and estimate  $L_0$  using

$$\frac{\|\nabla f(x^{(0)}) - \nabla f(\hat{x})\|_2}{\|x^{(0)} - \hat{x}\|_2} \leq L_0$$

for  $\hat{x} = \text{prox}_g(x^{(0)} - \nabla f(x^{(0)}))$ .



# Implementation – Initialization

Remember, derived bounds for  $\alpha_0$  and  $\beta_0$ :

$$\alpha_0 < \frac{2(1 - \beta_0)}{L_0};$$

$$\beta_0 \leq \frac{b_0 - 1}{b_0 - \frac{1}{2}} \quad \text{with} \quad b_0 := \frac{\delta_{-1} + \frac{L_n}{2}}{c_2 + \frac{L_n}{2}}.$$

Guessing an appropriate  $\beta_0$  is obviously easier than guessing  $\delta_{-1}$ , so fix  $\beta_0$  and estimate  $L_0$  using

$$\frac{\|\nabla f(x^{(0)}) - \nabla f(\hat{x})\|_2}{\|x^{(0)} - \hat{x}\|_2} \leq L_0$$

for  $\hat{x} = \text{prox}_g(x^{(0)} - \nabla f(x^{(0)}))$ .

# Implementation – Initialization

Remember, derived bounds for  $\alpha_0$  and  $\beta_0$ :

$$\alpha_0 < \frac{2(1 - \beta_0)}{L_0};$$
$$\beta_0 \leq \frac{b_0 - 1}{b_0 - \frac{1}{2}} \quad \text{with} \quad b_0 := \frac{\delta_{-1} + \frac{L_n}{2}}{c_2 + \frac{L_n}{2}}.$$

Guessing an appropriate  $\beta_0$  is obviously easier than guessing  $\delta_{-1}$ , so fix  $\beta_0$  and estimate  $L_0$  using

$$\frac{\|\nabla f(x^{(0)}) - \nabla f(\hat{x})\|_2}{\|x^{(0)} - \hat{x}\|_2} \leq L_0$$

for  $\hat{x} = \text{prox}_g(x^{(0)} - \nabla f(x^{(0)}))$ .

# Implementation – Initialization (cont'd)

In practice, fix  $K \gg 100$  and compute

$$\alpha_0^{(k)} := \alpha_0 - k \frac{a_0 - c_1}{K} \text{ with } a_0 := \frac{2(1 - \beta_0)}{(L_0 + 2c_2)} \text{ and } k = 1, \dots, K$$

until  $\alpha_0^{(k)}$  satisfies

$$\delta_0 := \frac{1}{\alpha_0^{(k)}} - \frac{L_0}{2} - \frac{\beta_0}{2\alpha_0^{(k)}} \geq \gamma_0 := \frac{1}{\alpha_0^{(k)}} - \frac{L_0}{2} - \frac{\beta_0}{\alpha_0^{(k)}} \geq c_2.$$

# Implementation – Initialization (cont'd)

In practice, fix  $K \gg 100$  and compute

$$\alpha_0^{(k)} := \alpha_0 - k \frac{a_0 - c_1}{K} \text{ with } a_0 := \frac{2(1 - \beta_0)}{(L_0 + 2c_2)} \text{ and } k = 1, \dots, K$$

until  $\alpha_0^{(k)}$  satisfies

$$\delta_0 := \frac{1}{\alpha_0^{(k)}} - \frac{L_0}{2} - \frac{\beta_0}{2\alpha_0^{(k)}} \geq \gamma_0 := \frac{1}{\alpha_0^{(k)}} - \frac{L_0}{2} - \frac{\beta_0}{\alpha_0^{(k)}} \geq c_2.$$

# Implementation – Initialization (cont'd)

In practice, fix  $K \gg 100$  and compute

$$\alpha_0^{(k)} := \alpha_0 - k \frac{a_0 - c_1}{K} \text{ with } a_0 := \frac{2(1 - \beta_0)}{(L_0 + 2c_2)} \text{ and } k = 1, \dots, K$$

until  $\alpha_0^{(k)}$  satisfies

$$\delta_0 := \frac{1}{\alpha_0^{(k)}} - \frac{L_0}{2} - \frac{\beta_0}{2\alpha_0^{(k)}} \geq \gamma_0 := \frac{1}{\alpha_0} - \frac{L_0}{2} - \frac{\beta_0}{2\alpha_0} \geq c_2.$$

# Implementation – Finding $\alpha_n$ and $\beta_n$

Given  $L_{n-1}$  and  $\eta > 1$ , find the smallest  $l \in \mathbb{N}$  such that

$$L_n := \eta^l L_{n-1}$$

satisfies

$$\begin{aligned} f(x^{(n+1)}) \leq & f(x^{(n)}) + \nabla f(x^{(n)})^T (x^{(n+1)} - x^{(n)}) \\ & + \frac{L_n}{2} \|x^{(n+1)} - x^{(n)}\|_2^2. \end{aligned}$$

Alternatively, instead of  $L_{n-1}$ , use

$$\frac{\|\nabla f(x^{(n-1)}) - \nabla f(\hat{x})\|_2}{\|x^{(n-1)} - \hat{x}\|_2} \leq L_n$$

with  $\hat{x} = \text{prox}_g(x^{(n-1)} - \nabla f(x^{(n-1)}))$  as starting point for Equation (??).

# Implementation – Finding $\alpha_n$ and $\beta_n$

Given  $L_{n-1}$  and  $\eta > 1$ , find the smallest  $l \in \mathbb{N}$  such that

$$L_n := \eta^l L_{n-1} \quad (2)$$

satisfies

$$\begin{aligned} f(x^{(n+1)}) \leq & f(x^{(n)}) + \nabla f(x^{(n)})^T (x^{(n+1)} - x^{(n)}) \\ & + \frac{L_n}{2} \|x^{(n+1)} - x^{(n)}\|_2^2. \end{aligned}$$

Alternatively, instead of  $L_{n-1}$ , use

$$\frac{\|\nabla f(x^{(n-1)}) - \nabla f(\hat{x})\|_2}{\|x^{(n-1)} - \hat{x}\|_2} \leq L_n$$

with  $\hat{x} = \text{prox}_g(x^{(n-1)} - \nabla f(x^{(n-1)}))$  as starting point for Equation (??).

# Implementation – Finding $\alpha_n$ and $\beta_n$

Given  $L_{n-1}$  and  $\eta > 1$ , find the smallest  $l \in \mathbb{N}$  such that

$$L_n := \eta^l L_{n-1} \tag{2}$$

satisfies

$$f(x^{(n+1)}) \leq f(x^{(n)}) + \nabla f(x^{(n)})^T (x^{(n+1)} - x^{(n)}) + \frac{L_n}{2} \|x^{(n+1)} - x^{(n)}\|_2^2.$$

Alternatively, instead of  $L_{n-1}$ , use

$$\frac{\|\nabla f(x^{(n-1)}) - \nabla f(\hat{x})\|_2}{\|x^{(n-1)} - \hat{x}\|_2} \leq L_n$$

with  $\hat{x} = \text{prox}_g(x^{(n-1)} - \nabla f(x^{(n-1)}))$  as starting point for Equation (??).



# Implementation – Finding $\alpha_n$ and $\beta_n$

Given  $L_{n-1}$  and  $\eta > 1$ , find the smallest  $l \in \mathbb{N}$  such that

$$L_n := \eta^l L_{n-1} \tag{2}$$

satisfies

$$\begin{aligned} f(x^{(n+1)}) \leq & f(x^{(n)}) + \nabla f(x^{(n)})^T (x^{(n+1)} - x^{(n)}) \\ & + \frac{L_n}{2} \|x^{(n+1)} - x^{(n)}\|_2^2. \end{aligned}$$

Alternatively, instead of  $L_{n-1}$ , use

$$\frac{\|\nabla f(x^{(n-1)}) - \nabla f(\hat{x})\|_2}{\|x^{(n-1)} - \hat{x}\|_2} \leq L_n$$

with  $\hat{x} = \text{prox}_g(x^{(n-1)} - \nabla f(x^{(n-1)}))$  as starting point for Equation (??).

# Implementation – Finding $\alpha_n$ and $\beta_n$ (cont'd)

Similar to initialization, fix  $J, K \gg 100$  and compute

$$\beta_n^{(j)} := \frac{b_n - 1}{b_n - \frac{1}{2}} - \frac{j}{J} \frac{b_n - 1}{b_n - \frac{1}{2}} \quad \text{with} \quad b_n := \frac{\delta_{n-1} + \frac{L_n}{2}}{c_2 + \frac{L_n}{2}} \quad \text{and } j = 0, \dots, J,$$

$$\alpha_n^{(k)} := a_n - k \frac{a_n - c_1}{K} \quad \text{with} \quad a_n := \frac{2(1 - \beta_n)}{(L_n + 2c_2)} \quad \text{and } k = 1, \dots, K$$

until  $\beta_n^{(j)}$  and  $\alpha_n^{(k)}$  satisfy

$$\delta_n := \frac{1}{\alpha_n^{(k)}} - \frac{L_n}{2} - \frac{\beta_n^{(j)}}{2\alpha_n^{(k)}} \geq \gamma_n := \frac{1}{\alpha_n^{(k)}} - \frac{L_n}{2} - \frac{\beta_n^{(j)}}{\alpha_n^{(k)}} \geq c_2$$

and

$$\delta_n \leq \delta_{n-1}.$$

# Implementation – Finding $\alpha_n$ and $\beta_n$ (cont'd)

Similar to initialization, fix  $J, K \gg 100$  and compute

$$\beta_n^{(j)} := \frac{b_n - 1}{b_n - \frac{1}{2}} - \frac{j}{J} \frac{b_n - 1}{b_n - \frac{1}{2}} \quad \text{with} \quad b_n := \frac{\delta_{n-1} + \frac{L_n}{2}}{c_2 + \frac{L_n}{2}} \quad \text{and} \quad j = 0, \dots, J,$$
$$\alpha_n^{(k)} := a_n - k \frac{a_n - c_1}{K} \quad \text{with} \quad a_n := \frac{2(1 - \beta_n)}{(L_n + 2c_2)} \quad \text{and} \quad k = 1, \dots, K$$

until  $\beta_n^{(j)}$  and  $\alpha_n^{(k)}$  satisfy

$$\delta_n := \frac{1}{\alpha_n^{(k)}} - \frac{L_n}{2} - \frac{\beta_n^{(j)}}{2\alpha_n^{(k)}} \geq \gamma_n := \frac{1}{\alpha_n^{(k)}} - \frac{L_n}{2} - \frac{\beta_n^{(j)}}{\alpha_n^{(k)}} \geq c_2$$

and

$$\delta_n \leq \delta_{n-1}.$$

# Implementation – Finding $\alpha_n$ and $\beta_n$ (cont'd)

Similar to initialization, fix  $J, K \gg 100$  and compute

$$\beta_n^{(j)} := \frac{b_n - 1}{b_n - \frac{1}{2}} - \frac{j}{J} \frac{b_n - 1}{b_n - \frac{1}{2}} \quad \text{with} \quad b_n := \frac{\delta_{n-1} + \frac{L_n}{2}}{c_2 + \frac{L_n}{2}} \quad \text{and} \quad j = 0, \dots, J,$$

$$\alpha_n^{(k)} := a_n - k \frac{a_n - c_1}{K} \quad \text{with} \quad a_n := \frac{2(1 - \beta_n)}{(L_n + 2c_2)} \quad \text{and} \quad k = 1, \dots, K$$

until  $\beta_n^{(j)}$  and  $\alpha_n^{(k)}$  satisfy

$$\delta_n := \frac{1}{\alpha_n^{(k)}} - \frac{L_n}{2} - \frac{\beta_n^{(j)}}{2\alpha_n^{(k)}} \geq \gamma_n := \frac{1}{\alpha_n^{(k)}} - \frac{L_n}{2} - \frac{\beta_n^{(j)}}{\alpha_n^{(k)}} \geq c_2$$

and

$$\delta_n \leq \delta_{n-1}.$$

# Implementation – Finding $\alpha_n$ and $\beta_n$ (cont'd)

Similar to initialization, fix  $J, K \gg 100$  and compute

$$\beta_n^{(j)} := \frac{b_n - 1}{b_n - \frac{1}{2}} - \frac{j}{J} \frac{b_n - 1}{b_n - \frac{1}{2}} \quad \text{with} \quad b_n := \frac{\delta_{n-1} + \frac{L_n}{2}}{c_2 + \frac{L_n}{2}} \quad \text{and } j = 0, \dots, J,$$

$$\alpha_n^{(k)} := a_n - k \frac{a_n - c_1}{K} \quad \text{with} \quad a_n := \frac{2(1 - \beta_n)}{(L_n + 2c_2)} \quad \text{and } k = 1, \dots, K$$

until  $\beta_n^{(j)}$  and  $\alpha_n^{(k)}$  satisfy

$$\delta_n := \frac{1}{\alpha_n^{(k)}} - \frac{L_n}{2} - \frac{\beta_n^{(j)}}{2\alpha_n^{(k)}} \geq \gamma_n := \frac{1}{\alpha_n^{(k)}} - \frac{L_n}{2} - \frac{\beta_n^{(j)}}{\alpha_n^{(k)}} \geq c_2$$

and

$$\delta_n \leq \delta_{n-1}.$$

# Table of Contents

# Denoising – Model

Given a noisy image  $u^{(0)} : \Omega = [0, 1]^2 \rightarrow [0, 1]$ , minimize

$$h(u; u^{(0)}, \lambda) = \int_{\Omega} \rho_1(u(x) - u^{(0)}(x)) dx + \lambda \int_{\Omega} \rho_2(\|\nabla u(x)\|_2) dx$$

with

$$\rho_{1,\text{abs}} = |x| \text{ and } \rho_{1,\text{sqr}}(x) = x^2;$$

$$\rho_2(x) = \log \left( 1 + \frac{x^2}{\sigma^2} \right).$$

# Denoising – Model

Given a noisy image  $u^{(0)} : \Omega = [0, 1]^2 \rightarrow [0, 1]$ , minimize

$$h(u; u^{(0)}, \lambda) = \int_{\Omega} \rho_1(u(x) - u^{(0)}(x)) dx + \lambda \int_{\Omega} \rho_2(\|\nabla u(x)\|_2) dx$$

with

$$\rho_{1,\text{abs}} = |x| \text{ and } \rho_{1,\text{sqr}}(x) = x^2;$$

$$\rho_2(x) = \log \left( 1 + \frac{x^2}{\sigma^2} \right).$$



# Denoising – Model

Given a noisy image  $u^{(0)} : \Omega = [0, 1]^2 \rightarrow [0, 1]$ , minimize

$$h(u; u^{(0)}, \lambda) = \int_{\Omega} \rho_1(u(x) - u^{(0)}(x)) dx + \lambda \int_{\Omega} \rho_2(\|\nabla u(x)\|_2) dx$$

with

$$\rho_{1,\text{abs}} = |x| \text{ and } \rho_{1,\text{sqr}}(x) = x^2;$$

$$\rho_2(x) = \log \left( 1 + \frac{x^2}{\sigma^2} \right).$$

# Denoising – Model

Given a noisy image  $u^{(0)} : \Omega = [0, 1]^2 \rightarrow [0, 1]$ , minimize

$$h(u; u^{(0)}, \lambda) = \int_{\Omega} \rho_1(u(x) - u^{(0)}(x)) dx + \lambda \int_{\Omega} \rho_2(\|\nabla u(x)\|_2) dx$$

with

$$\rho_{1,\text{abs}} = |x| \text{ and } \rho_{1,\text{sqr}}(x) = x^2;$$

$$\rho_2(x) = \log \left( 1 + \frac{x^2}{\sigma^2} \right).$$

$\rho_{1,\text{sqr}}$  and  $\rho_2$  are differentiable; the proximal mapping of  $\rho_{1,\text{abs}}(x - x^{(0)})$  is

$$\text{prox}_{\alpha\rho_{1,\text{abs}}}(x) = \max(0, |x| - \alpha) \cdot \text{sign}(x) - x^{(0)}.$$

# Denoising – Results




Figure : Signal denoising experiment; input signal shown on the left with the perturbed/noisy signal on its right. Results using  $\rho_{1,abs}$  and  $\rho_{1,sqr}$  with  $\lambda \in \{0.2, 0.4, 0.6, 0.8\}$  are shown.

# Denoising – Convergence




Figure : Convergence of iPiano. Shown is the value of the objective function  $h(x(n))$  for each iterate  $x(n)$ ,  $n \geq 0$ , as well as the corresponding parameters  $\alpha_n$ ,  $\beta_n$  and  $L_n$ . Furthermore,  $\Delta_n := \|x^{(n)} - x^{(n-1)}\|_2$  is shown.

# Denoising – Results (cont'd)



Figure : Image denoising experiment; noisy image in the top row,  $\rho_{1,abs}$  in the middle row and  $\rho_{1,sqr}$  in the bottom row.

# Binary Segmentation – Model

Binary segmentation based on an approximation of the Mumford-Shah model [?, ?];  $u : [0, 1]^2 \rightarrow [-1, 1]$ :

$$\begin{aligned} h_{\epsilon}(u; c_+, c_-, u^{(0)}, \lambda) = & \int_{\Omega} \left( 9\epsilon \|\nabla u(x)\|_2^2 + \frac{(1 - u(x)^2)^2}{64\epsilon} \right) dx \\ & + \lambda \int_{\Omega} \left( \frac{1 + u(x)}{2} \right)^2 (u^{(0)}(x) - c_+)^2 dx \\ & + \lambda \int_{\Omega} \left( \frac{1 - u(x)}{2} \right)^2 (u^{(0)}(x) - c_-)^2 dx. \end{aligned}$$

# Binary Segmentation – Model

Binary segmentation based on an approximation of the Mumford-Shah model [?, ?];  $u : [0, 1]^2 \rightarrow [-1, 1]$ :

$$\begin{aligned} h_{\epsilon}(u; c_+, c_-, u^{(0)}, \lambda) = & \int_{\Omega} \left( 9\epsilon \|\nabla u(x)\|_2^2 + \frac{(1 - u(x)^2)^2}{64\epsilon} \right) dx \\ & + \lambda \int_{\Omega} \left( \frac{1 + u(x)}{2} \right)^2 (u^{(0)}(x) - c_+)^2 dx \\ & + \lambda \int_{\Omega} \left( \frac{1 - u(x)}{2} \right)^2 (u^{(0)}(x) - c_-)^2 dx. \end{aligned}$$

(It can be shown, that for  $\epsilon \rightarrow 0$ ,

$$\int_{\Omega} \left( 9\epsilon \|\nabla u(x)\|_2^2 + \frac{(1 - u(x)^2)^2}{64\epsilon} \right) dx$$

approximates  $|u|_{BV}$ .)

# Binary Segmentation – Model

Binary segmentation based on an approximation of the Mumford-Shah model [?, ?];  $u : [0, 1]^2 \rightarrow [-1, 1]$ :

$$h_\epsilon(u; c_+, c_-, u^{(0)}, \lambda) = \int_{\Omega} \left( 9\epsilon \|\nabla u(x)\|_2^2 + \frac{(1 - u(x)^2)^2}{64\epsilon} \right) dx$$

$$+ \lambda \int_{\Omega} \left( \frac{1 + u(x)}{2} \right)^2 (u^{(0)}(x) - c_+)^2 dx$$
$$+ \lambda \int_{\Omega} \left( \frac{1 - u(x)}{2} \right)^2 (u^{(0)}(x) - c_-)^2 dx.$$

(It can be shown, that for  $\epsilon \rightarrow 0$ ,

$$\int_{\Omega} \left( 9\epsilon \|\nabla u(x)\|_2^2 + \frac{(1 - u(x)^2)^2}{64\epsilon} \right) dx$$

approximates  $|u|_{BV}$ .)



# Binary Segmentation – Results (cont'd)




Figure : Segmentation results for thresholds  $\tau = -0.2, 0.0, 0.2$  and using  $g_{\text{sqr}}$ ; the foreground segment  $S_f$  is depicted in white.

# Table of Contents

# Conclusion

We discussed the minimization of composite functions of the form

$$\min_{x \in \mathbb{R}^d} h(x) = \min_{n \in \mathbb{R}^d} (f(x) + g(x)).$$

# Conclusion

We discussed the minimization of composite functions of the form

$$\min_{x \in \mathbb{R}^d} h(x) = \min_{n \in \mathbb{R}^d} (f(x) + g(x)).$$

Ochs et al. [?] proposed the iPiano algorithm to solve this problem under to following requirements:

- $g$  proper closed convex and lower semi continuous;
- $f \in C^1$  with  $L$ -Lipschitz continuous  $\nabla f$ ;
- $h$  coercive and bounded below;
- and  $H_{\delta_n}(x^{(n)}, x^{(n-1)}) = h(x^{(n)}) + \delta_n \Delta_n$  satisfying the Kurdyka-Lojasiewicz property [?, ?] at a critical point.

# Conclusion

We discussed the minimization of composite functions of the form

$$\min_{x \in \mathbb{R}^d} h(x) = \min_{n \in \mathbb{R}^d} (f(x) + g(x)).$$

Ochs et al. [?] proposed the iPiano algorithm to solve this problem under to following requirements:

- $g$  proper closed convex and lower semi continuous;
- $f \in C^1$  with  $L$ -Lipschitz continuous  $\nabla f$ ;
- $h$  coercive and bounded below;
- and  $H_{\delta_n}(x^{(n)}, x^{(n-1)}) = h(x^{(n)}) + \delta_n \Delta_n$  satisfying the Kurdyka-Lojasiewicz property [?, ?] at a critical point.

The algorithm can be implemented efficiently in C++ and used to solve image processing tasks.

# Appendix – Kurdyka-Lojasiewicz Property

## Definition

$H$  has the Kurdyka-Lojasiewicz property at point  $\tilde{z} \in \text{dom}(\partial H)$  there exist  $\eta \in (0, \infty]$ , a neighborhood  $U$  of  $\tilde{z}$ , and a continuous concave function  $\phi : [0, \eta) \rightarrow \mathbb{R}_+$  such that

- $\phi \in C^1((0, \eta))$ ,  $\phi(0) = 0$ , and for all  $s \in (0, \eta)$ ,  $\phi'(s) > 0$ ;
- and for all  $z \in U \cap \{z \in \mathbb{R}^{2d} | H(\tilde{z}) < H(z) < H(\tilde{z}) + \eta\}$  the Kurdyka-Lojasiewicz inequality holds:

$$\phi'(H(z) - H(\tilde{z})) \inf_{\hat{z} \in \partial H(z)} \|\hat{z}\|_2 \geq 1.$$

# Appendix – Kurdyka-Lojasiewicz Property

## Definition

$H$  has the Kurdyka-Lojasiewicz property at point  $\tilde{z} \in \text{dom}(\partial H)$  there exist  $\eta \in (0, \infty]$ , a neighborhood  $U$  of  $\tilde{z}$ , and a continuous concave function  $\phi : [0, \eta) \rightarrow \mathbb{R}_+$  such that

- $\phi \in C^1((0, \eta))$ ,  $\phi(0) = 0$ , and for all  $s \in (0, \eta)$ ,  $\phi'(s) > 0$ ;
- and for all  $z \in U \cap \{z \in \mathbb{R}^{2d} | H(\tilde{z}) < H(z) < H(\tilde{z}) + \eta\}$  the Kurdyka-Lojasiewicz inequality holds:

$$\phi'(H(z) - H(\tilde{z})) \inf_{\hat{z} \in \partial H(z)} \|\hat{z}\|_2 \geq 1.$$

Intuitively, for  $H \in C^1$ , this means that  $\phi$  has to be steep around critical points  $\tilde{z}$  of  $H$  where  $\nabla H$  is flat.