

TOPOLOGY MATH4121A/9021
LECTURE 2

Definition. Let (X, \mathcal{T}) be a topological space, A be a subset of X , and p be a point in X . Then p is a **limit point** of A if and only if for each open set U containing p , $(U - \{p\}) \cap A \neq \emptyset$. Notice that p may or may not belong to A .

Theorem 1. Suppose $p \notin A$ in a topological space (X, \mathcal{T}) . Then p is not a limit point of A if and only if there exists a neighborhood U of p such that $U \cap A = \emptyset$.

Proof.

Definition. Let (X, \mathcal{T}) be a topological space, A be a subset of X , and p be a point in X . If $p \in A$ but p is not a limit point of A , then p is an **isolated point** of A .

Theorem 2. If p is an isolated point of a set A in a topological space X , then there exists an open set U such that $U \cap A = \{p\}$.

Proof.

Exercise 3. Give examples of sets A in topological spaces (X, \mathcal{T}) with

- (1) a limit point of A that is an element of A ;
- (2) a limit point of A that is not an element of A ;
- (3) an isolated point of A ;
- (4) a point not in A that is not a limit point of A .

Solution.

Definition. Let (X, \mathcal{T}) be a topological space, and $A \subset X$. Then the **closure** of A in X , denoted \bar{A} or $\text{Cl}(A)$ or $\text{Cl}_X(A)$, is the set A together with all its limit points in X . Let (X, \mathcal{T}) be a topological space and $A \subset X$. The subset A is **closed** if and only if $\bar{A} = A$, in other words, if A contains all its limit points.

The following theorem tells us that closed sets and open sets are complements of one another.

Theorem 4. *Let (X, \mathcal{T}) be a topological space. Then the set A is closed if and only if $X - A$ is open.*

Solution.

Similarly, one can show that removing a closed set from an open set leaves an open set. A theorem which may be helpful in proving this theorem is DeMorgans Law:

Theorem. (DeMorgan's Laws) Let X be a set, and let $\{A_k\}_{k=1}^N$ be a finite collection of sets such that $A_k \subset X$ for each $k = 1, 2, \dots, N$. Then

$$X - \left(\bigcup_{k=1}^N A_k \right) = \bigcap_{k=1}^N (X - A_k)$$

and

$$X - \left(\bigcap_{k=1}^N A_k \right) = \bigcup_{k=1}^N (X - A_k).$$

Theorem 5. Let (X, \mathcal{T}) be a topological space, and let U be an open set and A be a closed subset of X . Then the set $U - A$ is open and the set $A - U$ is closed.

Proof.

Theorem 6. $y \in \bar{A}$ iff for each open set U containing x , $U \cap A \neq \emptyset$.

Proof.

Theorem 7. *For any topological space (X, \mathcal{T}) and $A \subset X$, \bar{A} is closed. That is, for any set A in a topological space, $\overline{\bar{A}} = \bar{A}$. Hint: The previous theorem may be useful in the “ \subset ” direction.*

Proof.

Theorem 8. *For any set A in a topological space X , the closure of A equals the intersection of all closed sets containing A , that is,*

$$\overline{A} = \bigcap_{B \supset A, B \in \mathcal{C}} B$$

where \mathcal{C} is the collection of all closed sets in X .