

Confidence Limits (15-05-2025)

Exercise 1

For the following datasets on Bootstrap Sampling, generate a sampling distribution for the mean and median using bootstrapping:

- Bank Account Balance
- Cereal Sugar Content
- Female Student Heights

For each dataset this means you need to create at least 200 bootstrap derived datasets (the “bootstrap samples”) and the program will automatically generate a sampling distribution for the “Statistic of Interest”. You can then click on the tab “Get Confidence Interval” on the top title row next to “Build Sampling Distribution”. Comment on how the Central Limit Theorem is relevant in each case and how sampling distributions compare between the mean and the median.

Exercise 2

A sample of size $n = 100$ produced a sample mean of $\bar{x} = 16$. Assuming the population standard deviation is 3, compute a 95% confidence interval where we expect the population mean μ .

Exercise 3

Assuming a population standard deviation of $\sigma = 3$, how large should a sample be to estimate the population mean μ , using a 95% confidence interval, such that margin of error does not exceed 0.5? (This means that the confidence interval may contain only values between $\bar{x} - 0.5$ and $\bar{x} + 0.5$.) You can try to invert the formula for confidence intervals to get this value.

Exercise 4

The operations manager of a large production plant would like to estimate the mean amount of time a worker takes to assemble a new electronic component.

- After observing 120 workers assembling similar devices, the manager noticed that their average time was 16.2 minutes with a standard deviation of 3.6 minutes. Construct a 92 % confidence interval for the mean assembly time.
- How many workers should be involved in this study in order to have the mean assembly time estimated up to ± 15 seconds with 92 % confidence?

Exercise 5

In order to ensure efficient usage of a server, it is necessary to estimate the mean number of concurrent users. According to records, the sample mean and sample standard deviation of number of concurrent users at 100 randomly selected times is 37.7 and 9.2, respectively.

- Construct a 90% confidence interval for the mean number of concurrent users.
- (A question phrased as a chapter 10 question. You should be able to come up with a good answer without having studied chapter 10.) Do these data provide significant evidence, at 10 % significance level, that the mean number of concurrent users is greater than 35?

Exercise 6

Show that when calculating a confidence interval for an unknown population proportion p (say of left-handers) from some sample of people n , $E(\hat{p}) = p$ and $V(\hat{p}) = \frac{p(1-p)}{n}$. Hint: Write out the expectations in terms of the random variable X : Number of successes in the sample and use the expectations of the Binomial Distribution.

Exercise 7

Of 48 engineers that were interviewed about the topic, 36 think that the new reinforcement of a concrete building is not earthquake proof. Give a 95% confidence interval for the true proportion of engineers that thinks that the new reinforcement is not earthquake proof. [Note that we haven't yet covered this in lecture - but it is there in Lecture 7 and accompanying chapter.]

Exercise 8

The Normal Distribution can always be rescaled and shifted to a $N(0, 1)$ distribution. See slides in Lecture 7. Demonstrate that we can similarly transform a Uniform Distribution $U(a, b)$ to $U(0, 1)$.