


Calculus Lecture 3: Differentiation

- Tangent lines
- Definition of a derivative
- Calculation of derivatives
 - Product rule
 - Chain rule
 - Trigonometric functions
 - Exponential and logarithmic functions
- Higher order derivatives

Adams' Ch. 2.1-6, 3.3

Tangent lines and their slopes




slope: $\frac{\Delta y}{\Delta x} = \frac{f(x_0+h) - f(x_0)}{h}$

slope of tangent line

$$m = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$$


- Can the tangent line be vertical? *yes*



$$f(x) = \sqrt[3]{x}$$

$$\lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h} = \pm \infty \quad (\text{here } +\infty)$$

- Does the tangent line always exist?



$$f(x) = |x| \rightarrow f'(x) = \begin{cases} 1 & x > 0 \\ -1 & x < 0 \end{cases} = \text{sgn}(x)$$

no tangent line at $x_0 = 0$

$$f'(0) = \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h}$$

$$= \lim_{h \rightarrow 0^+} \frac{|0+h| - |0|}{h}$$

$$= \lim_{h \rightarrow 0^+} \frac{|h|}{h} = \lim_{h \rightarrow 0^+} \frac{h}{h} = 1$$

The derivative

The derivative of a function at a point c of the domain is defined as the slope of (the tangent of) the function at c .

$$f'(c) = \lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h} = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$$

The derivative only exists if this limit exists and is finite!

$$\text{domain}(f') \subseteq \text{domain}(f)$$

- Notations:

$$y', f'(x), \frac{d}{dx}(f(x)), D_x f, Df(x), \frac{dy}{dx}$$
$$f'(x_0), \left. \frac{d}{dx}(f(x)) \right|_{x=x_0}, \left. \frac{dy}{dx} \right|_{x=x_0}$$

- Differentiable:

$$f \text{ is differentiable at } x_0 = f'(x_0) \text{ exists}$$

- Singular point:

$$x_0 \text{ is a singular point} = f'(x_0) \text{ does not exist}$$

- Left and right derivatives: $f'_\pm(x_0) = \lim_{h \rightarrow 0^\pm} \frac{f(x_0+h) - f(x_0)}{h}$

Examples

$$\begin{aligned} \bullet f(x) = ax + b &\rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(a(x+h) + b) - (ax + b)}{h} \\ &= \lim_{h \rightarrow 0} \frac{ah}{h} = a \end{aligned}$$

$$\bullet f(x) = c \rightarrow f'(x) = 0$$

$$\begin{aligned} \bullet f(x) = x^2 &\Rightarrow f'(x) = 2x \quad \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{x^2 + 2hx + h^2 - x^2}{h} = 2x \end{aligned}$$

$$\frac{d}{dx}(x^n) = n \cdot x^{n-1} \quad (\forall n \in \mathbb{R})$$

Differentiation rules

- Differentiable \rightarrow continuous
- For functions f and g differentiable at x , and k constant,
 - $(f+g)'(x) = f'(x) + g'(x)$
 - $(kf)'(x) = k \cdot f'(x)$
- (The product rule) For functions f and g differentiable at x
 - $(f(x)g(x))' = f'(x) \cdot g(x) + f(x) \cdot g'(x)$
 - NOT: ~~$f'(x) \cdot g'(x)$~~

$$\begin{aligned}\frac{d}{dx}(f(x) \cdot g(x)) &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(f(x+h)g(x+h) - f(x)g(x+h)) + (f(x)g(x+h) - f(x)g(x))}{h} \\ &= \lim_{h \rightarrow 0} \left(g(x+h) \underbrace{\left(\frac{f(x+h) - f(x)}{h} \right)}_{f'(x)} + f(x) \underbrace{\left(\frac{g(x+h) - g(x)}{h} \right)}_{g'(x)} \right) \\ &= g(x) \cdot f'(x) + f(x) \cdot g'(x)\end{aligned}$$

Examples

$$f(x) = (x^2+1)(x^3+4)$$

$$f'(x) = (2x)(x^3+4) + (x^2+1)(3x^2) = 2x^4 + 8x + 3x^4 + 3x^2 = 5x^4 + 3x^2 + 8x$$

$$f(x) = x^5 + x^3 + 4x^2 + 4 \rightarrow f'(x) = 5x^4 + 3x^2 + 8x$$

- "power rule" $\frac{d}{dx}(x^n) = n \cdot x^{n-1}$ (valid for $n \in \mathbb{R}$)

proof for $n \in \mathbb{N}$: proof by induction :)

- Base case $n=1$: $\frac{d}{dx}(x^1) = 1$ ✓

- Induction step: $P(n) \rightarrow P(n+1)$

$$P(n): \frac{d}{dx}(x^n) = n \cdot x^{n-1}$$

$$P(n+1): \frac{d}{dx}(x^{n+1}) = (n+1) \cdot x^n$$

$$\frac{d}{dx}(x^{n+1}) = \frac{d}{dx}(x^n \cdot x) = \frac{d}{dx}(x^n) \cdot x + x^n \cdot \frac{d}{dx}(x)$$

$$= n \cdot \underbrace{x^{n-1} \cdot x}_{x^n} + x^n = (n+1) x^n \quad \square$$

- (The chain rule): If $f(u)$ is differentiable at $u=g(x)$, and $g(x)$ is differentiable at x , then $f(g(x))$ is differentiable and

$$\frac{d}{dx} (f(g(x))) = f'(g(x)) \cdot g'(x)$$

$$g(x)=u \quad \frac{d}{dx} (f(u(x))) = \frac{df}{du} \cdot \frac{du}{dx}$$

- (reciprocal rule)

$$\frac{d}{dx} \left(\frac{1}{f(x)} \right) = -\frac{1}{(f(x))^2} \cdot f'(x)$$

- (quotient rule) $\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{d}{dx} \left(f(x) \cdot \frac{1}{g(x)} \right) = f'(x) \cdot \frac{1}{g(x)} - \frac{f(x)g'(x)}{(g(x))^2}$


$$= \frac{f'(x) \cdot g(x) - f(x)g'(x)}{(g(x))^2}$$

Examples: • $\frac{d}{dx} \left(\frac{1}{1+x^2} \right) = \frac{-1}{(1+x^2)^2} \cdot \frac{d}{dx} (1+x^2) = \frac{-2x}{(1+x^2)^2}$

$$\begin{aligned} \bullet \frac{d}{dx} \left(\frac{x^2+1}{\sqrt{x}} \right) &= \frac{(2x \cdot \sqrt{x} - \frac{1}{2\sqrt{x}} (x^2+1))}{x} = \frac{2(2x \cdot)x - (x^2+1)}{2x\sqrt{x}} \\ &= \frac{3x^2 - 1}{2x\sqrt{x}} \end{aligned}$$

Derivatives of trigonometric functions

- $\lim_{x \rightarrow 0} \frac{\sin(x)}{x}$



$$\underline{x > 0} \quad \sin(x) < x < \tan(x)$$

$$\Leftrightarrow \boxed{\frac{\sin(x)}{x} < 1 < \frac{\tan(x)}{x}}$$

$$\frac{\tan(x)}{x} = \frac{\sin(x)}{\cos(x) \cdot x}$$

$$\Rightarrow \cos(x) < \frac{\sin(x)}{x}$$

$$\Rightarrow \cos(x) < \frac{\sin(x)}{x} < 1$$

\downarrow as $x \rightarrow 0$ \downarrow as $x \rightarrow 0$

$$\lim_{x \rightarrow 0^+} \frac{\sin(x)}{x} = 1$$

$$\rightarrow \lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1 \quad \text{since } \frac{\sin(x)}{x} \text{ is even}$$

$$\begin{aligned}
 \rightarrow \frac{d}{dx}(\sin(x)) &= \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{2 \sin\left(\frac{h}{2}\right) \cos\left(x + \frac{h}{2}\right)}{h} \\
 &\quad \text{approaches 1 as } h \rightarrow 0 \\
 &= \lim_{h \rightarrow 0} \underbrace{\frac{\sin\left(\frac{h}{2}\right)}{\frac{h}{2}}}_{\rightarrow 1} \cdot \lim_{h \rightarrow 0} \underbrace{\cos\left(x + \frac{h}{2}\right)}_{\cos(x)} \\
 &= \cos(x)
 \end{aligned}$$

(alternative calculation.)

$$\begin{aligned}
 \frac{d}{dx}(\sin(x)) &= \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin(x)}{h} = \\
 \lim_{h \rightarrow 0} \frac{(\sin(x)\cos(h) + \sin(h)\cos(x)) - \sin(x)}{h} &= \lim_{h \rightarrow 0} \left(\sin(x) \cdot \frac{\cos(h) - 1}{h} + \cos(x) \frac{\sin(h)}{h} \right) \\
 &\quad \text{to 0, see below} \\
 \lim_{h \rightarrow 0} \frac{\cos(h) - 1}{h} &= \lim_{h \rightarrow 0} \frac{(\cos(h) - 1)(\cos(h) + 1)}{h(\cos(h) + 1)} \\
 &= \lim_{h \rightarrow 0} \frac{\cos^2(h) - 1}{h(\cos(h) + 1)} = \lim_{h \rightarrow 0} \frac{-\sin^2(h)}{h(\cos(h) + 1)} \\
 &= \lim_{h \rightarrow 0} \left(\underbrace{\frac{\sin(h)}{h}}_{\rightarrow 1} \cdot \underbrace{\frac{-\sin(h)}{\cos(h) + 1}}_{\rightarrow 0} \right) = 0 \\
 &= \lim_{h \rightarrow 0} \left(\cos(x) \frac{\sin(h)}{h} \right) = \cos(x)
 \end{aligned}$$

trig. identities:

$$\sin(a+b) = \sin(a)\cos(b) + \sin(b)\cos(a)$$

$$\sin(a-b) = \sin(a)\cos(b) - \sin(b)\cos(a)$$

$$\rightarrow \sin(a+b) - \sin(a-b) = 2\sin(b)\cos(a)$$

$$\begin{aligned} \hookrightarrow \text{here: } a &= x + \frac{h}{2} \\ b &= \frac{h}{2} \end{aligned}$$

$$\begin{aligned} a+b &= x+h \\ a-b &= x \end{aligned}$$

other trig. limits:

$$\frac{d}{dx}(\cos(x)) = \frac{d}{dx}(\sin(\frac{\pi}{2} - x)) = -\cos(\frac{\pi}{2} - x) = -\sin(x)$$

$$\frac{d}{dx}(\tan(x)) = \frac{d}{dx}\left(\frac{\sin(x)}{\cos(x)}\right) = \frac{\cos(x) \cdot \cos(x) - (-\sin(x) \cdot \sin(x))}{\cos^2(x)} = \frac{1}{\cos^2(x)}$$

reciprocal rule

$$\bullet \frac{d}{dx}(e^x) = e^x$$

$$\lim_{h \rightarrow 0} \frac{e^{x+h} - e^x}{h} = e^x \cdot \lim_{h \rightarrow 0} \frac{e^h - 1}{h}$$

c. is chosen / determined
such, that this limit is 1

the exponential function $f(x) = e^x$ is the only function that is equal to its derivative!

$$\bullet \frac{d}{dx}(\ln(x)) = \frac{1}{x}$$

proof: $y = \ln(x) \Leftrightarrow e^y = x$ (def. of logarithm.)

$$\Rightarrow \frac{d}{dx}(e^y) = \frac{d}{dx}(x)$$

$$\Rightarrow e^y \frac{dy}{dx} = 1 \quad (\text{chain rule})$$

$$\Rightarrow x \frac{dy}{dx} = 1$$

$$\Rightarrow dy/dx = 1/x$$

Examples

$$\bullet \frac{d}{dx} \left(\frac{\cos(x)}{\sin(x)} \right) = \frac{-\sin(x) \cdot \sin(x) - \cos(x) \cdot \cos(x)}{\sin^2(x)} = \frac{-1}{\sin^2(x)}$$

$$\bullet \frac{d}{dx} (e^{-\sin(3x)}) = e^{-\sin(3x)} \frac{d}{dx} (-\sin(3x)) = e^{-\sin(3x)} (-\cos(3x)) \frac{d}{dx} (3x)$$

$$\text{chain rule: } \frac{d}{dx} (e^{f(x)}) = e^{f(x)} \cdot f'(x)$$

$$\frac{d}{dx} (-\sin(f(x))) = -\cos(f(x)) \cdot f'(x)$$

$$= -3 \cos(3x) e^{-\sin(3x)}$$

Higher order derivatives

$$y'' = f''(x) = \frac{d^2 f}{dx^2} = \frac{d}{dx} \left(\frac{d}{dx} f(x) \right) = D_x^2 f(x)$$

$$f(x) = x^3 \rightarrow f'(x) = 3x^2 \rightarrow f''(x) = 6 \cdot x \rightarrow f'''(x) = 6 \rightarrow f^{(4)} = 0$$