

MATH 223

Yang David Zhou

Winter 2015

0.1 Administrative

Professor Tiago Salvador

Website: <http://www.math.mcgill.ca/tsalvador/>

Office: Burnside Building 1036

Office Hours: M1:45-2:45PM W2:00-3:00PM, F3:30-4:30PM

Grading

Assignments	15%	15%
Midterm	25%	0%
Final	60%	85%

The midterm will be scheduled for the 7th week of class.

Chapter 1

Vectors

1.1 Vectors in \mathbb{R}^n

\mathbb{R}^n is the set of all n -tuples of real numbers $u = (a_1 \dots a_n) \mid a \in \mathbb{R}$ where a are the **components** or **entries**.

Remark 1. We use the term **scalar** to refer to an element in \mathbb{R} .

1.2 Basic Definitions

Definition 1. Addition

$$u, v \in \mathbb{R}^n$$

$$u = (a_1 \dots a_n)$$

$$v = (b_1 \dots b_n)$$

$$u + v = (a_1 + b_1 \dots a_n + b_n)$$

Definition 2. Scalar Multiplication

$$k \in \mathbb{R}$$

$$ku = (ka_1 \dots ka_n)$$

Definition 3. Two vectors u and v are said to be **equal** ($u = v$) if $a_i = b_i \forall i = 1 \dots n$.

Definition 4. The **zero vector** is defined as $0 = (0 \dots 0)$.

Definition 5. Suppose we are given m vectors $u_1 \dots u_m \in \mathbb{R}^n$ and m scalars $k_1 \dots k_m \in \mathbb{R}$.

$$\text{Let } u = k_1 u_1 + \dots + k_m u_m.$$

Such a vector u is called a **linear combination** of the vectors $u_1 \dots u_m$.

Definition 6. A vector u can be called a **multiple** of v if there is a scalar k such that $u = kv$ with $k \neq 0$. In the case $k > 0$ we say u is in the same direction as v . In the case $k < 0$ we say u is in the opposite direction of v .

1.3 The Dot Product

Definition 7. Let $u = (a_1 \dots a_n)$ and $v = (b_1 \dots b_n)$. The **dot product** or inner product is given by,

$$u \cdot v = a_1 b_1 + \dots a_n b_n =$$

Definition 8. The vectors u and v are **orthogonal** if $u \cdot v = 0$.

1.4 The Vector Norm

Definition 9. The **norm** or **length** of a vector is given by,

$$\|u\| = \sqrt{a_1^2 + \dots + a_n^2}$$

Thus $\|u\| \geq 0$ and $\|u\| = 0$ if and only if (iff) $u = 0$.

Definition 10. A vector is called a **unit vector** if $\|u\| = 1$.

Definition 11. For any non-zero vector v , the vector

$$\hat{v} = \frac{1}{\|v\|} v$$

is the only unit vector with the same direction of v . The process of finding \hat{v} is called **normalizing**.

1.5 Theorem: Cauchy-Schwarz Inequality

Theorem 1. Given any two vectors $u, v \in \mathbb{R}^n$, then,

$$|u \cdot v| \leq \|u\| \|v\|$$

Proof. Let $t \in \mathbb{R}$. So, $\|tu + v\|^2 \geq 0$.

$$\begin{aligned} \|tu + v\|^2 &= (tu + v)(tu + v) \\ &= (tu \cdot tu) + (tu \cdot v) + (v \cdot tu) + (v \cdot v) \\ &= t^2(u \cdot u) + t(v \cdot u) + t(u \cdot v) + (v \cdot v) \\ &= t^2\|u\|^2 + 2t(u \cdot v) + \|v\|^2 \end{aligned}$$

We can represent this in the form $at^2 + bt + c \geq 0$, so,

$$a = \|u\|^2, b = 2(u \cdot v), c = \|v\|^2$$

Take the Discriminant as $b^2 - 4ac \iff b^2 \leq 4ac$.

$$\begin{aligned} 4(u \cdot v)^2 &\leq 4\|u\|^2\|v\|^2 \\ |u \cdot v| &\leq \|u\| \|v\| \end{aligned}$$

□

1.6 Theorem: Minkowski Triangle Inequality

Theorem 2. Given $u, v \in \mathbb{R}^n$, then $\|u + v\| \leq \|u\| + \|v\|$.

Proof.

$$\begin{aligned}\|u + v\|^2 &= \|u\|^2 + 2(u \cdot v) + \|v\|^2 \\ &\leq \|u\|^2 + 2\|u\|\|v\| + \|v\|^2 \quad \text{by C-S inequality} \\ &= (\|u\| + \|v\|)^2\end{aligned}$$

So, $\|u + v\|^2 \leq (\|u\| + \|v\|)^2$. Take the square root and we are done. \square

1.7 Geometry with Vectors

Definition 12. The **distance** between vectors $u, v \in \mathbb{R}^n$ is given by,

$$d(u, v) = \|u - v\| = \sqrt{(a_1 - b_1)^2 + \dots + (a_n - b_n)^2}$$

Definition 13. The **angle** between vectors $u, v \in \mathbb{R}^n$ is given by,

$$\cos\theta = \frac{u \cdot v}{\|u\|\|v\|} \quad \theta \in [0, \pi]$$

Observe that in the previous definition, the angle is well defined.

$$-\|u\|\|v\| \leq -|u \cdot v| \leq u \cdot v \leq |u \cdot v| \leq \|u\|\|v\|$$

Dividing the entire inequality by $\|u\|\|v\|$ yields,

$$-1 \leq \frac{u \cdot v}{\|u\|\|v\|} \leq 1$$

Definition 14. A **hyperplane** \mathcal{H} in \mathbb{R}^n is the set of points $(x_1 \dots x_n)$ that satisfy $a_1x_1 + \dots + a_nx_n = b$ where $u = [a_1 \dots a_n] \in \mathbb{R}^n$ and $b \in \mathbb{R}$.

Definition 15. The **line** in \mathbb{R}^n passing through a point $P = (b_1 \dots b_n)$ and in the direction of $v \in \mathbb{R}^n$ with $v \neq 0$.

$$x = P + tv \quad t \in \mathbb{R}, \quad u = [a_1 \dots a_n]$$

$$\begin{cases} x_1 = a_1t + b_1 \\ x_n = a_nt + b_n \end{cases}$$

Chapter 2

Algebra of Matrices

2.1 Introduction

A matrix with n rows and m columns is written as,

$$A_{n \times m} = \begin{bmatrix} a_{11} & \dots & a_{1m} \\ \vdots & \ddots & \vdots \\ a_{n1} & \dots & a_{nm} \end{bmatrix}$$

Or,

$$A_{n \times m} = [a_{ij}]$$

Where a_{ij} is the entry in row i and column j .

2.2 Definitions and Properties of Matrices

Definition 16. Matrix Addition

$$A + B = [a_{ij} + b_{ij}] \quad \forall i = 1 \dots n, j = 1 \dots m$$

Definition 17. Scalar Multiplication

$$ka = [ka_{ij}] \quad \forall i = 1 \dots n, j = 1 \dots m$$

Definition 18. Zero Matrix

$$0 = [0]$$

Definition 19. Given a matrix $A_{m \times p}$ and a matrix $B_{p \times n}$, **matrix multiplication** is defined as,

$$AB = [c_{ij}] \quad c_{ij} = \sum_{k=1}^p a_{ik} b_{kj}$$

Definition 20. Given a matrix A , its **transpose** is $A^T = [a_{ji}]$ where $A = [a_{ij}]$.

Definition 21. A **square matrix** has the same number of rows as it does columns, i.e. $A_{n \times n}$ is a square matrix.

Definition 22. Given a matrix $A = [a_{ij}]$ the elements in the **diagonal** are $[a_{11}, \dots, a_{nn}]$.

Definition 23. The **trace** of a matrix A is given by,

$$\text{tr}(A) = a_{11} + \dots + a_{nn}$$

Definition 24. The **identity matrix** I_n is the matrix such that for any n -square matrix A ,

$$AI = IA = A$$

Definition 25. The **Kronecker delta** is defined by,

$$\delta_{ij} = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases}$$

Remark 2. Given the definitions for the identity matrix and the Kronecker delta, an alternative definition for the identity matrix is as follows,

$$I = [\delta_{ij}]$$

Definition 26. A matrix A is **invertible** if there is a matrix B such that $AB = BA = I$.

Remark 3. In general, for any matrices A and B , $AB \neq BA$.

Definition 27. A matrix D is **diagonal** if all the non-zero entries are in the diagonal.

$$D = \text{diag}(d_1, \dots, d_n)$$

Definition 28. A matrix A is **upper triangular** if,

$$a_{ij} = 0 \quad \forall i > j$$

2.3 Complex Numbers

The imaginary number i is defined as $i = \sqrt{-1}$ or equivalently, $i^2 = -1$.

Definition 29. A **complex number** z is given by,

$$z = a + bi \quad a, b \in \mathbb{R}$$

Where a is the real part and b is the imaginary part.

Real numbers are also complex numbers with no imaginary component, i.e.
 $a + 0i = a$.

Addition for two complex numbers $z = a + bi$ and $w = c + di$ is given by,

$$z + w = (a + c) + (b + d)i$$

Multiplication for the same two complex numbers is given by,

$$\begin{aligned} z \cdot w &= (a + bi)(c + di) \\ &= ac + adi + cbi - bd \\ &= (ac - bd) + (ad + bc)i \end{aligned}$$

Definition 30. The **conjugate** of $z = a + bi$ is $\bar{z} = a - bi$.

Definition 31. The **absolute value** or modulus of $z = a + bi$ is
 $|z| = \sqrt{a^2 + b^2}$.

Example 1.

$$z^{-1} = \frac{1}{z} = \frac{1}{a + bi} \cdot \frac{a - bi}{a - bi} = \frac{a - bi}{a^2 + b^2}$$

Observe that the following properties are true for conjugates and absolute values,

1.

$$z\bar{z} = |z|^2 = a^2 + b^2$$

2.

$$z \pm w = \bar{z} \pm \bar{w}$$

3.

$$z\bar{w} = \bar{z} \cdot \bar{w}$$

4.

$$(\bar{\bar{z}}) = z$$

5. z is real iff $z = \bar{z}$

6.

$$|zw| = |z||w|$$

7.

$$|z + w| \leq |z| + |w|$$

Chapter 3

Systems of Linear Equations

3.1 Representing Linear Systems with Matrices

Given a system of linear equations of the form,

$$\begin{cases} a_{11}x_1 + \dots + a_{1n}x_n = b_1 \\ \vdots \\ a_{m1}x_1 + \dots + a_{mn}x_n = b_m \end{cases}$$

- x_1, \dots, x_n are the unknowns, and
- a_{ij} and b_i are the constants.

The system can also be represented by matrices where,

- $A = [a_{ij}]$ is the matrix of coefficients
- $b = [b_i]$ is the column vector of constant
- $M = [A|b]$ is the matrix that represents the system.

Definition 32. A matrix A is in **echelon form** if

1. all zero rows are at the bottom, and
2. each leading non-zero entry in a row is to the right of the leading non-zero entry in the preceding row.

Example 2. This matrix is in echelon form. The boxed entries are the **pivots**.

$$A = \begin{bmatrix} 0 & \boxed{2} & 3 & 4 & 1 & 0 & 6 \\ 0 & 0 & 0 & \boxed{2} & 1 & 2 & 1 \\ 0 & 0 & 0 & 0 & 0 & \boxed{1} & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Definition 33. A matrix is said to be in the **row-reduced echelon form** if it is in the echelon form and,

1. each pivot is equal to 1, and
2. each pivot is the only non-zero entry in its column

Example 3. This matrix is in row-reduced echelon form. The boxed entries are the **pivots**.

$$A = \begin{bmatrix} 0 & \boxed{1} & 3 & 0 & 0 & 4 \\ 0 & 0 & 0 & \boxed{1} & 0 & -3 \\ 0 & 0 & 0 & 0 & \boxed{1} & 2 \end{bmatrix}$$

3.2 Elementary Row Operations

Suppose that A is a matrix with rows R_1, \dots, R_m . The elementary row operations that can be performed on A are as follows,

1. Row interchange, $R_i \leftrightarrow R_j$
2. Row scaling, $kR_i \rightarrow R_i$
3. Row addition, $kR_i + R_j \rightarrow R_j$

The method by which we find the (row-reduced) echelon form of a matrix is using the **Gaussian Elimination** algorithm.

Recall that every matrix is row equivalent to a unique matrix in the row-reduced echelon form.

Definition 34. The **rank** of a matrix $\text{rank}(A)$ is the number of pivots in the row-reduced echelon form. There are many other ways to define rank but they all have the same meaning.

The method by which we find the inverse of a square matrix A is as follows, Let $M = [A \mid I]$. Find the row-reduced echelon form of M . If there is a zero row in the resulting matrix then A is not invertible. Otherwise, $M \sim [I \mid B]$, $A^{-1} = B$.

Theorem 3. Let A be a square matrix. The following conditions are equivalent,

1. A is invertible
2. the row-reduced echelon form of A is I
3. the only solution to $Ax = 0$ is $x = 0$
4. the system $Ax = b$ has a solution for any choice of column b .

A partial proof is as follows,

Proof. (1) \Rightarrow (3) There is a matrix B such that $AB = I = BA$.
Let x be any solution of $Ax = [0]$.

$$\begin{aligned}BAx &= B[0] \\Ix &= [0] \\x &= [0]\end{aligned}$$

(1) \Rightarrow (4) Fix a column b ,

$$\begin{aligned}Ax &= b \\ \Leftrightarrow A^{-1}Ax &= A^{-1}b \\ \Leftrightarrow x &= A^{-1}b\end{aligned}$$

□

Definition 35. A linear system $Ax = b$ is **homogeneous** if $b = 0$. Otherwise, $Ax = b$ is said to be **non-homogeneous**.

Definition 36. A **particular solution** of $Ax = b$ is a vector x such that $Ax = b$. The set of all particular solutions is called the **general solution** of the solution set.

Definition 37. A system $Ax = b$ is **consistent** if it has one or more solutions and it is said to be **inconsistent** if it has no solutions.

Theorem 4. Any system $Ax = b$ has:

- (i) an unique solution,
- (ii) no solution, or
- (iii) an infinite number of solutions.

3.3 Examples

Example 4. The system,

$$\begin{aligned}x + y + 2z &= 1 \\ 3x - y + z &= -1 \\ -x + 3y + 4z &= 1\end{aligned}$$

is equivalent to,

$$\left[\begin{array}{ccc|c} 1 & 1 & 2 & 1 \\ 3 & -1 & 1 & -1 \\ -1 & 3 & 4 & 1 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 1 & 2 & 1 \\ 0 & 4 & 5 & 4 \\ 0 & 0 & 1 & -2 \end{array} \right]$$

Example 5. Back substitution:

$$z = -2$$

$$4y + 5z = 4 \Leftrightarrow y = \frac{7}{2}$$

$$x + y + 2z = 1 \Leftrightarrow x = \frac{3}{2}$$

$$\sim \left[\begin{array}{ccc|c} 1 & 0 & 0 & \frac{3}{2} \\ 0 & 1 & 0 & \frac{7}{2} \\ 0 & 0 & 1 & -2 \end{array} \right]$$

Example 6.

$$-2x + 3y + 3z = -9$$

$$3x - 4y + z = 5$$

$$-5x + 7y + 2z = -14$$

$$\sim \left[\begin{array}{ccc|c} -2 & 3 & 3 & -9 \\ 3 & -4 & 1 & 5 \\ -5 & 7 & 2 & -14 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & -1 & 4 & -4 \\ 0 & 1 & 11 & -17 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So there are infinitely many solutions.

Set $z = t$ since z is a free variable, then back substitute.

$$y = -17 - 11t \quad x = -21 - 15t \quad t \in \mathbb{R}$$

So the solution space is,

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -21 \\ -17 \\ 0 \end{bmatrix} + t \begin{bmatrix} -15 \\ -11 \\ 1 \end{bmatrix}$$

Where $(-21, -17, 0)$ is a particular solution and $(-15, -11, 1)$ is the set of basic solutions of the homogeneous system $Ax = 0$.

Example 7.

$$x + 2y - z = 2$$

$$2x + 5y - 3z = 1$$

$$x + 4y - 3z = 3$$

$$\left[\begin{array}{ccc|c} 1 & 2 & -7 & 2 \\ 2 & 5 & -3 & 1 \\ 1 & 4 & -3 & 3 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 2 & -1 & 2 \\ 0 & 1 & -1 & -3 \\ 0 & 0 & 0 & -1 \end{array} \right]$$

There are no solutions possible for this system.

Chapter 4

Vector Spaces

4.1 Introduction

Adding two vectors in \mathbb{R}^n produces a vector in \mathbb{R}^n . Similarly, multiplying by a scalar produces a vector in \mathbb{R}^n . These are some properties of a vector space, the following section is a formal list.

4.2 Basic Definitions

Definition 38. Let V be a non-empty set with two operations,

- i Vector addition: this assigns to any $u, v \in V$ the sum $u + v \in V$.
- ii Scalar multiplication: this assigns to any $u \in V$ and $k \in K$, a product $ku \in V$ where k is a field.

Then V is called a **vector** space (over the field K) if the following axioms hold for any $u, v, w \in V$.

- A1) $(u + v) + w = u + (v + w)$
- A2) there is a vector in V denoted by 0 called the zero vector such that $v + 0 = v$ for any $v \in V$.
- A3) for each $u \in V$, there is a vector in V denoted $-u$, such that $u + (-u) = 0$. $-u$ is called the negative of u .
- A4) $u + v = v + u$
- A5) $k(u + v) = ku + kv$ for any scalar $k \in K$
- A6) $(a + b)u = au + bu$ for any scalars $a, b \in K$
- A7) $(ab)u = a(bu)$ for any scalars $a, b \in K$

- A8) $1u = u$ for the unit scalar $k \in K$

Remark 4. A **field** K is a mathematical object with nice properties, with \mathbb{R} and \mathbb{C} being two examples. From now on, we will take it to be \mathbb{R} or \mathbb{C} .

4.3 Examples of Vector Spaces

Example 8. These are some examples of vector spaces,

1. \mathbb{R}^n

2. \mathbb{C}^n

3. The matrix space: $M_{m \times n}$

$M_{m \times n}$ denotes the set of all matrices with size m rows, n columns and real entries. $M_{m \times n}(\mathbb{C})$ permits the entries to be complex. The space of the real matrices are a subset of the space of complex matrices.

4. The polynomial space: $P(t)$

$P(t)$ denotes the set of all polynomials of the form,

$$P(t) = a_0 + a_1t + \dots + a_nt^n \mid a_i \in \mathbb{R}$$

5. The function space: $F(x)$

Let X be a non-empty set. Let $F(x)$ denote the set of all functions of X into \mathbb{R} . Then $F(x)$ is a vector space (over \mathbb{R}) with respect to the following operations,

i vector addition:

$$(f + g)(x) = f(x) + g(x) \mid \forall x \in X$$

ii scalar multiplication: for any $k \in K, f \in F(x)$

$$(kf)(x) = kf(x) \mid \forall x \in X$$

iii zero function: $\underline{0}(x) = 0$

Exercise 1. Consider the set \mathbb{R}^2 with the usual scalar multiplication, but with the following vector addition:

$$(a, b) \diamond (c, d) = (a + d, b + c)$$

Is this a vector space?

No because axiom 4 does not hold.

$$(1, 2) \diamond (-1, 1) = (2, 1)$$

$$(-1, 1) \diamond (1, 2) = (1, 2)$$

4.4 Vector Subspaces

Definition 39. Let V be a vector space and W be a subset of V . Then W is a **subspace** of V if W itself is a vector space with the operations of vector addition and scalar multiplication of V .

Example 9. $P(t)$ is a subspace of $F(\mathbb{R})$

The next theorem provides a simple criteria to show that a subset W of V is a subspace.

Theorem 5. Suppose that W is a subset of V , with V being a vector space. Then W is a subspace if the following two conditions hold:

- i The zero vector 0 belongs to W .
- ii For every two vectors $u, v \in W$ and $k \in R$
 - $u + v \in W$ (closed under vector addition)
 - $ku \in W$ (closed under scalar multiplication)

Proof. By (i), W is non-empty.

By (ii), the operations of vector addition and scalar multiplication are well defined.

It remains to prove each of the axioms of a vector space.

A1, 4, 5, 6, 7, and 8 hold in W because they hold in V .

A2 is true by (i).

A3: Let $v \in W$. We know that $-v \in V$ with $v + (-v) = 0$ by A3 for the vector space V . But W is closed under scalar multiplication (by (ii)) and so $v \in W$ and we are done. \square

4.5 Examples of Vector Subspaces

Example 10. These are some examples of vector subspaces,

1. $0, V$ are subspaces of V . These are called the trivial subspaces of V .
2. Subspaces of \mathbb{R}^3
 - i Line through the origin is a subspace.
 - ii Planes through the origin.
3. Subspaces of $P(t)$
 - i $P_m(t) = \{p(\cdot) \in P(t); \text{degree}(p(\cdot)) \leq m\}$
 - ii $Q(t)$ is the set of polynomials with only even powers
4. Subspaces of matrices $M_{m \times n}$

$$\text{i } W_1 = \{A \in M_{m \times n}; A \text{ is diagonal}\}$$

$$\text{ii } W_2 = \{A \in M_{m \times n}; A = A^T\}$$

5. Subspaces of $F(\mathbb{R})$

$$\text{i } C(\mathbb{R}) = \{f \in F(\mathbb{R}); f \text{ is continuous}\}$$

$$\text{ii } C'(\mathbb{R}) = \{f \in F(\mathbb{R}); f \text{ is differentiable}\}$$

4.6 More on Vector Spaces

Definition 40. Let $A \in M_{m \times n}$. The nullspace of A is $N(A)$ which is given by,

$$N(A) = \{x \in \mathbb{R}^n \mid Ax = 0\}$$

Proposition 1. $N(A)$ is a subspace of \mathbb{R}^n

Proof. Clearly $N(A)$ is a subset of \mathbb{R}^n

i $0 \in N(A)$. True because $A0 = 0$.

ii Let $u, v \in N(A)$, and $a, b \in \mathbb{R}$. We want to show that $au + bv \in N(A)$, which is the same as $A(au + bv) = 0$

$$\begin{aligned} A(au + bv) &= A(au) + A(bv) \\ &= a(Au) + b(Av) \quad \text{Since } u, v \in N(A), \text{ then } Au = 0, Av = 0 \\ &= a0 + b0 \\ &= 0 \end{aligned}$$

□

Remark 5. The solution set of a non-homogeneous system $\{x \in \mathbb{R}^n \mid Ax = b\}$ where $b \neq 0$ is not a subspace because the zero vector is not present.

Theorem 6. Let U and W be subspaces of a vector space V . Then $U \cap W$ is also a subspace.

Proof. Since $U \subseteq V$ and $W \subseteq V$ (U and W are subspaces),

$$U \cap W \subseteq V$$

i So, $0 \in V$ and $0 \in W$, therefore $0 \in U \cap W$

ii Let $u, v \in U \cap W$ and $a, b \in \mathbb{R}$

$$\begin{aligned} u, v \in U \cap W &\Rightarrow \begin{cases} u, v \in V \\ u, v \in W \end{cases} \\ &\Rightarrow \begin{cases} au + bv \in V \\ au + bv \in W \end{cases} \quad \text{both } U \text{ and } W \text{ are subspaces} \\ &\Rightarrow au + bv \in U \cap W \end{aligned}$$

□

Remark 6. In general, if U and W are subspaces, $U \cup W$ is **not** a subspace. An example would be two lines through the origin in \mathbb{R}^3 .

4.7 Linear Combinations

Observe that $au + bv$ is a linear combination.

Definition 41. Let U be a vector space. A vector $v \in V$ is a **linear combination** of u_1, \dots, u_m in V if there exists scalars a_1, \dots, a_m so that,

$$v = a_1u_1 + \dots + a_mu_m$$

Example 11. The following is an example of linear combinations in \mathbb{R}^3 . Is $v = (1, 5, 5) \in \mathbb{R}^3$ a linear combination of $u_1 = (1, 2, 3)$, $u_2 = (1, 0, 1)$, $u_3 = (0, 1, 0)$?

That is the same as asking, are there constants $a, b, c \in \mathbb{R}$ such that,

$$v = au_1 + bu_2 + cu_3$$

That is, are there $a, b, c \in \mathbb{R}$ such that,

$$\begin{bmatrix} 1 \\ 5 \\ 5 \end{bmatrix} = a \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + b \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + c \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

Yes. Take $a = 2, b = -1, c = 1$.

Definition 42. Let $A \in M_{m \times n}$. The column space of A is $C(A)$ which consists of all linear combinations of the columns of A . Alternatively,

$$C(A) = \{Ax \mid x \in \mathbb{R}^n\}$$

Proposition 2. The linear system $Ax = b$ is consistent iff $b \in C(A)$.

Example 12. The following is an example of linear combinations in $P(t)$. Is the polynomial $P(t) = t^2 + 5t + 5$ a linear combination of the polynomials $P_1(t) = t^2 + 2t + 3$, $P_2(t) = t^2 + 1$, $P_3(t) = t$? Equivalently, are there scalars $a, b, c \in \mathbb{R}$ such that,

$$p(\cdot) = ap_1(\cdot) + bp_2(\cdot) + cp_3(\cdot)$$

There are two ways of solving this,

1. Matching coefficients:

$$t^2 + 5t + 5 = (a + b)t^2 + (2a + c)t + 3a + b$$

$$\begin{cases} 1 = a + b \\ 5 = 2a + c \\ 5 = 3a + b \end{cases} \Leftrightarrow \begin{bmatrix} 1 & 1 & 0 \\ 2 & 0 & 1 \\ 3 & 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 1 \\ 5 \\ 5 \end{bmatrix}$$

2. "Trial approach": We set t in $P(t)$ equal to the three distinct values and each one provides a different equation,

$$\begin{array}{rcl} t = 0 & 5 & = 3a + b \\ t = 1 & 11 & = 6a + 2b + c \\ t = -1 & 1 & = 2a + 2b - c \end{array}$$

Then solve for a, b, c .

Example 13. The following are two examples of subspaces of \mathbb{R}^3

1. A line with direction $(1, 2, 3)$ through the origin,

$$\left\{ t \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \mid t \in \mathbb{R} \right\}$$

2. A plane through the origin,

$$\begin{aligned} \{(x, y, z) \in \mathbb{R}^3 \mid z = 0\} &= \{t(1, 0, 0) + s(0, 1, 0) \mid t, s \in \mathbb{R}\} \\ &= \{(t, s, 0) \mid t, s \in \mathbb{R}\} \end{aligned}$$

4.8 The Span of a Vector Space

Definition 43. Let u_1, \dots, u_m be vectors in V . The set of all linear combinations of u_1, \dots, u_m is called the **span** of u_1, \dots, u_m and is denoted by $\text{span}\{u_1, \dots, u_m\}$.

$$\text{span}\{u_1, \dots, u_m\} = \{t_1 u_1 + \dots + t_m u_m \mid t_1, \dots, t_m \in \mathbb{R}\}$$

Definition 44. The vectors $u_1, \dots, u_m \in V$ are said to span V or to form a **spanning set** of V if,

$$\text{span}\{u_1, \dots, u_m\} = V$$

The following are the properties of spans.

- If $\text{span}\{u_1, \dots, u_m\} = V$, then for any $v \in V$, $\text{span}\{v, u_1, \dots, u_m\} = V$.
- If $\text{span}\{0, u_1, \dots, u_m\} = V$, then $\text{span}\{u_1, \dots, u_m\} = V$.
- If $\text{span}\{u_1, \dots, u_m\} = V$ and u_k is a linear combination of $u_1, \dots, u_{k-1}, u_{k+1}, \dots, u_m$ then $\text{span}\{u_1, \dots, u_{k-1}, u_{k+1}, \dots, u_m\} = V$.

Proposition 3. Let u_1, \dots, u_m be vectors in V . Then $\text{span}\{u_1, \dots, u_m\}$ is a subspace.

Proof. Clearly $\text{span}\{u_1, \dots, u_m\} \subseteq V$.

We know $0 \in \text{span}\{u_1, \dots, u_m\}$ since,

$$0 = 0u_1 + \dots + 0u_m \in \text{span}\{u_1, \dots, u_m\}$$

Take any $u, v \in \text{span}\{u_1, \dots, u_m\}$ and $a, b \in \mathbb{R}$.

$$u = a_1u_1 + \dots + a_mu_m \quad a_1, \dots, a_m \in \mathbb{R} \text{ since } u \in \text{span}\{u_1, \dots, u_m\}$$

Likewise,

$$v = b_1u_1 + \dots + b_mu_m \quad b_1, \dots, b_m \in \mathbb{R}$$

So,

$$\begin{aligned} au + bv &= aa_1u_1 + \dots + aa_mu_m + bb_1u_1 + \dots + bb_mu_m \\ &= (aa_1 + bb_1)u_1 + \dots + (aa_m + bb_m)u_m \end{aligned}$$

Which shows that $au + bv$ is a linear combination of u_1, \dots, u_m with scalars $aa_1 + bb_1, \dots, aa_m + bb_m$. \square

Exercise 2. $W = \{(x, y, z) \in \mathbb{R}^3 \mid x = 2y = 3z\}$

Clearly, $W \subseteq \mathbb{R}^3$

$(0, 0, 0) \in W$ is true because $0 = 2(0) = 3(0)$.

$u, v \in W \quad a, b \in \mathbb{R}$

$u = (u_1, u_2, u_3)$ with (1) $u_1 = 2u_2$ and (2) $2u_2 = 3u_3$

$v = (v_1, v_2, v_3)$ with (3) $v_1 = 2v_2$ and (4) $2v_2 = 3v_3$

$$\begin{aligned} z &= (z_1, z_2, z_3) = au + bv \\ &= (au_1 + bv_1, au_2 + bv_2, au_3 + bv_3) \end{aligned}$$

We want to show $z \in W$, so,

$$\begin{aligned} z_1 &= 2z_2 = 3z_3 \\ z_1 &= au_1 + bv_1 \\ &= a(2u_2) + b(2v_2) \quad \text{by (1) and (3)} \\ &= 2(au_2 + bv_2) \\ &= 2z_2 \\ &= a(3u_3) + b(3v_3) \\ &= 3(au_3 + bv_3) \\ &= 3z_3 \end{aligned}$$

So, $z_1 = 2z_2 = 3z_3$.

It is also a line through the origin,

$$x = 2y = 3z \Rightarrow$$

$$\left\{ t \begin{bmatrix} 6 \\ 3 \\ 2 \end{bmatrix} \mid t \in \mathbb{R} \right\} = \text{span}\{(6, 3, 2)\}$$

Definition 45. The **span of a set** S is the set of all linear combinations of vectors in S . If $S \neq \{0\}$, $\text{span}(S) = \{0\}$.

Theorem 7. Let S be a subset of the vector space V . Then,

- i) $\text{span}(S)$ is a subspace of V .
- ii) if W is a subspace of V such that $S \subseteq W$, then $\text{span}(S) \subseteq W$.

Proof here

4.9 The Row Space of a Matrix

Definition 46. Let $A \in M_{m \times n}$. The **row space** of A , written as $\text{rowsp}(A)$, is the set of all linear combinations of rows of A .

$$\text{rowsp}(A) = \text{col}(A^T)$$

The notation for column space can also be $\text{colsp}(A^T)$. $A \in M_{m \times n}$. $\text{rowsp}(A)$ is a subspace of \mathbb{R}^n . $\text{col}(A)$ is a subspace of \mathbb{R}^m .

Example 14. Two matrices are row equivalent if you can get from one to the other with only elementary row operations.

$$A = \begin{bmatrix} 1 & 2 & -1 & 3 \\ 2 & 4 & -1 & 5 \\ 3 & 6 & -2 & 8 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 & -1 & 3 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix} = B$$

The following is true for A and B ,

- $\text{rowsp}(A) = \text{rowsp}(B)$
- $\text{rowsp}(A) = \text{span}(1, 2, -1, 3), (2, 4, -1, 5), (3, 6, -2, 8)$
- $\text{rowsp}(B) = \text{span}(1, 2, -1, 3), (0, 0, 1, -1)$

Observe that any basis of a subspace is not unique.

Theorem 8. Row equivalent matrices have the same row space.

4.10 Linear Dependence and Independence

Definition 47. The vectors v_1, \dots, v_n are **linearly independent** if the following condition is satisfied,

$$\text{if } a_1 v_1 + \dots + a_n v_n = 0, \text{ then } a_1 = \dots = a_n = 0$$

The vectors v_1, \dots, v_n are **linearly dependent** if they are not linearly independent.

Remark 7. Consider the vector equation,

$$x_1v_1 + \dots + x_nv_n = 0$$

where x_1, \dots, x_n are unknown scalars. If the only solution is $(0, \dots, 0)$, then the vectors are linearly independent. Otherwise they are linearly dependent.

Example 15. The following is an example of linear dependence in \mathbb{R}^3 . Geometrically, linearly dependent vectors run in the same direction.

- i) Two vectors in \mathbb{R}^3 are linearly dependent if they lie on the same line. i.e., $k \in \mathbb{R}, k \neq 0$

$$v_2 = kv_1 \Leftrightarrow kv_1 - v_2 = 0$$

- ii) Three vectors in \mathbb{R}^3 are linearly dependent if they lie on the same plane. i.e., $a_1, a_2 \in \mathbb{R}, a_1, a_2 \neq 0$

$$v_3 = a_1v_1 + a_2v_2 \Leftrightarrow a_1v_1 + a_2v_2 - v_3 = 0$$

Definition 48. An infinite set of vectors S is linearly dependent if there exist vectors $v_1, \dots, v_n \in S$ that are linearly dependent.

Proposition 4. Let V be a vector space.

- i) If $v \neq 0, \{v\}$ is linearly independent.
- ii) No independent set of vectors contains the zero vector. Any non-zero scalar multiplied by the zero vector will still yield the zero vector.
- iii) Two vectors are linearly dependent iff one of them is a multiple of the others. Let v_1, v_2 be linearly dependent and $a_1 \neq 0$.

$$\begin{aligned} a_1v_1 + a_2v_2 &= 0 \\ \Rightarrow v_1 + \frac{a_2}{a_1}v_2 &= 0 \\ \Leftrightarrow v_1 &= \frac{-a_2}{a_1}v_2 \end{aligned}$$

- iv) No independent set can contain two vectors that are multiples of each other.

Exercise 3. Show that $1 + t, 3t + t^2, 2 + t - t^2$ is linearly independent in $P_2(t)$.

Suppose,

$$a(1 + t) + b(3t + t^2) + c(2 + t - t^2) = 0 \quad \forall t \in \mathbb{R}$$

We want to show $a = b = c = 0$.

Substitute three different values for t to obtain three equations, then solve.

$$\begin{aligned} t = 0 & \quad a + b(0) + 2(c) = 0 \Leftrightarrow a + 2c = 0 \\ t = -1 & \quad -2b = 0 \Leftrightarrow b = 0 \\ t = 1 & \quad 2a + 0 + 2c = 0 \Leftrightarrow 2a + 2c = 0 \quad \text{zero term from } t = -1 \end{aligned}$$

So, $a + 2c = 0$ (1) and $2a + 2c = 0$ (2). (2)-(1) gives $a = 0$.

And we are done.

The alternate method is to match coefficients and solve that system as in the example in 4.7.

Proposition 5. The vectors v_1, \dots, v_n are linearly dependent iff one of them is a linear combination of another.

Proof. (\Rightarrow)

The vectors are linearly dependent. Then, $\forall a_1, \dots, a_n \exists a_i \neq 0$ such that $a_1 v_1 + \dots + a_n v_n = 0$.

Say $a_k \neq 0$.

$$\begin{aligned} a_1 v_1 + \dots + a_n v_n &= 0 \\ \Leftrightarrow \frac{a_1}{a_k} v_1 + \dots + \frac{a_{k-1}}{a_k} v_{k-1} + v_k + \frac{a_{k+1}}{a_k} v_{k+1} + \dots + \frac{a_n}{a_k} v_n &= 0 \\ \Leftrightarrow v_k &= -\frac{a_1}{a_k} v_1 - \dots - \frac{a_{k-1}}{a_k} v_{k-1} - \frac{a_{k+1}}{a_k} v_{k+1} - \dots - \frac{a_n}{a_k} v_n \end{aligned}$$

Observe that v_k can be any vector including the zero vector.

(\Leftarrow)

Say v_i is a linear combination of the other vectors. Then,

$$\begin{aligned} v_i &= b_1 v_1 + \dots + b_{i-1} v_{i-1} + b_{i+1} v_{i+1} + \dots + b_n v_n \\ \Leftrightarrow 0 &= b_1 v_1 + \dots + b_{i-1} v_{i-1} - v_i + b_{i+1} v_{i+1} + \dots + b_n v_n \end{aligned}$$

So, the scalar on v_i is -1 which is not zero.

And we are done. □

4.11 Basis

Definition 49. A set $B = \{u_1, \dots, u_n\}$ of vectors in V is a **basis** of V if two conditions are satisfied,

- i) B is a linearly independent set
- ii) $\text{span}(B) = V$

Example 16. The following are examples of basis,

1) \mathbb{R}^n

$$e_1 = (1, 0, \dots, 0), e_2 = (0, 1, 0, \dots, 0), e_n = (0, \dots, 0, 1)$$

$\{e_1, \dots, e_n\}$ is a basis for \mathbb{R}^n .

$(1, 2, 3) = e_1 + 2e_2 + 3e_3$. This is the canonical or standard basis.

2) $P_m(t)$

$$\{1, t, t^2, \dots, t^m\}$$

3) $M_{m \times n}$ For $M_{2 \times 3}$ it is,

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Observe that a basis can also be the minimum span of a vector space.

Example 17.

$$A = \begin{bmatrix} 1 & 2 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\{R_1, R_2, R_3\}$ is linearly independent as none of the rows are linear combinations of the others.

Observe that R_1, R_2, R_3 spans the row space of A .

Theorem 9. *The non-zero rows of a matrix in echelon form are linearly independent and form a basis for the row space.*

Proposition 6. A set $B = \{v_1, \dots, v_n\}$ is a basis of V iff every vector $v \in V$ can be uniquely written as a linear combination of v_1, \dots, v_n .

Proof. (\Rightarrow)

Suppose $v = a_1v_1 + \dots + a_nv_n$ $a_1, \dots, a_n \in \mathbb{R}$ and

$v = b_1v_1 + \dots + b_nv_n$ $b_1, \dots, b_n \in \mathbb{R}$.

We want to show that $a_i = b_i \quad \forall i = 1 \dots n$.

We have,

$$\begin{aligned} a_1v_1 + \dots + a_nv_n &= b_1v_1 + \dots + b_nv_n \\ \Leftrightarrow 0 &= (a_1 - b_1)v_1 + \dots + (a_n - b_n)v_n \end{aligned}$$

Since $\{v_1, \dots, v_n\}$ is linearly independent,

$$\begin{aligned} a_1 - b_1 &= 0, \dots, a_n - b_n = 0 \\ \Leftrightarrow a_1 &= b_1, \dots, a_n = b_n \end{aligned}$$

(\Leftarrow)

Suppose $a_1v_1 + \dots + a_nv_n = 0$. We have $a_1, \dots, a_n \in \mathbb{R}$. To show that B is linearly independent, we must show $a_1 = \dots = a_n = 0$,

$$0 = 0(v_1) + \dots + 0(v_n)$$

We assumed the linear combination is unique,

$$a_1 = \dots = a_n = 0$$

We assumed every vector can be written as a linear combination. B spans V by the assumption.

And we are done. \square

4.12 Coordinates

Definition 50. Let V be a vector space and $B = \{v_1, \dots, v_n\}$ a basis of V . Then for any $v \in V$,

$$v = a_1 v_1 + \dots + a_n v_n$$

where the a_1, \dots, a_n are unique for v .

We call these scalars the **coordinates** of v in the basis B and they form a vector $[a_1, \dots, a_n]$ called coordinate vectors of v relative to B and denoted by $[v]_B$.

Example 18. $t + 1, t - 1, (t - 1)^2$ form a basis of $P_2(t)$. This can be written as :

$$P(t) = (t + 1) - (t - 1) + (t - 1)^2 = t^2 - 2 + 2$$

or as coordinates: $[P(\cdot)]_B = [1, -1, 1]$.

4.13 Dimension

Next we will give a series of auxiliary lemmas and propositions to show that the size of any two basis for a vector space has the same number of vectors.

Proposition 7. Let V be a vector space and $S = \{v_1 + \dots + v_n\}$ be a spanning set of V . Then,

- i) if $w \in V$, $\{w, v_1, \dots, v_n\}$ is linearly dependent and spans V .
- ii) if v_i is a linear combination of the other vectors, S without v_i still spans V .

Proposition 1. Suppose $\{v_1, \dots, v_k\}$ is linearly dependent and all the vectors are non-zero. Then one of the vectors is a linear combination of the preceding vectors.

Proof. Then there are scalars, not all zero such that,

$$a_1 v_1 + \dots + a_n v_n = 0 \quad \text{where } a_1, \dots, a_k \in \mathbb{R}$$

Let i be the largest index such that $a_i \neq 0$. We claim $i > 1$. If $i = 1$, then $a_1 v_1 = 0$, which is a contradiction since $v_1 \neq 0$.

$$\begin{aligned} \Rightarrow \quad & a_1 v_1 + \dots + a_i v_i = 0 \\ \Leftrightarrow \quad & v_i = -\frac{a_1}{a_i} v_1 - \dots - \frac{a_{i-1}}{a_i} v_{i-1} \end{aligned}$$

□

Proposition 2. "Replacement Lemma".

Suppose $\{v_1, \dots, v_n\}$ spans V and $\{w_1, \dots, w_m\}$ is linearly independent. Then $m \leq n$ and V is spanned by a set of the form ,

$$\{w_1, \dots, w_m, v_{i_1}, \dots, v_{i_{n-m}}\}$$

Thus any $n + 1$ or more vectors in V are linearly dependent.

Proof. (General idea) In $\text{span}\{v_1, \dots, v_n\} = V$, we add w_1 .

$\text{span}\{w_1, v_1, \dots, v_n\} = V$ but it is now linearly dependent. Assume v_1, \dots, v_n are all non-zero.

By proposition and lemma from above, we can remove v_i so that

$$\text{span}\{w_1, v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n\} = V$$

Repeat the above steps when adding w_2 and removing v_k and we get,

$$\text{span}\{w_1, w_2, v_1, \dots, v_{k-1}, v_{k+1}, \dots, v_{i-1}, v_{i+1}, \dots, v_n\} = V$$

If $m \leq n$,

$$\text{span}\{w_1, \dots, w_m, v_{i_1}, \dots, v_{i_{n-m}}\} = V$$

Suppose $m > n$,

$$\text{span}\{w_1, \dots, w_m\} = V$$

If we add w_{n+1} , then $\text{span}\{w_1, \dots, w_n, w_{n+1}\} = V$ and is linearly dependent.

But this is a contradiction, w_1, \dots, w_m were all linearly independent. The only possible case is then $m \leq n$. □

Remark 8. The Replacement Lemma tells us that the size of any spanning set is at least as big as the size of any linearly independent set.

The preceding lemmas and propositions in this section were auxiliary to proving the following theorem.

Theorem 10. Let $\{u_1, \dots, u_m\}$ and $\{v_1, \dots, v_n\}$ be basis for V . Then $m = n$.

Proof. The proof is relatively simple as we have the Replacement Lemma.

$\{u_1, \dots, u_m\}$ is (1) linearly independent and (2) spans V .

$\{v_1, \dots, v_n\}$ (3) spans V and (4) is linearly independent.

Apply the remark on the Replacement Lemma twice to statements (1) with (3) and (2) with (4). Since, $m \leq n$ and $m \geq n$, then $n = m$. □

Definition 51. A vector space is said to be of **finite dimension** or **n-dimensional**, written $\dim(V) = n$ if V has a basis with n elements. The vector space $\{0\}$ has dimension 0. If a vector V doesn't have a finite basis, then V is said to be of infinite dimension or infinite dimensional.

Example 19. The following are examples of dimensions,

- 1) $\dim(\mathbb{R}^n) = n$
- 2) $\dim(M_{m \times n}) = mn$
- 3) $\dim(P_n(t)) = n + 1$ because of (t^0, t^1, \dots, t^n)
- 4) $\dim(P(t)) = \infty$

To find the basis of a set of vectors, put the vectors into matrix rows and find the echelon form.