

Higher Algebraic K-Theory

David Zhu

August 19, 2024

K -Theory started with Grothendieck's K_0 and projective modules/vector bundles. In this note, we present the various constructions for higher K -groups.

1 The Group Completion Theorem

First, we recall Grothendieck's definition of K_0 for an abelian monoid M

Definition 1.0.1. Let M be an abelian monoid. Then, the Grothendieck group $K_0(M)$ is an abelian group K with the inclusion map $i : M \rightarrow K$ satisfying the following universal property: given an abelian group A and a monoid morphism $f : M \rightarrow A$, we have the factorization

$$\begin{array}{ccc} M & \xrightarrow{f} & A \\ \downarrow i & \nearrow \exists! g & \\ K & & \end{array}$$

It is an easy exercise to show that K is unique up to isomorphism, which we will denote by $K_0(M)$. Explicitly, we can obtain $K_0(M)$ from the following "group completion" construction:

Proposition 1.0.1. Given an abelian monoid $K_0(M)$ is the abelian group generated by $[m]$ for each $m \in M$, modulo the relation $[x + y] = [x] + [y]$.

Example 1.0.1. (K_0 of a ring) Given a ring R , $K_0(R)$ is defined to be Grothendieck group over the abelian monoid of the isomorphism class of finitely generated projective modules over R , with monoid operation given by direct sum.

If M is a topological monoid, let BM be its classifying space (viewing M as a category with one object). Then, there is a natural map $M \rightarrow \Omega BM$, with $\pi_0(\Omega BM) = \pi_1(BM)$ an abelian group. When π_0 is a group, it can be shown that $M \rightarrow \Omega BM$ is a homotopy equivalence. Thus the map is referred to as the group completion of a topological monoid. When π_0 is not necessarily group, many can still be said: since M is an H -space, $H_*(M)$ is naturally a ring, and $H_0(M) = \mathbb{Z}[\pi_0(M)]$. Viewing $\pi_0(M)$ as a multiplicative subset of $H_0(M)$, the induced map $H_*(M) \rightarrow H_*(\Omega BM)$ sends $\pi_0(M)$ to units. McDuff and Segal proved the following result:

Theorem 1.1. If $\pi := \pi_0(M)$ is in the center of $H_*(M)$, then

$$H_*(M)[\pi^{-1}] \cong H_*(\Omega BM)$$

We will outline the idea of the proof following the original paper [MS76], as it provides many insights to the motivate the $S^{-1}S$ -construction.

The goal is to find an intermediate space M_∞ with prescribed homology $H_*(M_\infty) = H_*(M)[\pi^{-1}]$ and a homology equivalence $M_\infty \rightarrow \Omega BM$. The first step uses the Quillen's lemma, given in [Fri94], regarding localization:

Lemma 1.2. Let R be a ring (not necessarily commutative) and S a multiplicative subset. Let C be the category with objects elements of S and morphisms from s_1 to s_2 an element $t \in S$ such that $s_1 t = s_2$. Under the conditions that C is filtered, there is a canonical R -module morphism

$$u : \varinjlim_C R \rightarrow R[S^{-1}]$$

where the filtered colimit is defined on objects by the inclusion map, and on morphisms by right multiplication by t . If S acts on R by left multiplication bijectively, meaning

1. (injective) Given $r \in R$ and $s \in S$ such that $sr = 0$, then there exists $t \in S$ with $rt = 0$.
2. (bijective) Given $r \in R$ and $s \in S$, there exists $r' \in R$ and $t \in S$ such that $sr' = rt$

Then u is an isomorphism.

A simple example is colimit $\mathbb{Z} \xrightarrow{\times p} \mathbb{Z} \xrightarrow{\times p} \dots$ being isomorphic to $\mathbb{Z}[1/p]$ as \mathbb{Z} -modules. The hypothesis of the lemma is satisfied when π is in the center of $H_*(M)$, and we can define M_∞ to be the mapping telescope given by $M \xrightarrow{\times m} M \xrightarrow{\times m} \dots$, where m is any arbitrary element in the component of $1 \in \pi_0(M)$. Since homology commutes with filtered colimits, M_∞ will have the prescribed homology.

Given a topological group M , we may construct the universal bundle $EM \rightarrow BM$ by viewing M as a topological category. If M acts on a space X , let X_M denote the associated bundle to the universal bundle $X_M \rightarrow BM$ with fiber X . The construction still holds when M is only a monoid, and instead of the homotopy equivalent between the fiber and homotopy fiber, McDuff and Segal recovers the following proposition

Proposition 1.2.1. If M is a topological monoid which acts on a space X , and for each $m \in M$ the map $x \mapsto mx$ from X to itself is a homology equivalence, then the map $X_M \rightarrow BM$ is a homology fibration with fibre X , meaning the canonical map between the fiber and homotopy fiber induces isomorphism on homology.

By construction, M acts on M_∞ (which also induces an homology equivalence), and $(M_\infty)_M$ is also contractible since it will be a filtered colimit of the contractible $M_M = EM$. Thus, we have a map $(M_\infty)_M \rightarrow BM$ with fiber M_∞ and homotopy fiber ΩBM . The theorem then follows from proposition 1.2.1 and more colimit nonsense on components for the general case.

We now define group completion for general homotopy commutative, homotopy associative H -spaces.

Definition 1.2.1. Let X be a homotopy commutative, homotopy associative H -space. A group completion of X is an H -space map $f : X \rightarrow Y$ such that f induces the group completion on $\pi := \pi_0$, and the isomorphism

$$H_*(X)[\pi^{-1}] \cong H_*(Y)$$

2 The $S^{-1}S$ -construction

Given a symmetric monoidal category S , Quillen constructs the category $S^{-1}S$ such that $B(S^{-1}S)$ is the group completion of BS in the sense of definition 1.2.1. The group completion recovers the plus construction, and gives another more natural definition for higher K -theory. We shall note that the construction below is closely related to the group completion construction in chapter 1.

A left action of a monoidal category S on a category X is a functor $+$: $S \times X \rightarrow X$ with natural isomorphisms $A + (B + F) \cong (A + B) + F$ and $0 + F \cong F$, where $A, B \in S$ and $F \in X$, satisfying the associativity coherence conditions. An action is called invertible if each translation $F \mapsto A + F$ is a homotopy equivalence.

Definition 2.0.1. If S acts on X the category $\langle S, X \rangle$ has the same objects as X . A morphism is represented by an isomorphism class of the tuple $(s, sx \xrightarrow{\phi} t)$, where ϕ is a morphism in X . An isomorphism of tuples is given by an isomorphism $s \cong s'$, which induces the commutative diagram

$$\begin{array}{ccc} s + x & \xrightarrow{\cong} & s' + x \\ & \searrow & \swarrow \\ & t & \end{array}$$

We let $S^{-1}S := \langle S, S \times S \rangle$, where S acts on both factors of $S \times S$ by the monoidal operation. We also define an action of S on $S^{-1}S$ by $s + (s_1, s_2) = (s_1, s + s_2)$. Note that this action is invertible: the translation $(s_1, s_2) \mapsto (s_1, s + s_2)$ has homotopy inverse $(s_1, s_2) \mapsto (s + s_1, s_2)$, since there is the natural transformation $(s_1, s_2) \mapsto (s + s_1, s + s_2)$. Quillen then proves

Theorem 2.1. If every map in S is an isomorphism, and the translations are faithful in S , then the functor $S \rightarrow S^{-1}S$ given by $x \mapsto (0, x)$ induces isomorphism

$$H_*(S)[\pi^{-1}] \cong H_*(S^{-1}S)$$

In particular, $BS^{-1}S$ is the group completion of BS .

Proof. It is not hard to show that the projection functor $S^{-1}S \rightarrow \langle S, S \rangle$ onto the second factor is cofibered with fiber S . Then, there is a spectral sequence [Gra76]

$$E_{p,q}^2 = H_p(\langle S, S \rangle, H_q(S)) \Rightarrow H_{p+q}(S^{-1}S)$$

Localizing at $\pi_0(S)$ and noting the contractibility of $\langle S, S \rangle$ gives the desired degeneration. \square

We now show that the plus construction for $BGL(R)$ is a group completion.

Theorem 2.2. Let $S = \coprod GL_n(R)$ be the category of free R -modules. Then, $BS^{-1}S$ is the group completion of $BS = \coprod BGL_n(R)$, and

$$BS^{-1}S \cong \mathbb{Z} \times BGL(R)^+$$

Proof. We will construct a map $f : BGL(R) \rightarrow Y_S$, where Y_S is the connected component of $BS^{-1}S$ containing the basepoint, such that f induces isomorphism on integral homology. The universal property of the plus construction then implies a homotopy equivalence $BGL(R)^+ \rightarrow Y_S$. □

3 Exact Category

Definition 3.0.1. An **exact category** is an additive category \mathcal{M} equipped with a class \mathcal{E} of short exact sequences of the form

$$M' \xrightarrow{i} M \xrightarrow{j} M''$$

where the first arrow i is denoted an **admissible monomorphism**, and the second arrow j is denoted an **admissible epimorphism**. In addition, the class \mathcal{E} also satisfies the following properties:

1. (closed under trivial extension) For any M', M'' in $\text{ob}(\mathcal{M})$, the SES

$$M' \xrightarrow{(id, 0)} M' \oplus M'' \xrightarrow{pr_2} M''$$

is in \mathcal{E} .

2. The class of admissible epimorphism is closed under composition. Dually for admissible monomorphisms.
3. (closed under base-change) If $M \rightarrow M''$ is an admissible epimorphism and given $N \rightarrow M$, the pullback square exists

$$\begin{array}{ccc} N \times_{M''} M & \longrightarrow & M \\ \downarrow p & & \downarrow \\ N & \longrightarrow & M'' \end{array}$$

and the morphism p is an admissible epimorphism. Dually for admissible monomorphisms.

4. (admissible epimorphism is "epimorphism") Let $M \rightarrow M''$ be a map possessing a kernel in \mathcal{M} . If there exists a map $N \rightarrow M$ in \mathcal{M} such that $N \rightarrow M \rightarrow M''$ is an admissible epimorphism, then $M \rightarrow M''$ is an admissible epimorphism. Dually for admissible monomorphisms.

This is where we want to do K -theory. The motivation for exact categories is the following scenario: consider any additive category \mathcal{M} embedded as a full subcategory of an abelian category \mathcal{A} . Suppose further that \mathcal{M} is closed under taking extensions in \mathcal{A} , meaning if

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

is exact in \mathcal{A} and A, C is in \mathcal{M} , then B is also in \mathcal{M} . Then \mathcal{M} can be readily verified to be an exact category, with \mathcal{E} being the class of sequences in \mathcal{M} that is short exact in \mathcal{A} . The only non-trivial thing to check is axiom 3, which is a standard theorem regarding pullbacks. We now have a wealth of examples in algebra/algebraic geometry:

Example 3.0.1. The category $\mathbf{P}(\mathbf{R})$ of finitely generated projective modules over a commutative ring R is exact by its embedding in \mathbf{RMod} . Note that it is generally not abelian due to lacking of kernel/cokernels. For example $\mathbf{P}(\mathbb{Z})$ is not abelian, since $\mathbb{Z} \xrightarrow{\times 2} \mathbb{Z}$ does not have a cokernel.

Example 3.0.2. The category $\mathbf{VB}(X)$ over a paracompact space X is exact by its embedding in the category of family of vector spaces over X . It is also generally not abelian due to lacking kernels.

Definition 3.0.2. An exact functor $F : \mathcal{M} \rightarrow \mathcal{M}'$ between exact categories is an additive functor preserving exact sequences.

4 The Q -construction and Recovery of K_0

To build K -theory on a exact category such as $\mathbf{P}(\mathbf{R})$, we have to go through an intermediate category, which is called the Q -construction.

Definition 4.0.1. Given an exact category \mathcal{M} , let the category $Q\mathcal{M}$ have the same objects as \mathcal{M} , and morphisms from M to M' being isomorphisms classes of diagrams of the form

$$M \leftarrow N \rightarrow M'$$

where \leftarrow signifies an admissible epimorphism and \rightarrow an admissible monomorphism in \mathcal{M} . An isomorphism between diagrams of the form is one that induces identity on both M and M' . Composition of a morphisms is given by the pullback

$$\begin{array}{ccccc} N \times_{M'} N' & \xrightarrow{pr_2} & N' & \xrightarrow{i'} & M'' \\ \downarrow pr_1 & & \downarrow j' & & \\ N & \xrightarrow{i} & M' & & \\ \downarrow i & & & & \\ M & & & & \end{array}$$

It is clear that the composition is associative, and we have a well-defined category. Here are a few preliminary observations: 1. the classifying space $BQ\mathcal{M}$ is canonically a CW complex, and it is path-connected by the existence of a zero object, denoted 0, in \mathcal{M} . 2. If $i : M' \rightarrow M$ is an admissible monomorphism, then it induces a morphism in $Q\mathcal{M}$ denoted by $i_!$ given by

$$M' = M' \rightarrow M$$

which will be referred to as **injective maps**. Dually, If $j : M'' \rightarrow M$ is an admissible epimorphism, then it induces a morphism in $Q\mathcal{M}$ denoted by $j^!$ given by

$$M \leftarrow M'' = M''$$

which are called **surjective maps**. Note the superscript/subscript follows the contravariant/covariance convention. Then, each morphism u in $Q\mathcal{M}$ is the composition of $i_! \circ j^!$ for some i and j in \mathcal{M} (check the pullback diagram). We will abuse notation onwards and use the same arrows corresponding to admissible monomorphism/epimorphism to denote their induced maps when clear.

As of now, the structure of the intermediate category Q seems murky. We will motivate the definitions by proving it is the universal construction that recovers then well-accepted definition for K_0 .

Definition 4.0.2. given an exact category \mathcal{M} , $K_0(\mathcal{M})$ is defined to be the abelian group generated by $[C]$, one for each object C in \mathcal{M} , subjected to the relations $[B] = [A] + [C]$ whenever there is an short exact sequence $A \rightarrow B \rightarrow C$.

In some sense, K_0 "breaks up" short exact sequences, forcing them to split. We now examine how the Q -construction break up short exact sequences as well: recall the fact that every short exact sequence $A \rightarrow B \rightarrow C$ is equivalent to the bicartesian diagram

$$\begin{array}{ccc} A & \rightarrow & B \\ \downarrow & & \downarrow \\ 0 & \rightarrow & C \end{array}$$

On the other hand, we have the following proposition regarding bicartesian squares in $Q\mathcal{M}$, which occurs in definition for composition.

Proposition 4.0.1. Given a bicartesian square in \mathcal{M}

$$\begin{array}{ccc} N & \xrightarrow{i} & M' \\ \downarrow j & & \downarrow j' \\ M & \xrightarrow{i'} & N' \end{array}$$

we have $i_! j^! = j'^! i'_!$ in $Q\mathcal{M}$

The proof is simply tracing through the definitions. By the proposition, every short exact sequence $A \rightarrow B \rightarrow C$ in \mathcal{M} leads to the equivalence between the morphisms $0 \leftarrow A \rightarrow B$ and $0 \rightarrow C \leftarrow B$ in $Q\mathcal{M}$. We will see how this equivalence is exactly the splitting of exact sequences we desire in the proof of the following theorem:

Theorem 4.1. $\pi_1(BQ\mathcal{M}, 0) \cong K_0(\mathcal{M})$

Before proving the theorem, we first introduce the following lemma regarding trees.

Lemma 4.2. Suppose T is a maximal tree in a small connected category C . Then, $\pi_1(BC)$ is the group generated by $[f]$, one for each morphism not in T , modulo the relations that

1. $[t] = 1$ for every $t \in T$, and $[Id_c] = 1$ for each identity morphism.
2. $[f] \cdot [g] = [f \circ g]$ for every composable f, g .

Proof. Let X be the 1 skeleton of BC , which is a connected graph with maximal tree T . The proof for $\pi_1(X)$ being the free group generated by $X - T$ is given in Hatcher proposition 1A.2. The lemma is then a direct application of Van-Kampen's Theorem. \square

Proof of Theorem 2.1. We construct the isomorphism directly, which follows Weibel and slightly diverts from the original approach given by Quillen. Let T be the set of injective morphisms of the form $0 \rightarrow A$ in $Q\mathcal{M}$. Clearly, T contains all vertices (objects) in $Q\mathcal{M}$ and is thus a maximal tree.

Let $B' \rightarrow B$ be an injective morphism. Note that its left composition with $0 \rightarrow B'$ is the morphism induced by $0 \rightarrow B' \rightarrow B$, which is in T . Thus by lemma 2.2, all injective morphisms correspond to the identity element; given a surjective morphism $A \leftarrow A'$, note that its left composition with $0 \leftarrow A$ induces a surjective map $0 \leftarrow A'$, so we have the relation $[A \leftarrow A'] = [0 \leftarrow A]^{-1}[0 \leftarrow A']$. By the observation that every morphism in $Q\mathcal{M}$ factors as a surjective morphism followed by an injective one, we note that $\pi_1(BQ\mathcal{M})$ is generated by classes of the form $[0 \leftarrow A]$.

By the observation following proposition 2.0.1, each short exact sequence $A \rightarrow B \rightarrow C$ in \mathcal{M} leads to the equivalence between the morphisms $0 \leftarrow A \rightarrow B$ and $0 \rightarrow C \leftarrow B$ in $Q\mathcal{M}$, which in turn induces in the additivity relation in π_1

$$[0 \leftarrow C][0 \leftarrow A] = [0 \leftarrow B] = [0 \leftarrow A][0 \leftarrow C]$$

The composition rule follows the additivity relation, and the map $[0 \leftarrow A] \mapsto [A]$ is the desired isomorphism between $\pi_1(BQ\mathcal{M})$ and $K_0(\mathcal{M})$. \square

We can now define K -theory for exact categories.

Definition 4.2.1. For a small exact category \mathcal{M} , let $K\mathcal{M}$ be the loop space $\Omega BQ\mathcal{M}$, and set

$$K_i(\mathcal{M}) := \pi_{i+1}(BQ\mathcal{M}, 0) = \pi_i(K\mathcal{M}, 0)$$

For skeletally small categories, we define its K -groups to be those of its skeleton. It is not hard to see that K_i is a functor from the category of small exact categories and exact functors to \mathbf{Ab} , noting that isomorphic functors induce isomorphism of K -groups by the following proposition

Proposition 4.2.1. A natural transformation $\theta : f \rightarrow g$ of functors from C to C' induces an homotopy $BC \times I \rightarrow BC'$ between Bf and Bg . In particular, if a functor has a left/right adjoint, then it induces a homotopy equivalence between classifying spaces.

Proof. The key is to realize the data of the triple (f, g, θ) is exactly a functor $C \times 1 \rightarrow C$, where 1 is the ordered set $\{0 < 1\}$ with classifying space the unit interval. \square

With a bit more effort, one can show that K_i commutes with filtered colimits and products.

References

- [Fri94] E. M. 1944- author. (Eric M.) Friedlander. *Appendix Q. On the group completion of a simplicial monoid*. 1st ed. Vol. 110. Providence, Rhode Island, United States : American Mathematical Society, 1994. ISBN: 0-8218-2591-7.
- [Gra76] D. Grayson. "Higher algebraic K-theory: II". In: *Lecture Notes in Mathematics* 551 (1976). DOI: <https://doi-org.proxy.library.upenn.edu/10.1007/BFb0080003>.
- [MS76] D. McDuff and G. Segal. "Homology Fibrations and the "Group-Completion" Theorem". In: *Inventiones math.* 31 (1976), pp. 279–284. DOI: <https://doi.org/10.1007/BF01403148>.