

Equivariant Stable Homotopy Notes

David Zhu

October 26, 2024

For the entire note, we will assume a group G to be a compact Lie group, and subgroups $H \subset G$ are always closed.

1 Unstable Equivariant Homotopy Theory

1.1 G-CW Complexes

Fix a compact Lie group G acting on a space X . Similar to CW -complexes, we want to deconstruct X into cells, but this time with the additional data of the G -action along with each cell. The idea is that cells are of the form of a product $G/H \times D^n$, where G acts trivially on D^n , and G/H "represents" the orbits of D^n . To make this work, H must be the isotropy group of D^n .

Definition 1.0.1. A G-CW complex is the sequential colimit of spaces X_n , where X_{n+1} is a pushout:

$$\begin{array}{ccc} \coprod G/H \times S^n & \longrightarrow & X_n \\ \downarrow & & \downarrow \\ \coprod G/H \times D^{n+1} & \longrightarrow & X_{n+1} \end{array}$$

We will denote $G/H \times D^n$ as an n-cell.

Remark 1.0.1. Note that the topological dimension of an n -cell in a G -CW complex might be greater than n . For example, a 0-cell $S^1/e \times *$ is one dimensional.

Example 1.0.1. Let $G = C_2$ acting on S^2 by rotation by π along the Z-axis. It has a G -CW structure given by the following cells: 2 zero-cells $C_2/C_2 \times *$, which are the poles corresponding to the fixed points of the C_2 action. 1 one-cell $C_2/e \times D^1$, which are the two great circles joining the poles; 1 two-cell $C_2/C_2 \times D^2$, which are the two hemispheres.

Example 1.0.2. Let $G = C_2$ acting on S^2 by the antipodal map. It has a G -CW structure given by the following cells: 1 zero-cells $C_2/e \times *$, which are the poles; 1 one-cell $C_2/e \times D^1$, which are the two great circles joining the poles; 1 two-cell $C_2/C_2 \times D^2$, which are the two hemispheres.

Definition 1.0.2. Let H be a subgroup of G . Define $\pi_n^H(X) := \pi_n(X^H)$. A map $f : X \rightarrow Y$ of G -spaces is a weak equivalence if for all subgroups $H \subset G$,

$$f_* : \pi_n^H(X) \rightarrow \pi_n^H(Y)$$

is an isomorphism.

Let **GTop** be the category of G -spaces and G -maps. There is a cofibrantly-generated model structure that we can put on **GTop**:

Theorem 1.1. There is a cofibrantly-generated model structure on **GTop**, given by

1. A G -map $f : X \rightarrow Y$ is a fibration iff for all $H \subset G$, $f^H : X^H \rightarrow Y^H$ is a fibration.
2. A G -map $f : X \rightarrow Y$ is a weak equivalence iff for all $H \subset G$, $f^H : X^H \rightarrow Y^H$ is a weak equivalence.

An immediate consequence of the model category structure is the equivariant Whitehead's Theorem

Corollary 1.1.1. Let $f : X \rightarrow Y$ be a weak equivalence of cofibrant-fibrant objects in a model category. Then, f is a homotopy equivalence. In particular, every object in **GTop** is fibrant, and G -CW complexes are cofibrant.

1.2 Elmendorf's Theorem

From the model structure given in Theorem 1.1, we have a vague sense of the following "equivalence":

$$G\text{-Homotopy Type of } X \Leftrightarrow \{\text{ordinary homotopy type of } X^H : H \subset G\}$$

And Elmendorf's Theorem will make the equivalence precise. We start by introducing the orbit category:

Definition 1.1.1. The orbit category \mathcal{O}_G is the full subcategory of **GTop** on the objects $\{G/H : H \subset G\}$.

The following lemma will make the structure of \mathcal{O}_G clearer.

Lemma 1.2. $\text{Map}^G(G/H, G/K) \cong (G/K)^H$

Proof. Note that there exists a G -equivariant maps $\varphi : G/H \rightarrow G/K$, determined by $\varphi(H) = gK$ iff $gHg^{-1} \subseteq K$ iff $h(gK) = gK$ for all $h \in H$. \square

Let $\text{Fun}(\mathcal{O}_G^{\text{op}}, \text{Top})$ be the functor category. We have the following fact on the model structure on functor categories:

Theorem 1.3. Let \mathcal{D} be a model category and \mathcal{C} be a cofibrantly generated model category. Then, $\text{Fun}(\mathcal{C}, \mathcal{D})$ admits a model structure.

It is useful to know that the weak equivalences in $\text{Fun}(\mathcal{O}_G^{\text{op}}, \text{Top})$ is given pointwise: a natural transformation $\eta : \mathcal{F} \rightarrow \mathcal{G}$ is a weak equivalence iff $\eta_{G/H} : \mathcal{F}(G/H) \rightarrow \mathcal{G}(G/H)$ is a weak equivalence.

Definition 1.3.1. There is a functor $\psi : \mathbf{GTop} \rightarrow \mathbf{Fun}(\mathcal{O}_G^{op}, \mathbf{Top})$ given by

$$X \rightarrow (G/H \mapsto X^H)$$

It is easy to check the functoriality. Note that if we restrict ψ to \mathcal{O}_G , the functor is just the Yoneda embedding: $\text{Map}^G(G/H, G/K) \cong (G/K)^H$.

Proposition 1.3.1. There is a functor $\theta : \mathbf{Fun}(\mathcal{O}_G^{op}, \mathbf{Top}) \rightarrow \mathbf{GTop}$ given by $X \mapsto X(G/e)$, where $X(G/e)$ is equipped with the following G -action: note that every $g \in G$ defines an G -map $G/e \rightarrow G/e$, which we denote by R_g .

$$g \cdot x = X(R_g)(x)$$

It is easy to check that (θ, ψ) is an adjoint pair. In fact, more can be said:

Theorem 1.4. (Elmendorf's Theorem) $\mathbf{Fun}(\mathcal{O}_G^{op}, \mathbf{Top})$ and \mathbf{GTop} have the same homotopy category.

The original proof due to Elmendorf constructs the equivalence explicitly using the Bar construction to obtain a homotopy inverse to the embedding ψ . The theorem can now be put into a more modern framework:

Theorem 1.5. (θ, ψ) is an Quillen equivalence. ψ is an equivalence of $(\infty, 1)$ categories.

1.3 Bredon Cohomology

The goal is to construct a cohomology theory satisfying the Eilenberg-Steenrod axioms under the equivariant setting.

Definition 1.5.1. (Equivariant reduced generalized cohomology) Let GCW_* be the category of pointed G -CW complexes with equivariant maps. Then, a generalized cohomology theory on GCW_* is a sequence of contravariant functors

$$\tilde{H}^n := GCW_* \rightarrow \mathbf{Ab}$$

satisfying the following:

1. if f, g are equivariantly homotopic, then $\tilde{H}^n(f) = \tilde{H}^n(g)$.
2. There exists a sequence of natural isomorphisms

$$\tilde{H}^n(X) \rightarrow \tilde{H}^{n+1}(S^1 \wedge X)$$

where G acts trivially on S^1 in the smash.

3. The sequence

$$\tilde{H}^n(X/A) \rightarrow \tilde{H}^n(X) \rightarrow \tilde{H}^n(A)$$

is exact.

Remark 1.5.1. The above axioms is built upon pointed "single" spaces. It is in fact equivalent to the usual theory built upon pairs, and is justified in

For a non-equivariant reduced generalized cohomology theory \tilde{h}^* , the Atiyah-Hirzebruch spectral sequence tells us that knowing $\tilde{h}^*(pt)$ basically determines the cohomology theory on CW complexes. Heuristically,

the cohomology is determined by the building blocks, which are contractible open cells. However, in the equivariant setting, the building blocks are more complicated: the building blocks have become orbits of the form G/H . We are lead to the following definitions:

Definition 1.5.2. A coefficient system is a contravariant functor $\mathcal{F} : \mathcal{O}_G \rightarrow \text{Ab}$.

Recall that if a reduced cohomology theory is called ordinary if it satisfies the dimension axiom, i.e the zeroth reduced cohomology group (a.k.a the coefficient system) is trivial on a point. Our goal now is to construct such a theory in the equivariant setting, which is called Bredon cohomology.

Note that by the general theory of abelian categories, the functor category of coefficient systems $\mathcal{CS} := \text{Fun}(\mathcal{O}_G, \text{Ab})$ is abelian. It is now possible to define Bredon cohomology on a G -CW complex by explicitly defining the cochain complexes on cells, for example see . However, we may package the cochains into the following form

Definition 1.5.3. For each n , we may define a coefficient system $C_n(-)$, given by

$$G/H \mapsto H_n((X^H)_n, (X^H)_{n-1}; \mathbb{Z}) = C_n^{CW}(X^H)$$

The differential of the CW chain complex induces a chain complex of coefficient systems $C.(-)$.

It is easy to check that $\text{Hom}_{\mathcal{CS}}(C.(-), M)$ is a cochain complex whose differentials is induced by those of $C.()$.

Definition 1.5.4. The Bredon cohomology of X with coefficients in a system M is defined by

$$H_G^n(X; M) := H^n(\text{Hom}_{\mathcal{CS}}(C.(X), M))$$

proof of the axioms and computation of examples.

1.4 Classical Application

homotopy fixed points and actual fixed points: coarse vs fine Here are two questions we may ask: given a finite p -group G acting on X , can we recover the cohomology of X^G based on the cohomology of X , while somehow incorporating the G -action? The second point of interest is the Sullivan conjecture: in The study of étale homotopy theory allows one to determines algebraically the profinite completion of a variety given only its étale homotopy type. The same situation occurs as before: given a variety V over the complex numbers, there is a natural $\mathbb{Z}/2$ -action on the variety, with $V^G = V(\mathbb{R})$. How much can we say about $V(\mathbb{R})$ if we are given the étale homotopy type of V and the G -action? IN this case, the fixed points is not a homotopy invariant, so we should consider the homotopy fixed points instead, and the sullivan conjecture states that

Theorem 1.6. (Sullivan Conjecture) $V(\mathbb{C})^{\mathbb{Z}/2} \rightarrow V(\mathbb{C})^{h\mathbb{Z}/2}$ becomes an isomorphism after 2-adic completion. More generally, let G be a finite p -group, then $X^G \rightarrow X^{hG}$ becomes an equivalence after p -adic completion.

One possible application is a quick proof of a theorem by Smith:

Theorem 1.7. Let G be a finite p -group and X be a finite G -CW complex such that (the underlying topological space of) X is an p -cohomology sphere.¹⁶ Then, X^G is either empty or an p -cohomology sphere of smaller dimension.

2 Spectra and the Stable Category

2.1 Motivations

The first motivation is Brown Representability Theorem, which is about representing reduced cohomology theories

Theorem 2.1 (Brown Representability Theorem). Let \tilde{h}^* be a reduced cohomology theory on pointed CW-complexes. Then, for each $n \in \mathbb{Z}$, there exists a connected pointed CW complex K_n such that

$$\tilde{h}^n(X) \cong [X, K_n]$$

for all n . Moreover, the K_n are determined up to homotopy equivalence.

In fact, there are more structure to the set $\{K_n\}$: using the suspension axiom and loop-suspension adjunction, we see that there must be an isomorphism

$$[X, K_n] \cong \tilde{h}^n(X) \cong \tilde{h}^{n+1}(\Sigma X) \cong [\Sigma X, K_n] \cong [X, \Omega K_n]$$

Taking X to be K_n , we note that the identity map from the LHS corresponds uniquely to a map $\alpha_n : K_n \rightarrow \Omega K_n$, which we will call the structure map. By naturality and taking $A = S^k$, the structure map induces a weak equivalence

$$K_n \cong \Omega K_n$$

. This motivates the definition for Ω -Spectra

Definition 2.1.1. A spectrum is a sequence of pointed topological spaces $\{X_n\}$ with structure maps.

$$\Sigma X_n \rightarrow X_{n+1}$$

. An Ω spectrum is a spectrum whose adjoint structure maps

$$X_n \rightarrow \Omega X_{n+1}$$

are weak equivalences.

So we see that a reduced cohomology theory corresponds to an Ω -spectrum. In fact, it is not hard to show that an Ω -spectrum defines a reduced cohomology theory as well. We can then present Brown Representability Theorem in the following way:

Theorem 2.2 (Brown Representability Theorem). Every reduced cohomology theory on the category of basepointed CW complexes has the form $\tilde{h}^n(X) = [X, K_n]$ for some Ω -spectrum $\{K_n\}$

Example 2.2.1 (Eilenberg-MacLane Spectrum). The Ω -spectrum that represents reduced ordinary cohomology with coefficients in \mathbb{Z} is given by the Ω -spectrum that is the Eilenberg-MacLane spaces $H\mathbb{Z} := \{K(\mathbb{Z}, n)\}$.

Example 2.2.2. The Ω -spectrum that represents reduced complex topological K -theory is given by $\{KU_n\}$, where

$$KU_n = \begin{cases} BU \times \mathbb{Z} & n\text{-even} \\ \Omega BU & n\text{-odd} \end{cases}$$

In particular, Bott-periodicity shows that the structure maps are weak-equivalences.

Here is a natural question: how do we design a category of spectra, such that an "equivalence" of spectra will give equivalent reduced cohomology theories?

The second motivation is stable homotopy groups and stable maps:

Definition 2.2.1. Let X and Y be pointed CW-complexes. The set of stable homotopy classes of maps from X to Y is defined to be

$$[X, Y]^s := \varinjlim_k [\Sigma^k X, \Sigma^k Y]$$

One form of the Freudenthal suspension theorem says that if Y is n -connected and X has dimension less than $2n + 1$, then the suspension map $[X, Y] \rightarrow [\Sigma X, \Sigma Y]$ is bijective. In this case, we see that the colimit actual stabilizes after at a finite stage.

Definition 2.2.2. For a pointed CW-complex, the n -th stable homotopy group is defined to be

$$\pi_n^{st}(X) := \varinjlim_k \pi_{n+k}(\Sigma^k X) = \varinjlim_k [\Sigma^k S^n, \Sigma^k X] = [S^n, X]^s$$

By definition, we see that the stable homotopy group should be the homotopy group in some "stable category". Moreover, stable homotopy groups actually defines a reduced homology theory: the two axioms that are not trivial are the LES and the wedge axiom. The key point is that Blaker's Massey Theorem plus the LES of homotopy groups of a pair will give us the LES of stable homotopy groups of a pair; the wedge axiom follows from that $\Sigma^i X \wedge \Sigma^j Y$ is the $2i - 1$ skeleton of $\Sigma^i X \times \Sigma^j Y$. Generalizing this, we have

Theorem 2.3. Let K be a CW-complex. The sequence $h_i(X) = \pi_i^s(X \wedge K)$ forms a reduced homology theory on the category of basepointed CW-complexes and basepoint-preserving maps.

We can also define the homotopy groups of a spectrum as a generalization:

Definition 2.3.1 (Homotopy groups of a spectrum). Suppose $K = \{K_i\}$ is a spectrum. Then,

$$\pi_n(K) := \varinjlim_i \pi_{n+i}(K_i)$$

where the inductive limit is induced by the suspension structure maps.

Example 2.3.1. Given a topological space X , we have the associated suspension spectrum $\Sigma^\infty X = \{\Sigma^k X\}$. Then, the stable homotopy groups of X is given by the homotopy groups of its associated suspension spectrum.

Definition 2.3.2. Given a spectrum $K = \{K_n\}$ and a CW complex X , there is a associated smash spectrum $K \wedge X$, where $(K \wedge X)_n = K_n \wedge X$. The structure maps is given by

$$\Sigma(K \wedge X)_n = \Sigma K_n \wedge X \rightarrow K_{n+1} \wedge X = \Sigma(K \wedge X)_{n+1}$$

Proposition 2.3.1. Let K be a spectrum. Given a CW complex X , the sequence

$$h_n := \pi_n(X \wedge K)$$

is a reduced homology theory.

The proof mostly follows from the same tactics in proving the stable homotopy groups being a reduced homology theory.

Example 2.3.2 (Singular homology). If $K = HG$ is the Eilenberg-MacLane spectrum, then $h_i(X) := \pi_i(X \wedge K)$ is isomorphic to singular homology with coefficients in G . We only have to check the dimension axiom

$$h_i(S^0) := \varinjlim_n \pi_{n+i}(K(G, n))$$

which is trivial except $i = 0$.

Thus, we have seen that the singular homology is recovered from a spectrum; moreover, a spectrum gives a reduced homology theory by proposition 2.3.1.

2.2 Duality

We first recall a classifc duality theorems from singular cohomology:

Theorem 2.4 (Alexander Duality). If K is a compact, locally contractible, nonempty, proper subspace of S^n . Then we have the isomorphism

$$\tilde{H}^{n-i-1}(K; \mathbb{Z}) \cong \tilde{H}_i(S^n - K; \mathbb{Z})$$

The theorem in particular implies that different embeddings of some nice enough space in S^n has isomorphic homology. However, they do not necessarily have the same homotopy type, as there are tons of examples in knot theory. Nevertheless, Spanier and Whitehead prove the following theorem, which states that the homotopy types are the same after sufficiently many suspensions.

Theorem 2.5 (Spanier-Whitehead Duality). Let X be a compact simplicial complex. Let $f, g : X \rightarrow S^n$ be two simplicial embeddings. Then for some sufficiently large M the M -fold suspensions $\Sigma^M(S^n \setminus f(X))$ and $\Sigma^M(S^n \setminus g(X))$ are homotopy equivalent.

To study this phenomenon more formally, Spanier-Whitehead proposed the now-called Spanier-Whitehead category:

Definition 2.5.1 (Spanier-Whitehead Category). The **Spanier-Whitehead category**, or **S-category** for short, has objects pairs (X, n) , where X is a pointed finite CW complex, and n is an integer. The morphisms are defined by

$$\text{Hom}_S((X, n), (Y, m)) := \varinjlim_{q \geq \max(|m|, |n|)} [\Sigma^{q+n} X, \Sigma^{q+m} Y]$$

We can think of (X, m) as the m -fold suspension of X . If $m = n = 0$, we see that $\text{Hom}_S((X, 0), (Y, 0)) = \varinjlim_k [\Sigma^k X, \Sigma^k Y]$, which are simply the stable morphisms and that the different embeddings $X \rightarrow S^n$ become actually isomorphic in this category by theorem 2.5. If $m > 0$, we may denote $(X, -m)$ as $\Sigma^{-m} X$ to represent formal desuspensions. We can also think of this as "inverting smashing with spheres".

Definition 2.5.2. Let \mathcal{C} be a symmetric monoidal category with the tensor product denoted by \wedge , and unit object S^0 . Then, X, Y are **dual** if there are morphisms $X \wedge Y \rightarrow S^0$ and $S^0 \rightarrow Y \wedge X$ such that the following compositions

$$X \wedge S^0 \rightarrow X \wedge Y \wedge X \rightarrow S^0 \wedge Y$$

$$S^0 \wedge Y \rightarrow Y \wedge X \wedge Y \rightarrow Y \wedge S^0$$

are the swap isomorphisms in the symmetric monoidal category.

By the identity $\Sigma(X \wedge Y) \cong X \wedge \Sigma Y$, we see that the smash product defined by

$$(X, m) \wedge (Y, n) := (X \wedge Y, m + n)$$

endows the S -category with a symmetric monoidal structure. Let us see some explicit dualities

Example 2.5.1. S^n is dual to S^{-n} , the formal n -fold desuspension of S^0 . This follows directly from definition.

Example 2.5.2 (Alexander duality in S -category). Let X be a compact finite CW complex with an embedding $X \rightarrow S^{n+1}$. Let us denote the complement $S^n \setminus X$ as $D_n X$. Then, one may construct a map $X \wedge D_n X \rightarrow S^n$, and a desuspension of such map exhibits X and $\Sigma^{-n} D_n X$ as duals.

Let Y be a finite discrete set of points. Then, Y embeds in S^1 , and the complement is homotopy equivalent to Y again. This shows that Y is self-dual. This example will come up later when we deal with finite G -sets and equivariant embeddings.

Example 2.5.3 (Atiyah Duality). Suppose M is a closed smooth manifold with a smooth embedding $M \rightarrow \mathbb{R}^n$. Let Tv be the Thom space of the normal bundle of the embedding. The Pontrjagin-Thom construction yields a map

$$Tv \wedge M_+ \rightarrow \Sigma^n M_+ \xrightarrow{\text{collapse } M_+} S^n$$

which exhibits Tv and $\Sigma^{-n} M_+$ as duals.

2.3 The Desired Stable Homotopy Category

Building upon the motivations in the previous section, we want to build a category \mathcal{S} that captures the stable phenomena and cohomology theories/homology theories. In particular, we want it to satisfy the following

properties (not necessarily axioms):

1. There is a functor $\Sigma^\infty : \text{HoTop}_* \rightarrow \text{Sp}$, together with an adjoint $\Omega^\infty : \mathcal{S} \rightarrow \text{HoTop}_*$.
The motivation for this is the suspension spectrum.
2. Let A, B be CW complexes, where A has finite dimension. Then, there is a natural isomorphism

$$[\Sigma^\infty A, \Sigma^\infty B] \cong [A, B]^s$$

The motivation for this is to capture stable morphisms.

3. \mathcal{S} has a closed symmetric monoidal structure, with \wedge as the tensor product, and $F(-, -)$ as internal hom. $\Sigma^\infty S^0$ is the identity object.
4. The morphisms of \mathcal{S} has the structure of graded abelian groups, with bilinear composition. We denote that graded components as $[-, -]_*$. Moreover, given a reduced cohomology theory E^* there exists an object K in Sp such that

$$E^*(A) \cong [\Sigma^\infty A, K]_{-*}$$

The motivation for this is to represent cohomology theories.

5. For every object K in \mathcal{S} , one defines a reduced homology theory via

$$E_n(A) := \pi(K \wedge A)$$

The motivation for this is to determine homology theories.

6. \mathcal{S} has a closed symmetric monoidal structure, with \wedge as the tensor product and $F(-, -)$ as internal hom. The object $\Sigma^\infty S^0$ is the identity object.
The motivation for this is that smashing with the suspension functor induces isomorphism of stable homotopy groups, so it should induce an equivalence of spectra. The tensor-hom adjunction is going to help set up duality statements similar to Spanier-Whitehead duality and Poincaré duality..

In particular, property 6 was the central one in the sense that the entire theory began with the work of Spanier and Whitehead with their search for a category where one has the desired dualities in homotopy theory. It is also the structure that seems to be the hardest to obtain.

2.4 Attempts

2.4.1 The Spanier-Whitehead Category

The first attempt to construct such a category is the Spanier-Whitehead category, where the objects are finite CW complexes, and the morphisms are stable maps. The category was originally set up to perform Spanier-Whitehead duality (chronologically it was), but the objects "almost" represents cohomology theories. We use the word almost because since the objects are only finite complexes, we do not have arbitrary products and coproducts, and the wedge axiom do not hold.

2.4.2 An Attempt for Sp

By Brown representability, it should be natural to consider a category with objects spectra, and the natural choice for morphisms between spectra is a collection of level-wise maps that is compatible with the structure maps. A natural candidate for homotopies in this category is a map $H : X \wedge I_+ \rightarrow Y$. However, it turns out there are not enough maps in this category, such that non-homotopy equivalent spectra might represent isomorphic cohomology theories. Adams then in his 74 paper resolves the issue by consider equivalences classes of "cofinal spectra," which turns out to be quite a complicated a construction.

2.4.3 Ho(Sp) and Structured Spectra

The first construction in line with modern treatment of the subject is Bousfield and Friedlander's construction, which puts a model structure on the naive category of spectra, and considers its homotopy category. This category turns out satisfies almost all the properties that people wanted. (We say "almost" since Lewis in 91 showed that it is impossible to put a symmetric monoidal structure on the category of spectra that also satisfies a list of desirable axioms). However, people still searched for a point-set model that does so without going through the homotopy category. People later constructed more structured spectra, such as orthogonal spectra and symmetric spectra, whose additional structure then provided a fix to such a problem.

2.5 Orthogonal Spectra

Definition 2.5.3. An **orthogonal spectrum** consists of the following data:

1. A sequence of pointed topological spaces X_n
2. A basepoint-preserving left $O(n)$ -action on X_n .
3. Structure maps $\sigma_n : X_n \wedge S^1 \rightarrow X_{n+1}$
4. The iterated structure maps

$$X_n \wedge S^m \rightarrow X_{n+m}$$

define by the composition

$$X_n \wedge S^m = (X_n \wedge S^1) \wedge S^{m-1} \xrightarrow{\sigma_n \wedge Id_{S^{m-1}}} X_{n+1} \wedge S^{m-1} \rightarrow \dots \rightarrow X_{n+m}$$

is $O(n) \times O(m)$ -equivariant. Specifically, $O(m)$ -acts on S^m canonically since S^m is the one-point compactification of \mathbb{R}^m ; $O(n) \times O(m)$, through orthogonal sum, is identified as a subgroup of $O(n+m)$ and acts by restriction on X_{n+m} .

Definition 2.5.4. A morphism of orthogonal spectra $f : X \rightarrow Y$ is a sequences of maps $f_n : X_n \rightarrow Y_n$ that are $O(n)$ -equivariant, and compatible with the structure maps such that

$$\begin{array}{ccc} X_n \wedge S^1 & \xrightarrow{f_n \wedge S^1} & Y_n \wedge S^1 \\ \downarrow \sigma_n & & \downarrow \sigma_n \\ X_{n+1} & \xrightarrow{f_{n+1}} & Y_{n+1} \end{array}$$

Our goal now is to show that there exists a tensor product structure. We now recall a few constructions:

Definition 2.5.5. Given subgroup $H \subset G$ and a pointed H -space X , the **balanced smash product** $G_+ \wedge_H X$ is the quotient space of $G_+ \wedge H$ by the relation $gh \wedge x \sim g \wedge hx$

Note that $G_+ \wedge_H X$ is canonically a G -space by the action on the G_+ -coordinate. One can think of this as the base-change from H to G .

Proposition 2.5.1. The functor $\text{Htop}_* \rightarrow \text{GTop}_*$ given by the balanced smash product is left adjoint to the forgetful functor. Specifically, we have the natural isomorphism

$$\text{Hom}_H(X, Y) \cong \text{Hom}_G(G \wedge_H X, Y)$$

Recall the universal property of the tensor product in **RMod** is given by the universal object from which a R -balanced map from the product factors through. In our situation, the ring R is more or less the sphere

spectrum, and in order to define the tensor product, we should first give an appropriate definition of a balanced map.

Definition 2.5.6. Given orthogonal spectra X, Y, Z , a **bimorphism** from the pair (X, Y) to Z is given by a collection of $O(p) \times O(q)$ -equivariant maps

$$b_{p,q} : X_p \wedge Y_q \rightarrow Z_{p+q}$$

subjected to the following commutative diagram

$$\begin{array}{ccc} X_p \wedge Y_q \wedge S^1 & \xrightarrow{X_p \wedge \sigma_Y} & X_p \wedge Y_{q+1} \\ b_{p,q} \wedge S^1 \downarrow & & \downarrow b_{p,q+1} \\ Z_{p+q} \wedge S^1 & \xrightarrow{\sigma_Z} & Z_{p+q+1} \end{array}$$

and

$$\begin{array}{ccccc} X_p \wedge Y_q \wedge S^1 & \xrightarrow{X_p \wedge \sigma_Y} & X_p \wedge Y_{q+1} & \xrightarrow{b_{p,q+1}} & Z_{p+q+1} \\ X_p \wedge twist \downarrow & & & & \uparrow Id \times \chi_{1,q} \\ X_p \wedge S^1 \wedge Y_q & \xrightarrow{\sigma_X \wedge Y_q} & X_{p+1} \wedge Y_q & \xrightarrow{b_{p+1,q}} & Z_{p+1+q} \end{array}$$

The map $\chi_{p,q}$ correspond to the element in $O(p+q)$ that swaps the first q coordinates with the remaining q -coordinates. In analogy to **RMod**, the first diagram says that $(x \otimes y) \cdot r = x \otimes (y \cdot r)$, and the second diagram says that $x \otimes (y \cdot r) = (x \cdot r) \otimes y$.

Recall that the tensor product of two R -modules A, B is the coequalizer of the diagram

$$\alpha_1, \alpha_2 : A \otimes R \otimes B \rightarrow A \otimes B$$

where the tensor is in **Ab** and the two maps are given by the action on A and B , respectively. We shall recreate this in our category **Sp**:

Definition 2.5.7. Given orthogonal spectra X, Y , define $(X \wedge Y)_n$ to be the coequalizer of the maps

$$\alpha_X, \alpha_Y : \bigvee_{p+1+q=n} O(n)_+ \wedge_{O(p) \times Id \times O(q)} X_p \wedge S^1 \wedge Y_q \rightarrow \bigvee_{p+q=n} O(n)_+ \wedge_{O(p) \times O(q)} X_p \wedge Y_q$$

in the category of $O(n)$ -spaces, where α_X is induced by the maps

$$X_p \wedge S^1 \wedge Y_q \xrightarrow{\sigma_X \wedge Y_q} X_{p+1} \wedge Y_q$$

on each summand, and α_Y is induced by the maps

$$X_p \wedge S^1 \wedge Y_q \xrightarrow{X_p \wedge twist} X_p \wedge Y_p \wedge S^1 \rightarrow X_p \wedge Y_{q+1} \xrightarrow{X_p \wedge \chi_{q,1}} X_p \wedge Y_{1+q}$$

Note that the coequalizer in $GTop$ is created by the coequalizer in the underlying Top , so in our case, $(X \wedge Y)_n$ is a quotient space of $\bigvee_{p+q=n} O(n)_+ \wedge_{O(p) \times O(q)} X_p \wedge Y_q$. Then, the maps

$$O(n)_+ \wedge_{O(p) \times O(q)} X_p \wedge Y_q \wedge S^1 \rightarrow O(n+1)_+ \wedge_{O(p) \times O(q)} X_p \wedge Y_q$$

given by $Id \wedge \sigma_Y$ induces a morphism $\varphi_n : (X \wedge Y)_n \wedge S^1 \rightarrow (X \wedge Y)_{n+1}$. It follows that

Proposition 2.5.2. The collection $(X \wedge Y)_n$, with structure maps φ_n forms an orthogonal spectra.

As remarked before, the smash product is equipped with a universal bimorphism $i : (X, Y) \rightarrow X \wedge Y$ given by

$$X_p \wedge Y_q \xrightarrow{x \wedge y \mapsto e \wedge x \wedge y} \bigvee_{p+q=n} O(n)_+ \wedge_{O(p) \times O(q)} X_p \wedge Y_q \xrightarrow{\text{projection}} (X \wedge Y)_{p+q}$$

The punchline is that we also have an internal hom orthogonal spectra $F(X, Y)$ for orthogonal spectra X, Y , which makes \mathbf{Sp} a closed symmetric monoidal category in which we have the adjunction

$$\mathrm{Hom}_{\mathbf{Sp}}(X \wedge Y, Z) \cong \mathrm{Hom}_{S_p}(X, F(Y, Z))$$

which we will show later in the equivariant case below.

2.6 The Equivariant Case

One way to motivate the correct definition of equivariant stable homotopy category is to represent the equivariant cohomology theories. In particular, we want a form of the suspension isomorphism

$$H^*(X) \cong H^{*+V}(\Sigma^V X)$$

In the non-equivariant case, we only take V to be trivial representations, which means we are smashing with a sphere with trivial G -action. However, it turns out that the cohomology would carry more useful data if we want to keep track all possible representations V .

Definition 2.5.8. A complete G -universe U is an infinite dimensional real inner product space with a G -action, such that it is a direct sum of countably many copies of each irreducible representations of G .

Remark 2.5.1. If we drop the adjective "complete" from the definition, we would only require that the universe containing some subset of G -representations and the trivial representation instead of all possible representations. However, if we want dualizable orbit, a convenient assumption is requiring U to be complete.

Definition 2.5.9. A G -prespectrum X is a collection of pointed G -spaces $X(V)$, one for each finite dimensional subspace $V \subset U$ of a given G universe. The collection is equipped with structure maps

$$\Sigma^W X(V) \cong X(V \oplus W)$$

The structure maps must be associative. A G -prespectrum is a G -Spectrum (or $\Omega - G$ -spectrum) if the adjoint structure maps

$$X(V) \rightarrow \Omega^W X(V \oplus W)$$

are homeomorphisms.

This definition enlarges the usual \mathbb{Z} -graded notion of spectra. Sometimes in literature this representation graded spectrum is referred to as the genuine G -spectra.

Definition 2.5.10. The homotopy groups of G -prespectrum are

$$\pi_q^H(X) := \operatorname{colim}_{V \subset U} \pi_q^H \Omega^V X(V)$$

$$\pi_{-q}^H(X) := \operatorname{colim}_{\mathbb{R}^m \subset V} \pi_q^H \Omega^{V - \mathbb{R}^m} X(V)$$

where $V - \mathbb{R}^m$ denotes the orthogonal complement.

A weak-equivalence between G -spectra is a π_* equivalence.

Theorem 2.6. There is a model structure on G -prespectra where the weak equivalences are the stable equivalences.

Again, we want the symmetric monoidal structure on the category of genuine G -spectra, and we are going to use the orthogonal G -spectra model again.

3 $RO(G)$ -graded cohomology Theories

3.1 Mackey Functors

Mackey Functors will serve the role of coefficient system in $RO(G)$ -graded coefficient theories. In particular, we take into account of the transfer maps, which is not present in the orbit category \mathcal{O}_G . We first give a functorial definition for Mackey functors in terms of the Burnside category, and then present an alternative definition with axioms instead, which hopefully further elucidates the structure of a Mackey functor. For the following discussions, G will be a finite discrete group.

Definition 3.0.1. Let C be a category with finite limits and colimits. Then, $\mathbf{Span}(C)$ is the category with the same objects as C , and morphisms equivalence classes of spans. Composition of morphisms is given by pullback of spans.

Given two morphisms in $\mathbf{Span}(C)(X, Y)$ $X \leftarrow Z \rightarrow Y$ and $X \leftarrow Z' \rightarrow Y$, the disjoint union operation $X \leftarrow Z \amalg Z' \rightarrow Y$ endows the homsets a monoid structure. Let $\mathbf{Span}^+(C)$ denote the preadditive completion of $\mathbf{Span}(C)$.

Definition 3.0.2. Let G be a finite group. Then, the Burnside Category is defined to be $\mathcal{A}(G) := \mathbf{Span}^+(\mathbf{GSet}^{\text{Fin}})$.

Recall in the orbit category, we have a morphism $G/H \rightarrow G/K$ iff H is subconjugate to K . The Burnside category can be viewed as formally adding the "wrong way maps" to each morphisms in the orbit category.

Definition 3.0.3. A Mackey Functor is an additive functor $F : \mathcal{A}(G)^{op} \rightarrow \mathbf{Ab}$.

And Mackey functors are coefficients systems with wrong way maps (transfers) added.

we can then define a functor $i_* : \mathcal{O}_G \rightarrow \mathcal{A}(G)$ by sending a morphism $f : G/H \rightarrow G/K$ to the span $G/H \xleftarrow{Id} G/H \xrightarrow{f} G/K$; similarly, we have a functor $i^* : \mathcal{O}_G^{op} \rightarrow \mathcal{A}(G)$ by sending a morphism $f : G/H \rightarrow G/K$ to the span $G/K \xleftarrow{f} G/H \xrightarrow{Id} G/H$. Given a Mackey functor $\underline{M} : \mathcal{A}(G) \rightarrow \mathbf{Ab}$, precomposition with i_* gives a

coefficients system, which we will denote M^* , and precomposition with i^* is a covariant functor $\mathcal{O}_G \rightarrow \text{Ab}$, which we will denote as M_* . By construction, M^* and M_* agree on objects, and they determine the value of the Mackey functor on objects: a Mackey functor is additive, its data is determined by the disjoint orbits. We would also like to know how the morphisms commute: if we have the pullback square of orbits

$$\begin{array}{ccc} W & \xrightarrow{\beta} & Y \\ \alpha \downarrow & & \downarrow \delta \\ X & \xrightarrow{\gamma} & Z \end{array}$$

then $M_*(\beta)M^*(\alpha) = M^*(\delta)M_*(\gamma)$, as illustrated by the following composition diagram.

$$\begin{array}{ccccc} & & W & & \\ & \swarrow \alpha & \downarrow \vee & \searrow \beta & \\ & X & & Y & \\ \swarrow Id & & \searrow \gamma & \swarrow \delta & \searrow Id \\ X & & Z & & Y \end{array}$$

From the above discussion, we have another equivalent definition for Mackey functors, which is sometimes more useful

Definition 3.0.4. A **Mackey Functor** is a pair of functors $M^* : \text{GSet}^{\text{Fin}} \rightarrow \text{Ab}$ and $M_* : \text{GSet}^{\text{Fin}} \rightarrow \text{Ab}$ such that the following axioms hold:

1. M^* and M_* agree on objects;
2. For every pullback square of orbits

$$\begin{array}{ccc} W & \xrightarrow{\beta} & Y \\ \alpha \downarrow & & \downarrow \delta \\ X & \xrightarrow{\gamma} & Z \end{array}$$

then $M_*(\beta)M^*(\alpha) = M^*(\delta)M_*(\gamma)$;

3. Both functors maps disjoint unions to direct sums.

Before describing concrete examples, we want to clarify some structure inherent in the burnside categories and Mackey Functors: suppose we have a G -Mackey functor \underline{M} ; since it is additive, its data is completely determined by its value on orbits, and the restriction/transfer morphisms induced by morphisms between orbits. Further, recall that a G -map between orbits $G/H \rightarrow G/K$ given by $H \mapsto gK$ must satisfy $gHg^{-1} \subseteq K$. Thus, the map can be decomposed into

$$G/H \xrightarrow{c_g} G/gHg^{-1} \xrightarrow{\pi} G/K$$

where $c_x : G/H \rightarrow G/gHg^{-1}$ is the conjugation map defined by $H \mapsto g(gHg^{-1})$, and π is the canonical projection that sends $gHg^{-1} \mapsto K$. herefore, the data of a Mackey functor is completely determined by the conjugation morphisms and the canonical projection. We will see how these morphisms commute with each other.

Let $c_g^* := M^*(c_g)$ and $c_g^g := M_*(c_g)$.

We have the following pullback square

$$\begin{array}{ccc} G/H & \xrightarrow{Id} & G/H \\ Id \downarrow & \lrcorner & \downarrow c_g \\ G/H & \xrightarrow{c_g} & G/gHg^{-1} \end{array}$$

So in fact we have $c_g^* = (c_g^g)^{-1}$.

Given a canonical projection $\pi : G/H \rightarrow G/K$, denote $\text{Res}_H^K := M^*(\pi)$ and $\text{Tr}_H^K := M_*(\pi)$. Now consider the following commutative diagram:

$$\begin{array}{ccc} G/H & \xrightarrow{\pi} & G/G \\ c_g \downarrow & & \downarrow Id \\ G/gHg^{-1} & \xrightarrow{\pi} & G/G \end{array}$$

which implies $\text{Tr}_H^G \circ c_g^g = \text{Tr}_H^G$ and $\text{Res}_H^G(x) = c_g^* \circ \text{Res}_H^G(x)$ for all g . Moreover, consider the pullback diagram

$$\begin{array}{ccc} G/H \times G/K & \xrightarrow{\beta} & G/K \\ \alpha \downarrow & & \downarrow \delta \\ G/H & \xrightarrow{\gamma} & G/G \cong * \end{array}$$

By the G -isomorphism $G/H \times G/K \cong G \times_H G/K$, it is easy to see the the class $(1, xK)_H$ has stablizer $H \cap xKx^{-1}$, and the orbits are indexed by a class in the double coset $H \backslash G/K$. Thus,

$$G/H \times G/K \cong \coprod_{H \backslash G/K} G/H \cap xKx^{-1}$$

. We may use this decomposition to compute the composition of tranfers and restrictions: the composition $\text{Res}_K^G \circ \text{Tr}_H^G : M(G/H) \rightarrow M(G/K)$ is given by the value of the Mackey functor on the diagram

$$\begin{array}{ccccc} & & \coprod_{H \backslash G/K} G/H \cap xKx^{-1} & & \\ & \swarrow & \downarrow & \searrow & \\ G/H & & & & G/K \\ \swarrow & \searrow & & \swarrow & \searrow \\ G/H & & G/G & & G/K \end{array}$$

so by the additivity, it is the direct sum of the value of the morphisms

$$\begin{array}{ccc} & G/H \cap xKx^{-1} & \\ \swarrow & & \searrow \\ G/H & & G/K \end{array}$$

which is the composition $\text{Tr}_{x^{-1}Hx \cap K}^K \circ c_x \circ \text{Res}_{H \cap xKx^{-1}}^H$. We now have the double coset formula

Theorem 3.1 (Double Coset Formula).

$$\text{Res}_K^G \circ \text{Tr}_H^G = \oplus_{x \in H \backslash G / K} \text{Tr}_{x^{-1}Hx \cap K}^K \circ c_x \circ \text{Res}_{H \cap xKx^{-1}}^H$$

By the above discussion, we have a very concrete alternative definition for Mackey functors

Definition 3.1.1. A Mackey functor $\underline{M} : \text{GSet}^{\text{Fin}} \rightarrow \text{Ab}$ consists of the following data:

1. An abelian group $\underline{M}(G/H)$ for every subgroup $H \leq G$;
2. For every subgroup $H \leq K \leq G$, a restriction map $\text{Res}_H^K : \underline{M}(G/K) \rightarrow \underline{M}(G/H)$ and a transfer map $\text{Tr}_H^K : \underline{M}(G/H) \rightarrow \underline{M}(G/K)$.
3. For every subgroup $H \leq G$ and $g \in G$, a conjugation homomorphism $c_g : \underline{M}(G/H) \rightarrow \underline{M}(G/H)$.

The three types of morphisms must also satisfy the following relations:

1. Tr_H^H and Res_H^H are both identity maps for all subgroups $H \leq G$.
2. $\text{Res}_H^K \circ \text{Res}_K^L = \text{Res}_H^L$ and $\text{Tr}_K^L \circ \text{Tr}_H^K = \text{Tr}_H^L$ for $K \leq H \leq L$.
3. $c_g \circ c_h = c_{gh}$ for all $g, h \in G$.
4. $\text{Res}_{gHg^{-1}}^{gKg^{-1}} \circ c_g = c_g \circ \text{Res}_H^K$ and $\text{Tr}_{gHg^{-1}}^{gKg^{-1}} \circ c_g = c_g \circ \text{Tr}_H^K$ for $H \leq K$.
5. For subgroups $H \leq K \leq L$

$$\text{Res}_K^L \circ \text{Tr}_H^L = \oplus_{x \in H \backslash L / K} \text{Tr}_{x^{-1}Hx \cap K}^K \circ c_x \circ \text{Res}_{H \cap xKx^{-1}}^H$$

For any group G , we can now give an example of a constant Mackey functor:

Example 3.1.1 (Constant Mackey functor). For every abelian group A , we have the constant Mackey functor \underline{A} , which assigns every orbit the abelian group A , and restriction maps the identity morphism. One may check that this forces the transfer maps $\text{Tr}_H^K : \mathbb{Z} \rightarrow \mathbb{Z}$ to be multiplication by $|K/H|$, and the conjugation maps to be the identity.

In the case where $G = C_2$ and $A = \mathbb{Z}$, the Mackey functor is described by the following diagram:

$$\begin{array}{c} \mathbb{Z} \\ \text{Id} \left(\begin{array}{c} \nearrow \\ \searrow \end{array} \right) \times 2 \\ \mathbb{Z} \\ \text{Id} \left(\begin{array}{c} \curvearrowright \end{array} \right) \end{array}$$

In the case where $G = C_2$ and $A = \mathbb{Z}/2$, the Mackey functor is described by the following diagram:

$$\begin{array}{c} \mathbb{Z}/2 \\ \text{Id} \left(\begin{array}{c} \nearrow \\ \searrow \end{array} \right) \times 0 \\ \mathbb{Z}/2 \\ \text{Id} \left(\begin{array}{c} \curvearrowright \end{array} \right) \end{array}$$

Remark 3.1.1. Note that the "constant" Mackey functor $\underline{\mathbb{Z}}$ breaks up to a constant coefficient system \mathbb{Z} , but the covariant part is **NOT** the constant functor. We will run into this problem later.

Definition 3.1.1 seems bulky, but when we are dealing with $G = C_p$ Mackey functors, the axioms can be

reduced to the following:

Proposition 3.1.1. A C_p -Mackey functor \underline{M} consists of the following data:

$$\begin{array}{ccc} & \underline{M}(C_p/C_p) & \\ \text{Res}_e^{C_p} \swarrow & & \searrow \text{Tr}_e^{C_p} \\ & \underline{M}(C_p/e) & \\ & \text{\scriptsize c_g} \nearrow & \end{array}$$

where we ignore the some of all of the obvious identity morphisms. The morphisms satisfies the following:

1. $c_g \circ \text{Res}_e^{C_p} = \text{Res}_e^{C_p}$.
2. $\text{Tr}_e^{C_p} \circ c_g = \text{Tr}_e^{C_p}$
3. $\text{Res}_e^{C_p} \circ \text{Tr}_e^{C_p} = \sum_{\gamma \in C_p} c_\gamma$

Here are a few more examples of Mackey functors:

Example 3.1.2 (Burnside Mackey Functor). The Burnside Mackey Functor A_G assigns every orbit G/H the Grothendieck group of the symmetric monoidal category of finite H -sets under coproduct. The transfer and restriction maps come from induction and the forgetful functor, respectively, between H -Sets and K -Sets.

When $G = C_p$, the Burnside Mackey Functor A_{C_p} is given by the following: there are two subgroups e and C_p ; the corresponding Grothendieck groups are $A_{C_p}(e) = \mathbb{Z}$, generated by the singleton with trivial action, and $A_{C_p}(C_p) = \mathbb{Z} \oplus \mathbb{Z}$, generated by the singleton with trivial action and a C_p orbit. The induced G set $C_p \times_e *$ is the C_p orbit, and forgetting the G action on the C_p -orbit gives us a disjoint union of p singletons with trivial action. Diagrammatically, we have the following

$$\begin{array}{ccc} & \mathbb{Z} & \\ (0, Id) \swarrow & & \searrow (Id, p) \\ & \mathbb{Z} \oplus \mathbb{Z} & \\ & \text{\scriptsize Id} \nearrow & \end{array}$$

Example 3.1.3. The homotopy groups of a genuine G -spectra X , $\pi_n^*(X)$, is naturally a Mackey functor: we have the assignment $\pi_n^*(X)(G/H) := \pi_n^H(X)$; by the Wirthmuller isomorphism and the fact that orbits are self dual, we have

$$\pi_n^H(X) \cong [S^n \wedge G/H_+, X] \cong [S^n, X \wedge G/H_+]$$

since we have availabilities on both variances, the tranfers and restrictions maps are easy to define.

Definition 3.1.2. $(\text{RO}(G))$

Theorem 3.2. (Eilenberg-MacLane Spectra) Given a Mackey functor \underline{M} , there exists a Eilenberg MacLane Spectra HM that represents the ordinary $RO(G)$ -graded cohomology theories on Sp^G .

3.2 The $RO(C_2)$ cohomology of a point

3.2.1 Computing $H_{C_2}^{p,q}(*; \mathbb{Z})$

Proposition 3.2.1. The representation ring $RO(C_2) \cong \mathbb{Z}[\sigma]/(\sigma^2 - 1)$.

Thus as abelian group, $RO(C_2) \cong \mathbb{Z}^2$, generated by the trivial representation and the sign representation. We now calculate the ordinary $RO(C_2)$ -cohomology of a point, $H^{p,q}(*; \mathbb{Z})$ where p is the dimension of the trivial representation and q is the dimension of the sign representation.

First, we recall the important fact that the $RO(C_2)$ -cohomology at a trivial representation reduces to the Bredon cohomology with constant coefficient system induced from the Mackey functor. Our game plan is then to first abuse the suspension isomorphism and reduce to Bredon cohomology. We now break the calculation to two parts:

If $q \leq 0$, then by the suspension isomorphism we have

$$H^{p,q}(*; \mathbb{Z}) := \tilde{H}^{p,q}(S^0; \mathbb{Z}) \cong \tilde{H}^{p,0}(S^{-q\sigma}; \mathbb{Z}) \cong \tilde{H}_{\text{Bredon}}^p(S^{-q\sigma}; \mathbb{Z})$$

When the coefficient system is constant, we have the following reduction:

Proposition 3.2.2. If the coefficient system is constant, then

$$\tilde{H}_{\text{Bredon}}^*(X; \underline{M}) \cong \tilde{H}_{\text{Sing}}^*(X/G; M)$$

Recall that $S^{-q\sigma}$ is the one-point compactification of the sign-representation on \mathbb{R}^{-q} . It is not hard to see that $S^{-q\sigma}/C_2$ is homotopy equivalent to $\Sigma\mathbb{RP}^{-q-1}$. (Insert picture). Using suspension isomorphism, we have

$$H^{p,q}(*; \mathbb{Z}) \cong \tilde{H}^{p-1}(\mathbb{RP}^{-q-1})$$

which we know how to compute.

When $q > 0$, we still want to transform the $RO(G)$ -graded cohomology to \mathbb{Z} -graded, so we apply the Spanier-Whitehead duality

Proposition 3.2.3 (Spanier-Whitehead Duality). Note that the sphere spectrum is self-dual, and in particular S^V is dual to S^{-V} .

$$\tilde{H}^{p,q}(S^0; \mathbb{Z}) := [S^0, S^{p+q\sigma} \wedge H\mathbb{Z}] \cong [S^{-p-q\sigma}, S^0 \wedge H\mathbb{Z}] =: \tilde{H}_{-p,-q}(S^0; \mathbb{Z})$$

and we reduce the problem again to computing $\tilde{H}_{-p,-q}(S^0; \mathbb{Z}) \cong \tilde{H}_{-p}^{\text{Bredon}}(S^{q\sigma}; \mathbb{Z})$. However, we should note that the constant Mackey functor does not restrict to the constant covariant coefficient system! Thus, we do not get the analog of proposition 3.2.2 for homology. However, we still have the following simplification available:

Proposition 3.2.4. Suppose the G -action on X is free, then

$$H_*^{\text{Bredon}}(X, \underline{M}) \cong H_*^{\text{Bredon}}(X/G, \underline{M}(e))$$

The C_2 -action on $S^{q\sigma}$ is not free: it has two fixed points at the poles. All hope is not lost, since we have the following cofiber sequence

$$S(V)_+ \rightarrow D(V)_+ \rightarrow S^V$$

where $S(V)$ and $D(V)$ are the unit sphere and unit disk in the representation V , respectively. Moreover, $D(n)_+$ is C_2 -homotopy equivalent to S^0 , and the C_2 -action on $S(V)_+$ is free. Then by the cofiber long exact sequence, we have

$$0 \longrightarrow \tilde{H}_{p+1}(S^{q\sigma}; \mathbb{Z}) \longrightarrow \tilde{H}_p(S(q\sigma)_+; \mathbb{Z}) \longrightarrow 0$$

for $p \geq 1$, and we have the same reduction again. When $p = 0$, we open up the LES

$$0 \longrightarrow \tilde{H}_1(S^{q\sigma}; \mathbb{Z}) \longrightarrow \tilde{H}_0(S(q\sigma)_+; \mathbb{Z}) \cong \mathbb{Z} \xrightarrow{\gamma} \tilde{H}_0(S^0; \mathbb{Z}) \cong \mathbb{Z} \longrightarrow \tilde{H}_0(S^{q\sigma}; \mathbb{Z}) \longrightarrow 0$$

Note that the middle map is induced by the inclusion

$$\tilde{H}_0(S(q\sigma)_+/C_2; \mathbb{Z}) \cong \mathbb{Z} \rightarrow \tilde{H}_0(D(q\sigma)_+/C_2; \mathbb{Z}) \cong \mathbb{Z}$$

which is multiplication by 2.

3.2.2 Computing $H^{p,q}(C_2)(*; A_{C_2})$

First, we will reduce the problem to computing the reduced cohomology of S^0 : $H_{C_2}^{p,q}(*; A_{C_2}) := \tilde{H}_{C_2}^{p,q}(S^0; A_{C_2})$. From now on, we will suppress the coefficient A_{C_2} unless otherwise specified. Our primary strategy is going to be abusing the cofiber sequence

$$S(\sigma)_+ \cong C_{2+} \rightarrow S^0 \rightarrow S^\sigma$$

So the first step is understanding $\tilde{H}_{C_2+}^{p,q}(C_{2+})$.

Proposition 3.2.5. The Mackey functor $\tilde{H}_{C_2+}^{p,q}(C_{2+})$ is represented by the following diagram

$$\begin{array}{ccc} & \tilde{H}_{C_2}^{p,q}(C_{2+}) & \\ \text{Res}_e^{C_2} \swarrow & & \searrow \text{Tr}_e^{C_2} \\ \tilde{H}_{C_2}^{p,q}(C_{2+}) \oplus \tilde{H}_{C_2}^{p,q}(C_{2+}) & & \\ \uparrow \text{swap} & & \end{array}$$

By adjunction, $\tilde{H}_{C_2}^{p,q}(C_{2+}) := [C_{2+}, \Sigma^{p+q\sigma} H A_{C_2}]_{C_2} = [S^0, H A_{C_2}]_e = \tilde{H}_{\text{Sing}}^{|p+q|}(S^0; \mathbb{Z})$.

As a result, we have $\tilde{H}_{C_2}^{p,q}(C_{2+}) = 0$ if $p + q \neq 0$. Then, we can apply the long exact sequence of the cofiber

$$\dots \longrightarrow H_{C_2}^{p-1,q}(C_{2+}) \longrightarrow H_{C_2}^{p,q}(S^\sigma) \longrightarrow H_{C_2}^{p,q}(S^0) \longrightarrow H_{C_2}^{p,q}(C_{2+}) \longrightarrow \dots$$

Thus, if $p + q \neq 0, 1$, we have the isomorphism $\tilde{H}_{C_2}^{p,q-1}(S^0) \cong \tilde{H}_{C_2}^{p,q}(S^\sigma) \cong \tilde{H}_{C_2}^{p,q}(S^0)$. If we plot the Mackey functors onto the plane, where X -axis represents index p , and Y -axis represents index $p+q$, then every lattice point not on $y = 0, 1$ is isomorphic to the lattice point below it. Moreover, by dimension axiom, we have $\tilde{H}_{C_2}^{p,0}(S^0) = 0$ if $p \neq 0$. Thus, all lattice points on the diagonal will be 0, and by the previous observation, all of the 1st and 3rd quadrant will be 0 Mackey functors.

For the rest of the missing pieces, it suffices to compute the values at $p + q = 0, 1, -1$.

Proposition 3.2.6. When $p + q = -1$, we have

$$\tilde{H}_{C_2}^{p,q}(S^0) \cong \ker(\tilde{H}_{C_2}^{p,q}(S^0) \rightarrow \tilde{H}_{C_2}^{p,q+1}(C_{2+}))$$

When $p + q = 1$, we have

$$\tilde{H}_{C_2}^{p,q}(S^0) \cong \operatorname{coker}(\tilde{H}_{C_2}^{p,q-1}(C_{2+}) \rightarrow \tilde{H}_{C_2}^{p,q-1}(S^0))$$

The first isomorphism comes from the cohomology LES of cofiber sequence, and the second isomorphism comes from the homology LES of the cofiber sequence after applying Spanier-Whitehead duality. Note that as Mackey functors, the kernel and cokernel are both trivial at e , since the cohomologies will reduce to singular cohomology, and the maps are injective/surjective. Thus, it suffices to determine the value of the Mackey functors at C_2 , which amounts to computing the value of $\tilde{H}_{C_2}^{p,q}(S^0)$, $\tilde{H}_{C_2}^{p,q}(S^0)$ when $p + q = 1, -1$. These computations can be done again using the cofiber sequence

$$S(q\sigma)_+ \rightarrow D(q\sigma)_+ \rightarrow S^{q\sigma}$$

Proposition 3.2.7. If $p \geq 0$ and $p + q = -1$, then

$$\tilde{H}_{C_2}^{p,q}(S^0) \cong \begin{cases} \mathbb{Z} & p = 0 \\ \mathbb{Z}/2 & p \text{ odd and greater or equal to } 3 \\ 0 & \text{otherwise} \end{cases}$$

Proof. Since $p \geq 0$ and $p + q = -1$, then $\tilde{H}_{C_2}^{p,q}(S^0) \cong \tilde{H}_{C_2}^{p,-p-1}(S^0) \cong \tilde{H}_{C_2}^{p,0}(S^{(p+1)\sigma})$. When $p \geq 2$, the LES gives the isomorphism $\tilde{H}_{C_2}^{p-1,0}(S^{(p+1)\sigma}) \cong \tilde{H}_{C_2}^{p,0}(S^{(p+1)\sigma})$. Again by proposition 3.2.4, $\tilde{H}_{C_2}^{p-1,0}(S^{(p+1)\sigma}) \cong \tilde{H}_{\text{Sing}}^{p-1}(\mathbb{RP}^p)$, which is $\mathbb{Z}/2$ iff p is odd. When $p = 1$, we open up the LES

$$\tilde{H}^{0,0}(D(2\sigma)_+) \rightarrow H^{0,0}(S(2\sigma)_+) \rightarrow \tilde{H}_{C_2}^{1,0}(S^{2\sigma})$$

where the first arrow is easily see to be surjective, and $H_{C_2}^{1,0}(S^{2\sigma}) \cong 0$. For $p = 0$, we have

$$\tilde{H}^{0,0}(S^\sigma) \rightarrow \tilde{H}^{0,0}(D(\sigma)_+) \rightarrow \tilde{H}_{C_2}^{0,0}(S(\sigma)_+)$$

where the second arrow is a projection $\mathbb{Z} \oplus \mathbb{Z} \rightarrow \mathbb{Z}$, so $\tilde{H}_{C_2}^{0,0}(S^\sigma) \cong \mathbb{Z}$.

Proposition 3.2.8. If $p \leq 0$ and $p + q = 1$, then

$$\tilde{H}_{C_2}^{p,q}(S^0) \cong \begin{cases} \mathbb{Z} & p = 0 \\ \mathbb{Z}/2 & p \text{ even and less or equal to } -2 \\ 0 & \text{otherwise} \end{cases}$$

The argument is entirely symmetric by first using duality to turn cohomology into homology, and then apply the LES to the same cofiber sequence.

We are left with what happens when $p + q = 0$. At $p = q = 0$, we know by dimension axiom that the Mackey functor is the burnside Mackey functor A_{C_2} . If $p < 0$, then the LES

$$\tilde{H}_{C_2}^{p,q-1}(S^0) \cong \tilde{H}_{C_2}^{p,q}(S^\sigma) \rightarrow H_{C_2}^{p,q}(S^0) \rightarrow H_{C_2}^{p,q}(S(\sigma)_+) \rightarrow H_{C_2}^{p+1,q-1}(S^0)$$

If further that $p \neq -1$, the two terms on the end are trivial since they land in the 3rd quadrant, and we have the middle arrow an isomorphism. If $p = -1$, one also note that there are no non-trivial maps from $\tilde{H}_{C_2}^{-1,1}(S(\sigma)_+) \rightarrow \tilde{H}_{C_2}^{0,0}(S^0)$. \square

If $p \geq 2$, then we have the long exact sequence

$$\tilde{H}_{C_2}^{p-1,q+1}(S^0) \rightarrow \tilde{H}_{C_2}^{p-1,q+1}(S(\sigma)_+) \rightarrow \tilde{H}_{C_2}^{p,q+1}(S^\sigma) \cong \tilde{H}_{C_2}^{p,q+1}(S^0) \rightarrow \tilde{H}_{C_2}^{p,q-1}(S^0)$$

and the two terms on the end are trivial again since they are in the 1st quadrant, so the middle is an isomorphism.

The last step is to determine the specific case where $p = 1, q = -1$, and it is given by the LES

$$\tilde{H}_{C_2}^{0,0}(S^0) \rightarrow \tilde{H}_{C_2}^{0,0}(C_{2+}) \rightarrow \tilde{H}_{C_2}^{1,0}(S^\sigma) \cong \tilde{H}_{C_2}^{1,-1}(S^0) \rightarrow \tilde{H}_{C_2}^{1,0}(S^0)$$

where the last term is 0 and the first term is the Burnside Mackey functor by dimension axiom, so we realize $\tilde{H}_{C_2}^{1,-1}(S^0)$ as the cokernel.