

MATH 624 HW2

David Zhu

September 22, 2024

Problem 2b

A representative of \tilde{O}_a is given by a pair $(W_1, \frac{f_1}{g_1})$, with $g_1 \neq 0$ on W_1 , and $(W_1, \frac{f_1}{g_1}) \sim (W_2, \frac{f_2}{g_2})$ iff there exists a open $U_{h'} \subset W_1 \cap W_2$ such that $\frac{f_1}{g_1} = \frac{f_2}{g_2}$ on $U_{h'}$. On the other hand, a representative of $k[V]_{\mathfrak{p}_a}$ is given by some $\frac{f}{g}$, where $g(a) \neq 0$. By continuity, there exists a basic open U_h containing a on which g does not vanish. We define the k -algebra homomorphism:

$$i : k[V]_{\mathfrak{p}_a} \rightarrow \tilde{O}_a \quad \frac{f}{g} \mapsto (U_h, \frac{f}{g})$$

Surjectivity is obvious by construction, so there are two things to check: well-definedness (it is clearly that this will be a k -algebra morphism once we check well-definedness) and injectivity.

Well-definedness: suppose $\frac{f}{g} \sim \frac{f'}{g'}$ in $k[V]_{\mathfrak{p}_a}$, which means there exists some $h' \in K[V]$ such that $h'(fg' - f'g) = 0$, which implies $\frac{f}{g} = \frac{f'}{g'}$ on $U_{h'}$. Thus, both will be mapped to the equivalence class $(U_{h'}, \frac{f}{g})$.

Injectivity: suppose $i(\frac{f}{g}) = (U_h, \frac{f}{g})$ represents the 0 element. WLOG, we may assume that f vanishes on U_h , for otherwise we may replace U_h with a smaller basic open. Then, $\frac{f}{g} \sim \frac{0}{1}$ in $k[V]_{\mathfrak{p}_a}$ since $h(f \cdot 1 - g \cdot 0)$ is identically 0 on V .