

# The monodromy determination and McMullen's theorem

Albert Jinghui Yang

Department of Mathematics  
University of Pennsylvania

Nov 2025

- 1 Motivation: Finite Fermat
- 2 Monodromy Representation
- 3 Parshin's Trick

- 1 Motivation: Finite Fermat
- 2 Monodromy Representation
- 3 Parshin's Trick

# Classical Problem in Number Theory

## Problem

For  $n \geq 1$ , how many integral solutions to the equation

$$x^n + y^n = z^n,$$

with  $\gcd(x, y, z) = 1$ ?

# Classical Problem in Number Theory

## Problem

For  $n \geq 1$ , how many integral solutions to the equation

$$x^n + y^n = z^n,$$

with  $\gcd(x, y, z) = 1$ ?

- Turn out to be extremely hard (except for  $n = 1, 2$ , of course).

# Classical Problem in Number Theory

## Problem

For  $n \geq 1$ , how many integral solutions to the equation

$$x^n + y^n = z^n,$$

with  $\gcd(x, y, z) = 1$ ?

- Turn out to be extremely hard (except for  $n = 1, 2$ , of course).
- This is known as the **Fermat's last theorem**, which is proved by Wiles in 1995.

# Classical Problem in Number Theory

## Problem

For  $n \geq 1$ , how many integral solutions to the equation

$$x^n + y^n = z^n,$$

with  $\gcd(x, y, z) = 1$ ?

- Turn out to be extremely hard (except for  $n = 1, 2$ , of course).
- This is known as the **Fermat's last theorem**, which is proved by Wiles in 1995.
- We have a nice way to deal with the weakened version of this problem.

# Finite Fermat



## Theorem (Finite Fermat)

*For  $n \geq 4$ , the equation*

$$x^n + y^n = z^n$$

*has only a finite number of integral solutions with  $\gcd(x, y, z) = 1$ .*

## Theorem (Finite Fermat)

*For  $n \geq 4$ , the equation*

$$x^n + y^n = z^n$$

*has only a finite number of integral solutions with  $\gcd(x, y, z) = 1$ .*

Here's the ideas:

## Theorem (Finite Fermat)

For  $n \geq 4$ , the equation

$$x^n + y^n = z^n$$

has only a finite number of integral solutions with  $\gcd(x, y, z) = 1$ .

Here's the ideas:

- Consider the Riemann surface  $C \subset \mathbb{P}^2(\mathbb{C})$  given by  $x^n + y^n = z^n$ .

# Finite Fermat

## Theorem (Finite Fermat)

For  $n \geq 4$ , the equation

$$x^n + y^n = z^n$$

has only a finite number of integral solutions with  $\gcd(x, y, z) = 1$ .

Here's the ideas:

- Consider the Riemann surface  $C \subset \mathbb{P}^2(\mathbb{C})$  given by  $x^n + y^n = z^n$ .
- Treat this as a family  $C/B$  spread out over the prime  $p \in \mathbb{Z}$ , with fibers  $C_p \simeq C \bmod p$ . Also  $g(C) \geq 2$ . Actually  $B = \operatorname{Spec}(\mathbb{Z}) - \{\text{primes}\}$ .

## Theorem (Finite Fermat)

For  $n \geq 4$ , the equation

$$x^n + y^n = z^n$$

has only a finite number of integral solutions with  $\gcd(x, y, z) = 1$ .

Here's the ideas:

- Consider the Riemann surface  $C \subset \mathbb{P}^2(\mathbb{C})$  given by  $x^n + y^n = z^n$ .
- Treat this as a family  $C/B$  spread out over the prime  $p \in \mathbb{Z}$ , with fibers  $C_p \simeq C \bmod p$ . Also  $g(C) \geq 2$ . Actually  $B = \operatorname{Spec}(\mathbb{Z}) - \{\text{primes}\}$ .
- An integral solution gives a coherent family of points on  $C_p$ , and hence a section of  $C/B$ .

## Theorem (Finite Fermat)

For  $n \geq 4$ , the equation

$$x^n + y^n = z^n$$

has only a finite number of integral solutions with  $\gcd(x, y, z) = 1$ .

Here's the ideas:

- Consider the Riemann surface  $C \subset \mathbb{P}^2(\mathbb{C})$  given by  $x^n + y^n = z^n$ .
- Treat this as a family  $C/B$  spread out over the prime  $p \in \mathbb{Z}$ , with fibers  $C_p \simeq C \bmod p$ . Also  $g(C) \geq 2$ . Actually  $B = \operatorname{Spec}(\mathbb{Z}) - \{\text{primes}\}$ .
- An integral solution gives a coherent family of points on  $C_p$ , and hence a section of  $C/B$ .
- Finite Fermat follows from an arithmetic version of the finiteness of sections for families  $C/B$ .

# Finite Fermat

## First Step

Need to understand the classical version of the finiteness of sections for families  $C/B$ .



## First Step

Need to understand the classical version of the finiteness of sections for families  $C/B$ .

## Theorem (McMullen, a.k.a. Geometric Mordell Conjecture)

*A truly varying family  $C/B$  of genus  $g \geq 2$  has only a finite number of sections  $s : B \rightarrow C$ .*

## First Step

Need to understand the classical version of the finiteness of sections for families  $C/B$ .

## Theorem (McMullen, a.k.a. Geometric Mordell Conjecture)

*A truly varying family  $C/B$  of genus  $g \geq 2$  has only a finite number of sections  $s : B \rightarrow C$ .*

## Goal

Make sense of this theorem!

- 1 Motivation: Finite Fermat
- 2 Monodromy Representation
- 3 Parshin's Trick

# Basics in Riemann Surfaces

Let  $S$  be a smooth oriented Riemann surface of genus  $g \geq 0$ .

# Basics in Riemann Surfaces

Let  $S$  be a smooth oriented Riemann surface of genus  $g \geq 0$ .

## Definition

The **mapping class group**  $\text{Mod}(S)$  is the group of isotopy classes of orientation-preserving diffeomorphism of  $S$ , with group action being the composition.

# Basics in Riemann Surfaces

Let  $S$  be a smooth oriented Riemann surface of genus  $g \geq 0$ .

## Definition

The **mapping class group**  $\text{Mod}(S)$  is the group of isotopy classes of orientation-preserving diffeomorphism of  $S$ , with group action being the composition.

Let  $\mathcal{S}$  be the set of isotopy classes of simple loops (i.e. cannot be shrunk to a point) in  $S$ , and

$$i : \mathcal{S} \times \mathcal{S} \rightarrow \{0, 1, 2, \dots\}$$

be the minimal number of transverse intersections between the given  $\alpha, \beta \in \mathcal{S}$ .

# Basics in Riemann Surfaces

Let  $S$  be a smooth oriented Riemann surface of genus  $g \geq 0$ .

## Definition

The **mapping class group**  $\text{Mod}(S)$  is the group of isotopy classes of orientation-preserving diffeomorphism of  $S$ , with group action being the composition.

Let  $\mathcal{S}$  be the set of isotopy classes of simple loops (i.e. cannot be shrunk to a point) in  $S$ , and

$$i : \mathcal{S} \times \mathcal{S} \rightarrow \{0, 1, 2, \dots\}$$

be the minimal number of transverse intersections between the given  $\alpha, \beta \in \mathcal{S}$ .

There is a natural group action  $\text{Mod}(S)$  on  $(\mathcal{S}, i)$ .

# Classification of Surface Diffeomorphisms

Recall that we learned in Ellis' talk:



# Classification of Surface Diffeomorphisms

Recall that we learned in Ellis' talk:

## Theorem (Nielsen-Thurston)

Any  $f \in \text{Mod}(S)$  is either

- 1 **finite order:**  $f_* : H_1(S; \mathbb{R}) \rightarrow H_1(S; \mathbb{R})$  has complex eigenvalues, and  $f^n = \text{id}$  for some  $n > 0$ ;

# Classification of Surface Diffeomorphisms

Recall that we learned in Ellis' talk:

## Theorem (Nielsen-Thurston)

Any  $f \in \text{Mod}(S)$  is either

- 1 **finite order**:  $f_* : H_1(S; \mathbb{R}) \rightarrow H_1(S; \mathbb{R})$  has complex eigenvalues, and  $f^n = \text{id}$  for some  $n > 0$ ;
- 2 **reducible**: there exists a finite set  $\alpha_1, \dots, \alpha_n \in \mathcal{S}$  permuted by  $f$  with  $i(\alpha_i, \alpha_j) = 0$ ;

# Classification of Surface Diffeomorphisms

Recall that we learned in Ellis' talk:

## Theorem (Nielsen-Thurston)

Any  $f \in \text{Mod}(S)$  is either

- 1 **finite order**:  $f_* : H_1(S; \mathbb{R}) \rightarrow H_1(S; \mathbb{R})$  has complex eigenvalues, and  $f^n = \text{id}$  for some  $n > 0$ ;
- 2 **reducible**: there exists a finite set  $\alpha_1, \dots, \alpha_n \in \mathcal{S}$  permuted by  $f$  with  $i(\alpha_i, \alpha_j) = 0$ ;
- 3 **pseudo-Anosov**: there exists an expansion factor  $K > 1$  such that  $i(f^n(\alpha), \beta)$  grows like  $K^n$  for all  $\alpha, \beta \in \mathcal{S}$ .

To know the McMullen's theorem, we need to understand the finiteness theorem.

To know the McMullen's theorem, we need to understand the finiteness theorem.

## Theorem (Finiteness Theorem, a.k.a. Geometric Shafaravich Conjecture)

*For a given base  $B$  with genus  $g \geq 2$ , there are only finitely many truly varying families  $C/B$  with fibers of genus  $g$ .*

# Finiteness

## Definition

A family  $C/B$  consists of data:

- 1  $B$  is the base Riemann surface.

## Definition

A family  $C/B$  consists of data:

- ①  $B$  is the base Riemann surface.
- ②  $C$  is the 2-dimensional complex manifold with holomorphic fibration

$$\pi : C \rightarrow B,$$

satisfying

- each fiber  $C_t = \pi^{-1}(t)$  is a compact Riemann surface of genus  $g$ , and



## Definition

A family  $C/B$  consists of data:

- ①  $B$  is the base Riemann surface.
- ②  $C$  is the 2-dimensional complex manifold with holomorphic fibration

$$\pi : C \rightarrow B,$$

satisfying

- each fiber  $C_t = \pi^{-1}(t)$  is a compact Riemann surface of genus  $g$ , and
- $B \cong \bar{B} - P$  is the complement of a finite set  $P$  in a compact Riemann surface  $\bar{B}$ .

## Definition

A family  $C/B$  consists of data:

- ①  $B$  is the base Riemann surface.
- ②  $C$  is the 2-dimensional complex manifold with holomorphic fibration

$$\pi : C \rightarrow B,$$

satisfying

- each fiber  $C_t = \pi^{-1}(t)$  is a compact Riemann surface of genus  $g$ , and
- $B \cong \bar{B} - P$  is the complement of a finite set  $P$  in a compact Riemann surface  $\bar{B}$ .

It is **truly varying** if there are  $t, u \in B$  such that  $C_t \not\cong C_u$ .

## Definition

A family  $C/B$  consists of data:

- ①  $B$  is the base Riemann surface.
- ②  $C$  is the 2-dimensional complex manifold with holomorphic fibration

$$\pi : C \rightarrow B,$$

satisfying

- each fiber  $C_t = \pi^{-1}(t)$  is a compact Riemann surface of genus  $g$ , and
- $B \cong \overline{B} - P$  is the complement of a finite set  $P$  in a compact Riemann surface  $\overline{B}$ .


It is **truly varying** if there are  $t, u \in B$  such that  $C_t \not\cong C_u$ .

- How to understand this?

# Monodromy Representation


# Monodromy Representation

Let  $f : X \rightarrow Y$  be a non-constant holomorphic map of Riemann surfaces.



# Monodromy Representation


Let  $f : X \rightarrow Y$  be a non-constant holomorphic map of Riemann surfaces.



For any  $x \in X$ , there is a local chart such that the local expression of  $f$  is  $f(z) = z^{m(x)}$  for some integer  $m(x) \geq 1$ .

# Monodromy Representation

Let  $f : X \rightarrow Y$  be a non-constant holomorphic map of Riemann surfaces.



For any  $x \in X$ , there is a local chart such that the local expression of  $f$  is  $f(z) = z^{m(x)}$  for some integer  $m(x) \geq 1$ . Call  $f(x)$  the **branched points** if  $m(x) > 1$ , and denote  $\mathcal{B}$  the set of branched points (**branched locus**).

# Monodromy Representation



# Monodromy Representation


Let  $f : X \rightarrow Y$  be a non-constant holomorphic map of compact Riemann surfaces of degree  $d$ , and  $\mathcal{B} = \{b_1, \dots, b_n\} \subset Y$  be the branched locus (finiteness follows from Riemann-Hurwitz theorem).

# Monodromy Representation


Let  $f : X \rightarrow Y$  be a non-constant holomorphic map of compact Riemann surfaces of degree  $d$ , and  $\mathcal{B} = \{b_1, \dots, b_n\} \subset Y$  be the branched locus (finiteness follows from Riemann-Hurwitz theorem). Let  $y_0 \notin \mathcal{B}$ , and  $\gamma : [0, 1] \rightarrow Y - \mathcal{B}$  be a loop at  $y_0$ .

# Monodromy Representation


Let  $f : X \rightarrow Y$  be a non-constant holomorphic map of compact Riemann surfaces of degree  $d$ , and  $\mathcal{B} = \{b_1, \dots, b_n\} \subset Y$  be the branched locus (finiteness follows from Riemann-Hurwitz theorem). Let  $y_0 \notin \mathcal{B}$ , and  $\gamma : [0, 1] \rightarrow Y - \mathcal{B}$  be a loop at  $y_0$ .



# Monodromy Representation




# Monodromy Representation



Every  $x \in f^{-1}(y_0)$ ,  $\gamma$  lifts to a path  $\tilde{\gamma}_x$  with the other endpoint  $\tilde{\gamma}_x(1) \in f^{-1}(y_0)$ .

# Monodromy Representation



Every  $x \in f^{-1}(y_0)$ ,  $\gamma$  lifts to a path  $\tilde{\gamma}_x$  with the other endpoint  $\tilde{\gamma}_x(1) \in f^{-1}(y_0)$ . This gives rise to a function

$$\sigma_\gamma : f^{-1}(y_0) \rightarrow f^{-1}(y_0)$$

sending  $x$  to  $\tilde{\gamma}_x(1)$ . This  $\sigma_\gamma$  is actually a permutation of elements in  $f^{-1}(y_0)$ .

# Monodromy Representation

# Monodromy Representation

For any  $y_0 \in Y - \mathcal{B}$ , one obtains a group homomorphism

$$\begin{aligned}\Phi : \pi_1(Y - \mathcal{B}, y_0) &\rightarrow S_d \\ \gamma &\mapsto \sigma_\gamma\end{aligned}$$



# Monodromy Representation

For any  $y_0 \in Y - \mathcal{B}$ , one obtains a group homomorphism

$$\begin{aligned}\Phi : \pi_1(Y - \mathcal{B}, y_0) &\rightarrow S_d \\ \gamma &\mapsto \sigma_\gamma\end{aligned}$$

## Definition

The group homomorphism  $\Phi$  above is called the **monodromy representation** of  $f : X \rightarrow Y$ .

# Examples of Monodromy Representation

# Examples of Monodromy Representation

- Inspired by the nice note by Jennifer Li's note on "Hurwitz Numbers and Monodromy Representation".

# Examples of Monodromy Representation

- Inspired by the nice note by Jennifer Li's note on "Hurwitz Numbers and Monodromy Representation".

Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be  $f(z) = z^3$ . Clearly  $\deg f = 3$ . Choose  $y_0 = 1 \in \mathbb{C}$  and the loop  $\gamma(z) = e^{2\pi iz}$ .

# Examples of Monodromy Representation


- Inspired by the nice note by Jennifer Li's note on "Hurwitz Numbers and Monodromy Representation".

Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be  $f(z) = z^3$ . Clearly  $\deg f = 3$ . Choose  $y_0 = 1 \in \mathbb{C}$  and the loop  $\gamma(z) = e^{2\pi iz}$ . Easy to check  $f^{-1}(1) = \{1, e^{2\pi i/3}, e^{4\pi i/3}\}$ .

# Examples of Monodromy Representation

- Inspired by the nice note by Jennifer Li's note on "Hurwitz Numbers and Monodromy Representation".

Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be  $f(z) = z^3$ . Clearly  $\deg f = 3$ . Choose  $y_0 = 1 \in \mathbb{C}$  and the loop  $\gamma(z) = e^{2\pi iz}$ . Easy to check  $f^{-1}(1) = \{1, e^{2\pi i/3}, e^{4\pi i/3}\}$ .




Lifts

$$\gamma_1(t) = e^{2\pi i/3 \cdot t}$$


Permutation

$$\rightsquigarrow \sigma_{\gamma}(1) = e^{2\pi i/3}$$



$$\gamma_2(t) = e^{2\pi i/3 \cdot (t+1)}$$

$$\rightsquigarrow \sigma_{\gamma}(e^{2\pi i/3}) = e^{4\pi i/3}$$



$$\gamma_3(t) = e^{4\pi i/3 \cdot (t + \frac{2\pi i}{3})}$$

$$\rightsquigarrow \sigma_{\gamma}(e^{4\pi i/3}) = 1$$

$$\Rightarrow \sigma_{\gamma} = (1 \ 2 \ 3) \in S_3.$$

Let  $S$  be a Riemann surface of genus  $g$ .

Let  $S$  be a Riemann surface of genus  $g$ .

## Definition

Let  $X$  be another Riemann surface and  $f : S \rightarrow X$  be any orientation-preserving diffeomorphism.



Let  $S$  be a Riemann surface of genus  $g$ .

## Definition

Let  $X$  be another Riemann surface and  $f : S \rightarrow X$  be any orientation-preserving diffeomorphism. Two such diffeomorphism are equivalent, if

$$f_2 \circ f_1^{-1} : X_1 \rightarrow X_2$$

is isotopic to a biholomorphic map  $h : X_1 \rightarrow X_2$ . In other word, there is an isotopy between  $f_1$  and  $f_2$ .

Let  $S$  be a Riemann surface of genus  $g$ .

## Definition

Let  $X$  be another Riemann surface and  $f : S \rightarrow X$  be any orientation-preserving diffeomorphism. Two such diffeomorphism are equivalent, if

$$f_2 \circ f_1^{-1} : X_1 \rightarrow X_2$$

is isotopic to a biholomorphic map  $h : X_1 \rightarrow X_2$ . In other word, there is an isotopy between  $f_1$  and  $f_2$ . The space

$$T(S) := \{(X, f) \mid f : S \rightarrow X\} / \sim$$

is called a **Teichmüller space** of  $S$ .

# Moduli Spaces

## Definition

The **moduli space** of  $S$ , denoted  $\mathcal{M}_g(S)$ , is given by

$$\mathcal{M}_g(S) := T(S)/\text{Mod}(S).$$

# Moduli Spaces

## Definition

The **moduli space** of  $S$ , denoted  $\mathcal{M}_g(S)$ , is given by

$$\mathcal{M}_g(S) := T(S)/\text{Mod}(S).$$

## Remark

One has  $\mathcal{M}_g(S) \simeq B \text{Diff}^+(S)$  as stacks. In this case,

$$\pi_1(\mathcal{M}_g(S)) = \pi_0(\text{Diff}^+(S)) = \text{Mod}(S).$$

# Moduli Spaces

## Definition

The **moduli space** of  $S$ , denoted  $\mathcal{M}_g(S)$ , is given by

$$\mathcal{M}_g(S) := T(S)/\text{Mod}(S).$$

## Remark

One has  $\mathcal{M}_g(S) \simeq B \text{Diff}^+(S)$  as stacks. In this case,

$$\pi_1(\mathcal{M}_g(S)) = \pi_0(\text{Diff}^+(S)) = \text{Mod}(S).$$

Something of personal interest: Take  $S = S^n$ , then

$$\pi_0(\mathcal{M}_0(S^n)) = \Theta_n,$$

where the latter is the group of smooth structures on  $S^n$ . This is deeply intertwined with the celebrated Kervaire invariant one problem – one of my all-time mathematical obsessions!

# Summary

# Summary

If we don't care about a specific  $S$ , we denote  $M_g$  by the moduli space of Riemann surfaces of genus  $g$ , which is the set of isomorphism classes of Riemann surfaces of genus  $g$ .

# Summary

If we don't care about a specific  $S$ , we denote  $M_g$  by the moduli space of Riemann surfaces of genus  $g$ , which is the set of isomorphism classes of Riemann surfaces of genus  $g$ .

## Remark

To summarize:

- 1 We have the data of family  $C/B$  in the monodromy determination theorem.



# Summary

If we don't care about a specific  $S$ , we denote  $M_g$  by the moduli space of Riemann surfaces of genus  $g$ , which is the set of isomorphism classes of Riemann surfaces of genus  $g$ .

## Remark

To summarize:

- 1 We have the data of family  $C/B$  in the monodromy determination theorem.
- 2 We discuss the monodromy representation

$$\Phi : \pi_1(Y, y_0) \rightarrow S_d.$$

# Summary

If we don't care about a specific  $S$ , we denote  $M_g$  by the moduli space of Riemann surfaces of genus  $g$ , which is the set of isomorphism classes of Riemann surfaces of genus  $g$ .

## Remark

To summarize:

- 1 We have the data of family  $C/B$  in the monodromy determination theorem.
- 2 We discuss the monodromy representation

$$\Phi : \pi_1(Y, y_0) \rightarrow S_d.$$

- 3 We introduce the Teichmüller spaces and the moduli spaces.

# Summary

If we don't care about a specific  $S$ , we denote  $M_g$  by the moduli space of Riemann surfaces of genus  $g$ , which is the set of isomorphism classes of Riemann surfaces of genus  $g$ .

## Remark

To summarize:

- 1 We have the data of family  $C/B$  in the monodromy determination theorem.
- 2 We discuss the monodromy representation

$$\Phi : \pi_1(Y, y_0) \rightarrow S_d.$$

- 3 We introduce the Teichmüller spaces and the moduli spaces.

# Summary

# Summary

To put them in the same picture: there is a natural complex structure on  $\mathcal{M}_g$  (inherited from  $T_g$ , the **universal covering of  $\mathcal{M}_g$** , see Tom's talk), such that for all family  $C/B$ , it determines a holomorphic classifying map

$$\begin{aligned} F : B &\rightarrow \mathcal{M}_g \\ t &\mapsto [C_t = \pi^{-1}(t)] \end{aligned}$$

with monodromy representation

$$F_* : \pi_1(B, t) \rightarrow \text{Mod}(C_t) = \pi_1(\mathcal{M}_g).$$

# Summary

To put them in the same picture: there is a natural complex structure on  $\mathcal{M}_g$  (inherited from  $T_g$ , the **universal covering of  $\mathcal{M}_g$** , see Tom's talk), such that for all family  $C/B$ , it determines a holomorphic classifying map

$$\begin{aligned} F : B &\rightarrow \mathcal{M}_g \\ t &\mapsto [C_t = \pi^{-1}(t)] \end{aligned}$$

with monodromy representation

$$F_* : \pi_1(B, t) \rightarrow \text{Mod}(C_t) = \pi_1(\mathcal{M}_g).$$

This is the generalized monodromy representation in the new setting  $C/B$ , instead of a simple covering  $C \rightarrow B$  of fixed degree.

# Proof of Finiteness: Baby Case

# Proof of Finiteness: Baby Case

Recall in Tom' talk:

- $\dim T_g = 3g - 3$ . So  $T_g \cong$  open bounded domain in  $\mathbb{C}^{3g-3}$ .
- **Uniformization Theorem:** Up to biholomorphism, there are just 3 simply connected Riemann surface:  $\mathbb{C}, \hat{\mathbb{C}}, \mathbb{D}$ .



# Proof of Finiteness: Baby Case

Recall in Tom' talk:

- $\dim T_g = 3g - 3$ . So  $T_g \cong$  open bounded domain in  $\mathbb{C}^{3g-3}$ .
- **Uniformization Theorem:** Up to biholomorphism, there are just 3 simply connected Riemann surface:  $\mathbb{C}, \hat{\mathbb{C}}, \mathbb{D}$ .

Let  $\tilde{B}$  be the universal cover of  $B$  in the family  $C/B$ , we have a commutative diagram

$$\begin{array}{ccc} B & \xrightarrow{F} & \mathcal{M}_g \\ \uparrow & & \uparrow \\ \tilde{B} & \xrightarrow{\tilde{F}} & T_g \end{array}$$

# Proof of Finiteness: Baby Case

Recall in Tom' talk:

- $\dim T_g = 3g - 3$ . So  $T_g \cong$  open bounded domain in  $\mathbb{C}^{3g-3}$ .
- **Uniformization Theorem:** Up to biholomorphism, there are just 3 simply connected Riemann surface:  $\mathbb{C}, \hat{\mathbb{C}}, \mathbb{D}$ .

Let  $\tilde{B}$  be the universal cover of  $B$  in the family  $C/B$ , we have a commutative diagram

$$\begin{array}{ccc} B & \xrightarrow{F} & \mathcal{M}_g \\ \uparrow & & \uparrow \\ \tilde{B} & \xrightarrow{\tilde{F}} & T_g \end{array}$$

So in the case of  $g(B) < 2$ ,  $\tilde{B} \cong \mathbb{C}$  ( $g(B) = 1$ ) or  $\hat{\mathbb{C}}$  ( $g(B) = 0$ ), and thus  $\tilde{F}$  is trivial. Hence, all  $C/B$  is trivial.

# Modular Schwarz Lemma

# Modular Schwarz Lemma

Consider the case  $B$  is hyperbolic, i.e.  $g(B) \geq 2$ . To decide the fate of  $C/B$ , we need to modular Schwarz lemma.

# Modular Schwarz Lemma

Consider the case  $B$  is hyperbolic, i.e.  $g(B) \geq 2$ . To decide the fate of  $C/B$ , we need to modular Schwarz lemma.

## Theorem

*Any holomorphic  $F : B \rightarrow \mathcal{M}_g$  is distance-decreasing from the hyperbolic metric on  $B$  to the Teichmüller metric on  $\mathcal{M}_g$ . More precisely,*

$$d_{\text{Teichmüller}}(F(s), F(t)) \leq \frac{d_{\text{hyperbolic}}(s, t)}{2}.$$

# Modular Schwarz Lemma

Consider the case  $B$  is hyperbolic, i.e.  $g(B) \geq 2$ . To decide the fate of  $C/B$ , we need to modular Schwarz lemma.

## Theorem

*Any holomorphic  $F : B \rightarrow \mathcal{M}_g$  is distance-decreasing from the hyperbolic metric on  $B$  to the Teichmüller metric on  $\mathcal{M}_g$ . More precisely,*

$$d_{\text{Teichmüller}}(F(s), F(t)) \leq \frac{d_{\text{hyperbolic}}(s, t)}{2}.$$

## Remark

- 1 The standard hyperbolic metric on  $B$  is

$$ds^2 = \frac{4|dz|^2}{(1 - |z|^2)^2}.$$

# Modular Schwarz Lemma

Consider the case  $B$  is hyperbolic, i.e.  $g(B) \geq 2$ . To decide the fate of  $C/B$ , we need to modular Schwarz lemma.

## Theorem

*Any holomorphic  $F : B \rightarrow \mathcal{M}_g$  is distance-decreasing from the hyperbolic metric on  $B$  to the Teichmüller metric on  $\mathcal{M}_g$ . More precisely,*

$$d_{\text{Teichmüller}}(F(s), F(t)) \leq \frac{d_{\text{hyperbolic}}(s, t)}{2}.$$

## Remark

- 1 The standard hyperbolic metric on  $B$  is

$$ds^2 = \frac{4|dz|^2}{(1 - |z|^2)^2}.$$

- 2 The Teichmüller metric on  $\mathcal{M}_g$  is

## Remark

- ① The standard hyperbolic metric on  $B$  is

$$ds^2 = \frac{4|dz|^2}{(1 - |z|^2)^2}.$$



## Remark

- ① The standard hyperbolic metric on  $B$  is

$$ds^2 = \frac{4|dz|^2}{(1 - |z|^2)^2}.$$

- ② The Teichmüller metric on  $\mathcal{M}_g$  is

$$d(X, Y) = \frac{1}{2} \log \inf \{ K \geq 1 : \text{there exists } K\text{-quasiconformal map} \\ f : X \rightarrow Y \text{ respecting markings} \}.$$

# Example of Monodromy over Punctured Disk

# Example of Monodromy over Punctured Disk

Consider  $C/B$  a family with  $B = \Delta^* = \{z : 0 < |z| < 1\}$  a punctured disk.

# Example of Monodromy over Punctured Disk

Consider  $C/B$  a family with  $B = \Delta^* = \{z : 0 < |z| < 1\}$  a punctured disk. Equipped  $B$  with the standard hyperbolic metric. Then 0 is infinitely far away in this metric. This is because for any point  $p$ , the hyperbolic line  $z(t) = t$  for  $t \in [\epsilon, r]$  has

$$ds = \frac{2dt}{1 - t^2},$$

# Example of Monodromy over Punctured Disk

Consider  $C/B$  a family with  $B = \Delta^* = \{z : 0 < |z| < 1\}$  a punctured disk. Equipped  $B$  with the standard hyperbolic metric. Then 0 is infinitely far away in this metric. This is because for any point  $p$ , the hyperbolic line  $z(t) = t$  for  $t \in [\epsilon, r]$  has

$$ds = \frac{2dt}{1-t^2},$$

implying

$$\int_{\epsilon}^r \frac{2dt}{1-t^2} = \log \frac{1+r}{1-r} - \log \frac{1+\epsilon}{1-\epsilon}.$$

When  $\epsilon \rightarrow 0$ , the length tends to  $\log \frac{1+r}{1-r}$ , which tends to 0 as  $r \rightarrow 0$ . So  $\Delta^*$  has a cusp at 0.

# Example of Monodromy over Punctured Disk

# Example of Monodromy over Punctured Disk

Similarly, the circle  $S^1(r) = \{z : |z| = r\}$  has the hyperbolic metric (by setting  $z = re^{i\theta}$  for  $\theta \in [0, 2\pi]$ )

$$ds = \frac{d\theta}{\log(1/r)},$$

implying the length of  $S^1(r) \rightarrow 0$  as  $r \rightarrow 0$ .

# Example of Monodromy over Punctured Disk

Similarly, the circle  $S^1(r) = \{z : |z| = r\}$  has the hyperbolic metric (by setting  $z = re^{i\theta}$  for  $\theta \in [0, 2\pi]$ )

$$ds = \frac{d\theta}{\log(1/r)},$$

implying the length of  $S^1(r) \rightarrow 0$  as  $r \rightarrow 0$ . By the modular Schwarz lemma,  $F : \Delta^* \rightarrow \mathcal{M}_g$  shrinks the distance. It follows that

$$\begin{array}{ccc} F_*(\pi_1(B)) & \xlongequal{\quad} & [f] \in \mathcal{M}_g \\ \uparrow & & \uparrow \\ \tilde{\gamma} \in \tilde{B} & \longrightarrow & \tilde{\gamma}_f \in T_g \end{array}$$

and  $\tau(f) = \inf \text{length}(\tilde{\gamma}_f) = 0$  because the hyperbolic metric for  $\gamma$  tends to 0.



## Example of Monodromy over Punctured Disk

Similarly, the circle  $S^1(r) = \{z : |z| = r\}$  has the hyperbolic metric (by setting  $z = re^{i\theta}$  for  $\theta \in [0, 2\pi]$ )

$$ds = \frac{d\theta}{\log(1/r)},$$

implying the length of  $S^1(r) \rightarrow 0$  as  $r \rightarrow 0$ . By the modular Schwarz lemma,  $F : \Delta^* \rightarrow \mathcal{M}_g$  shrinks the distance. It follows that

$$\begin{array}{ccc} F_*(\pi_1(B)) & \xlongequal{\quad} & [f] \in \mathcal{M}_g \\ \uparrow & & \uparrow \\ \tilde{\gamma} \in \tilde{B} & \longrightarrow & \tilde{\gamma}_f \in T_g \end{array}$$

and  $\tau(f) = \inf \text{length}(\tilde{\gamma}_f) = 0$  because the hyperbolic metric for  $\gamma$  tends to 0. By Nielsen-Thurston,  $f$  is reducible or of finite order. (If  $\tau(f) > 0$ , then pseudo-Anosov since  $f$  moves every surface some distance.)

# Proof Idea of Finiteness

# Proof Idea of Finiteness

Idea: To show only finitely many truly varying families  $C/B$  by showing the space  $\mathcal{F}$  of all classifying maps  $F : B \rightarrow \mathcal{M}_g$  is compact and discrete.

# Proof Idea of Finiteness

Idea: To show only finitely many truly varying families  $C/B$  by showing the space  $\mathcal{F}$  of all classifying maps  $F : B \rightarrow \mathcal{M}_g$  is compact and discrete.

- I won't go into the proof, since it is rather heavy on analysis for a homotopy theorist. 🤖

# Proof Idea of Finiteness

Idea: To show only finitely many truly varying families  $C/B$  by showing the space  $\mathcal{F}$  of all classifying maps  $F : B \rightarrow \mathcal{M}_g$  is compact and discrete.

- I won't go into the proof, since it is rather heavy on analysis for a homotopy theorist. 🤖

## Theorem (Rigidity)

*A truly varying family  $C/B$  is determined up to finitely many choices by its monodromy*

$$F_* : \pi_1(B) \rightarrow \text{Mod}_g = \pi_1(\mathcal{M}_g).$$

# Outline

- 1 Motivation: Finite Fermat
- 2 Monodromy Representation
- 3 Parshin's Trick

# Parshin's Argument

# Parshin's Argument

**Upshot**: Given a section  $s : B \rightarrow C$  for a fibration  $C \rightarrow B$ , one can use  $s$  to build a new family by taking a branched covering of  $C$  along  $s(B)$ .



# Parshin's Argument

**Upshot**: Given a section  $s : B \rightarrow C$  for a fibration  $C \rightarrow B$ , one can use  $s$  to build a new family by taking a branched covering of  $C$  along  $s(B)$ . The new constructed family  $D/B$  has fibers of higher genus  $g \geq 2$ . Then we can use the Finiteness Theorem to deal with  $D/B$ .

# Parshin's Argument

**Upshot:** Given a section  $s : B \rightarrow C$  for a fibration  $C \rightarrow B$ , one can use  $s$  to build a new family by taking a branched covering of  $C$  along  $s(B)$ . The new constructed family  $D/B$  has fibers of higher genus  $g \geq 2$ . Then we can use the Finiteness Theorem to deal with  $D/B$ .

## Theorem

*Given a genus  $g \geq 1$  and a base  $B$ , there exists a genus  $h \geq 2$  and a finite-to-one map*

$$\{C/B \text{ with fibers of genus } g + \text{sections } s : B \rightarrow C\} \rightarrow \{D/B \text{ with fibers of genus } h\}.$$

*For each  $t \in B$ , the surface  $D_t$  is a covering of  $C_t$  branched over the single point  $s(t) \in C(t)$ .*

# McMullen's Theorem

We are ready to prove the McMullen's theorem, as our ultimate goal:

## Theorem (McMullen)

*A truly varying family  $C/B$  of genus  $g \geq 2$  has only a finite number of sections  $s : B \rightarrow C$ .*

# McMullen's Theorem

We are ready to prove the McMullen's theorem, as our ultimate goal:

## Theorem (McMullen)

*A truly varying family  $C/B$  of genus  $g \geq 2$  has only a finite number of sections  $s : B \rightarrow C$ .*

## Proof

For each section  $s : B \rightarrow C$ , we can form the family  $D/B$  of genus  $h \geq 2$  branched over  $s(B)$  by Parshin's trick.

# McMullen's Theorem

We are ready to prove the McMullen's theorem, as our ultimate goal:

## Theorem (McMullen)

*A truly varying family  $C/B$  of genus  $g \geq 2$  has only a finite number of sections  $s : B \rightarrow C$ .*

## Proof

For each section  $s : B \rightarrow C$ , we can form the family  $D/B$  of genus  $h \geq 2$  branched over  $s(B)$  by Parshin's trick. The number of truly varying families  $D/B$  is finite by the Finiteness Theorem, so the number of pairs  $(C, s)$  is finite up to automorphism over  $B$ .

# McMullen's Theorem

We are ready to prove the McMullen's theorem, as our ultimate goal:

## Theorem (McMullen)

*A truly varying family  $C/B$  of genus  $g \geq 2$  has only a finite number of sections  $s : B \rightarrow C$ .*

## Proof

For each section  $s : B \rightarrow C$ , we can form the family  $D/B$  of genus  $h \geq 2$  branched over  $s(B)$  by Parshin's trick. The number of truly varying families  $D/B$  is finite by the Finiteness Theorem, so the number of pairs  $(C, s)$  is finite up to automorphism over  $B$ . However,  $\text{Aut}(C/B)$  is finite (Theorem 2.2 in the paper, also in Ellis' talk).

# McMullen's Theorem

We are ready to prove the McMullen's theorem, as our ultimate goal:

## Theorem (McMullen)

*A truly varying family  $C/B$  of genus  $g \geq 2$  has only a finite number of sections  $s : B \rightarrow C$ .*

## Proof

For each section  $s : B \rightarrow C$ , we can form the family  $D/B$  of genus  $h \geq 2$  branched over  $s(B)$  by Parshin's trick. The number of truly varying families  $D/B$  is finite by the Finiteness Theorem, so the number of pairs  $(C, s)$  is finite up to automorphism over  $B$ . However,  $\text{Aut}(C/B)$  is finite (Theorem 2.2 in the paper, also in Ellis' talk). It follows that the number of sections  $s : B \rightarrow C$  is also finite.

- ① C. T. McMullen, *From dynamics on surfaces to rational points on curves*. Bulletin of the American Mathematical Society, volume 37 (2), pages 119–140, 1999.



*Thank you!*