

## The use of He's variational iteration method for solving the telegraph and fractional telegraph equations

Mehdi Dehghan<sup>1,\*</sup>, S. A. Yousefi<sup>2</sup> and A. Lotfi<sup>2</sup>

<sup>1</sup>*Department of Applied Mathematics, Faculty of Mathematics and Computer Sciences,  
Amirkabir University of Technology, No. 424 Hafez Avenue, Tehran, Iran*

<sup>2</sup>*Department of Mathematics, Shahid Beheshti University, Tehran, Iran*

### SUMMARY

In this paper the variational iteration method is used to compute the solution for the linear, variable coefficient, fractional derivative and multi space telegraph equations. The method constructs a convergent sequence of functions, to approximate the exact solution with a few number of iterations without discretization. Numerical results and comparison with exact solutions are given for some examples in order to show its applicability and efficiency. Copyright © 2009 John Wiley & Sons, Ltd.

Received 5 March 2009; Accepted 22 May 2009

KEY WORDS: telegraph equation; variational iteration method; semi-analytical solution

### 1. INTRODUCTION

Telegraph equations are hyperbolic partial differential equations that are applicable in several fields such as wave propagation [1], signal analysis [2], random walk theory [3], etc. In this paper we consider four different types of linear telegraph equations.

The first one is the classical linear telegraph equation:

$$\begin{aligned}u_{tt} + \alpha u_t + \beta u &= u_{xx} + f(x, t) \\ u(x, 0) &= \phi, \quad u_t(x, 0) = \psi\end{aligned}\tag{1}$$

where  $u$  can be considered as a function depending on distance ( $x$ ) and time ( $t$ ),  $\alpha$  and  $\beta$  are constants depending on given problem and  $f$ ,  $\phi$ ,  $\psi$  are known continuous functions. This type of telegraph equation applies to high-frequency transmission lines such as telegraph wires and radio frequency conductors. It is also applicable for designing high voltage transmission lines. Several numerical methods exist for solving this equation. Authors of [4] used an analytical approach to solve it.

\*Correspondence to: Mehdi Dehghan, Department of Applied Mathematics, Faculty of Mathematics and Computer Sciences, Amirkabir University of Technology, No. 424 Hafez Avenue, Tehran, Iran.

†E-mail: mdehghan@aut.ac.ir

The second equation is the following linear telegraph equation with variable coefficients [5]:

$$\begin{aligned} r(x)u_{tt}(x, t) &= [p(x)u_x(x, t)]_x - q(x)u(x, t) + F(x, t), \quad 0 \leq x \leq 1 \\ au_x(0, t) + bu(0, t) &= 0, \quad t > 0 \\ cu_x(1, t) + du(1, t) &= 0, \quad t > 0 \\ u(x, 0) &= f(x), \quad 0 \leq x \leq 1 \\ u_t(x, 0) &= g(x), \quad 0 \leq x \leq 1 \end{aligned} \quad (2)$$

where  $|a| + |b| > 0$ ,  $|c| + |d| > 0$ ,  $r(x) > 0$ ,  $p(x) > 0$  and  $p, r, q, f, g, F$  are continuous functions. This equation is solved in [5] by means of discrete eigenfunction method.

The third equation that we are going to solve numerically is the telegraph equation with fractional derivatives:

$$\begin{aligned} D_x^\alpha u(x, t) &= u_{tt}(x, t) + u_t(x, t) + u(x, t) + f(x, t), \quad t \geq 0, \quad 1 < \alpha \leq 2 \\ u(0, t) &= l_1(t), \quad t \geq 0 \\ u_x(0, t) &= l_2(t), \quad t \geq 0 \\ u(x, 0) &= L(x), \quad 0 < x < 1 \end{aligned} \quad (3)$$

where  $\alpha$  is the parameter of order of fractional derivative and  $D_x^\alpha$  is the Caputo fractional differential operator defined as

$$D_x^\alpha u(x, t) = \begin{cases} \frac{1}{\Gamma(m-\alpha)} \int_0^x (x-\theta)^{m-\alpha-1} \frac{\partial^m}{\partial \theta^m} u(\theta, t) d\theta, & m-1 < \alpha \leq m \\ \frac{\partial^m}{\partial x^m} u(x, t), & \alpha = m \in \mathbb{N} \end{cases}$$

This fractional type of telegraph equation is solved in [6, 7] by the Adomian decomposition method. Applications of this equation are pointed out in [6] and the references therein.

Finally, we discuss on the multi-dimensional telegraph equations such as the three-space-dimensional equation with variable coefficients [8, 9]. This equation can be shown by

$$\begin{aligned} u_{tt}(x, y, z, t) + p(x, y, z, t)u_t + q(x, y, z, t)u(x, y, z, t) \\ = A(x, y, z, t)u_{xx}(x, y, z, t) + B(x, y, z, t)u_{yy}(x, y, z, t) \\ + C(x, y, z, t)u_{zz}(x, y, z, t) + f(x, y, z, t), \quad 0 < x, y, z < 1, \quad t > 0 \end{aligned} \quad (4)$$

with initial conditions,

$$\begin{aligned} u(x, y, z, 0) &= \phi(x, y, z), \quad 0 \leq x, y, z \leq 1 \\ u_t(x, y, z, 0) &= \psi(x, y, z), \quad 0 \leq x, y, z \leq 1 \end{aligned}$$

and boundary conditions,

$$\begin{aligned} u(x, y, z, t) &= g(x, y, z, t), \quad (x, y, z, t) \in \partial\Omega \times [0, T] \\ \Omega &= \{(x, y, z) | 0 \leq x, y, z \leq 1\} \end{aligned}$$

where  $A, B, C, p$  and  $q$  are in  $\Omega \times [0, T]$  and  $\phi, \psi$  and their partial derivatives up to order two are continuous and  $T$  is a positive integer. The two-dimensional version of this equation is solved numerically in [9]. It is worth pointing out that authors of [10] used the Adomian decomposition method [11] to obtain solutions of the fourth-order fractional diffusion-wave equation that is defined in a bounded space domain. The one-dimensional telegraph equation is investigated in [12].

A numerical technique is presented [12] for the solution of this equation that uses the Chebyshev cardinal functions. The method consists of expanding the required approximate solution as the elements of the Chebyshev cardinal functions. Using the operational matrix of derivative, the problem is reduced to a set of algebraic equations [12]. Authors of [13] developed a numerical scheme to solve this hyperbolic telegraph equation using collocation points and approximating directly the solution using the thin plate splines radial basis function. The scheme works in a similar fashion as finite difference methods. Authors of [14] combined a high-order compact finite difference scheme to approximate the spatial derivative and collocation technique for the time component to numerically solve the one-space-dimensional linear hyperbolic telegraph equation. A numerical technique is presented in [15] to solve this equation. This method is based on the shifted Chebyshev tau method. This method consists of expanding the required approximate solution as the elements of shifted Chebyshev polynomials. Using the operational matrices of integral and derivative, the problem can be reduced [15] to a set of linear algebraic equations. The two-dimensional version of the telegraph equation is investigated in [9] and a high-order accurate method is proposed for solving it. The proposed technique is based on a compact finite difference approximation of fourth order for discretizing spatial derivatives and collocation method for the time component. The resulted method is unconditionally stable and solves the two-dimensional linear telegraph equation with high accuracy. In Dehghan and Mohebbi's approach [9], the solution is approximated by a polynomial at each grid point whose coefficients are determined by solving a linear system of equations.

In the current investigation, the approach is different as we use the variational iteration method to solve four kinds of telegraph equations. In this paper we are going to find the solution of the telegraph equations by using this method. With demonstrating examples we will see that by few iterations we get approximate solutions that are very accurate in comparison with exact or other solutions, which are achieved by other methods, and it shows the efficiency of this method.

The article is organized as follows: In Section 2 we describe the variational iteration method. In Section 3 we use the method for Problems (1)–(4) in order to construct an iteration formula. In Section 4 we demonstrate the accuracy of method by considering numerical examples. Finally Section 5 concludes this article with a brief summary.

## 2. VARIATIONAL ITERATION METHOD

The variational iteration method is a method for solving linear and nonlinear problems. This method is introduced by the Chinese researcher He [16] by modifying the general Lagrange multiplier method [17]. The method constructs an iterative sequence of functions converging to exact solution. In the case of linear problems by determining exact Lagrange multiplier, approximate solution turns into exact solution and is available by only one iteration. The method works without discretization of problem and round off error does not affect it. The method is used in several well-known problems successfully such as delay differential equations [18], differential equations with fractional derivatives [19–21], autonomous differential equations [22], Helmholtz equations [23], Burgers and coupled Burgers equations [24], non-linear wave and diffusion equations [25], parabolic integro-differential equations arising in heat conduction in materials with memory [26] and some other problems such as [16, 21, 27, 28]. In [29] a convergence criterion based on Banach fixed point is given for this method. The variational iteration method is used in [30] for solving the Lane–Emden equation, in [31] to solve several problems in calculus of variations, in [32] to find the solution of an inverse problem in partial differential equations, in [33] to solve a system of nonlinear integro-differential equations arising in biology, in [34] to solve the Kline–Gordon equation, in [35] to find solution of the Fokker–Planck equation, in [36] to solve the Cauchy–reaction–diffusion partial differential equation, in [37] to solve the biological population equation. In addition the main advantage of the new approach in comparison with the mesh point techniques [38] is highlighted in these works. Wazwaz [39] employed this method to solve the nonlinear Goursat problem. In [40] the variational iteration method is compared with

the Adomian decomposition method for homogeneous and nonhomogeneous advection problems. Author of [41] used this scheme to determine rational solutions for the KdV,  $k(2, 2)$ , Burgers and cubic Boussinesq equations. Abbasbandy and Shirzadi [42] employed the variational iteration method to solve some eighth-order boundary-value problems. Some special cases of the mentioned equations that govern scientific and engineering experimentations are tested. In addition the initial approximation was selected as a polynomial with unknown constants, which was determined by considering the boundary conditions. Nonlinear Volterra's integro-differential equations are solved in [43] using the variational iteration technique. The variational iteration method is applied to find the solution of the wave equation subject to an integral conservation condition [44]. Authors of [45] modified the variational iteration method to solve a system of differential equations. The variation iteration technique is used in [46] to solve the generalized pantograph equation [47]. This scheme is employed in [48] to solve several variational problems.

Consider the following general nonlinear problem:

$$L(u(t)) + N(u(t)) = g(t) \quad (5)$$

where  $L$  is a linear operator,  $N$  is a nonlinear operator and  $g(t)$  is a known analytical function. Variational iteration method constructs an iterative sequence called correction functional as

$$u_{n+1}(t) = u_n(t) + \int_0^t \lambda(L(u_n(s)) + N(\tilde{u}_n(s)) - g(s)) ds \quad (6)$$

where  $\lambda$  is general Lagrange multiplier, which can be identified optimally via the variational theory [16, 24],  $\tilde{u}_n(s)$  is considered as restricted variation [16, 49], i.e.  $\delta\tilde{u}_n = 0$ , and the index  $n$  denotes the  $n$ th iteration.

### 3. ANALYSIS OF TELEGRAPH EQUATIONS

Considering Equation (1) with respect to Equation (6), we construct an iteration formula in  $t$ -direction, which is as follows:

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t \lambda(s)((u_n(x, s))_{ss} + \alpha(u_n(x, s))_s + \beta\widetilde{u_n(x, s)} - (\widetilde{u_n(x, s)})_{xx} - f(x, s)) ds$$

We now determine the Lagrange multiplier.

$$\begin{aligned} \delta u_{n+1}(x, t) &= \delta u_n(x, t) + \delta \int_0^t \lambda(s)((u_n(x, s))_{ss} + \alpha(u_n(x, s))_s + \beta\widetilde{u_n(x, s)} \\ &\quad - ((\widetilde{u_n(x, s)})_{xx}) - f(x, s)) ds = 0 \\ \delta u_{n+1} &= \delta u_n + \alpha\lambda(s)\delta u_n|_{s=t} + \lambda(s)\delta((u_n)_s)|_{s=t} - \lambda'(s)\delta(u_n)|_{s=t} \\ &\quad + \int_0^t (\lambda''(s) - \alpha\lambda'(s) + \beta\lambda(s))\delta u_n ds = 0 \\ \delta u_n : (1 - \lambda'(s) + \alpha\lambda(s))|_{s=t} &= 0 \\ \delta((u_n)_s) : \lambda(s)|_{s=t} &= 0 \\ \delta u_n : (\lambda''(s) - \alpha\lambda'(s)) &= 0 \end{aligned}$$

So we have

$$\lambda(s) = \frac{1}{\alpha}(e^{\alpha(s-t)} - 1)$$

and we obtain the following iteration formula:

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t \frac{1}{\alpha} (e^{\alpha(s-t)} - 1) ((u_n(x, s))_{ss} + \alpha(u_n(x, s))_s + \beta u_n(x, s) - ((u_n(x, s))_{xx} - f(x, s)) ds \quad (7)$$

with initial approximation  $u_0(x, t)$  such that  $u_0(x, 0) = \phi$  and  $u_{0t}(x, 0) = \psi$ , which guarantees satisfaction of boundary conditions in each iteration. Considering Equation (2) we construct an iteration sequence the same as the above:

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t \lambda(s) (r(x) u_{ss}(x, s) - ([p(x) \widetilde{u_x(x, s)}]_x) + q(x) \widetilde{u(x, s)} - F(x, s)) ds$$

By taking variation with respect to  $u_n$ , we determine the Lagrange multiplier.

$$\begin{aligned} \delta u_{n+1}(x, t) &= \delta u_n(x, t) + \delta \int_0^t \lambda(s) (r(x) u_{ss}(x, s) \\ &\quad - ([p(x) \widetilde{u_x(x, s)}]_x) + q(x) \widetilde{u(x, s)} - F(x, s)) ds = 0 \end{aligned}$$

$$\delta u_n : 1 - r(x) \lambda'(s) |_{s=t} = 0$$

$$\delta((u_n)_s) : r(x) \lambda(s) |_{s=t} = 0$$

$$\delta u_n : r(x) \lambda''(s) = 0$$

So we have

$$\lambda(s) = \frac{1}{r(x)} (s - t)$$

and we get the iteration formula:

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t \frac{1}{r(x)} (s - t) (r(x) u_{ss}(x, s) - ([p(x) \widetilde{u_x(x, s)}]_x) + q(x) u(x, s) - F(x, s)) ds \quad (8)$$

We start the iteration formula with initial approximation

$$u_0(x, t) = f(x) + tg(x)$$

For Equation (3) we construct the following iteration formula:

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^x \lambda(s) (D_s^\alpha u(s, t) - \widetilde{u_{tt}(s, t)} - \widetilde{u_t(s, t)} - \widetilde{u(s, t)} - f(s, t)) ds$$

By taking variation with respect to  $u_n$  when  $\alpha=2$ , we determine the Lagrange multiplier approximately.

$$\delta u_{n+1}(x, t) = \delta u_n(x, t) + \delta \int_0^x \lambda(s) (D_s^\alpha u(s, t) - \widetilde{u_{tt}(s, t)} - \widetilde{u_t(s, t)} - \widetilde{u(s, t)} - f(s, t)) ds = 0$$

Then we have

$$\begin{aligned} \lambda(s) &= (s - t) u_{n+1}(x, t) = u_n(x, t) + \int_0^x (s - t) (D_s^\alpha u(s, t) - u_{tt}(s, t) \\ &\quad - u_t(s, t) - u(s, t) - f(s, t)) ds \end{aligned} \quad (9)$$

with initial approximation  $u_0(x, t) = l_1(t) + x l_2(t)$ . It is obvious that by setting  $\alpha = 2$  the fractional telegraph equation reduces to the classical form equation (1), which is discussed above. Considering Equation (4) we construct the iteration formula for it. Thus we get

$$\begin{aligned} u_{n+1}(x, y, z, t) = & u_n(x, y, z, t) + \int_0^t \lambda(s) (u_{ss}(x, y, z, s) + p(x, y, z, s) u_s(x, y, z, s) \\ & + q(x, y, z, s) u(x, y, z, s) - A(x, y, z, s) u_{xx}(x, y, z, s) - B(x, y, z, s) u_{yy}(x, y, z, s) \\ & - C(x, y, z, s) u_{zz}(x, y, z, s) - f(x, y, z, s)) ds \end{aligned}$$

By taking variation with respect to  $u_n$  we determine the Lagrange multiplier.

$$\begin{aligned} \delta u_{n+1}(x, y, z, t) = & \delta u_n(x, y, z, t) + \delta \int_0^t \lambda(s) (u_{ss}(x, y, z, s) + p(x, y, z, s) u_s(x, y, z, s) \\ & + q(x, y, z, s) \widetilde{u(x, y, z, s)} - A(x, y, z, s) (\widetilde{u_{xx}})(x, y, z, s) \\ & - B(x, y, z, s) (\widetilde{u_{yy}})(x, y, z, s) \\ & - C(x, y, z, s) (\widetilde{u_{zz}})(x, y, z, s) - f(x, y, z, s)) ds = 0 \end{aligned} \quad (10)$$

$$\begin{aligned} \delta u_n(s) : 1 + p\lambda(s) - \lambda'(s)|_{s=t} &= 0 \\ \delta((u_n)_s) : \lambda(s)|_{s=t} &= 0 \\ \delta u_n(s) : [\lambda''(s) - p(x, y, z, s)\lambda'(s)] &= 0 \end{aligned} \quad (11)$$

By solving the above equations in each case we determine the Lagrange multiplier.

#### 4. ILLUSTRATIVE EXAMPLES

We applied the method presented in this paper and solved five test problems to show the efficiency of the new approach.

##### 4.1. Example 1

Let

$$\alpha = \beta = 1, \quad f(x, t) = 0, \quad \phi = e^x, \quad \psi = -e^x, \quad 0 \leq x \leq 4$$

In Equation (1) then we have

$$\begin{aligned} u_{tt} + u_t + u &= u_{xx} \\ u(x, 0) &= e^x, \quad u_t(x, 0) = -e^x \end{aligned}$$

Now we have the iteration formula from (7) as

$$\begin{aligned} u_{n+1}(x, t) &= u_n(x, t) + \int_0^t (e^{(s-t)} - 1) ((u_n(x, s))_{ss} + (u_n(x, s))_s + u_n(x, s) - ((u_n(x, s))_{xx})) ds \\ u_0(x, t) &= (1-t)e^x \\ u_1(x, t) &= (1-t)e^x + \int_0^t (e^{(s-t)} - 1) (0 - e^x + (1-s)e^x - (1-s)e^x) ds = e^{(x-t)} \end{aligned}$$

which is the exact solution of the given equation.

#### 4.2. Example 2

Let

$$\alpha = 6, \quad \beta = 2, \quad f(x, t) = (2 - \alpha + \beta) e^{-t} \sin(x)$$

$$\phi = \sin(x), \quad \psi = -\sin(x), \quad 0 \leq x \leq \pi$$

In Equation (1) then we have

$$u_{tt} + 6u_t + 2u = u_{xx} - 2e^{-t} \sin(x)$$

$$u(x, 0) = \sin(x), \quad u_t(x, 0) = -\sin(x)$$

In addition we have the iteration formula from (6) that can be written as

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t \frac{1}{6} (e^{6(s-t)} - 1) ((u_n(x, s))_{ss} + 6(u_n(x, s))_s + 2u_n(x, s) - ((u_n(x, s))_{xx}) + 2e^{-s} \sin(x)) ds$$

$$u_0(x, t) = (1 - t) \sin(x)$$

$$u_1(x, t) = (1 - t) \sin(x) + \frac{1}{360} (e^{-6t} + 144e^{-t} + 5(-29 + 6t(5 + 3t))) \sin(x)$$

and so on. The exact solution  $u(x, t) = e^{-t} \sin(x)$  and also  $u_1(x, t)$  are plotted in Figure 1.

#### 4.3. Example 3

Considering Equation (2) with

$$a = 1, \quad b = -1, \quad c = 1, \quad d = -1, \quad f(x) = g(x) = e^x, \quad r(x) = 1$$

$$p(x) = x^2, \quad q(x) = x, \quad F(x, t) = (1 - x - x^2) e^{x+t}$$

Then we have

$$u_{tt} - x^2 u_{xx} - 2x u_x + x u = (1 - x - x^2) e^{t+x}$$

and using (8) we get the following iteration formula:

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t (s - t) ((u_n(x, s))_{ss} - x^2 (u_n(x, s))_{xx} - 2x (u_n(x, s))_x + x u_n(x, s) - (1 - x - x^2) e^{t+x}) ds$$




Figure 1. Exact and approximate solutions.




Figure 2. Exact and approximate solutions.

We start with

$$u_0(x, t) = e^x(1+t)$$

and by the above iteration formula we will have

$$\begin{aligned} u_1(x, t) &= e^x(1+t) + \left(\frac{1}{6}\right)e^x(t^2x(1+x) - 6e^t(-1+x+x^2) + 6(t+1)(-1+x+x^2)) \\ u_2(x, t) &= u_1(x, t) + 1/120e^x x(-120e^t(1+x(2+x)(4+x)) + 120(1+t)(1+x(2+x)(4+x)) \\ &\quad + t^2(5t^2(2+x)(1+x(4+x)) + t^3(2+x)(1+x(4+x)) \\ &\quad + 60(1+x(2+x)(4+x)) + 20t(1+x(2+x)(4+x)))) \end{aligned}$$

and so on. In Figure 2, the exact solution  $u(x, t) = e^{(x+t)}$  and the approximate solution  $u_2(x, t)$  are plotted.

#### 4.4. Example 4

Consider Equation (3) with

$$\alpha = 3/2, \quad l_1(t) = l_2(t) = e^{-t}, \quad L(x) = e^x, \quad f(x, t) = 0$$

Then we have

$$\begin{aligned} D_x^\alpha u(x, t) &= u_{tt}(x, t) + u_t(x, t) + u(x, t) \\ u(0, t) &= e^{-t}, \quad t \geq 0 \\ u_x(0, t) &= e^{-t}, \quad t \geq 0 \\ u(x, 0) &= e^x, \quad 0 < x < 1 \end{aligned}$$

With this assumption and using (9) the following iteration formula will be obtained:

$$\begin{aligned} u_{n+1}(x, t) &= u_n(x, t) + \int_0^x (s-x)(D_s^\alpha u(s, t) - u_{tt}(x, t) - u_t(x, t) - u(x, t)) ds \\ u_0(x, t) &= e^{-t}(1+x) \end{aligned}$$



$$u_1(x, t) = e^{-t}(1+x) + 1/6e^{-t}x^2(3+x)$$

$$u_2(x, t) = e^{-t}(1+x) + 1/6e^{-t}x^2(3+x) + (e^{-t}x^2(-64\sqrt{x}(7+2x) + 7\sqrt{\pi}(60+20x+5x^2+x^3)))/(840\sqrt{\pi})$$

and so on. The approximate solution  $u_3(x, t)$  and the solution obtained by the Adomian decomposition method proposed in [6]

$$u(x, t) = e^{-t}(1+x) + e^{-t}((4x^{3/2})/(3\sqrt{\pi}) + (8x^{5/2})/(15\sqrt{\pi}) + x^3/6 + x^4/24)$$

are plotted in Figure 3.

#### 4.5. Example 5

Consider Equation (4) with

$$p, q = 1, \quad A = x, \quad B = y, \quad C = z, \quad f(x, y, z, t) = e^{(x+y+z+t)}(3-x-y-z)$$

$$\phi(x, y, z) = e^{(x+y+z)}, \quad \psi(x, y, z) = e^{(x+y+z)}, \quad T = 1$$

$$u_{tt} + u_t + u = xu_{xx} + yu_{yy} + zu_{zz} + e^{(x+y+z+t)}(3-x-y-z)$$

$$u(x, y, z, 0) = e^{(x+y+z)}, \quad u_t(x, y, z, 0) = e^{(x+y+z)}$$

$$u(0, y, z, t) = e^{(y+z)}, \quad u(x, 0, z, t) = e^{(x+z)}, \quad u(x, y, 0, t) = e^{(x+y)}$$

By using Equations (10)–(11), we get  $\lambda(s) = e^{(s-t)} - 1$  and we have the iteration formula:

$$\begin{aligned} u_{n+1}(x, y, z, t) = & u_n(x, y, z, t) + \int_0^t (e^{(s-t)} - 1)(u_{ss}(x, y, z, s) \\ & + u_s(x, y, z, s) + u(x, y, z, s) - xu_{xx}(x, y, z, s) \\ & - yu_{yy}(x, y, z, s) - zu_{zz}(x, y, z, s) - e^{(x+y+z+s)}(3-x-y-z)) ds \end{aligned}$$




Figure 3. Exact and approximate solutions.




Figure 4. Exact and approximate solutions for  $x = y = z = 0.25$  and  $0 \leq t \leq 1$ .




Figure 5. Exact and approximate solutions for  $x = y = z = 0.5$  and  $0 \leq t \leq 1$ .




Figure 6. Exact and approximate solutions for  $x = y = z = 0.75$  and  $0 \leq t \leq 1$ .

We start with initial approximation

$$u_0(x, y, z, t) = (1+t)e^{(x+y+z)}$$

The approximate solution  $u_2(x, y, z, t)$  and the exact solution  $u(x, y, z, t) = e^{(x+y+z+t)}$  for various values of  $x$ ,  $y$  and  $z$  are plotted in Figures 4–7.




Figure 7. Exact and approximate solutions for  $x = y = z = 1$  and  $0 \leq t \leq 1$ .

## 5. CONCLUSION

In this paper the variational iteration method was used for solving the linear, variable coefficient, fractional derivative telegraph equations and also for the three-space-dimensional telegraph equation. We described the method and used it in some test examples in order to show its applicability and validity in comparison with exact and other numerical solutions. We achieved accurate approximations by using only a few number of iterations, which reveals efficiency of the new method.

## ACKNOWLEDGEMENTS

The authors are very grateful to both reviewers for carefully reading this paper.

## REFERENCES

1. Weston VH, He S. Wave splitting of the telegraph equation in  $R^3$  and its application to inverse scattering. *Inverse Problems* 1993; **9**:789–812.
2. Jordan PM, Puri A. Digital signal propagation in dispersive media. *Journal of Applied Physics* 1999; **85**(3): 1273–1282.
3. Banasiak J, Mika JR. Singular perturbed telegraph equations with applications in the random walk theory. *Journal of Applied Mathematics and Stochastic Analysis* 1998; **11**:9–28.
4. Gao F, Chi C. Unconditionally stable difference schemes for a one-space-dimensional linear hyperbolic equation. *International Journal of Applied Mathematics and Computation* 2007; **187**:1272–1276.
5. Aloy R, Cassaban MC, Caudillo-Mata LA, Jodar L. Computing the variable coefficient telegraph equation using a discrete eigenfunctions method. *Computers and Mathematics with Applications* 2007; **54**:448–458.
6. Momani S. Analytical and approximate solutions of the space-and-time-fractional telegraph equations. *Applied Mathematics and Computation* 2005; **170**:1126–1134.
7. Momani S, Odibat Z. Analytical approach to linear fractional partial differential equations arising in fluid mechanics. *Physics Letters, A* 2006; **355**:271–279.
8. Mohanty RK. An operator splitting technique for an unconditionally stable difference method for a linear three space dimensional hyperbolic equation with variable coefficient. *Applied Mathematics and Computation* 2005; **162**:549–557.
9. Dehghan M, Mohebbi A. High order implicit collocation method for the solution of two-dimensional linear hyperbolic equation. *Numerical Methods for Partial Differential Equations* 2009; **25**:232–243.
10. Jafari H, Dehghan M, Sayevand Kh. Solving a fourth-order fractional diffusion-wave equation in a bounded domain by decomposition method. *Numerical Methods for Partial Differential Equations* 2008; **24**:1115–1126.
11. Dehghan M, Shakourifar M, Hamidi A. The solution of linear and nonlinear systems of Volterra functional equations using Adomian–Pade technique. *Chaos, Solitons and Fractals* 2009; **39**:2509–2521.
12. Dehghan M, Lakestani M. The use of Chebyshev cardinal functions for solution of the second-order one-dimensional telegraph equation. *Numerical Methods for Partial Differential Equations* 2009; **25**:931–938.

13. Dehghan M, Shokri A. A numerical method for solving the hyperbolic telegraph equation. *Numerical Methods for Partial Differential Equations* 2008; **24**:1080–1093.
14. Mohebbi A, Dehghan M. High order compact solution of the one-space-dimensional linear hyperbolic equation. *Numerical Methods for Partial Differential Equations* 2008; **24**:1222–1235.
15. Saadatmandi A, Dehghan M. Numerical solution of hyperbolic telegraph equation using the Chebyshev tau method. *Numerical Methods for Partial Differential Equations* 2009; DOI: 10.1002/num.
16. He JH. Variational iteration method—a kind of nonlinear analytical technique: some examples. *International Journal of Non-Linear Mechanics* 1999; **34**:699–708.
17. Inokuti M, Sekine H, Mura T. General use of the Lagrange multiplier in nonlinear mathematical physics. In *Variational Method in the Mechanics of Solids*, Nemat-Nasser S (ed.). Pergamon Press: Oxford, 1978; 156–162.
18. He JH. Variational iteration method for delay differential equations. *Communications in Nonlinear Science and Numerical Simulation* 1997; **2**(4):235–236.
19. He JH. Approximate analytical solution for seepage flow with fractional derivatives in porous media. *Computational Methods in Applied Mechanical Engineering* 1998; **167**:57–68.
20. Odibat Z, Momani S. Numerical approach to differential equations of fractional order. *Journal of Computational and Applied Mathematics* 2007; **207**:96–110.
21. He JH, Wu XH. Construction of solitary solution and compaction-like solution by variational iteration method. *Chaos, Solitons and Fractals* 2006; **29**:108–113.
22. He JH. Variational iteration method for autonomous ordinary differential systems. *Applied Mathematics and Computation* 2000; **114**:115–123.
23. Momani S, Abuasad S. Application of He's variational iteration method to Helmholtz equation. *Chaos, Solitons and Fractals* 2006; **27**(5):1119–1123.
24. Abdou MA, Soliman AA. Variational iteration method for solving Burgers and coupled Burgers equations. *Journal of Computational and Applied Mathematics* 2005; **181**(2):245–251.
25. Abbasbandy S. Numerical method for non-linear wave and diffusion equations by the variational iteration method. *International Journal for Numerical Methods in Engineering* 2008; **73**:1836–1843.
26. Dehghan M, Shakeri F. Solution of parabolic integro-differential equations arising in heat conduction in materials with memory via He's variational iteration technique. *Communications in Numerical Methods in Engineering* 2008; DOI: 10.1002/cnm.
27. He JH. Approximate solution of nonlinear differential equations with convolution product nonlinearities. *Computational Methods in Applied Mechanical Engineering* 1998; **167**:69–73.
28. He JH. Some asymptotic methods for strongly nonlinear equations. *International Journal of Modern Physics B* 2006; **20**(10):1141–1199.
29. Tatari M, Dehghan M. On the convergence of He's variational iteration method. *Journal of Computational and Applied Mathematics* 2007; **207**:121–128.
30. Dehghan M, Shakeri F. Approximate solution of a differential equation arising in astrophysics using the variational iteration method. *New Astronomy* 2008; **13**:53–59.
31. Tatari M, Dehghan M. Solution of problems in calculus of variations via He's variational iteration method. *Physics Letters A* 2007; **362**:401–406.
32. Dehghan M, Tatari M. Identifying an unknown function in a parabolic equation with overspecified data via He's variational iteration method. *Chaos, Solitons and Fractals* 2008; **36**:157–166.
33. Shakeri F, Dehghan M. Solution of a model describing biological species living together using the variational iteration method. *Mathematical and Computer Modelling* 2008; **48**:685–699.
34. Shakeri F, Dehghan M. Numerical solution of the Klein–Gordon equation via He's variational iteration method. *Nonlinear Dynamics* 2008; **51**:89–97.
35. Dehghan M, Tatari M. The use of He's variational iteration method for solving the Fokker–Planck equation. *Physica Scripta* 2006; **74**:310–316.
36. Dehghan M, Shakeri F. Application of He's variational iteration method for solving the Cauchy reaction–diffusion problem. *Journal of Computational and Applied Mathematics* 2008; **214**:435–446.
37. Dehghan M, Shakeri F. Numerical solution of a biological population model using He's variational iteration method. *Computers and Mathematics with Applications* 2007; **54**:1197–1209.
38. Dehghan M. Finite difference procedures for solving a problem arising in modeling and design of certain optoelectronic devices. *Mathematics and Computers in Simulation* 2006; **71**:16–30.
39. Wazwaz AM. The variational iteration method for a reliable treatment of the linear and the nonlinear Goursat problem. *Applied Mathematics and Computation* 2007; **193**:455–462.
40. Wazwaz AM. A comparison between the variational iteration method and Adomian decomposition method. *Journal of Computational and Applied Mathematics* 2007; **207**:129–136.
41. Wazwaz AM. The variational iteration method for solutions for KdV, K(2,2), Burgers and cubic Boussinesq equations. *Journal of Computational and Applied Mathematics* 2007; **207**:18–23.
42. Abbasbandy S, Shirzadi A. The variational iteration method for a class of eighth-order boundary-value differential equations. *Zeitschrift für Naturforschung* 2008; **63a**:745–751.
43. Abbasbandy S, Shivanian E. Application of the variational iteration method for nonlinear Volterra's integro-differential equations. *Zeitschrift für Naturforschung* 2008; **63a**:538–542.
44. Dehghan M, Saadatmandi A. Variational iteration method for solving the wave equation subject to an integral conservation condition. *Chaos, Solitons and Fractals* 2009; **41**:1448–1453.

45. Tatari M, Dehghan M. Improvement of He's variational iteration method for solving systems of differential equations. *Computers and Mathematics with Applications* 2009; DOI: 10.1016/j.camwa.2009.03.081.3.
46. Saadatmandi A, Dehghan M. Variational iteration method for solving a generalized pantograph equation. *Computers and Mathematics with Applications* 2009; DOI: 10.1016/j.camwa.2009.03.017.
47. Dehghan M, Shakeri F. The use of the decomposition procedure of Adomian for solving a delay differential equation arising in electrodynamics. *Physica Scripta* 2008; **78**:1–11. Article No. 065004.
48. Yousefi SA, Dehghan M. The use of He's variational iteration method for solving variational problems. *International Journal of Computer Mathematics* 2008; in press.
49. Finlayson BA. *The Method of Weighted Residuals and Variational Principles*. Academic Press: New York, 1972.