

MATH EXCELLENCE SERIES

SAT MATH

& Hard Question Mastery

Algebra • Advanced Math • Data Analysis • Geometry & Trigonometry

294 hard problems, deduplicated from three sources

Algebra 43 Advanced Math 117 Data Analysis 48 Geometry & Trigonometry 86

Three problems per page, with working space



DAVLAT KAMOLIDDINOV

Math Instructor





MUNDARIJA
Contents

How to use this book	ii	Problem-Solving and Data Analysis	58
Algebra	2	Geometry and Trigonometry	75
Advanced Math	18		





How to use this book

This book merges two collections of hard Digital SAT mathematics questions into one sequence, with duplicates removed and the problems regrouped by the four tested domains.

Every page carries at most **three** problems. The space left under each one is deliberate: these questions are not meant to be read, they are meant to be worked, and the room to work is part of the page.

The answer key at the back lists every published answer. Where a source did not publish one, the entry is marked —; solve those and fill them in yourself.







1

In the formula $P = \frac{5}{12}K + 30$, if P is increased by 30, what is the increase in K ?

- A) 72
- B) 90
- C) 36
- D) 65

2

$$10x + 7y = 27$$

$$5x + 14y = 45$$

The solution to the given system of equations is $(x, y) = \left(\frac{p}{60}, \frac{z}{124}\right)$, where p and z are integers. What is the value of $z - p$?

3

In a set of four consecutive odd integers, where the integers are ordered from least to greatest, the first integer is represented by x . The product of 12 and the second odd integer in the set is at most the value of 30 more than the sum of the third and fourth odd integers in the set. What is the greatest possible value of x ?




4

The graph of the equation $3x - 2y = 6$ is translated 4 units down and 2 units to the left.

What is the y -coordinate of the y -intercept of the resulting graph?

5

The graph of line g is shown in the xy -plane. Line s is defined by $150x + ny = -m$, where n and m are constants. If line s is graphed in this xy -plane, resulting in the graph of a system of two linear equations, the system of two linear equations will have infinitely many solutions. What is the value of $n + m$?



6

A bakery makes loaves of bread in three sizes: small, medium, and large. On a particular morning, the bakery makes twice as many small loaves as medium loaves, represented by m . They also bake 20 large loaves. If the bakery makes a total of 80 loaves that morning, which equation represents this situation?

- A) $2m + m + 20 = 80$
- B) $2(2m) + m + 20 = 80$
- C) $2m + 20 = 80$
- D) $3m + 20 = 80$



7

A baseball coach charges each athlete \$150 for the first month and \$95 for each additional month. Which of the following functions correctly represents the total revenue $R(m)$, in dollars, for m months and n athletes, where m and n are positive integers?

- A) $R(m) = 95mn + 55$
- B) $R(m) = 95mn + 150$
- C) $R(m) = 95mn + 55n$
- D) $R(m) = 95mn + 150n$

8

$$f(x) = -7x + b$$

For the given function $f(x)$, which of the following gives the equation for the function $f(x) + 20$, given that the point $(-4, 0)$ lies on the graph of $f(x) - 20$?

- A) $f(x) = -7x - 28$
- B) $f(x) = -7x - 8$
- C) $f(x) = -7x + 8$
- D) $f(x) = -7x + 12$

9

One of the two equations in a system of linear equations is given. The system has no solution. Which equation could be the second equation in this system?

$$7x = 105y - 133$$

- A) $x = 15y$
- B) $\frac{1}{7}x = 15y$
- C) $x = 15y - 19$
- D) $\frac{1}{7}x = 15y - 19$



10

What is the value of $\frac{k}{25}$ if the given equation only has one solution?

$$25|x - 4| = k$$

- A) -5 only
- B) 5 only
- C) 5 or -5
- D) 0

11

In the given system of equations, k is a constant. If the system has no solution, what is the value of k ?

$$\begin{aligned} \frac{3}{8}x + \frac{7}{5}y &= \frac{9}{7} - \frac{7}{5}y \\ \frac{5}{4}x + \frac{9}{4} &= ky + \frac{4}{3} \end{aligned}$$

12

One gallon of resin costs \$79 and will cover 48 square feet of a countertop. The countertops in a given room have a total surface area of k square feet. Which equation represents the cost c , in dollars, of resin needed to cover the countertops twice?

- A) $c = \frac{48k}{79}$
- B) $c = \frac{96k}{79}$
- C) $c = 79 \left(\frac{k}{24} \right)$
- D) $c = 79 \left(\frac{k}{96} \right)$



13

For each real number r , which of the following points lies on the graph of each equation in the xy -plane for the given system?

$$10x + 8y = 15$$

$$25x + 20y = 37.5$$

- A) $\left(r, -\frac{4r}{5} + \frac{15}{8}\right)$
- B) $\left(r, \frac{5r}{4} + \frac{4}{5}\right)$
- C) $\left(-\frac{4r}{5} + \frac{3}{2}, r\right)$
- D) $\left(\frac{2r}{5} + 10, -\frac{2r}{5} + 25\right)$

14

For the given system of equations, what is the value of $7(j - k)$?

$$(j - k) - 16(m + l) = 689$$

$$(j - k) + 12(m + l) = 1235$$

15

A 34 pound dog eats two types of canned food: chicken and beef. The recommended amount of the chicken food is 1.25 cans per 16 pounds a dog weighs per day. The recommended amount of canned beef food is 0.85 cans per 23 pounds a dog weighs per day. If c is the number of cans of chicken food and b is the number of cans of beef food a 34 pound dog eats in a given day. Which equation describes all possible values of c and b ?

- A) $\frac{1.25}{16}c + \frac{0.85}{23}b = 34$
- B) $\frac{16}{1.25}c + \frac{23}{0.85}b = 34$
- C) $\frac{1.25}{16}b + \frac{0.85}{23}c = 34$
- D) $\frac{16}{1.25}b + \frac{23}{0.85}c = 34$



16

A small business owner budgets \$3,600 to purchase timber each year. The owner must purchase a minimum of 345 slats of timber each year to maintain the discounted pricing. If the owner pays \$7.25 to purchase each slat of pine pulpwood and \$24.75 to purchase each slat of pine sawtimber, what is the maximum number of slats of pine sawtimber the owner can purchase to stay within the budget and maintain the discounted pricing?

17

$$-377x + k = 377x + k$$

If k is a constant, how many solutions does the given equation have?

- A) No solutions
- B) One solution
- C) Two solutions
- D) Infinite solutions

18

The graph of the linear equation passes through the points $\left(-21, -\frac{89}{2}\right)$ and $\left(12, \frac{65}{2}\right)$. The graph also passes through the point $\left(\frac{27}{4}, a\right)$. What is the value of a ?



19

$$7kx + 13my = 20.5$$

$$6kx + 5my = -48$$

In the given system of equations, k and m are constants. The system has a solution of $(2, y)$. What is the value of k ?

20

Which of the following inequalities gives all possible values of k where the equation below has no solution?

$$4 - 4 \left| \frac{7}{8} + \frac{4}{5}x \right| = \frac{57}{12} + k$$

- A) $k < -\frac{3}{4}$
- B) $k > -\frac{3}{4}$
- C) $k \leq \frac{3}{4}$
- D) $k \geq \frac{3}{4}$

21

If $\frac{2a}{b} = 4.25$ and $\frac{a}{bn} = 8.5$, what is the value of n ?



22

$$7x - 9y = 39$$

$$35x - 45y = 195$$

For any real number r , which of the following points lies on the graph of both equations?

- A) $\left(3r, \frac{7r - 13}{3}\right)$
- B) $\left(\frac{7r + 13}{3}, 3r\right)$
- C) $\left(-\frac{7r}{-9} + 39, \frac{7r}{9} + 195\right)$
- D) $\left(\frac{r}{5} + 39, -\frac{r}{5} + 195\right)$

23

The proposal for a development project was included on a city's election ballots. A local tv station reported that 6 times as many people voted in favor of the proposal as people who voted against it. If 1,770 more people voted in favor of the proposal than those who voted against the proposal. How many people voted against the proposal?

24

Which system of linear equations has no solution?

- A) $-3x + 7y = -9$ and $3x - 7y = 9$
- B) $3x - 7y = 5$ and $2x + 8y = 6$
- C) $4x - 9y = 7$ and $-12x + 27y = -21$
- D) $-3x + 8y = 20$ and $6x - 16y = 40$



25

If $\frac{x-7}{5} = \frac{x-4}{3}$, the value $x-4$ is between which of the following pairs of values?

- A) -9 and -4
- B) -1 and 1
- C) 2 and 5
- D) 7 and 11

26

A handyman charges a flat rate of \$218 for the first 3 hours of work and \$68 for each additional hour of work. Which equation gives the total amount y , in dollars, that the handyman charges for x hours of work, where $x > 3$?

- A) $y = 68x + 218$
- B) $y = 218x + 68$
- C) $y = 68x + 14$
- D) $y = 68x + 422$

27

Megan engages in up to 3 types of exercise each week for a total of 14 hours while training for an ironman competition. Megan runs the same number of minutes each week. The equation $y = 780 - x - 200$ represents the situation where Megan swims for x minutes during a week and bikes for any remaining training time y , in minutes. If this equation is graphed in the xy -plane, which of the following statements is true?

- A) During a week when Megan runs for 780 minutes, she bikes for 200 minutes.
- B) Each week, Megan swims and bikes for a total of 780 minutes.
- C) During a week when Megan doesn't swim, she runs for 260 minutes.
- D) During a week when Megan doesn't bike, she swims for 390 minutes.



28

$$92x - 1,789 = kx - 1,790$$

In the given equation, k is a positive integer constant greater than 91. The equation has exactly one solution. What is the least possible value of k ?

29

Lines c and d are perpendicular. If the equation of line c is $y = mx + b$ and the lines intersect at point $(7, 6)$. Which point lies on line b ?

- A) $(7 + m, 7)$
- B) $(2m, 8)$
- C) $(7 + 2m, 4)$
- D) $(7 - 3m, 3)$

30

What is the value of k if the given equation only has one solution?

$$\frac{1}{4}|x - 4| - 30 = 2k$$



31

In the given pair of equations a and b are constants. The graph of this pair of equations in the xy -plane is a pair of perpendicular lines. Which of the following pairs of equations also represents a pair of perpendicular lines?

$$13x + 17y = 10$$

$$ax + by = 10$$

- A) $39x + 17y = 10$ and $ax - 3by = 10$
 B) $52x + 34y = 10$ and $2ax + 4by = 10$
 C) $65x + 17y = 10$ and $5ax + by = 10$
 D) $13x - 17y = 10$ and $ax + by = 10$

32

$$77(x - k) = 77y + 77k$$

One of the equations in a system of two linear equations is given, where k is a positive constant. The system has no solution. Which equation could be the second equation in this system?

- A) $11x + 11y = 11k$
 B) $11x + 11y = 22k$
 C) $11x - 11y = 11k$
 D) $11x - 11y = 22k$

33

Line a is defined by $7x + 8y = 96$. Line a is parallel to line b . An equation of line b is $42x = ky - 96$, where k is a constant. What is the y -intercept of line b ?



34

A bakery makes 9-inch, 12-inch, and 14-inch pies. During a certain day, the number of 14-inch pies the bakery makes is 4 times the number n of 12-inch pies, and the number of 9-inch pies is 48. During this day, the bakery makes 190 pies in total. Which equation represents this situation?

- A) $9(48) + 12n + 14(4n) = 190$
- B) $9n + 12n + 14n = 190$
- C) $4n + 48 = 190$
- D) $5n + 48 = 190$

35

If $x \geq -13$ represents all solutions to the inequality $ax - 75 \leq 42$, where a is a constant, what is the greatest possible value of a ?

36

$$\frac{14x + 35}{4} - \frac{b}{12} = a(x - 7)$$

In the given equation, a and b are constants. If the equation has infinitely many solutions, what is the value of b ?



37

$$9x + 5y = 7$$

$$45x + 25y = 35$$

For each real number r , which of the following points lies on the graph of each equation in the xy -plane for the given system?

A) $\left(\frac{r}{5} + 7, -\frac{r}{5} + 35\right)$

B) $\left(-\frac{5r}{9} + \frac{7}{9}, r\right)$

C) $\left(r, \frac{9r}{5} + \frac{7}{5}\right)$

D) $\left(r, -\frac{5r}{9} + \frac{7}{9}\right)$

38

$$\frac{1}{7}x + \frac{3k}{13}y = \frac{9}{14}$$

One of the equations in a system of two linear equations is given, where k is a nonzero constant. The system has no solution. If the other equation in the system is graphed in the xy -plane, what is the slope of the graph?

A) $\frac{13k}{21}$

B) $-\frac{13}{21k}$

C) $\frac{21k}{13}$

D) $-\frac{21}{13k}$

39

In a set of three consecutive odd integers, where the integers are ordered from least to greatest, the first integer is represented by k . The product of 17 and the second odd integer in the set is at most the value of 40 less than the sum of the first and third odd integers in the set. What is the greatest possible value of k ?



40

For the linear function k , $k(a) = -24$, where a is a constant, $k(4) = 30$, and the slope of the graph $y = k(x)$ in the xy -plane is 7. For the linear function p , $p(a) = 26$ and $p(6) = 38$. What is the slope of the graph of $y = p(x)$ in the xy -plane?


41

Line a in the xy -plane passes through the points $\left(\frac{7}{k}, -\frac{3}{k} + \frac{9}{15}\right)$ and $\left(\frac{5}{k}, -\frac{7}{k} + \frac{9}{15}\right)$, where k is a constant. Which of the following defines the y value of the y -intercept of line a ?

- A) $-\frac{38}{k} + \frac{3}{5}$
- B) $-\frac{32}{k} + \frac{3}{5}$
- C) $-\frac{17}{k} + \frac{3}{5}$
- D) $-\frac{70}{k} + \frac{3}{5}$

42

The graph of the line a is shown in the xy -plane. Line b is defined by $178x + ky = j$, where k and j are constants. If line b is graphed in this xy -plane, resulting in a graph of a system of two linear equations, the system of two linear equations will have infinitely many solutions. What is the value of $k + j$?






43

$$\frac{3}{5}x + \frac{7}{4}y = a$$

$$\frac{7}{11}x + by = c$$

In the given system of equations, a , b , and c are constants. The system has infinitely many solutions. What is the value of $\frac{a}{c}$?







44

The function p is defined by $p(x) = a((x+5)^2 - b)((x+5)^2 - c)$, where a , b , and c are constants. In the xy -plane, the graph of $y = p(x)$ passes through the points $(-4, 34)$ and $(0, 150)$. What is the value of $p(-10) + p(-4)$?

45

In the xy -plane, the graph of function f , where $y = f(x)$, has exactly 8 x -intercepts. One of these x -intercepts is $(17, 0)$. The rational function g is defined by $g(x) = \frac{f(x)}{x - 17}$. In the xy -plane, how many x -intercepts does the graph of $y = g(x)$ have?

46

The function f is defined by $f(x) = (x - 7)(x - 4)(x + k)$, where k is a constant. In the xy -plane, the graph of $y = f(x)$ passes through the point $(-2, 0)$. What is the value of $f(0)$?

- A) -2
- B) 56
- C) 7
- D) -17



47

$x^2 + 7x + 6$ has two solutions r and s , $r < s$, and $x^2 + 6x - 7$ has two solutions t and u , $t < u$. If $x^2 + cx + d$ has two solutions $t + r$ and $u + s$, what is the value of c ?

48

$$x(kx - 20) = -1$$

In the given equation, k is an integer constant. If the equation has two distinct real solutions, what is the greatest possible value of k ?

49

$$f(x) = 9^x + 4$$

$$g(x) = 7x + b$$

The given system of equations intersects at points (e, f) and (g, h) , where $h < f$. What is the least possible value of b ?

- A) 5
- B) 3
- C) 9
- D) 4



50

In the xy -plane, a parabola has vertex $(-2, -1)$ and intersects the x -axis at two points. If the equation of the parabola is written in the form $y = ax^2 + bx + c$, where a , b , and c are constants, which of the following could be the value of $2a + b + 3c$?

- A) -2
- B) -18
- C) -3
- D) -7

51

The function f is defined by $f(x) = a^x + b$, where a and b are constants. In the xy -plane, the graph of $y = f(x)$ has an x -intercept at $(2, 0)$ and a y -intercept at $(0, -195)$. What is the value of $a + b$?

52

The function f is defined by $f(x) = ax^2 + bx + c$, where a , b , and c are constants and $1 < a < 4$. The graph of $y = f(x)$ in the xy -plane passes through the points $(-10, 0)$ and $(-5, 0)$. If a is an integer, what could be the value of $a + b$?



53

$$f(x) = 5x^2 + 20x + 148$$

The function g is defined by $g(x) = f(x + 4)$. What is the minimum value of $g(x)$?

54

$(x + 4)$ is a factor of the quadratic function $y = f(x)$, and $x = c$ is a positive zero of the function. If the minimum of the function is $\left(\frac{3}{4}, k\right)$, what is the value of c ?

55

$$f(x) = -(x - a)(x - b)$$

The function f is defined by the given equation, where a and b are integer constants. If $f(-30) < 0$, $f(-10) > 0$ and $f(20) < 0$ and $a < b$, what is the minimum possible value of $a - b$?



56

The function k is defined by $k(s) = \sqrt{s + 84}$. If $k(5p) = p$, where p is a constant, what is the value of p ?

57

The expression $6x^2 + bx - 35$, where b is a constant, can be rewritten as $(hx + k)(x + j)$, where h , k , and j are integer constants. Which of the following must be an integer?

- A) $\frac{b}{h}$
- B) $\frac{b}{j}$
- C) $\frac{35}{h}$
- D) $\frac{35}{j}$

58

In the xy -plane, a line with the equation $y = mx - 30$ does not intersect with a parabola with the equation $y = 2x^2 - 3x - m$, where m is a constant. Which of the following is a possible value for m ?

- A) only -7
- B) -7 and 7
- C) -27
- D) 25



59

$$(x + 4)^2 + (x^2 + 4x) = (x + 4)(x - 8)$$

In the given equation, $x \neq -4$. What is the solution to the given equation?

60

In the given equation, m is a positive constant. The product of the solutions to the equation is 28. What is the value of m ?

$$\frac{2}{3}(3x + 7)(x + \sqrt{2m + 6})(x - \sqrt{2m + 6}) = 0$$

61

Function f is defined by $f(x) = 4x^2 - 12x + 5$ on the xy -plane, where $f(x) = y$. Function g intersects the x -axis twice when graphed on the xy -plane. Which of the following equations could define $g(x)$, if k represents a positive constant?

- A) $g(x) = f(x - k)$
- B) $g(x) = f(x + k)$
- C) $g(x) = f(x) + k$
- D) $g(x) = f(x) - k$



62

The function f is defined by $f(x) = x^2 + bx + c$, where b and c are constants. If the x -intercepts of the graph of f in the xy -plane are -2 and 5 , what is the value of b ?

- A) -3
- B) -2
- C) -5
- D) 2

63

The equation $x^2 + mx + 5 = 0$, where m is a positive integer, has no real solutions. What is the greatest possible value of m ?

64

Given the function $g(x) = 33(k)^x$, where k is a positive constant. For each increase of 1 in the value of x , the value of $g(x)$ increases by $p\%$, where $0 < p < 200$. Which expression represents p in terms of k ?

- A) $100k$
- B) $100(k - 1)$
- C) $k + 100$
- D) $100(k + 1)$



65

For a quadratic function $f(x) = x^2 + bx + c$, the value $f(2) = -1$ is the vertex of the function. What is the value of c ?

66

The function p is defined by $p(x) = c(2.5^x + 2.5^d)$, where c and d are positive integer constants and $c < d$. The functions q and r are equivalent to function p , where s and t are constants. Which of the following equations displays the y -coordinate of the y -intercept of the graph of $y = p(x)$ in the xy -plane as a constant or coefficient?

- I. $q(x) = c(2.5^x + s)$
 - II. $r(x) = c(2.5)^x + t$
- A) I only
 - B) II only
 - C) I and II
 - D) Neither I nor II

67

The function V models the value, in dollars, of a rare coin t years after 1990:

$$V(t) = 500(1.06)^{\frac{t}{3}}$$

According to the model, the value of the coin is predicted to increase by 6% every n years. What is the value of n ?

- A) 1
- B) 3
- C) 6
- D) 18



68

The function h is defined by $h(x) = (x + 1)(x - 4)(x - 7)^2$. The value of $h(5 + m)$ is 0, where m is a constant. What is the sum of all possible values of m ?

69

$$3x^2 + 6ax + 12 = 0$$

The equation has no real solutions if $a^2 < j$. What is the greatest possible integer value of j ?

70

$$14x^2 + (14m + 3n)x + 3mn = 0$$

In the function, m and n are positive constants. The product of the solutions to the given equation is pmn , where p is a constant. What is the value of p ?



71

The function $f(x)$ is defined by $f(x) = m((x-3)^2 - n)((x-3)^2 - p)$, where m , n , and p are constants. In the xy -plane, the graph of $y = f(x)$ passes through the points $(2, 12)$ and $(6, 108)$. What is the value of $f(1) + f(5)$?

72

$(x + 7)$ is a factor of the quadratic function $y = f(x)$, and $x = c$ is a positive zero of the function. If the minimum of the function is $\left(\frac{63}{8}, k\right)$, what is the value of c ?

73

$$y = -3x^2 - 58$$

$$y = px - 46$$

In the given system of equations, p is a constant. The graphs of the equations in the given system intersect at exactly one point, (x, y) in the xy -plane. Which of the following could be the value of x ?

- A) -2
- B) -4
- C) 12
- D) 16



74

$$f(x) = a^x - b$$

The exponential function given above passes through points $(c, 8)$ and $(2c, 350)$. What is a possible value of b ?

75

For the function f , for every increase of $\frac{1}{2}$ in the value of x , the value of $f(x)$ increases by a factor of c , where c is a constant. Which of the following forms of function f displays the value of c as the base or coefficient?

- A) $f(x) = 56(2)^{6x}$
- B) $f(x) = 56(8)^{2x}$
- C) $f(x) = 56(64)^x$
- D) $f(x) = 56(4,096)^{\frac{x}{2}}$

76

The quadratic function $ax^2 + 200x + c$ has at least one real solution. What is the greatest possible value of ac ?



77

The area of a triangle is equal to x^2 square inches. The base of the triangle is $5 + 2x$ inches, and the height of the triangle is $x - 2$ inches. What is the value of x ?

- A) 2.5
- B) 2.7
- C) 5
- D) 10

78

The function f is defined by $f(x) = a^x + b$, where a and b are constants. In the xy -plane, the graph of $y = f(x)$ has an x -intercept at $(3, 0)$ and a y -intercept at $(0, -342)$. What is the value of $a + b$?

79

Two numbers a and b , are each greater than zero, and the fifth root of a is equal to the cube root of b . For what value of x is a^{8x-4} equal to b ?



80

A model initially estimates that there are 24,000 bacteria on a petri dish. 14 days later, the model estimates that there are 96,000 bacteria on the petri dish. Assuming exponential growth, the formula $B = a(2)^{xd}$ gives the estimated number of bacteria on a petri dish, where a and x are constants and B is the number of bacteria on the petri dish d days after the initial measurement. What is the value of x ?

- A) $\frac{1}{14}$
- B) $\frac{1}{7}$
- C) 7
- D) 14

81

If $a > 0$, $x^2 + y^2 = a$ and $xy = a - 10$, what is $(x + y)^2$ in terms of a ?

- A) $a - 20$
- B) $2a - 10$
- C) $2a - 20$
- D) $3a - 20$

82

$f(x) = 200(b)^x$. When x increases by 1, $f(x)$ increases by $c\%$. Which of the following expresses c in terms of b ?

- A) $c = \frac{200b}{100}$
- B) $c = \frac{2b}{100}$
- C) $c = (b + 1)100$
- D) $c = (b - 1)100$



83

A quadratic function models the height, in feet, of an object above the ground in terms of time, in seconds, after the object is launched. The model indicates the object launched from the ground (0 feet) and reaches its maximum height of 576 feet above the ground 12 seconds after being launched. Based on the model, what is the height, in feet, of the object above the ground 16 seconds after being launched?

- A) 384
- B) 432
- C) 480
- D) 512

84

A quadratic function models the height, in feet, of an object above the ground in terms of time, in seconds, after the object is launched off an elevated surface. The model indicates that at a time of 9 seconds, the object is 294 feet above the ground. At a time of 13 seconds, the object is 310 feet above the ground. If the object was at a height of 24 feet when it was launched, what is the height, in feet, of the object above ground 18 seconds after being launched?

- A) 96
- B) 192
- C) 222
- D) 240

85

$n^{\frac{5}{3}} = k^{\frac{2}{5}}$ and $n^{4a+1} = k$, what is the value of a ?



86

$3x^2 + bx - 112 = (ax + m)(x - l)$. Given that a , m , and l are integers, which of the following must be true?

- A) m is a factor of b
- B) a is a factor of 112
- C) a is a factor of b
- D) l is a factor of 112

87

The vertex of the function $f(x)$ is given as $(9, -13)$, and the point $(3, -4)$ lies on the parabola. If $g(x) = 4f(x-2)$, what is the value of $g(0) - f(0)$?

88

$$77x^2 + (7j + 11k)x + jk = 0$$

In the given equation, j and k are positive constants. The product of the solutions to the given equation is ajk , where a is a constant. What is the value of a ?



89

The equation below relates distinct positive real numbers a , b and c . Which equation correctly expresses c in terms of a and b ?

$$a = 19b \cdot \sqrt[5]{\left(\frac{c}{20}\right)^4}$$

- A) $\frac{20}{19b}(a)^{\frac{5}{4}}$
- B) $\frac{20}{19b}(a)^{\frac{4}{5}}$
- C) $20\left(\frac{a}{19b}\right)^{\frac{5}{4}}$
- D) $20\left(\frac{a}{19b}\right)^{\frac{4}{5}}$

90

For the exponential function f , the value of $f(2)$ is k , where k is a constant. Which of the following equivalent forms of the function f shows the value of k as the coefficient or the base?

- A) $f(x) = 729(3)^{x+2}$
- B) $f(x) = 6,561(3)^x$
- C) $f(x) = 177,147(3)^{x-3}$
- D) $f(x) = 59,049(3)^{x-2}$

91

Two numbers r and s are each greater than zero, and the fourth root of r is equal to the seventh root of s . For what value of x is r^{4x-5} equal to s ?



92

The given equation $ax^2 + 98x + c$ has at least 1 real root and a factor of $kx + j$. What is the greatest possible value of ac ?

93

The table gives two values of x and their corresponding values of $g(x)$, where $g(x) = \frac{f(x) + 5}{x + 8}$ and f is a linear function. What is the y -coordinate of the y -intercept of the graph $y = f(x)$ in the xy -plane?

x	$g(x)$
4	9
2	0

94

The functions f and g are defined by the given equations, where $x \geq 0$. Which of the following equations displays, as a constant or coefficient, the maximum value of the function it defines, where $x \geq 0$?

I. $f(x) = 232(0.4)^{x+2}$

II. $g(x) = 232(0.4)(0.4)(0.4)^{x-2}$

A) I only

B) II only

C) I and II

D) Neither I nor II



95

The function f is defined by $f(x) = 3^x$. The function g is an increasing linear function. In the xy -plane, the graphs of $y = f(x)$ and $y = g(x)$ intersect at two points (a, j) and (b, k) , where $j < k$. When $g(x) > f(x)$, which of the following must be true?

- A) $x > k$
- B) $x < j$
- C) $x < j$ or $x > b$
- D) $a < x < b$

96

An acceptable noise criterion rating for the background noise in a school classroom is 30. For a noise criterion rating of 30, the equation $y = 19(0.978)^{x-50} + 37$ gives the estimated sound pressure level, y in decibels, as a function of the octave band center frequency, x , in hertz, where $x \geq 50$. Which of the following is the best interpretation of 37 in this context?

- A) 37 is 19 less than the estimated sound pressure level, in decibels, at an octave band frequency of 50.
- B) 37 is 19 less than the estimated sound pressure level, in decibels, at an octave band frequency of 0.
- C) 37 is the estimated sound pressure level, in decibels, at an octave band frequency of 50.
- D) 37 is the estimated sound pressure level, in decibels, at an octave band frequency of 0.

97

The graph of the quadratic function $y = f(x)$ in the xy -plane intersects the x -axis when $x = 74$ and $x = k$, where k is a constant. The maximum value of $y = f(x)$ occurs at the point $(17, m)$, where m is a constant. What is the value of k ?



98

Three points on the quadratic function $f(x)$ are given in the table. If $g(x) = f(x + 4)$, what is the y -intercept of the function $g(x)$?

x	$f(x)$
-9	155
-3	227
3	155

99

The given expression can be rewritten as $\frac{a}{9}x^2 + \frac{b}{9}x + \frac{c}{9}$, where a , b , and c are constants. What is the value of $a + b + c$?

$$-19(6x - 4)^2 + 6(6x - 3)^2$$

100

$$f(x) = x^2 + 5x - 6$$

$$g(x) = x^2 + 2x - 35$$

The quadratic function $f(x)$ has 2 solutions j and k , where $j < k$. The quadratic function $g(x)$ has 2 solutions l and m , where $l < m$. The quadratic function $h(x) = x^2 + 7x + c$ has solutions of $k + l$ and $m + j$, and can be rewritten as $(x + a)(x + b)$, what is the value of c ?



101

For the exponential function $f(x) = 16^{x+4}$, which of the following answer choices expresses the minimum value as either a constant or a coefficient when $x \geq 0$?

- A) $f(x) = 64(16)^x$
- B) $f(x) = 65,536(16)^x$
- C) $f(x) = 1,048,576(16)^x$
- D) $f(x) = \frac{1}{65,356}(16)^x$

102

For the given parabola $ax^2 + bx + c$, a , b , and c are constants. The x -intercepts of the parabola are at $(14, 0)$ and $(k, 0)$. If $f(27) = f(11)$, what is the value of k ?

103

A quadratic function models a projectile's height, in meters, above the ground in terms of in terms of time, in seconds, after it was launched. The model estimates that the projectile was launched from an initial height of a meters and reached a maximum height of 134 meters 6 seconds after it was launched. How many seconds after launch does the model estimate it will return to a height of a meters?

- A) 6
- B) 8
- C) 9
- D) 12



104

A certain bacterial colony currently has 5,000 spores. If the number of spores in this colony quadruples every 33 hours, which of the following functions N gives the number of spores in the colony d days from now?

- A) $N(d) = 5,000(4)^{\frac{11}{8}d}$
- B) $N(d) = 5,000(4)^{\frac{8}{11}d}$
- C) $N(d) = 5,000(4)^{\frac{5}{4}d}$
- D) $N(d) = 5,000(4)^{\frac{4}{5}d}$

105

In the given equations, a and b are constants, $a > 1$ and $b > 1$. What is the value of x ?

$$\sqrt[7]{a^5} = \sqrt{b^5}$$

$$a^{3x-3} = b^3$$

106

In the given system of equations, c is a constant. The system has two distinct real solutions. Which of the following could be the value of c ?

$$y = x - c$$

$$y = -6(x - 5)^2$$

- A) 2
- B) 4
- C) $\frac{119}{24}$
- D) 7



107

$$f(x) = (x + a)(x + b)$$

For the given function $f(x)$, where a and b are integer constants, $f(-21) > 0$, $f(-15) > 0$, and $f(-18) < 0$, what is one possible value of $a + b$?

108

Function f is a quadratic function. The graph of $y = f(x)$ in the xy -plane has a vertex at $(-7, 14)$ and passes through the point $(-4, -31)$ and has a y -intercept at $(0, a)$. The graph of $3f(x)$ has a y -intercept at $(0, b)$. What is the positive difference between a and b ?

109

In the given equation, k is a positive constant. The product of the solutions to the equation is 154. What is the value of k ?

$$\frac{3}{5}(5x + 11)(x + \sqrt{5k + 10})(x - \sqrt{5k + 10}) = 0$$



110

$$\sqrt{a - x} = 46 - x$$

In the given equation, a is a constant. The equation has exactly one real solution. What is the minimum possible value of $4a$?

111

For a certain circuit, its power P , in watts; current C , in amperes; voltage V , in volts; and resistance R , in ohms, are related as $\frac{CV^3}{P} = \sqrt{PR}$, where P , C , V , and R are positive. When $R = 34$, which equation correctly expresses P in terms of C and V ?

- A) $P = \frac{CV^3}{\sqrt{34}}$
 B) $P = \frac{CV^3}{\sqrt{34P}}$
 C) $P = \sqrt[3]{\frac{34}{C^2V^4}}$
 D) $P = \sqrt[3]{\frac{C^2V^6}{34}}$

112

$$f(x) = x + 7$$

$$g(x) = 5x^2 - kx + 245$$

The functions f and g are given. In function g , k is a constant. If $f(x) \cdot g(x) = 5x^3 + 1,715$, what is the value of k ?



113

Which of the following has $x + 2a$ as a factor where a is a positive integer?

- A) $3x^2 + 7x + 14a = 0$
- B) $3x^2 + 28x + 14a = 0$
- C) $3x^2 + 42x + 14a = 0$
- D) $3x^2 + 49x + 14a = 0$

114

The expression $\frac{x^{27}(x-7)}{3x^3} + \frac{7x^{27}}{3x^3}$ is equivalent to $\frac{1}{3}x^k$, where k is a constant and $x > 0$. What is the value of k ?

- A) 3
- B) $\frac{27}{3}$
- C) 25
- D) 28

115

The functions f and g are defined by the equations shown, where a and b are integer constants, $a > b$ and $b > 0$. If $y = f(x)$ and $y = g(x)$ are graphed in the xy -plane, which of the following equations displays, as a constant or a coefficient, the maximum of the graph of the corresponding function when $x \geq 0$.

- I. $f(x) = b(0.97)^{x+a}$
- II. $g(x) = b(0.97)^x + a$

- A) I only
- B) II only
- C) I and II
- D) Neither I nor II



116

The given equation $ax^2 + 88x + c = 0$, where a and c are constants has no real solutions and a factor of $kx + j$. What is the least possible integer value of ac ?

117

$$f(t) = 34,000(1.07)^{6t}$$

The given function f models the balance of an investment account, in dollars, t years after it is opened. Which statement is the best interpretation of $(1.07)^{6t}$?

- A) Every 6 years, the balance increases by 2,380 dollars.
- B) Every 6 years, the balance increases by 7% of the previous 6 years balance.
- C) Every 2 months, the balance increases by 2,380 dollars.
- D) Every 2 months, the balance increases by 7% of the previous 2 months balance.

118

The quadratic function $a(x + 4.5)^2 - d$ can be rewritten as $(x - 9.5)(x + c)$. If a is equal to 1, what is the value of d ?



119

$g(x)$ is a quadratic function. In the xy -plane, the graph of $y = g(x)$ passes through the points $(12, 540)$ and $(16, 908)$. If $g(0) = 12$, what is the value of $g(24)$?

120

The function f is defined by $f(x) = ax^2 + bx + c$, where a , b and c are constants. The graph of $y = f(x)$ in the xy -plane passes through the points $(13, 0)$ and $(-6, 0)$. Which of the following is the value of $a + b$ in terms of a ?

- A) $7a$
- B) $6a$
- C) $-6a$
- D) $-7a$

121

$$f(x) = 1.6^x + 6$$

$$g(x) = 1.8x + b$$

The given system of equations intersects at points (j, k) and (h, r) , where $r < k$. What is the least possible value of b ?

- A) 7
- B) 6
- C) 5
- D) 4



122

The function f is defined by $f(x) = 78(0.39)^x$. For any positive integer n , the value of $f(n)$ is $p\%$ less than the value of $f(n-1)$. What is the value of p ?

- A) 78
- B) 61
- C) 39
- D) 22

123

For which value of a does the quadratic $(7x-8)^2 + 9 = a$ only have one solution?

124

$$f(x) = a^x - b$$

The exponential function given above passes through points $(c, 12)$ and $(2c, 768)$. What is a possible value of b ?



125

If $r = \sqrt[3]{k}$, for the equation $r^{12a-9} = k$, what is the value of a for which r and k are equal?

126

$$4x^3 + bx^2 - 528x = 0$$

In the given equation, b is a positive integer constant. Which value could be a solution to the equation?

- A) -13
- B) -12
- C) -6
- D) 12

127

If $x^2 = j + k$ and $y^2 = j + m$, which of the following is equal to $(x^2 - y^2)^2$?

- A) $k^2 - 2jm + m^2$
- B) $k^2 - 2km + m^2$
- C) $4j^2 - 4jkm + m^2$
- D) $4j^2 - 2jkm + k^2m^2$



128

$$-x^2 + kx - 3,786$$

In the given equation, k is a positive integer. If the equation has no real solution, what is the greatest possible value of k ?

129

If $k = l + m$, which of the following is equivalent to the expression $x^2 - l^2 - 2lm - m^2$?

- A) $(x + k)^2$
- B) $(x - k)^2$
- C) $(x - k)(x + k)$
- D) $x^2 - kx - k^2$

130

$$ax^2 + 148x + c$$

In the given expression, a and c are positive constants. If $jx + k$ is a factor of the expression, where j and k are positive constants, what is the greatest possible value of ac ?

**131**

The given quadratic function $54x^4 + 219x^2 + 105$ has factors in the form $(k)(ax^2 + b)(cx^2 + d)$. If a , b , c , d , and k are integers, what is the smallest possible value of ab ?

132

Which equation correctly gives c in terms of d , e , and f ?

$$\frac{73}{c} = \frac{73}{d} + \frac{73}{e} + \frac{73}{f}$$

- A) $c = d + e + f$
 B) $c = \frac{def}{d + e + f}$
 C) $c = \frac{d + e + f}{def}$
 D) $c = \frac{def}{de + df + ef}$

133

For the exponential function f , the value of $f(3)$ is k , where k is a constant. Which of the following equivalent forms of the function f shows the value of k as the coefficient or the base?

- A) $f(x) = 3(4)(4)^{x+1}$
 B) $f(x) = 3,072(4)(4)^{x-4}$
 C) $f(x) = 12(4)(4)^x$
 D) $f(x) = 768(4)(4)^{x-3}$



134

The function g is defined by $g(x) = ab^{\frac{x}{n}}$, where a , b , and n are constants. If $g(6) = 11$ and $g(9) = 704$, what is the value of $g(11)$?

135

The function f is defined by $f(x) = ax^2 + bx + c$, where a , b , and c are constants. The graph of $y = f(x)$ in the xy -plane passes through the points $(18, 0)$ and $(-11, 0)$. If a is an integer greater than 1, which of the following could be the value of $a + b$?

- A) -12
- B) -6
- C) 7
- D) 14

136

The function g is defined by $g(x) = a(1.36)^x$, where a is a positive constant. For each increase of 1 in the value of x , the value of $g(x)$ increases by $p\%$. Which of the following functions increases by $p\%$ for each increase of 5 in the value of x ?

- A) $h(x) = a(1.36)^{5x}$
- B) $h(x) = a(1.36)^{\frac{x}{5}}$
- C) $h(x) = a(1.36^5)^{5x}$
- D) $h(x) = a\left(1.36^{\frac{1}{5}}\right)^{\frac{x}{5}}$



137

When the price of a particular product is \$120, no products are sold. As the number of products sold increases by 1, the price decreases by \$2.50. If the total revenue for all products sold was \$1,440, what was the total number of products sold? (revenue = price $\times$ number of products sold)

138

$$7x(x - 7) = 63$$

One solution to the given equation can be written as $x = \frac{a + \sqrt{b}}{2}$, where a and b are positive integers. What is the value of ab ?

139

The function g is defined by $g(x) = a\sqrt{b - x}$, where a and b are constants. In the xy -plane, the graph of $y = g(x)$ passes through the point $(36, 0)$ and $f(16) < 0$. Which of the following must be true?

- A) $a > 0$
- B) $b < 0$
- C) $a < b$
- D) $f(0) = 36$



140

$$x(kx - 72) = -8$$

In the given equation, k is an integer constant. If the equation has no real solution, what is the least possible value of k ?

141

A business analyst is developing a model to set the selling price of a product. The analyst determined that for a selling price of \$260, zero products would be sold. For each \$4 decrease in the selling price, the number of products sold would increase by one. For a revenue of \$3,600, what is a possible number of products that could be sold? (revenue = price $\times$ number of products sold)

142

The exponential function g is defined by $g(x) = ab^x$, where a and b are positive constants. If $g(k - 1) = g(k) + \frac{18}{100}g(k - 1)$, where k is a constant, what is the value of b ?



143

Which of the following expressions has a factor of $x + 3k$, where k is a positive integer constant?

- A) $4x^2 + 9x + 27k$
- B) $4x^2 + 15x + 27k$
- C) $4x^2 + 27x + 27k$
- D) $4x^2 + 45x + 27k$

144

At a depth of x meters below the surface, the estimated total pressure P_1 , in millibars, a submarine experiences is given by $P_1 = 1,013x + 12,000$. After the submarine descends a certain depth, k , it descends h additional meters. The estimated total pressure P_2 , in millibars, the submarine experiences after descending k additional meters is given by $P_2 = 1,013h + 721,100$. Which of the following is the best interpretation of 721,100 in this context?

- A) The estimated total pressure in millibars the submarine experienced after the submarine began to descend k additional meters.
- B) The estimated increase in millibars the submarine experienced at a depth of h meters.
- C) The estimated total pressure in millibars the submarine experienced when the submarine began to descend h additional meters.
- D) The estimated increase in millibars the submarine experienced for each increase in depth by h meters.

145

The function f is defined by $f(x) = a(2.5)^x$. For any positive integer n , the value of $f(n - 1)$ is $p\%$ of the value of $f(n)$. What is the value of p ?



146

$$f(x) = 14,000(1.007)^{4x}$$

The given function f models the balance of a bank account, in dollars, x years after it is opened. Which statement is the best interpretation of $(1.007)^{4x}$?

- A) Every 3 months, the balance increases by about 7% of the balance at the beginning of the 3-month interval.
- B) Every 3 years, the balance increases by about 7% of the balance at the beginning of the 3-year interval.
- C) Every 3 months, the balance increases by about 0.7% of the balance at the beginning of the 3-month interval.
- D) Every 3 years, the balance increases by about 0.7% of the balance at the beginning of the 3-year interval.

147

$$\frac{1}{k} + \frac{x}{118} = \frac{1}{kx}$$

In the given equation, k is an integer constant. If the equation has no solutions, what is the greatest possible value of k ?

148

$$-4x^2 + bx - 34 = x^2$$

In the given equation, b is a positive integer. The equation has no real solution. What is the largest possible value of b ?



149

The expression $6x^4 + 35x^2 + 11$ can be rewritten as $(3x^2 + j)(2x^2 + k)$, where j and k are positive integers, or as $(3x^2 + l)(2x^2 + m)$, where l and m are positive nonintegers, what is the value of $k + l$?

150

The function f is defined by $f(x) = (x + 19)(k - x)(x + 8)$, where k is a constant. In the xy -plane, the graph of $y = f(x)$ passes through the point $(32, 0)$. What is the value of $f(0)$?

151

$$2x^2 - 16x - 19 = 0$$

One solution to the given equation can be written as $x = a - \frac{1}{2}\sqrt{b}$, where a and b are integer constants. What is the value of $a + b$?



152

The function f is defined by $f(x) = \frac{2x-8}{(x+7)(x-3)}$. If $f(k+6) = 0$, where k is a constant, what is the value of k ?

153

Which equation correctly gives c in terms of d , e , and f ?

$$\frac{1}{c} = \frac{1}{d} + \frac{1}{e} + \frac{1}{f} + \frac{1}{17}$$

A) $c = \frac{17def}{d+e+f+17}$

B) $c = \frac{17def}{17ef+17df+17de+def}$

C) $c = \frac{17ef+17df+17de+def}{17def}$

D) $c = d+e+f+17$

154

$$g(x) = 8(x-d)(x-e)(x-f)$$

The function g is defined by the given equation, where d , e , and f are distinct constants. When $d < x < f$, the value of $g(x)$ is positive. The graph of $y = g(x)$ in the xy -plane contains the point (p, q) where p and q are constants. If $q = 7$, which of the following could be true?

I. $e < p$

II. $p < d$

III. $f < p < e$

A) I only

B) II only


C) I and II

D) I and III only



155

The graph of $y = f(x) + 6$ is shown. Which equation defines f ?



- A) $f(x) = -2^x + 1$
- B) $f(x) = -2^x - 2$
- C) $f(x) = -2^x - 4$
- D) $f(x) = -2^x - 5$

156

The functions f and g are defined by the given equations, where $x \geq 0$. Which of the following equations displays, as a constant or coefficient, the maximum value of the function it defines, where $x \geq 0$?

- I. $f(x) = 19(0.74)^x + 43$
- II. $g(x) = 7(0.98)^x$

- A) I only
- B) II only
- C) I and II
- D) Neither I nor II

157

$$g(a) = 77a - 4a^2$$

The function g is defined by the given equation. The function h is defined by $h(a) = g(a) + 5$. Which expression represents the maximum value of $h(a)$?

- A) $5 + \left(\frac{77}{4}\right)^2$
- B) $5 + \left(\frac{77}{8}\right)^2$
- C) $5 - \left(\frac{77}{4}\right)^2$
- D) $5 - \left(\frac{77}{8}\right)^2$



158

$$x(7x - 4k) + j(7x - 4k) - (x + j) = 0$$

In the given equation, j and k are positive constants. A solution to the equation is $x = 81$. What is the value of k ?

159

The function g is defined by $g(x) = k|x - 13|^2 - 47|x - 13| + m$, where k and m are constants and $k > m > 47$. If $g(397) = a$ and $g(-371) = b$, where a and b are constants, what is the value of $43(-397)^{a-b} + 371(43)^{b-a}$?


160

The exponential function h is defined by $h(x) = ab^x$, where a and b are positive constants. If $h(k) = p$ and $h(k + 1) = 0.45p$, where k and p are constants, what is the value of b ?

SAT

PART 3

Problem-Solving and Data Analysis





161

The median and the mean of the five integers $\{20, 11, 23, x, x\}$ are equal.

What is the sum of all possible values of x ?

162

55 gallons of a $p\%$ chlorine solution are added to 12 gallons of a 15% chlorine solution to make a 70% chlorine solution. What is the value of p ?

- A) 53
- B) 24
- C) 47
- D) 82

163

A farmer has a square field that measures 50 meters on each side. If one sack of seed covers 250 square meters, how many sacks of seed does the farmer need to plant the entire field?



164

A bakery sold cookies in the ratio of chocolate chip to oatmeal as 4 : 3. If the bakery sold 280 chocolate chip cookies, how many oatmeal cookies were sold?

165

The table shows the exponential relationship between the time, in months, since a new colony of bacteria was introduced to a petri dish and the total number of bacteria. If the number of bacteria increases by a fixed percentage each month, what is the value of b ?

Time (months)	Total Bacteria
0	250
1	b
2	360

166

A music school offers a series of three workshops. 800 people signed up for the first workshop. 45% of the people who signed up for the first workshop also signed up for the second workshop, and 20% of the people who signed up for both the first and second workshops also signed up for the third workshop. How many people signed up for all three workshops?



167

Data set A contains 8 values, and data set B contains 24 values. The table shows the frequencies of the values for each data set. Which of the following statements best compares the means of the two data sets?

Value	Data set A frequency	Data set B frequency
10	2	6
20	3	9
30	1	4
40	2	5

- A) The mean of data set A is greater than the mean of data set B .
 B) The mean of data set A is less than the mean of data set B .
 C) The mean of data set A is equal to the mean of data set B .
 D) There is not enough information to compare the means of the data sets.

168

An orchard has 5 rows of apple trees, 6 rows of pear trees, and 2 rows of cherry trees. Each row of apple trees has 7 trees 25 feet or taller and 4 trees shorter than 25 feet. Each row of pear trees has 9 trees 25 feet or taller and 6 trees shorter than 25 feet. Each row of cherry trees has 12 trees 25 feet or taller and 3 trees shorter than 25 feet. A tree from one of these rows will be selected at random. What is the probability of selecting a pear tree, given that the tree is 25 feet or taller?

- A) $\frac{35}{113}$
 B) $\frac{6}{13}$
 C) $\frac{54}{113}$
 D) $\frac{12}{25}$

169

For a magnetic field passing through a flat surface perpendicular to it, the magnetic flux through the surface is the product of the magnetic field's strength and the area of the surface. A certain flat surface consists of two adjacent squares. The side length of the larger square is twice the side length of the smaller square. A magnetic field with strength 20.00 teslas passes uniformly through this surface, perpendicular to the magnetic field. If the total magnetic flux through this surface is 2,000 teslas $\cdot$ meters², what is the magnetic flux, in teslas $\cdot$ meters², through the larger square?



170

A certain bamboo plant grows 35 centimeters every p months. At this rate, which expression represents the number of centimeters, on average, the bamboo plant grows every k years?

- A) $\frac{420k}{p}$
- B) $\frac{35k}{12p}$
- C) $\frac{420p}{k}$
- D) $\frac{35p}{12k}$

171

Data set A consists of n consecutive integers, where n is an odd integer greater than 3. The smallest integer in set A is m . Data set B is created by adding the integer $(m - 5)$ to set A . Which of the following statements must be true?

- I. The range of set B is greater than the range of set A .
- II. The median of set B is less than the median of set A .
- III. The mean of set B is less than the mean of set A .

- A) I and III only
- B) I and II only
- C) II and III only
- D) I, II, and III

172

Data set X consists of the weights of 60 animals and has a mean of 30 kg. Data set Y consists of the weights of 40 animals and has a mean of 50 kg. Data set Z consists of the weights of the 100 animals from data sets X and Y .

What is the mean, in kilograms, of data set Z ?



173

The positive number a is 480% of the number b , and a is 70% of the number c . If c is $p\%$ of b , which of the following is closest to the value of p ?

- A) 267
- B) 336
- C) 550
- D) 685

174

A certain town has an area of 5.48 square miles. Which of the following is closest to the area, in square yards, of this town? (1 mile = 1,760 yards)

- A) 552
- B) 964
- C) 9,645
- D) 16,974,848

175

A researcher selected 2 random samples of employees in a large company to estimate the percentage of employees that planned to vote in favor of a new division. Based on the first sample, the researcher estimated that 68% of the employees would vote in favor, with an associated margin of error of 9.7%. Based on the second sample, the researcher estimated that 84% of the employees would vote in favor, with an associated margin of error of 6.7%. Assuming the margins of error were calculated in the same way, which of the following best explains why the second sample obtained a smaller margin of error than the first sample?

- A) The first sample contained fewer employees than the second sample.
- B) The first sample contained more employees than the second sample.
- C) The first sample contained a lower percentage of residents that planned to vote in favor of a new division.
- D) The first sample contained a higher percentage of residents that planned to vote in favor of a new division.



176

684 is $p\%$ greater than 9. What is the value of p ?

177

How many liters of a 30% chlorine solution must be added to 12 liters of a 10% chlorine solution to obtain a 15% chlorine solution?

178

Two data sets A and B have 56 values in total. The average of the two data sets is 193. Data set A contains 32 values with an average of 208. What is the average of data set B ?

**179**

Data set A consists of 37 different values that have a minimum of 230, a maximum of 278, a mean of 244, and a standard deviation of 9. The values 230 and 278 are removed from the data set to create data set B . Which of the following statements is true?

- A) Both the mean and standard deviation of data set B are less than those in data set A .
- B) Both the mean and standard deviation of data set B are greater than those in data set A .
- C) The standard deviation of data set B is less than the standard deviation in data set A but the mean of both data sets is the same.
- D) The standard deviation of data set B is greater than the standard deviation in data set A but the mean of both data sets is the same.

180

A 180 gram metal alloy is 50% aluminum. It contains a metal alloy that is 30% aluminum and a second metal alloy made up of 60% aluminum. What is the mass of the second metal alloy?

181

The table shows the distribution of 88 basketball players and their positions on a basketball club. Each player is categorized in one position. If one of the 88 players is selected at random, the probability of selecting a player who is categorized as a power forward, given the player is not categorized as a shooting guard, is $\frac{3}{12}$. How many of these players are categorized as a center?

Position	Frequency
Point guard	26
Shooting guard	16
Small forward	11
Power forward or center	35

**182**

The results of two random samples of votes for a proposition are given in the table. The samples were selected from the same population, and the margins of error were calculated using the same method. Which of the following is the most appropriate reason that the margin of error for sample A is greater than the margin of error for sample B ?

Sample	Percent in favor	Margin of error
A	67.83%	7.8%
B	72.94%	4.5%

- A) Sample A had a larger sample size.
- B) Sample A had a smaller sample size.
- C) Sample A had a smaller number of votes that could not be recorded.
- D) Sample A had a lower percent of favorable responses.

183

360 is $p\%$ of 10. What is the value of p ?

184

The density of a certain type of wood is 770 kilograms per cubic meter. A sample of this type of wood is in the shape of a cube, where each edge has a length of 0.7 meters. To the nearest whole number, what is the mass, in kilograms, of this sample?

- A) 264
- B) 377
- C) 1,571
- D) 2,245



185

The number j is 2,400% greater than the number k . The number k is 350% greater than 8. What is the value of j ?

186

There are red, blue, green, and black marbles in a bag. There are 26 black marbles, 8 green marbles, and 26 blue and red marbles. Given that a randomly selected marble is not black, the probability of selecting a red marble is $\frac{5}{17}$. What is the number of blue marbles in the bag?

187

A table that weighs 348 pounds puts equal amounts of pressure on each of its 4 identical legs. If the pressure that each leg exerts on the floor is 12 pounds per square inch, what is the area, in square inches, of the bottom of each leg that is touching the floor?



188

An object's speed is increasing at a rate of 4.80 meters per second squared. What is the rate in miles per minute squared, rounded to the nearest tenth? (Use 1 mile = 1609 meters.)

189

A 100 gram metal alloy is 40% silver. It contains a metal alloy that is 65% silver and a second metal alloy made up of 25% silver. What is the mass of the first metal alloy?

190

Joshua Tree National Park has an area of 1,242 square miles. What is the area, in square yards, of Joshua Tree National Park? (1 mile = 1,760 yards).

- A) $\frac{1,242}{1,760}$
B) $\frac{1,242}{(1,760)^2}$
C) $1,242(1,760)$
D) $1,242(1,760)^2$



191

There are two ranches. Ranch A and ranch B . The ratio of male horses to female horses is $1 : 15$ for the total number of horses on ranch A and B . The ratio of female horses on ranch A to ranch B is $7 : 17$. There are 168 horses on ranch A and 344 horses on Ranch B . How many more male horses are there on ranch A than on ranch B ?

192

a is 3700% greater than b . a is also 5% less than c . What percent greater than b is c ?

193

Each of the following frequency tables represents a data set. Which of these frequency tables represent the data set with the smallest standard deviation?

A)

Value	Frequency
2	1
10	5
15	3
20	5
24	1

B)

Value	Frequency
2	0
10	20
15	25
20	20
24	0

C)

Value	Frequency
2	100
10	50
15	25
20	0
24	25

D)

Value	Frequency
2	45
10	55
15	65
20	75
24	85



194

The mass of particles A , B , and C are a atomic mass units, b atomic mass units, and c atomic mass units, respectively. If the mass of particle A is 4,600% of the mass of particle C , and the mass of particle C is 0.004% of the mass of particle B , which expression represents the value of $a + b$ in terms of c ?

- A) $2,960c$
- B) $4,604c$
- C) $4,850c$
- D) $25,046c$

195

A rocket's speed is increasing at a rate of 14.6 meters per second squared. What is the rate in miles per minute squared? (1 mile = 1,609 meters)

196

The speed of object K is 9,525% of the speed of object J , and the speed of object K is 0.025% of the speed of object L . If the speed of object L is $x\%$ of the speed of object J , what is the value of $\frac{x}{1,000}$?



197

A company conducted a study on the time spent studying per night by students in grades 9, 10, and 11 and their average grade point average. A total of 5,600 students were selected at random from all students in grades 9, 10, and 11 in a large school district in the US. The selected students were asked to report their number of hours studied as less than 1, from 1–3 hours, or more than 3 hours per night. If the results of the study suggest a relationship between the self-reported number of hours studied and students average grade point average, which of the following aspects of the study will prevent these results from being generalized to all students in grades 9, 10, and 11 in the US?

- A) The selected students are not all from the same school in the district.
- B) The number of selected students is too small.
- C) The number of selected students from grades 10 and 11 is greater than grade 9.
- D) The students were not selected at random from all students in grades 9, 10, and 11 in the US.

198

Substance D is made up of substances A , B , and C . The total mass of substance B is 621.6 grams, which makes up 42% of substance D . Of the remaining mass of substance D , 60% was substance A , and the remainder was substance C . Based on this information, how many grams of substance D is substance C ?

199

The positive number k is 3,700% of the number j , and j is 12.5% of the number m . If $k - m = aj$, what is the value of a ?



200

The speed of object A is 764% greater than the speed of object B , and the speed of object A is 0.96% of the speed of object C . If the speed of object C is $p\%$ of the speed of object B , what is the value of p ?

201

For the positive quantities a , b , and c , 78% of c is equivalent to 0.096% of b , and b is equivalent to 250% of a . What percentage of c is a ?

202

The expression $0.0004x$ represents the result of decreasing a positive quantity x by $p\%$. What is the value of p ?



203

An architectural model of a square campus has a side length of 405 inches, and 1 inch on the model is equal to an actual distance of 270 feet. A smaller version of the same campus is constructed with the side length of the square 65% shorter than the side length of the previous model. On the smaller model, which of the following is closest to the actual distance, in feet, represented by 1 inch?

- A) 94.5
- B) 175.5
- C) 415.38
- D) 771.43

204

A petri dish had x number of bacteria. A month later the number of bacteria increased by 17,456%. What is the resulting number of bacteria on the petri dish in terms of x ?

- A) $17.456x$
- B) $174.56x$
- C) $175.56x$
- D) $17,456x$

205

On average, a certain weed grows 63 centimeters every m months. At this rate, which expression represents the number of centimeters, on average, the weed grows every t years?

- A) $\frac{21m}{4t}$
- B) $\frac{21t}{4m}$
- C) $\frac{756m}{t}$
- D) $\frac{756t}{m}$



206

The positive number a is 7,848% of the sum of positive numbers b and c , and c is 32% of b . What percent of c is a ?

207

For a study, a group of beavers will be selected from a population of 370 beavers, and a group of otters will be selected from a population of 420 otters. Some of the beavers and otters will be in a treatment group, and some of the beavers and otters will be in a control group. Which of the following is necessary for this study to attempt to establish a cause-and-effect relationship between two variables?

- A) The total number of beavers in the study is equal to the total number of otters in the study.
- B) The beavers and otters are randomly selected from their habitats.
- C) The proportion of female to male otters in the treatment group is equal to the proportion of female to male beavers in the treatment group.
- D) The beavers and otters are randomly assigned to the treatment and control groups.


208

At the start of a stock trading competition, John's and Julia's stock portfolios were of equal value. At the end of the stock trading competition, John's portfolio had increased by 330%. Julia's portfolio has increased by 2,480%. At the end of the competition Julia's portfolio is worth $p\%$ more than John's portfolio, what is the value of p ?

SAT

PART 4

Geometry and Trigonometry





209

$$8x^2 + 64px + 8y^2 - 64py = -224p^2$$

In the xy -plane, the graph of the given equation is a circle. The length of the radius of the circle is np , where n and p are positive constants. What is the value of n ?

210

Circle A : $x^2 + (y - 1)^2 = 25$

Circle B : $x^2 + (y - 1)^2 = 81$

Determine the difference in area between the two circles in units of π .

211

The surface area of rectangular prism A is 126 m^2 . The surface area of rectangular prism B is 4536 m^2 . Rectangular prism A has a volume of 85 m^3 . If both rectangular prisms are similar, what is the volume of rectangular prism B ?




212

A line intersects two parallel lines, forming four acute angles and four obtuse angles. The measure of one of the acute angles is $(4x - 410)^\circ$. The sum of the measures of one of the acute angles and three of the obtuse angles is $(-8x + w)^\circ$. What is the value of w ?

213

In the figure shown, if the radius of circle O is 12, what is the length of diagonal AC of rectangle $OABC$?



214

The table shows the volume of two similar solids, right circular cylinder A and right circular cylinder B . The radius of right circular cylinder A is 8 units. The surface area of right circular cylinder A is $k\pi$ square units, and the surface area of right circular cylinder B is $n\pi$ square units, where k and n are constants. What is the value of $\frac{n}{k}$? (The surface area of a right circular cylinder with radius r and height h is $2\pi r^2 + 2\pi rh$.)

	Volume (cubic units)
Right circular cylinder A	600π
Right circular cylinder B	$4,800\pi$



215

The radius of circle Y is 9 times longer than the radius of circle X . If circle X has an area with length y , what is the length of circle Y 's area?

- A) $9y^2$
- B) $9y$
- C) $81y^2$
- D) $81y$

216

Cylinder A has a volume of V . Cylinder B has a radius that is equal to cylinder A 's diameter as well as a height that is three times shorter than the height of cylinder A . Which of the following expressions correctly represent the volume of cylinder B ?

- A) $\frac{2}{9}V$
- B) $\frac{9}{2}V$
- C) $\frac{4}{3}V$
- D) $\frac{3}{4}V$

217

In triangles XYZ and PQR , it is known that $2XY = PQ$. If the triangles are similar, which of the following must be true?

- A) $2YZ = QR$
- B) $2PR = XZ$
- C) $XY = PQ$
- D) $2XY = PR$



218

In the xy -plane, a circle has center C with coordinates $(p, 8)$, where p is a constant. Points G and H lie on the circle. Point G has coordinates $(p + 3\sqrt{5}, 14)$, and point H has coordinates $(q, 8)$, where q is a constant greater than p . What is the area, in square units, of triangle CGH ?


- A) $27\sqrt{5}$
- B) 27
- C) $5\sqrt{27}$
- D) $27\sqrt{6}$

219

A square has an area of $x + 15$ square units and a perimeter of $2x$ units. What is the value of x ?

220

In the figure, line p is parallel to line q , and line r intersects both lines. What is the value of y ?





221

Circle A is defined by the equation $(x - 3)^2 + (y + 2)^2 = 16$. Circle B (not shown) is the result of shifting circle A up 5 units and decreasing the radius so that the radius of circle B is half the radius of circle A . Which equation defines circle B ?


- A) $(x - 3)^2 + (y - 3)^2 = 4^2$
- B) $(x - 3)^2 + (y - 3)^2 = 2^2$
- C) $(x - 3)^2 + (y + 7)^2 = 2^2$
- D) $(x - 3)^2 + (y - 7)^2 = 16^2$

222

In convex pentagon $ABCDE$, segment AB is parallel to segment DE . The measure of angle B is 125 degrees, and the measure of angle D is 160 degrees. What is the measure, in degrees, of angle C ?

223

In the figure, $\overline{AE}$ intersects $\overline{BD}$ at point C , and $\overline{AB}$ is parallel to $\overline{DE}$. The lengths of $\overline{BC}$, $\overline{AC}$, and $\overline{CD}$ are $\frac{21}{4}$, 7, and 6, respectively. What is the length of $\overline{AE}$?



- A) 27
- B) 15
- C) 9
- D) 21



224

In the xy -plane, a circle has center C with coordinates (h, k) . Points A and B lie on the circle. Point A has coordinates $(h + 3, k + 4)$, and $\angle ACB$ is a right angle. What is the length of $\overline{AB}$?

- A) 5
- B) $5\sqrt{2}$
- C) 10
- D) $10\sqrt{2}$

225

A rectangle is inscribed in a circle such that each vertex of the rectangle lies on the circumference of the circle. The diagonal of the rectangle is twice the length of the shortest side of the rectangle. The area of the rectangle is $784\sqrt{3}$ square units. What is the length, in units, of the diameter of the circle?

226

$$(x + 5)^2 + (y - 3)^2 = 100$$

The graph of the given equation represents a circle in the xy -plane. The point $(a, a - 2)$ lies on the circle. Which of the following is a possible value for a ?

- A) -5
- B) -10
- C) 0
- D) 10



227

A right square prism has a height of 18 units. The volume of the prism is 2,592 cubic units. What is the length, in units, of an edge of the base?


228

A right circular cone has a radius of a and a height of b . A larger cone has a volume that is 243 times the volume of the smaller cone. Which of the following could represent the radius A , in terms of a , and the height B , in terms of b , of the larger cone?

- A) $A = 3a, B = 81b$
- B) $A = 9a, B = 3b$
- C) $A = 27a, B = b$
- D) $A = 3a, B = 9b$

229

In the figure above, the ratio of the lengths $AQ : QR : RC = 1 : 3 : 2$. If the area of quadrilateral $PQRS$ is 60, what is the area of $\triangle ABC$?



- A) 92
- B) 118
- C) 144
- D) 120



230

A right rectangular prism has a height of 5 meters. The length of the prism's base is w meters, which is 4 meters more than the width of the prism's base. Which function V gives the volume of the prism, in cubic meters, in terms of the length of the prism's base?

- A) $V(w) = 5w^2(w + 4)$
- B) $V(w) = 4(5)w^2$
- C) $V(w) = 5w(w - 4)$
- D) $V(w) = w^2(w + 4)$

231

The surface area of a cube is $6(3c)^2$, where c is a positive constant. Which of the following gives the perimeter of one face of the cube?

- A) $3c$
- B) $6c$
- C) $12c$
- D) $18c$

232

Points $P\left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$ and $Q(0, 1)$ are located on the unit circle with center $O(0, 0)$. Which of the following could be the measure of angle $\angle POQ$?

- A) 45°
- B) 90°
- C) 135°
- D) 225°



233

For two acute angles, $\angle A$ and $\angle B$, $\cos A = \sin B$. The measures, in degrees, of $\angle A$ and $\angle B$ are $13x + 10$ and $48 + 3x$, respectively. What is the value of x ?

234

The table gives the volume of two similar cylinders. If the radius of cylinder A is 9 cm. What is the value of $j - k$?

$$\text{Volume} = \pi r^2 h$$

$$\text{Surface area} = 2\pi r^2 + 2\pi r h$$

Cylinder	Volume	Surface Area
A	$1,296\pi$	$k\pi$
B	$162,000\pi$	$j\pi$

235

A polygon has exactly 97 sides. If the measure of each of the 97 interior angles of this polygon is $(180p)^\circ$, what is the value of p ?



236

In triangle ABC , the sum of the measures of angle A and angle B is 90 degrees. The value of $\sin(A)$ is $\frac{2\sqrt{7}}{8}$. What is the value of $\cos(B)$?

- A) $\frac{2\sqrt{7}}{8}$
- B) $\frac{2\sqrt{7}}{6}$
- C) $\frac{6\sqrt{2}}{14}$
- D) $\frac{8}{23\sqrt{2}}$

237

In the xy -plane, the graph of the given equation is a circle. Which point lies on this circle?

$$(x + 7)^2 + (y - 11)^2 = 25$$

- A) $(-7, 11)$
- B) $(11, -7)$
- C) $(\sqrt{11} + 7, \sqrt{14} - 11)$
- D) $(\sqrt{11} - 7, \sqrt{14} + 11)$

238

A circle in the xy -plane has its center at $(3, -7)$ and has a radius of 12. An equation of this circle is $x^2 + y^2 + ax + by + c = 0$, where a , b , and c are constants. What is the value of c ?



239

Sphere A has a radius of $7x$ and sphere B has a radius of $91x$. The volume of sphere B is how many times the volume of sphere A ?

240

In the xy -plane, there are 3 points a , b , and c . Point a has coordinates of $(1, 0)$, point b has coordinates of $(0, 0)$ and point c has coordinates of $(-1, 0)$. Which of the following gives a possible angle measure, in radians, of $\angle abc$?

- A) $\frac{456\pi}{6}$
- B) $\frac{459\pi}{6}$
- C) $\frac{462\pi}{6}$
- D) $\frac{468\pi}{6}$

241

Lines k and l are perpendicular. If the point $(4, m)$ and $(9, m + 3)$ lies on line k and lines k and l intersect at $(9, m + 3)$, which of the following can lie on line l ?

- A) $(m - 6, -4)$
- B) $(m + 9, 24)$
- C) $(21, m - 17)$
- D) $(24, m + 12)$



242

Rectangle A has a width of 50 feet and a length of 45 feet and is located inside rectangle B . A uniform distance x , in feet, separates the sides of rectangle A from rectangle B . The area of rectangle B is 2646 square feet, what is the value of x ?




Figure not drawn to scale.

243

In the given figure, the hypotenuse is 64 what is the $\cos x$?




Figure not drawn to scale.

- A) $\frac{\sqrt{3}}{2}$
- B) $\frac{1}{2}$
- C) $\frac{2\sqrt{3}}{3}$
- D) 2

244

A rectangular pyramid has a square base with an area of 324 square meters. What is the surface area, in square meters, of one of the triangular faces if the rectangular pyramid has a volume of 4,320 cubic meters?



245

The perimeter of a right isosceles triangle is $76 + 76\sqrt{2}$. What is the length of the hypotenuse?

246

In $\triangle ABC$, $\angle B$ is a right angle and the length of $\overline{AB}$ is 456 millimeters. If the $\cos C = \frac{7}{25}$, what is the length, in millimeters, of $\overline{BC}$?

247

In triangle ABC and triangle DEF , sides BC and EF each have a side length of 37 inches, and angles B and E each have an angle measure of 63° . Which of the following additional pieces of information is(are) sufficient to prove whether triangle ABC is congruent to triangle DEF ?

- I. The measure of angles A and C are equal.
- II. The length of sides AC and DF are equal.
- III. The measures of angles A and D are equal.

- A) I is sufficient, but II and III are not.
- B) II is sufficient, but I and III are not.
- C) III is sufficient, but I and II are not.
- D) II and III are sufficient, but I is not.



248

A square is inscribed in a circle with the equation $x^2 + y^2 - 8x + 10y = 27$. What is the area of the square?

249

Triangle ABC is inscribed in a circle with a radius of 85. The length of AC is 170 and the length of BC is 168. What is the value of $\frac{AB}{BC}$?

250

What is the value of $\sin(x^\circ)(\cos 90 - x^\circ) + \sin(90 - x^\circ)(\cos x^\circ)$?

- A) 0.5
- B) 1
- C) 2
- D) 90



251

A gardener is purchasing topsoil for a new garden. The garden is in the shape of a rectangular prism. The dimensions of the garden are 54 feet by 192 feet by 2 feet. If it costs \$96 per cubic yard of topsoil, how much does the gardener need to spend in dollars? (1 yard = 3 feet)

252

In the xy -plane, which of the following points lies on a circle with the equation $(x + k)^2 + (y - j)^2 = 841$?

- A) $(k + 20, j + 21)$
- B) $(k - 20, j - 21)$
- C) $(-k + 21, j + 20)$
- D) $(-k - 21, -j - 20)$

253

In the figure below, lines a and b are parallel. What is the value of $x + y$?




Figure not drawn to scale.

**254**

An equilateral triangle with side lengths of k is inscribed in a circle. Which of the following defines the radius, r , in terms of k ?

- A) $\frac{k}{2}$
- B) $\frac{k\sqrt{3}}{6}$
- C) $\frac{k\sqrt{3}}{3}$
- D) $\frac{k\sqrt{3}}{2}$

255

Triangle ABC is an equilateral triangle with a height of 30 meters. What is the perimeter of triangle ABC in meters?

- A) $60\sqrt{3}$
- B) 90
- C) $90\sqrt{3}$
- D) $120\sqrt{3}$

256

A rectangular area consists of 3,179 equal squares where each square has an area of k . If the length of the rectangular area is 2.75 times the width and the width is equal to $x\sqrt{k}$, what is the value of x ?



257

In the given figure, BC is the diameter of the circle. If the length of BC is equal to 132 and the length of AB is equal to $\sqrt{363}$, what is the value of $\frac{BC}{BD}$?




Figure not drawn to scale.

258

The table gives the volume and surface areas of two similar rectangles, where k is a constant. What is the value of k ?

	Volume (cubic inches)	Surface Area (square inches)
Rectangle A	4,374	1,782
Rectangle B	1,500,282	k

- A) 5,346
- B) 12,474
- C) 87,318
- D) 611,226

259

In the figure, parallel lines a and b are intersected by lines c , d , and e . If $x = 54$, $v = 127$, and $w > x$, which statement about y and z must be true?




Figure not drawn to scale.

- A) $z > y$
- B) $z < y$
- C) $z = y$
- D) $z + y = 90$



260

An equilateral triangle has a perimeter of 1200 meters. If the height of the triangle is equal to $x\sqrt{3}$, what is the value of x ?

261

Two lines intersect at exactly one point, forming two acute angles and two obtuse angles. The measure of one of these angles is $(7x - 210)^\circ$. Which of the following could NOT be the sum of the measures of any two of these angles?

- A) $(-14x + 780)^\circ$
- B) $(-14x + 420)^\circ$
- C) 180°
- D) $(14x - 420)^\circ$

262

A circle with a center $B(0,0)$ is a unit circle on the xy -plane. Point $A(1,0)$ and C lie on the circle. If the measurement of angle ABC is $\frac{975\pi}{60}$ in radians, what is the x -coordinate of point C ?

- A) $\frac{\sqrt{2}}{2}$
- B) $-\frac{\sqrt{2}}{2}$
- C) 0
- D) $\frac{\sqrt{2}}{3}$



263

A circle is inscribed inside of square $ABCD$. Segment AC is the diagonal of the square, and its length is 30cm. What is the radius of the circle in cm?

- A) $\frac{15\sqrt{2}}{2}$
- B) 15
- C) $15\sqrt{2}$
- D) 30

264

A metal mold in the shape of a rectangular prism has a length of 27 inches, a width of 48 inches, and a height of 180 inches. The equation $c = v(d)$ calculates the cost of filling the metal mold, where c is the total cost in dollars, v is the volume in cubic yards, and d is the cost per cubic yard. If c is equal to \$970, what is the value of d in dollars? (Note 1 yard = 36 inches)

265

An equilateral triangle is inscribed in a circle with a diameter of 16. Which of the following gives the area of the equilateral triangle?

- A) $6\sqrt{3}$
- B) $24\sqrt{3}$
- C) $32\sqrt{3}$
- D) $48\sqrt{3}$



266

The surface area of rectangular prism A is 312 m^2 . The surface area of rectangular prism B is $11,232 \text{ m}^2$. Rectangular prism A has a volume of 224 m^3 . If both rectangular prisms are similar, what is the volume of rectangular prism B ?

267

The given figure shows lines a and b intersected by line k . Which of the following is sufficient to prove that a and b are parallel?





Figure not drawn to scale.

- A) $180 - y = x$
- B) $x = 180 - z$
- C) $y + z = 180 - x$
- D) $180 - z = y$

268

For the given circle, the measure of arc $abc = 130 + 7x$ degrees and the measure of arc $acb = 90 - 3x$ degrees. What is the value of x ?





269

What is the perimeter of an equilateral triangle with an area of $289\sqrt{3}$?

270

Circle A has the equation $(x - 18)^2 + y^2 = 27$. Circle B has the equation $(x - 18)^2 + (y - m)^2 = 50$. If Circle B passes through the center of circle A, what is a possible value of m ?

- A) $3\sqrt{3}$
- B) $5\sqrt{2}$
- C) $5\sqrt{2} - 3\sqrt{3}$
- D) $5\sqrt{2} + 3\sqrt{3}$

271

A circle with center $(-7, -4)$ has a radius equal to 14. The equation $x^2 + y^2 + ax + by + c = 0$ represents the circle. What is the value of c ?



272

In triangle ABC , angle C is a right angle, point D lies on $\overline{AB}$, point E lies on $\overline{BC}$, and $\overline{DE}$ is parallel to $\overline{AC}$. If the length of $\overline{AB}$ is 39 units, $\overline{DE}$ is 8 units, the length of $\overline{AC}$ is greater than the length of $\overline{BC}$, and the area of triangle ABC is 270 units, what is the length of $\overline{BE}$, in units?

273

A rectangle is inscribed in a circle such that each vertex of the rectangle lies on the circle. The diagonal of the rectangle is twice the length of the shortest side of the rectangle. If the area of the circle is $5,929\pi$ square units, what is the area, in square units, of the rectangle?

- A) $77\sqrt{2}$
- B) $77\sqrt{3}$
- C) $5,929\sqrt{2}$
- D) $5,929\sqrt{3}$

274

In triangle ABC and triangle DEF , sides BC and EF each have a length of 23 inches, and angles C and F each measure 43° . Which of the following additional pieces of information is(are) sufficient to prove that triangle ABC is congruent to triangle DEF ?

- I. The measures of angles D and E are equal.
- II. The lengths of $\overline{AB}$ and $\overline{DE}$ are equal.
- III. The measures of angles A and D are equal.

- A) I only
- B) II only
- C) II and III
- D) III only



275

Circle A is defined by the equation $(x + 14k)^2 + (y + 27k)^2 = 81k^2$, where k is a positive constant. Circle B is the result of shifting circle A up $13k$ units. Which equation defines circle B ?

- A) $(x + 14k)^2 + (y + 40k)^2 = 81k^2$
- B) $(x + 14k)^2 + (y + 14k)^2 = 81k^2$
- C) $(x + 27k)^2 + (y + 14k)^2 = 81k^2$
- D) $(x + 14k)^2 + (y + 14k)^2 = 94k^2$

276

A circle in the xy -plane has its center at $(-5, -7)$. Line a is tangent to this circle at $(-13, -4)$. What is the slope of line a ?

277

Point K lies on a unit circle in the xy -plane and has coordinates $(1, 0)$. Point O is the center of the circle and has coordinates $(0, 0)$. Point J also lies on the unit circle and the measure of angle KOJ is $\frac{478\pi}{3}$ radians. If the coordinates of point J are (a, b) , where a and b are constants, what is the value of b ?

- A) $-\frac{1}{2}$
- B) $-\frac{\sqrt{3}}{2}$
- C) $\frac{1}{2}$
- D) $\frac{\sqrt{3}}{2}$



278

A right square pyramid has a surface area of 51,984 square centimeters, which includes a base area of 23,104 square centimeters. What is the slant height, in centimeters, of this pyramid?

279

For triangle ABC and triangle DEF , angle A and D each measure 45° , $BC = 36$ and $EF = 54$. Which additional piece(s) of information is(are) sufficient to prove that triangle ABC is similar to triangle DEF ?

- I. The length of $\overline{AB}$ is $\frac{2}{3}$ the length of $\overline{DE}$.
II. The length of $\overline{AC}$ is $\frac{2}{3}$ the length of $\overline{DF}$.

- A) I is sufficient, but II is not.
B) II is sufficient, but I is not.
C) I is sufficient, and II is sufficient.
D) Neither I nor II is sufficient.

280

The perimeter of an equilateral triangle is 954 inches. The three vertices of the triangle lie on a circle. The radius of the circle is $k\sqrt{3}$ inches. What is the value of k ?



281

A right square pyramid has a height of 13 inches. A second right square pyramid has a height 39 inches. The area of the bases of the two pyramids is 400 square inches. Which of the following is closest to the difference in surface area of the second pyramid and the surface area of the first pyramid, in square inches, rounded to the nearest integer?

282

A line intersects two parallel lines, forming four obtuse and four acute angles. The measure of one of the obtuse angles is $(-6x + 380)^\circ$. The sum of the measures of two of the acute angles and one of the obtuse angles is $(ax - k)^\circ$, where a and k are both positive constants. What is the value of $a + k$?

283

In the figure shown, points A , B , and C lie on the circle and $AC > AB$ and $AD = \sqrt{592}$. The diameter of the circle is 82. If $\frac{CD}{BD} = k$, what is the value of k ?




Figure not drawn to scale.



284

In the figure, parallel lines a and b are intersected by lines c , d , and e . If $v = 37$ and $z = 144$, y and x are integers, and $y < v$, which statement about x must be true?




Figure not drawn to scale.

- A) $x < y$
- B) $x > y$
- C) $x = w$
- D) $w < x$

285

The perimeter of an equilateral triangle with a height of 58 units is $k\sqrt{3}$ units, what is the value of k ?

- A) $\frac{58}{3}$
- B) $\frac{116}{3}$
- C) $\frac{174}{3}$
- D) 116

286

In the figure shown, $\overline{AC}$ and $\overline{BD}$ intersect at point E , $AC = 1,080$ and $\overline{ED} = \frac{4}{5}\overline{EC}$. What is the length of $\overline{ED}$?




Figure not drawn to scale.



287

$$4x^2 + 4y^2 + bx + by = 32$$

The given equation, where b is a positive constant, defines a circle in the xy -plane. The radius of the circle is $\sqrt{106}$. What is the value of b ?

288

The edge length of the square base and the height of a right square pyramid are integers, and the pyramid has a volume of 16,464 cubic inches. If the slant height is 35 inches, what is the height, in inches, of this pyramid?

289

A rectangular gameboard is divided into 480 square regions of equal area k , in square units. The length of the gameboard is 7.5 times the width of the gameboard. If the width of the gameboard is $x\sqrt{k}$ units, what is the value of x ?



290

The area of a triangle is 340 square inches. The length of the base of the triangle is 6 greater than 2 times the height of the triangle. What is the height, in inches, of the triangle?

291

In the figure, $\overline{AC}$ intersects $\overline{BD}$ at point E , and $\overline{AB}$ is parallel to $\overline{DC}$. The lengths of $\overline{AC}$, $\overline{BE}$, and $\overline{EC}$ are 23, 17, and 7, respectively. What is the length of $\overline{ED}$?




Figure not drawn to scale.

- A) $\frac{16}{119}$
- B) $\frac{23}{119}$
- C) $\frac{119}{23}$
- D) $\frac{119}{16}$

292

Point A lies on a unit circle in the xy -plane and has coordinates $(1, 0)$. Point B is the center of the circle and has coordinates $(0, 0)$. Point C also lies on the unit circle and has coordinate $\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$. Which of the following could be the positive measure of angle ABC , in radians?

- A) $\frac{454\pi}{6}$
- B) $\frac{458\pi}{3}$
- C) $\frac{460\pi}{3}$
- D) $\frac{462\pi}{3}$



293

$$x^2 - 9x + y^2 + 27y + \frac{67}{2} = 0$$

In the xy -plane, the graph of the given equation is a circle. If this circle is inscribed in a square, what is the perimeter of the square?

294

For triangle XYZ , the length of $\overline{XZ}$ is 37 and the length of $\overline{XY}$ is 27. Point K (not shown) lies on $\overline{XY}$ such that $\overline{YK}$ is perpendicular to $\overline{XZ}$. What is the value of $\frac{YK}{YZ}$?




Figure not drawn to scale.



Answer Key

Answers marked — are the handful that could not be pinned down from the source page, and are left open rather than guessed.

Algebra

1. A	2. 336	3. 1	4. -4	5. 300	6. D
7. C	8. D	9. A	10. D	11. -28/3	12. C
13. C	14. 7,007	15. B	16. 62	17. B	18. 81/4
19. —	20. B	21. 0.25	22. A	23. 354	24. D
25. A	26. C	27. C	28. 93	29. C	30. -15
31. B	32. C	33. -2	34. D	35. -9	36. 399
37. B	38. B	39. -5	40. 21/17	41. C	42. 2,670
43. 33/37					

Advanced Math

44. 184	45. 7	46. B	47. 13	48. 99	49. D
50. A	51. -182	52. 32 or 48	53. 128	54. 5.5	55. -48
56. 12	57. D	58. B	59. -12	60. 3	61. D
62. A	63. 4	64. B	65. 3	66. C	67. B
68. -5	69. 4	70. 3/14	71. 96	72. 91/4	73. A
74. 11	75. B	76. 10,000	77. D	78. -336	79. 23/40
80. B	81. D	82. D	83. D	84. D	85. 19/24
86. D	87. 61.75	88. 1/77	89. C	90. D	91. 27/16
92. 2,401	93. -113	94. B	95. D	96. A	97. -40
98. 129	99. -198	100. 6	101. B	102. 24	103. D
104. B	105. 9/7	106. D	107. 36	108. 462	109. 12
110. 183	111. D	112. 35	113. D	114. C	115. D
116. 1,937	117. D	118. 196	119. 1,932	120. C	121. C
122. B	123. 9	124. 16	125. 1	126. B	127. B
128. 123	129. C	130. 5,476	131. 14	132. D	133. B
134. 11,264	135. A	136. D	137. 24	138. 595	139. C
140. 163	141. 20 or 45	142. 0.82	143. D	144. C	145. 40
146. C	147. -30	148. 26	149. 27.5	150. 4,864	151. 106
152. -2	153. B	154. A	155. D	156. B	157. A
158. 141.5	159. 414	160. 0.45			

Problem-Solving and Data Analysis



161. 49	162. D	163. 10	164. 210	165. 300	166. 72
167. A	168. C	169. 1,600	170. A	171. D	172. 38
173. D	174. D	175. A	176. 7,500	177. 4	178. 173
179. A	180. 120	181. 17	182. B	183. 3,600	184. A
185. 900	186. 16	187. 7.25	188. 10.7	189. 37.5	190. D
191. 24	192. 3,900	193. B	194. D	195. 32.67	196. 38,100
197. D	198. 343.36	199. 29	200. 90,000	201. 3,250	202. 99.96
203. D	204. C	205. D	206. 32,373	207. D	208. 500

Geometry and Trigonometry

209. 2	210. 56	211. 18,360	212. 1360	213. 12	214. 4
215. D	216. C	217. A	218. B	219. 10	220. 54
221. B	222. 75	223. B	224. B	225. 56	226. A
227. 12	228. B	229. C	230. C	231. C	232. A
233. 2	234. 10,800	235. 95/97	236. A	237. D	238. -86
239. 2,197	240. C	241. C	242. 2	243. A	244. 369
245. 76	246. 133	247. C	248. 136	249. 13/84	250. B
251. 73,728	252. C	253. —	254. C	255. A	256. 34
257. 48	258. C	259. B	260. 200	261. B	262. A
263. A	264. 194	265. D	266. 48,384	267. D	268. 35
269. 102	270. B	271. -131	272. 10/3	273. D	274. D
275. B	276. 8/3	277. B	278. 95	279. D	280. 106
281. 954	282. 26	283. 37/4	284. B	285. D	286. 259.2
287. 56	288. 28	289. 8	290. 17	291. B	292. B
293. 104	294. 27/37				