

Modèle Monte Carlo du transport dans l'atmosphère des électrons relativistes et des photons gamma en relation avec les Flashes Gamma Terrestres (TGF)

Soutenance de thèse

David Sarria

Sous la direction de Pierre-Louis Blelly

14 Septembre 2015

En vue de l'obtention du grade de docteur de l'Université Paul Sabatier



Outline

1 Introduction : Transient events from thunderstorms, environment

2 Monte-Carlo model

3 Applications of the model

- Simulation of a typical Terrestrial Gamma-ray Flash
- Simulation of Fermi e^-/e^+ events

4 TARANIS


5 Conclusion



Introduction


- Transient events from thunderstorms, environment

TLE and TGF




TLE and TGF

TLE = Transient Luminous Event (1990)
= Sprites, Elves, Jets, etc ...




TLE and TGF

TGF = Terrestrial Gamma-ray Flash (1994)





TGF : Production sketch





- Ingredients : high energy electrons ("seeds"), strong electric field
- Relativistic electron avalanche
- Bremsstrahlung radiation
- Cold Runaway VS Relativistic Feedback

TGF : atmospheric filtering



- Reference model : MSIS (empirical)
- Density falls exponentially with increasing altitude, scale height $\sim 7 \text{ km}$
- Gamma ray flux too faint to be detected from ground \rightarrow detected from space

TGF : detections



First detections by BATSE on-board NASA-CGRO,
published in Science (*Fishman et al., 1994*)


Also detected by RHESSI, Fermi and AGILE (Not their primary goal)

Fermi : > 400 TGF/day (*Briggs et al.*). Maybe 50,000 TGF/day ! (*Ostgaard et al.*)


Primary objective of TARANIS and ASIM



TGF : morphology




Fermi
Briggs et al.
(2011)



AGILE
Marisaldi et al.
(2014)

- Typical duration of ~ 0.5 ms
- Bremsstrahlung energy spectrum filtered by the atmosphere.
- ~ 1 photon/cm² at satellite's altitude (~ 600 km)
- Simulations $\rightarrow \gtrsim 10^{16}$ photons at source (~ 15 km)

TGF : source altitude




- The energy spectrum seen at satellite altitude is changed if the photons are emitted at different altitudes
- Monte-Carlo simulations for different source altitudes
- Comparison with satellite data
- Early estimation : 15-21 km
- Later estimation : 12-15 km

TEB


- During their propagation in the atmosphere, photons will produce a lot of electrons and positrons.

Once escaping the atmosphere :

- Electrons and Positrons are guided by magnetic field lines
→Terrestrial Electron Beam (**TEB**)
- TGF (=photons) follow straight trajectories



TEBs : Electron motion





Conservation of the first adiabatic invariant


→ Magnetic mirror

→ Possible detection of electrons on both sides

TEB example : 091214 Fermi event



(Briggs et al., 2011)



- Double peaked
 - Much longer than a TGF
 - TGF detection above the egyptian desert ?
 - 511 keV line : positron annihilation
 - Electrons/positron beams ! TGF was actually over Zambia

Monte-Carlo model

Generalities

**MC-PEPTITA = Monte-Carlo Photon Electron Positron Tracking
In Terrestrial Atmosphere**

3D


Involved particles:

Photons Electrons Positrons

- Energy range : 10 keV to 100 MeV
- External models :
 - MSIS for the atmosphere
 - IGRF-12 for the magnetic field
- 11 processes included


Processes : photons

Coherent (Rayleigh) scattering




- Only deviation, no energy change

Incoherent (Compton) scattering




- Photon is deviated loses energy
- Electron is added

Photo-electric absorption




- Photon is removed
- Electron is added

Electron/positron pair production




- Photon is removed
- Electron and positron are added

Processes : photons, relative probability




Processes : Electrons and positrons

Elastic scattering




- Only deviation, no energy change

Inelastic scattering



- e-/e+ is deviated and loses energy E_s
- Electron is added

Bremsstrahlung




- e-/e+ loses energy
- Photon is added

Processes : Positrons

One extra process :

Positron annihilation




In a dense medium, annihilation only after para-positronium phase

$\rightarrow e^-/e^+$ annihilation into 2 photons with $E_{\gamma+} \approx E_{\gamma-} \approx 511$ keV



Processes : Electrons/Positrons, relative probability



Particles' displacement

- s = distance between 2 interactions
- σ = total cross section, ρ = number density, λ = mean free path
- $P(s)$ = probability of not interacting after reaching a distance s


$$\lambda^{-1} = \sigma(E) \rho(h)$$

$$P(s) = 1 - \exp \left[- \int_0^s \frac{ds'}{\lambda(s')} \right]$$

- γ trajectories : straight lines between two interactions
- e^-/e^+ trajectories :
 - Solving $\frac{d\vec{p}}{ds} = \frac{\vec{v} \times \vec{\Omega}}{v} - \nu \frac{\vec{v}}{v}$
 - Different regimes function of ν/Ω , a lot of care taken

Validation : Comparison with GEANT4

- GEANT4 is a reference Monte-Carlo code developed by CERN
- Testing similar set-ups with GEANT4 and MC-PEPTITA
- Test of the collision model



- Excellent agreement between GEANT4 and MC-PEPTITA



MC-PEPTITA : assets

- Full tracking of all the particle
- Particle's outputs with new information, including :
 - Process of production (see next)
 - Production altitude (see next)
 - Pitch angles of e^- , e^+ (see next)
 - Larmor radii of e^- , e^+
 - Number of interactions
- Accurate resolution of e^- , e^+ transport
- Fast (e.g. compared to G4), optimized for Earth's environment
- Parallelized, with very good scalability





Applications of the model

- 1 Simulation of a typical TGF : γ and e^-e^+
- 2 Simulation of Fermi e^-e^+ events, focus on 091214




Initial conditions for a typical TGF

- Source at 15 km altitude, southern hemisphere.
- Energy spectrum $\propto \frac{1}{E} \exp\left(\frac{-E}{\epsilon}\right)$, $\epsilon = 7.3$ MeV.
- Time distribution $\propto \exp\left(\frac{-t^2}{2\sigma_{t,src}^2}\right)$, $\sigma_{t,src} = 0.15$ ms.
- Angle beaming $\propto \exp\left(\frac{-\theta^2}{2\sigma_\theta^2}\right)$, $\sigma_\theta = 35^\circ$.




- Every particle crossing satellite's altitude is saved.

γ , e^- , e^+ energy spectra at satellite altitude



- Photon spectrum roughly agree with previous works (e.g. Dwyer *et al.*)
- e^-, e^+ more noisy than γ (Monte-Carlo), $\frac{N(\gamma)}{N(e^-)} \sim 100$; $\frac{N(e^-)}{N(e^+)} \sim 8$
- Positron spectrum determined for the first time.
- MC-PEPTITA permits to decompose these spectra : focus on electrons

e⁻ energy spectra decomposition




- > 100 keV : domination of Compton, < 100 keV : domination of inelastic
 - Pair production : same spectrum as positrons, not dominant, but $\sim 12\%$ of total

Simulation of a typical Terrestrial Gamma-ray Flash


e^-, e^+ : production profile

- Determined for the first time with MC-PEPTITA.




- $\sim 10^{19}$ e^- produced (above 10 keV),
Effects of this massive production ?
 - $\sim 10^{14}$ can escape, all produced
above ≈ 30 km
 - e^- production non-negligible
up to 100 km, neglected above 60 km
in previous works.
 - Below 70 km, e^- production
dominated by Compton scattering
 - Above 70 km : 50% Compton
scattering, 50% inelastic scattering
 - e^+ lower average production altitude
(≈ 48 km vs ≈ 53 km)

e^- , e^+ fluxes at satellite altitude : definition



e^- , e^+ fluxes at satellite altitude




Applications of the model

- 1 Simulation of a typical TGF : γ and e^-e^+
- 2 Simulation of Fermi e^-e^+ events, focus on 091214


Simulating 3 Fermi e^-e^+ events (1/2)

- Briggs et al. (2011) : Three e^-e^+ events detected by Fermi-GBM





Simulating 3 Fermi e^-e^+ events (2/2)


080807




- Accurate modelling from simple TGF assumption
- N = Fermi detectors total count
- Estimation of the photon number required at source $n_{\gamma,\text{src}}$: strong events
- 090813 needs adjusting $\sigma_{t,\text{src}}$
- 091214 : X 5 statistics and double peak




Fermi 091214 event : pitch angle definition




Simulation of Fermi e^-/e^+ events

Fermi 091214 event : pitch angle time decomposition



Simulation of Fermi e^-/e^+ events


Fermi 091214 event : pitch angle time decomposition



- Full TEB pitch angle distribution resolved for the first time !


Simulation of Fermi e^-/e^+ events

Fermi 091214 event : pitch angle time decomposition

Pitch angle of $\sim 120^\circ$ → Had mirroring altitude of ~ 100 km



Fermi 091214 event : pitch angle time decomposition



Three populations :

- 1 Coming from south
- 2 Mirroring with negligible interactions
- 3 Mirroring with non-negligible interactions


Applications of the model : summary

- e^+ spectrum (new).
- Decomposition of e^- spectrum (new).
- Production profile of e^-, e^+ (new).
- Spatial properties of e^-, e^+ beams, $\sim 40\%$ wider than previously expected.
- Accurate simulation of Fermi events.
- Pitch angle decomposition of e^-/e^+ time histogram (new).
- Method for estimating the satellite's position within the TEB (not discussed).
- Determining geographical positions for mirror pulse of TEBs (not discussed).
- Gives important information for understanding the data of the TARANIS XGRE and IDEE instruments, and makes predictions



TARANIS

General features



Orbit:

- Sun-synchronous
- Inclination: 98°
- Altitude: 700 km


Principal Investigator : J.L. Pinçon (LPC2E Orleans, France)
Project manager : C. Bastien-Thiry (CNES Toulouse, France)

© CNES - Septembre 2007 / illus D. Ducros




Expected launch : 2018

TARANIS : Main scientific objectives




- To advance the physical understanding of the links between TLEs, TGFs and environmental conditions
- To identify the signatures associated with these phenomena and to provide inputs to test generation mechanisms.
- To provide inputs for the modelling of the effects of TLEs, TGFs and bursts of precipitated and accelerated electrons on the Earth's atmosphere.

TARANIS : Instrumentation




TARANIS : XGRE





Rough estimation :
~ 850 TGF/year
each with ~ 280 counts

- Sandwich design of plastic/crystal scintillators
- Photons : 20 keV to 10 MeV (LaBr₃)
- Electrons : 1 MeV to 10 MeV (plastic)
- Dead time < 350 ns
- Total detection area ~800 cm²
- 3 detectors, 20° tilts
→ Estimation of the direction of photons



TARANIS : IDEE



- 2 detectors
- Electrons : 60 keV to 5 MeV
- < 2 ms time resolution
- Pitch angle estimated from coincidence between Si and CdTe cells.

Conclusions

- Transient events related to thunderstorms, including TGFs and associated electron and positron emissions :
 - Recently discovered (1990+)
 - Generation mechanism not well understood
 - as well as their effects on the environment
- We built a model of propagation of γ , e^- , e^+ on Earth :
 - Accounts for all the relevant physics.
 - Accurate, fast.
 - Access to a lot of new information.
- Will be helpful for the TARANIS data analysis (2018), in particular :
 - XGRE : X/ γ photons and high-energy electrons.
 - IDEE : Electrons on a large energy range, with pitch angle estimation.

Future work

- Applying our model to other events (Fermi,TARANIS,ASIM ?).
- Collaboration with *M.S. Briggs* at Univ. of Alabama :
 - Effects of pitch angles of the e^-/e^+ , interactions with satellite and detectors.
 - Important for TARANIS' IDEE and XGRE data.
- Coupling MC-PEPTITA with others models
 - Ionospheric models : effects of TGF, e.g. production of excited states and optical emissions.
 - Radio emission
- Spectral response of the XGRE detector (Post-doc at APC).
- 2018 : TARANIS data analysis.



Merci pour votre attention

Un grand merci à mon directeur de thèse et à tous mes collègues,
ma famille et mes amis pour leur soutien durant ces trois ans

Extra slides


Process : Positron annihilation

- In-flight annihilation is very unlikely
- Positrons will lose most of their energy (=slow down) catch an electron to form a positronium

Two ways of annihilation :



Para-Positronium





Ortho-Positronium


→ 2 photons,
each at 511 keV

→ 3 photons

- InterStellar Medium density $\sim 30 \text{ cm}^{-3}$, Air density at 40 km altitude $\sim 7 \times 10^{16} \text{ cm}^{-3}$
- PP lifetime $= 1.2 \times 10^{-10} \text{ s} << \text{OP lifetime} = 1.4 \times 10^{-7} \text{ s}$
- In ISM, if any of these two state forms, it will annihilate a long time before encountering another electron from the medium.
- In a dense medium, if OP forms, its long lived lifetime permits "pick-off" annihilation where an opposite spin electron from some surrounding material will annihilate in para-orientation before the ortho-bound electron can collapse and annihilate with the positron.

091214 : Larmor radii





- Full TEB pitch angle distribution resolved for the first time !

Three populations :


- 1 Coming from south
- 2 Mirroring with negligible interactions
- 3 Mirroring with non-negligible interactions

- Accurate analytical model for the corresponding time distributions
 - Permits estimation of position of the satellite within the TEB
- Both time and pitch angle will be estimated by TARANIS-IDEE




- Accurate modelling from simple TGF assumption
- N = Fermi detectors total count
- Estimation of the photon number required at source $n_{\gamma,src}$: strong events
- 090813 needs adjusting $\sigma_{t,src}$
- 091214 : X 5 statistics and double peak


Detection at 400 km




Detection at 550 km




Detection at 620 km



Detection at 700 km




Detection at 850 km




- $\sim 10^{19}$ e^- produced (above 10 keV),
- Effects of this massive production ?
- $\sim 10^{14}$ can escape, all produced above ≈ 30 km
- e^- production non-negligible up to 100 km, neglected above 60 km in previous works.
- Below 70 km, e^- production dominated by Compton scattering
- Above 70 km : 50% Compton scattering, 50% inelastic scattering
- e^+ lower average production altitude (≈ 48 km vs ≈ 53 km)


RREA : Relativistic Runaway Electron Avalanche (2/2)



RREA : Relativistic Runaway Electron Avalanche (2/2)



RREA : Relativistic Runaway Electron Avalanche (1/2)





Conclusions

- We built the MC-PEPTITA model of propagation of γ , e^- , e^+ on Earth :
 - Accounts for all the relevant physics
 - Access to a lot of new information
 - Accurate, fast
- Applications of the model :
 - Energetic decomposition of γ , e^- , e^+
 - Spatial properties of e^- , e^+ beams
 - Simulation of Fermi events
 - Pitch angle decomposition of e^- , e^+ time histogram.
 - Method of estimating the satellite's position within the TEB (not discussed here).
 - Determining geographical positions where we may or may not see the mirror pulse for TEBs (not discussed here).
- Made in preparation for the TARANIS mission, featuring :
 - Simultaneous measurement of transient events in X/ γ , optical, and radio.
 - X/ γ photons and high-energy electrons (XGRE)
 - Electrons on a large energy range, with pitch angle estimation (IDEE).