

# Phil/LPS 31 Introduction to Inductive Logic

## Lecture 13

David Mwakima

[dmwakima@uci.edu](mailto:dmwakima@uci.edu)

Department of Logic and Philosophy of Science  
University of California, Irvine

May 17th 2023

# Topics

- ▶ Odds
- ▶ Expected Value (Discrete Case Only)

## Recap: Probability Rules

Rule 1  $P(S) = 1$

Rule 2  $0 \leq P(E) \leq 1$

Rule 3 If  $E \cap F = \emptyset$ , then  $P(E \cup F) = P(E) + P(F)$  (Special Disjunction Rule)

Rule 4  $P(E \cup E^c) = 1$

Rule 5  $P(E^c) = 1 - P(E)$

Rule 6  $P(\emptyset) = 0$

Rule 7  $P(E) = P(E \cap F) + P(E \cap F^c)$

Rule 8  $P(E \cup F) = P(E) + P(F) - P(E \cap F)$  (General Disjunction Rule)

Rule 9  $P(E \cap F) = P(E) \times P(F)$  if  $E$  and  $F$  are independent.

Rule 10  $P(E \cap F) = P(E) \times P(F|E)$  if  $E$  and  $F$  are dependent.

## Recap: Conditional Probability & Bayes' Theorem

- ▶ Conditional Probability. This is a consequence of Rule 10:

$$P(E | F) = \frac{P(E \cap F)}{P(F)}$$

## Recap: Conditional Probability & Bayes' Theorem

- ▶ Conditional Probability. This is a consequence of Rule 10:

$$P(E | F) = \frac{P(E \cap F)}{P(F)}$$

- ▶ Bayes' Theorem

$$P(E | F) = \frac{P(F|E)P(E)}{P(F)}$$

## Recap: Conditional Probability & Bayes' Theorem

- ▶ Conditional Probability. This is a consequence of Rule 10:

$$P(E | F) = \frac{P(E \cap F)}{P(F)}$$

- ▶ Bayes' Theorem

$$P(E | F) = \frac{P(F|E)P(E)}{P(F)}$$

- ▶ If  $\{E_1, E_2, \dots, E_n\}$  is a partition of  $E$ , then from the theorem of total probability:

$$\begin{aligned} P(E | F) &= \frac{P(F | E)P(E)}{P(F | E_1)P(E_1) + \dots + P(F | E_n)P(E_n)} \\ &= \frac{P(F | E)P(E)}{\sum_{i=1}^n P(F | E_i)P(E_i)} \end{aligned}$$

## Alternative ways of talking about chance

- ▶ There are other ways that people talk about chances that are related to **but different** from probability. Let's talk about that today.

## Alternative ways of talking about chance

- ▶ There are other ways that people talk about chances that are related to **but different** from probability. Let's talk about that today.
- ▶ Consider the following sentences:



# Alternative ways of talking about chance

- ▶ There are other ways that people talk about chances that are related to **but different** from probability. Let's talk about that today.
- ▶ Consider the following sentences:
  - ▶ The **odds** are 3 to 2 in favor of Mage winning the 2023 Kentucky Derby.

## Alternative ways of talking about chance

- ▶ There are other ways that people talk about chances that are related to **but different** from probability. Let's talk about that today.
- ▶ Consider the following sentences:
  - ▶ The **odds** are 3 to 2 in favor of Mage winning the 2023 Kentucky Derby.
  - ▶ The **odds are fifty fifty** that Real Madrid will beat Manchester City in the Champions League Semi-Final Today.

# Alternative ways of talking about chance

- ▶ There are other ways that people talk about chances that are related to **but different** from probability. Let's talk about that today.
- ▶ Consider the following sentences:
  - ▶ The **odds** are 3 to 2 in favor of Mage winning the 2023 Kentucky Derby.
  - ▶ The **odds are fifty fifty** that Real Madrid will beat Manchester City in the Champions League Semi-Final Today.
- ▶ Odds are a useful way of talking about the chances of an event and many people often go back and forth between probability and odds as if they mean the same thing. They don't!

# Alternative ways of talking about chance

- ▶ There are other ways that people talk about chances that are related to **but different** from probability. Let's talk about that today.
- ▶ Consider the following sentences:
  - ▶ The **odds** are 3 to 2 in favor of Mage winning the 2023 Kentucky Derby.
  - ▶ The **odds are fifty fifty** that Real Madrid will beat Manchester City in the Champions League Semi-Final Today.
- ▶ Odds are a useful way of talking about the chances of an event and many people often go back and forth between probability and odds as if they mean the same thing. They don't!
- ▶ In this lecture we will see why and also introduce ways of going (1) from probability to odds and (2) from odds to probability.

## From Probabilities to Odds

- ▶ If the probability of an event  $E$  is  $p$ , then we know by Rule 5, that the probability of  $E^c$  is  $1 - p$ .

## From Probabilities to Odds

- ▶ If the probability of an event  $E$  is  $p$ , then we know by Rule 5, that the probability of  $E^c$  is  $1 - p$ .
- ▶ We define the **odds in favor** of  $E$ , or simply just **odds** of  $E$ , as:

$$\mathcal{O}(E) = \frac{p}{1 - p}$$

## From Probabilities to Odds

- ▶ If the probability of an event  $E$  is  $p$ , then we know by Rule 5, that the probability of  $E^c$  is  $1 - p$ .
- ▶ We define the **odds in favor** of  $E$ , or simply just **odds** of  $E$ , as:

$$\mathcal{O}(E) = \frac{p}{1 - p}$$

- ▶ We write  $\mathcal{O}(E) = p : 1 - p$  and read this as “Odds in favor of  $E$  are  $p$  to  $1 - p$ ”.

## From Probabilities to Odds

- ▶ If the probability of an event  $E$  is  $p$ , then we know by Rule 5, that the probability of  $E^c$  is  $1 - p$ .
- ▶ We define the **odds in favor** of  $E$ , or simply just **odds** of  $E$ , as:

$$\mathcal{O}(E) = \frac{p}{1 - p}$$

- ▶ We write  $\mathcal{O}(E) = p : 1 - p$  and read this as “Odds in favor of  $E$  are  $p$  to  $1 - p$ ”.
- ▶ Examples. Calculate the odds in favor of the following events.



# From Probabilities to Odds

- ▶ If the probability of an event  $E$  is  $p$ , then we know by Rule 5, that the probability of  $E^c$  is  $1 - p$ .
- ▶ We define the **odds in favor** of  $E$ , or simply just **odds** of  $E$ , as:

$$\mathcal{O}(E) = \frac{p}{1 - p}$$

- ▶ We write  $\mathcal{O}(E) = p : 1 - p$  and read this as “Odds in favor of  $E$  are  $p$  to  $1 - p$ ”.
- ▶ Examples. Calculate the odds in favor of the following events.
  - (1) The odds of Manchester City winning if the probability that Manchester City will win after being ahead 2 - 0 at half time is 0.8.

## From Probabilities to Odds

- ▶ If the probability of an event  $E$  is  $p$ , then we know by Rule 5, that the probability of  $E^c$  is  $1 - p$ .
- ▶ We define the **odds in favor** of  $E$ , or simply just **odds** of  $E$ , as:

$$\mathcal{O}(E) = \frac{p}{1 - p}$$

- ▶ We write  $\mathcal{O}(E) = p : 1 - p$  and read this as “Odds in favor of  $E$  are  $p$  to  $1 - p$ ”.
- ▶ Examples. Calculate the odds in favor of the following events.
  - (1) The odds of Manchester City winning if the probability that Manchester City will win after being ahead 2 - 0 at half time is 0.8.
  - (2) The odds of having two girls, if the probability that a family has two children who are both girls is  $\frac{1}{4}$ .

# From Probabilities to Odds

- ▶ Some properties of odds:

# From Probabilities to Odds

- ▶ Some properties of odds:
  - (1)  $\mathcal{O}(E)$  can be any real number in the interval  $[0, +\infty)$ .

# From Probabilities to Odds

- ▶ Some properties of odds:
  - (1)  $\mathcal{O}(E)$  can be any real number in the interval  $[0, +\infty)$ .
  - (2) If  $\mathcal{O}(E) > 1$ , then  $P(E) > P(E^c)$

# From Probabilities to Odds

- ▶ Some properties of odds:
  - (1)  $\mathcal{O}(E)$  can be any real number in the interval  $[0, +\infty)$ .
  - (2) If  $\mathcal{O}(E) > 1$ , then  $P(E) > P(E^c)$
  - (3) If  $\mathcal{O}(E) < 1$ , then  $P(E) < P(E^c)$

# From Probabilities to Odds

- Some properties of odds:
  - (1)  $\mathcal{O}(E)$  can be any real number in the interval  $[0, +\infty)$ .
  - (2) If  $\mathcal{O}(E) > 1$ , then  $P(E) > P(E^c)$
  - (3) If  $\mathcal{O}(E) < 1$ , then  $P(E) < P(E^c)$
  - (4) If  $\mathcal{O}(E) = 1$ , then  $P(E) = P(E^c)$

# From Probabilities to Odds

- Some properties of odds:
  - (1)  $\mathcal{O}(E)$  can be any real number in the interval  $[0, +\infty)$ .
  - (2) If  $\mathcal{O}(E) > 1$ , then  $P(E) > P(E^c)$
  - (3) If  $\mathcal{O}(E) < 1$ , then  $P(E) < P(E^c)$
  - (4) If  $\mathcal{O}(E) = 1$ , then  $P(E) = P(E^c)$
  - (5)  $\mathcal{O}(E)$  is invariant to scaling.



# From Probabilities to Odds

► Some properties of odds:

- (1)  $\mathcal{O}(E)$  can be any real number in the interval  $[0, +\infty)$ .
- (2) If  $\mathcal{O}(E) > 1$ , then  $P(E) > P(E^c)$
- (3) If  $\mathcal{O}(E) < 1$ , then  $P(E) < P(E^c)$
- (4) If  $\mathcal{O}(E) = 1$ , then  $P(E) = P(E^c)$
- (5)  $\mathcal{O}(E)$  is invariant to scaling.
- (6)  $\mathcal{O}(E^c)$ , read as "odds against  $E$ " is given by the reciprocal of  $\mathcal{O}(E)$

# From Probabilities to Odds

- ▶ Some properties of odds:
  - (1)  $\mathcal{O}(E)$  can be any real number in the interval  $[0, +\infty)$ .
  - (2) If  $\mathcal{O}(E) > 1$ , then  $P(E) > P(E^c)$
  - (3) If  $\mathcal{O}(E) < 1$ , then  $P(E) < P(E^c)$
  - (4) If  $\mathcal{O}(E) = 1$ , then  $P(E) = P(E^c)$
  - (5)  $\mathcal{O}(E)$  is invariant to scaling.
  - (6)  $\mathcal{O}(E^c)$ , read as "odds against  $E$ " is given by the reciprocal of  $\mathcal{O}(E)$
- ▶ Exercise. What are the ways that probabilities are different from odds given these properties of odds?

## From Probabilities to Odds

- ▶ Odds can be extremely useful if what you are interested in is the relative likelihood of an event  $E$  and its complement  $E^c$ .

# From Probabilities to Odds

- ▶ Odds can be extremely useful if what you are interested in is the relative likelihood of an event  $E$  and its complement  $E^c$ .
- ▶ Examples of complementary events are usually binary events like “Success” or “Failure”; “Win” or “Lose”; “Adverse Health Effect” or “No adverse Effect”.

# From Probabilities to Odds

- ▶ Odds can be extremely useful if what you are interested in is the **relative likelihood** of an event  $E$  and its complement  $E^c$ .
- ▶ Examples of complementary events are usually **binary events** like “Success” or “Failure”; “Win” or “Lose”; “Adverse Health Effect” or “No adverse Effect”.
- ▶ Later in the course we will see that on the Bayesian interpretation of probability, odds provide a useful way of going from fair bets (based on odds) to probabilities as **degrees of belief**.

# From Probabilities to Odds

- ▶ Odds can be extremely useful if what you are interested in is the **relative likelihood** of an event  $E$  and its complement  $E^c$ .
- ▶ Examples of complementary events are usually **binary events** like “Success” or “Failure”; “Win” or “Lose”; “Adverse Health Effect” or “No adverse Effect”.
- ▶ Later in the course we will see that on the Bayesian interpretation of probability, odds provide a useful way of going from fair bets (based on odds) to probabilities as **degrees of belief**.
- ▶ We will see later that the **Odds Ratio** provides a good measure of association (or dependence) between two groups in the same way that we used joint and marginal probabilities to assess independence between groups.

## From odds to probabilities

- ▶ Going from odds to probabilities we just reverse the operations.

## From odds to probabilities

- ▶ Going from odds to probabilities we just reverse the operations.
- ▶ Recall that:

$$\mathcal{O}(E) = \frac{p}{1-p}$$



## From odds to probabilities

- ▶ Going from odds to probabilities we just reverse the operations.
- ▶ Recall that:

$$\mathcal{O}(E) = \frac{p}{1-p}$$

- ▶ Solving this equation for  $p$  we get:

$$p = \frac{\mathcal{O}(E)}{1 + \mathcal{O}(E)}$$

## From odds to probabilities

- ▶ Exercise. Given the following odds find the probability of the following events.

# From odds to probabilities

- ▶ Exercise. Given the following odds find the probability of the following events.
  - (1) The odds of Real Madrid making it to the final after being 3 - 0 down at the 80th minute is 1:9.

# From odds to probabilities

- ▶ Exercise. Given the following odds find the probability of the following events.
  - (1) The odds of Real Madrid making it to the final after being 3 - 0 down at the 80th minute is 1:9.
  - (2) There are even odds of getting a head after tossing a coin once.

# From odds to probabilities

- ▶ Exercise. Given the following odds find the probability of the following events.
  - (1) The odds of Real Madrid making it to the final after being 3 - 0 down at the 80th minute is 1:9.
  - (2) There are even odds of getting a head after tossing a coin once.
- ▶ See **Homework 7** for more exercises.

## Expected Value (Discrete Case)

- The number of telephone calls received at a remote call center per minute show the following frequencies:

Number of Calls	Count
1	10
2	8
3	4
4	4
5	2
6	2
7	1
8	1
TOTAL	32

## Expected Value (Discrete Case)

- ▶ The number of calls received per minute at a remote call center is called **a random variable**.

## Expected Value (Discrete Case)

- ▶ The number of calls received per minute at a remote call center is called **a random variable**.
- ▶ What is a random variable? A random variable has a technical definition that we shall not consider in this class. But! Let me mention that a random variable is neither a variable nor “random”, whatever that means.



## Expected Value (Discrete Case)

- ▶ The number of calls received per minute at a remote call center is called **a random variable**.
- ▶ What is a random variable? A random variable has a technical definition that we shall not consider in this class. But! Let me mention that a random variable is neither a variable nor “random”, whatever that means.
- ▶ For our purposes it is enough to think that whenever we have an event that can occur with different probabilities, we can **model** the **number of occurrences** of that event using a **probability distribution**

## Expected Value (Discrete Case)

- ▶ The number of calls received per minute at a remote call center is called **a random variable**.
- ▶ What is a random variable? A random variable has a technical definition that we shall not consider in this class. But! Let me mention that a random variable is neither a variable nor “random”, whatever that means.
- ▶ For our purposes it is enough to think that whenever we have an event that can occur with different probabilities, we can **model** the **number of occurrences** of that event using a **probability distribution**
- ▶ Using the probability distribution, we can then ask useful questions such as: what is the **expected value** or **expectation** of the random variable.

## Expected Value (Discrete Case)

- ▶ The expected value  $E(X)$ , “read as the expected value of a random variable  $X$ , can be helpfully thought of as the **average** or **mean** of  $X$ .

## Expected Value (Discrete Case)

- ▶ The expected value  $E(X)$ , “read as the expected value of a random variable  $X$ ”, can be helpfully thought of as the **average** or **mean** of  $X$ .
- ▶ In fact  $E(X)$  is a **weighted average**, where we weigh each possible value of  $X$ ,  $x_i$  say, by its probability  $p_i$ .

## Expected Value (Discrete Case)

- ▶ The expected value  $E(X)$ , “read as the expected value of a random variable  $X$ , can be helpfully thought of as the **average** or **mean** of  $X$ .
- ▶ In fact  $E(X)$  is a **weighted average**, where we weigh each possible value of  $X$ ,  $x_i$  say, by its probability  $p_i$ .
- ▶ For example, in the call center example, we can let the number of calls received at the center be a random variable  $X$ . We can calculate its expected value of the number of calls received per minute in two ways. Let's do that.

## Expected Value (Discrete Case)

- ▶ The expected value  $E(X)$ , “read as the expected value of a random variable  $X$ , can be helpfully thought of as the **average** or **mean** of  $X$ .
- ▶ In fact  $E(X)$  is a **weighted average**, where we weigh each possible value of  $X$ ,  $x_i$  say, by its probability  $p_i$ .
- ▶ For example, in the call center example, we can let the number of calls received at the center be a random variable  $X$ . We can calculate its expected value of the number of calls received per minute in two ways. Let's do that.
- ▶ Here is the data for your convenience.

## Expected Value (Discrete Case)

Number of Calls	Count
1	10
2	8
3	4
4	4
5	2
6	2
7	1
8	1
TOTAL	32

## Expected Value (Discrete Case)

- ▶ Let  $\{a_1, a_2, a_3, \dots, a_n\}$  be the possible values that a random variable  $A$  can take and let  $\{p_1, p_2, p_3, \dots, p_n\}$  be the probabilities that it takes on these values.



## Expected Value (Discrete Case)

- ▶ Let  $\{a_1, a_2, a_3, \dots, a_n\}$  be the possible values that a random variable  $A$  can take and let  $\{p_1, p_2, p_3, \dots, p_n\}$  be the probabilities that it takes on these values.
- ▶ The **expected value** of  $A$  is given by:

$$\begin{aligned} E(A) &= a_1p_1 + a_2p_2 + a_3p_3 + \cdots + a_np_n \\ &= \sum_{i=1}^n a_i p_i \end{aligned}$$

## Expected Value (Discrete Case)

- ▶ Let  $\{a_1, a_2, a_3, \dots, a_n\}$  be the possible values that a random variable  $A$  can take and let  $\{p_1, p_2, p_3, \dots, p_n\}$  be the probabilities that it takes on these values.
- ▶ The **expected value** of  $A$  is given by:

$$\begin{aligned} E(A) &= a_1p_1 + a_2p_2 + a_3p_3 + \dots + a_np_n \\ &= \sum_{i=1}^n a_i p_i \end{aligned}$$

- ▶ Commit this formula to memory!