Phil/LPS 31 Introduction to Inductive Logic Lecture 2

David Mwakima dmwakima@uci.edu Department of Logic and Philosophy of Science University of California, Irvine

April 5th 2023

Topics

- Sentential Logic: Formulas
- ► Sentential Logic: Main connectives
- Sentential Logic: Paraphrasing English
- Truth functions and Truth tables
- Uses of Truth Tables

Sentential Logic: Formulas

Remember that the formal symbols of sentential logic are:

- 1. p, q, r, s and t as symbols for sentences. If we need more than 5 symbols (rarely!), then add the following countably many symbols p₁, p₂, p₃,
- 2. ∨ for "or", ¬ for "not" since the other symbols for "and" and "if..., then..." can be defined from these. (More of this later)
- 3. (for left bracket and) for right bracket.

The transformation rules (also known as "syntactic rules") are:

- 1. Any sentence p is a formula.
- 2. If F is a formula, then $\neg F$ is formula.
- 3. If F is a formula and G is a formula, then $(F \vee G)$ is a formula.

The closure condition states that a formula is anything that results from 1 or from a finite number of applications of rule 2 or rule 3. Nothing else is a formula of sentential logic.

Sentential Logic: Formulas

Are the following formulas in sentential logic?

- 1. q
- 2. t
- **3**. ¬*q*
- **4**. ¬¬*r*
- 5. $(\neg\neg\neg p \lor q)$
- 6. $(p \lor (r \lor \neg q))$
- 7. $((q \lor (t \lor s)) \lor (r \lor \neg p))$

Sentential Logic: Formulas

Are the following formulas in sentential logic?

- 1. a
- 2. *b*
- 3. *q*¬
- 4. $\neg r \neg$
- 5. $(\neg \lor \neg p \lor q)$
- 6. $p \lor (r \lor \neg q)$
- 7. $((q \lor (t \lor s)) \lor r \lor \neg p)$

► Consider:

Consider:

1.
$$((q \lor (t \lor s)) \lor (r \lor \neg p))$$

- Consider:
 - 1. $((q \lor (t \lor s)) \lor (r \lor \neg p))$
 - 2. $((q \lor (t \lor s)) \lor r \lor \neg p)$

- Consider:
 - 1. $((q \lor (t \lor s)) \lor (r \lor \neg p))$
 - 2. $((q \lor (t \lor s)) \lor r \lor \neg p)$
- ► We said that (1) is a formula while (2) is not because brackets matter.

- Consider:
 - 1. $((q \lor (t \lor s)) \lor (r \lor \neg p))$
 - 2. $((q \lor (t \lor s)) \lor r \lor \neg p)$
- ► We said that (1) is a formula while (2) is not because brackets matter.
- ► Think about similar uses of brackets in algebra. Brackets tell us which operations to do first.

- Consider:
 - 1. $((q \lor (t \lor s)) \lor (r \lor \neg p))$
 - 2. $((q \lor (t \lor s)) \lor r \lor \neg p)$
- ► We said that (1) is a formula while (2) is not because brackets matter.
- ► Think about similar uses of brackets in algebra. Brackets tell us which operations to do first.
 - 1. $((24 \div 2) + 2) 4 = 10$

- Consider:
 - 1. $((q \lor (t \lor s)) \lor (r \lor \neg p))$
 - 2. $((q \lor (t \lor s)) \lor r \lor \neg p)$
- ► We said that (1) is a formula while (2) is not because brackets matter.
- ► Think about similar uses of brackets in algebra. Brackets tell us which operations to do first.
 - 1. $((24 \div 2) + 2) 4 = 10$
 - 2. $(24 \div (2+2)) 4 = 2$

- Consider:
 - 1. $((q \lor (t \lor s)) \lor (r \lor \neg p))$
 - 2. $((q \lor (t \lor s)) \lor r \lor \neg p)$
- ► We said that (1) is a formula while (2) is not because brackets matter.
- ► Think about similar uses of brackets in algebra. Brackets tell us which operations to do first.
 - 1. $((24 \div 2) + 2) 4 = 10$
 - 2. $(24 \div (2+2)) 4 = 2$
- ▶ So using brackets is not just a stipulation of our logic, it turns out that having brackets helps us parse the formula correctly in order to identify the main connective. In (1) we can identify the main connective but in (2) we cannot. This is why brackets matter.

▶ We identified the main connective in (1) by thinking about how the formula was built up from simpler formulas.

- ▶ We identified the main connective in (1) by thinking about how the formula was built up from simpler formulas.
- ► The main connective is the connective that is added last when forming complex formulas from simple ones and closing everything up with brackets, where applicable.

- ▶ We identified the main connective in (1) by thinking about how the formula was built up from simpler formulas.
- ► The main connective is the connective that is added last when forming complex formulas from simple ones and closing everything up with brackets, where applicable.
- Identify the main connective in these formulas:

- ▶ We identified the main connective in (1) by thinking about how the formula was built up from simpler formulas.
- ► The main connective is the connective that is added last when forming complex formulas from simple ones and closing everything up with brackets, where applicable.
- Identify the main connective in these formulas:
 - 1. $\neg q$

- ▶ We identified the main connective in (1) by thinking about how the formula was built up from simpler formulas.
- ► The main connective is the connective that is added last when forming complex formulas from simple ones and closing everything up with brackets, where applicable.
- ▶ Identify the main connective in these formulas:
 - 1. $\neg q$
 - 2. ¬¬*r*

- ▶ We identified the main connective in (1) by thinking about how the formula was built up from simpler formulas.
- ► The main connective is the connective that is added last when forming complex formulas from simple ones and closing everything up with brackets, where applicable.
- ▶ Identify the main connective in these formulas:
 - 1. $\neg q$
 - 2. ¬¬*r*
 - 3. $(\neg\neg\neg p \lor q)$

- ▶ We identified the main connective in (1) by thinking about how the formula was built up from simpler formulas.
- ► The main connective is the connective that is added last when forming complex formulas from simple ones and closing everything up with brackets, where applicable.
- ▶ Identify the main connective in these formulas:
 - 1. $\neg q$
 - 2. ¬¬*r*
 - 3. $(\neg\neg\neg p \lor q)$
 - 4. $(p \lor (r \lor \neg q))$

▶ We use sentential logic to represent the truth-functional sentential structure of a natural language such as English. This process of representing is called paraphrasing (also sometimes called translating) natural language into sentential logic.

- ▶ We use sentential logic to represent the truth-functional sentential structure of a natural language such as English. This process of representing is called paraphrasing (also sometimes called translating) natural language into sentential logic.
- Here are the steps for paraphrasing:

- ▶ We use sentential logic to represent the truth-functional sentential structure of a natural language such as English. This process of representing is called paraphrasing (also sometimes called translating) natural language into sentential logic.
- Here are the steps for paraphrasing:
 - 1. Identify the English sentence truth-functional connectives.

- ▶ We use sentential logic to represent the truth-functional sentential structure of a natural language such as English. This process of representing is called paraphrasing (also sometimes called translating) natural language into sentential logic.
- Here are the steps for paraphrasing:
 - 1. Identify the English sentence truth-functional connectives.
 - 2. Replace them with symbolic connectives.

- ▶ We use sentential logic to represent the truth-functional sentential structure of a natural language such as English. This process of representing is called paraphrasing (also sometimes called translating) natural language into sentential logic.
- Here are the steps for paraphrasing:
 - 1. Identify the English sentence truth-functional connectives.
 - 2. Replace them with symbolic connectives.
 - 3. Identify the simplest (atomic) sentential components of the English sentences.

- ▶ We use sentential logic to represent the truth-functional sentential structure of a natural language such as English. This process of representing is called paraphrasing (also sometimes called translating) natural language into sentential logic.
- Here are the steps for paraphrasing:
 - 1. Identify the English sentence truth-functional connectives.
 - 2. Replace them with symbolic connectives.
 - 3. Identify the simplest (atomic) sentential components of the English sentences.
 - 4. Replace each distinct component with a distinct sentence symbol.

- ▶ We use sentential logic to represent the truth-functional sentential structure of a natural language such as English. This process of representing is called paraphrasing (also sometimes called translating) natural language into sentential logic.
- ► Here are the steps for paraphrasing:
 - 1. Identify the English sentence truth-functional connectives.
 - 2. Replace them with symbolic connectives.
 - Identify the simplest (atomic) sentential components of the English sentences.
 - 4. Replace each distinct component with a distinct sentence symbol.
 - 5. Use the structure of the English sentence to determine where the parentheses are to be placed.

- ▶ We use sentential logic to represent the truth-functional sentential structure of a natural language such as English. This process of representing is called paraphrasing (also sometimes called translating) natural language into sentential logic.
- ► Here are the steps for paraphrasing:
 - 1. Identify the English sentence truth-functional connectives.
 - 2. Replace them with symbolic connectives.
 - 3. Identify the simplest (atomic) sentential components of the English sentences.
 - 4. Replace each distinct component with a distinct sentence symbol.
 - 5. Use the structure of the English sentence to determine where the parentheses are to be placed.
- ➤ Since the focus of this class is not sentential logic, we will not spend a lot of time doing these paraphrases in class. But! See Homework 1.

Once we have paraphrased the sentences of natural language, the next step is to specify an interpretation of each paraphrase.

- Once we have paraphrased the sentences of natural language, the next step is to specify an interpretation of each paraphrase.
- ▶ Specifying an interpretation takes us from the syntax of sentential logic to its semantics, i.e., the content of the symbols or what the symbols mean.

- Once we have paraphrased the sentences of natural language, the next step is to specify an interpretation of each paraphrase.
- Specifying an interpretation takes us from the syntax of sentential logic to its semantics, i.e., the content of the symbols or what the symbols mean.
- ➤ Since we are interested in the truth-functional structure of the sentences in natural language, specifying an interpretation of our symbols will involve truth functions (also called "truth value assignments").

- Once we have paraphrased the sentences of natural language, the next step is to specify an interpretation of each paraphrase.
- Specifying an interpretation takes us from the syntax of sentential logic to its semantics, i.e., the content of the symbols or what the symbols mean.
- Since we are interested in the truth-functional structure of the sentences in natural language, specifying an interpretation of our symbols will involve truth functions (also called "truth value assignments").
- ➤ To understand what truth functions are, remember that we agreed that sentences in sentential logic will be unambiguously true or false (not both).

- Once we have paraphrased the sentences of natural language, the next step is to specify an interpretation of each paraphrase.
- Specifying an interpretation takes us from the syntax of sentential logic to its semantics, i.e., the content of the symbols or what the symbols mean.
- Since we are interested in the truth-functional structure of the sentences in natural language, specifying an interpretation of our symbols will involve truth functions (also called "truth value assignments").
- ➤ To understand what truth functions are, remember that we agreed that sentences in sentential logic will be unambiguously true or false (not both).
- So truth functions will take as input the truth value (true or false) of the paraphrased sentences in sentential logic and return as output a unique a truth value (true or false).

► Let us use the symbol 1 to represent the truth value "True" and the symbol 0 to represent the truth value "False".

- ► Let us use the symbol 1 to represent the truth value "True" and the symbol 0 to represent the truth value "False".
- Other authors use *T* and *F* for these truth values. But remember that these are just formal symbols, which we can use to mean anything we want. There is no difference between using *T* for "True" as opposed to 1 for "True".

- ► Let us use the symbol 1 to represent the truth value "True" and the symbol 0 to represent the truth value "False".
- ▶ Other authors use *T* and *F* for these truth values. But remember that these are just formal symbols, which we can use to mean anything we want. There is no difference between using *T* for "True" as opposed to 1 for "True".
- But! We shall see that using 0 and 1 will help us see the difference between deductive logic and inductive logic more clearly.

➤ Truth tables are a useful way to visualize how truth functions work.

- Truth tables are a useful way to visualize how truth functions work.
- ► From a truth table, one can see the unique output of a truth function for all of the possible inputs.

- Truth tables are a useful way to visualize how truth functions work.
- From a truth table, one can see the unique output of a truth function for all of the possible inputs.
- ▶ Here is the truth table for the simplest truth function. This is the truth function takes the truth value of *p* and returns its truth value.

р	
1	1
0	0

▶ Here is the truth table for the truth function that takes the truth value of p and returns the truth value of $\neg p$.

р	$\neg p$
1	0
0	1

▶ Here is the truth table for the truth function that takes the truth value of p and returns the truth value of $\neg p$.

p	$\neg p$
1	0
0	1

▶ Here is the truth table for the truth function that takes the truth value of p and q and returns the truth value of $(p \lor q)$.

р	q	$(p \lor q)$
1	1	1
1	0	1
0	1	1
0	0	0

▶ Here is the truth table for the truth function that takes the truth value of p and returns the truth value of $\neg p$.

p	$\neg p$
1	0
0	1

▶ Here is the truth table for the truth function that takes the truth value of p and q and returns the truth value of $(p \lor q)$.

р	q	$(p \lor q)$
1	1	1
1	0	1
0	1	1
0	0	0

► The simplicity of the idea of truth tables, understates just how very very powerful truth tables are!

▶ Here are the steps for constructing a truth table.

- ► Here are the steps for constructing a truth table.
 - 1. Identify all the simple sentences in the paraphrase.

- Here are the steps for constructing a truth table.
 - 1. Identify all the simple sentences in the paraphrase.
 - 2. Assign all the possible truth values to these sentences.

- Here are the steps for constructing a truth table.
 - 1. Identify all the simple sentences in the paraphrase.
 - 2. Assign all the possible truth values to these sentences.
 - 3. Use **known** truth functions to write down the truth value of the other **unknown** truth functions.

- Here are the steps for constructing a truth table.
 - 1. Identify all the simple sentences in the paraphrase.
 - 2. Assign all the possible truth values to these sentences.
 - Use known truth functions to write down the truth value of the other unknown truth functions.
 - 4. The truth value of the entire formula will be assigned under the **main connective** using the truth values from step 3.

- ▶ Here are the steps for constructing a truth table.
 - 1. Identify all the simple sentences in the paraphrase.
 - 2. Assign all the possible truth values to these sentences.
 - Use known truth functions to write down the truth value of the other unknown truth functions.
 - 4. The truth value of the entire formula will be assigned under the **main connective** using the truth values from step 3.
- Use truth tables to determine the truth value of the following formulas:

- ▶ Here are the steps for constructing a truth table.
 - 1. Identify all the simple sentences in the paraphrase.
 - 2. Assign all the possible truth values to these sentences.
 - Use known truth functions to write down the truth value of the other unknown truth functions.
 - 4. The truth value of the entire formula will be assigned under the **main connective** using the truth values from step 3.
- Use truth tables to determine the truth value of the following formulas:
 - 1. $(\neg p \lor q)$

- Here are the steps for constructing a truth table.
 - 1. Identify all the simple sentences in the paraphrase.
 - 2. Assign all the possible truth values to these sentences.
 - Use known truth functions to write down the truth value of the other unknown truth functions.
 - 4. The truth value of the entire formula will be assigned under the **main connective** using the truth values from step 3.
- Use truth tables to determine the truth value of the following formulas:
 - 1. $(\neg p \lor q)$
 - 2. $\neg(\neg p \lor \neg q)$

- Here are the steps for constructing a truth table.
 - 1. Identify all the simple sentences in the paraphrase.
 - 2. Assign all the possible truth values to these sentences.
 - Use known truth functions to write down the truth value of the other unknown truth functions.
 - 4. The truth value of the entire formula will be assigned under the **main connective** using the truth values from step 3.
- Use truth tables to determine the truth value of the following formulas:
 - 1. $(\neg p \lor q)$
 - 2. $\neg(\neg p \lor \neg q)$
- ► The focus of this class will not be on truth tables. But! See Homework 1.

► Truth tables can be useful because they provide a quick way for checking certain properties of the formulas of sentential logic.

- ► Truth tables can be useful because they provide a quick way for checking certain properties of the formulas of sentential logic.
- ▶ The properties that we will be interested in are: the property of being a tautology, the property of being contradictory, and the property of two formulas being truth-functionally equivalent.

- Truth tables can be useful because they provide a quick way for checking certain properties of the formulas of sentential logic.
- The properties that we will be interested in are: the property of being a tautology, the property of being contradictory, and the property of two formulas being truth-functionally equivalent.
- ▶ A formula is a tautology if its truth function always returns the value 1. If a complex formula is a tautology, then it always has the value 1 under every truth-value assignment to its component sentences.

- ► Truth tables can be useful because they provide a quick way for checking certain properties of the formulas of sentential logic.
- ► The properties that we will be interested in are: the property of being a tautology, the property of being contradictory, and the property of two formulas being truth-functionally equivalent.
- ➤ A formula is a tautology if its truth function always returns the value 1. If a complex formula is a tautology, then it always has the value 1 under every truth-value assignment to its component sentences.
- ▶ A formula is contradictory if its truth function always returns the value 0. If a complex formula is a tautology, then it always has the value 0 under every truth-value assignment to its component sentences.

- Truth tables can be useful because they provide a quick way for checking certain properties of the formulas of sentential logic.
- ► The properties that we will be interested in are: the property of being a tautology, the property of being contradictory, and the property of two formulas being truth-functionally equivalent.
- ➤ A formula is a tautology if its truth function always returns the value 1. If a complex formula is a tautology, then it always has the value 1 under every truth-value assignment to its component sentences.
- ▶ A formula is contradictory if its truth function always returns the value 0. If a complex formula is a tautology, then it always has the value 0 under every truth-value assignment to its component sentences.
- ► Two formulas are truth-functionally equivalent if their truth functions are identical.

▶ Use truth tables to verify that (1) $(\neg p \lor p)$ is a tautology and (2) $(\neg \neg p \lor \neg p)$ is a tautology.

- Use truth tables to verify that (1) $(\neg p \lor p)$ is a tautology and (2) $(\neg \neg p \lor \neg p)$ is a tautology.
- Use truth tables to verify that $(1) \neg (\neg p \lor p)$ is contradictory and $(2) \neg (\neg \neg p \lor \neg p)$ is contradictory.

- Use truth tables to verify that (1) $(\neg p \lor p)$ is a tautology and (2) $(\neg \neg p \lor \neg p)$ is a tautology.
- Use truth tables to verify that $(1) \neg (\neg p \lor p)$ is contradictory and $(2) \neg (\neg \neg p \lor \neg p)$ is contradictory.
- ▶ Use truth tables to verify that $(\neg p \lor p)$ is truth-functionally equivalent to $(\neg \neg p \lor \neg p)$.

- Use truth tables to verify that (1) $(\neg p \lor p)$ is a tautology and (2) $(\neg \neg p \lor \neg p)$ is a tautology.
- Use truth tables to verify that $(1) \neg (\neg p \lor p)$ is contradictory and $(2) \neg (\neg \neg p \lor \neg p)$ is contradictory.
- Use truth tables to verify that $(\neg p \lor p)$ is truth-functionally equivalent to $(\neg \neg p \lor \neg p)$.
- See Homework 1 for more exercises.