

TRINITY COLLEGE DUBLIN THE UNIVERSITY OF DUBLIN

Faculty of Engineering, Mathematics and Science

School of Computer Science & Statistics

Integrated Computer Science Programme
B.A. (Mod.) Business and Computing

Year 3 Annual Examinations

Trinity Term 2015

Computational Mathematics

Fergal Shevlin

Friday May 15th 2015

Sports Centre

14:00–16:00h

Instructions to Candidates:

Answer **both** questions in **Part A** and **two** questions out of three in **Part B**.
All questions carry equal marks.

Suggestion: Take 20 minutes to read all five questions. This leaves
100 minutes for answering questions worth 100 marks in total. So if a part of a
question is worth five marks, spend five minutes answering it.

Materials permitted for this examination:

Mathematical formulae and tables booklet—available from the invigilators.

Graph paper—available from the invigilators.

Non-programmable calculator—indicate make and model.

Part A

- Question 1. (i) In the IEEE 754 standards for *half*, *single* and *double* precision binary floating point number systems, how many binary digits are specified for sign p , significand s and exponent e ? [6 marks]
- (ii) Calculate (in decimal) the approximate minimum non-zero positive values that can be expressed in the above *half* and *single* systems with an exponent bias of $2^{e-1} - 1$, when the numbers are:
(a) normalized; (b) not normalized. [8 marks]
- (iii) Write a program fragment to plot the differences between division by 10_{10} and multiplication by 0.1_{10} , implemented as double precision floating point numbers, for a subset of the range of integers between 1×10^0 and 1×10^{30} . Explain your choice of starting value and sampling interval such that the program is relatively efficient. Assume an appropriate plotting method is available. [6 marks]
- (iv) Derive an expression for the relative error of subtraction. [5 marks]

- Question 2. (i) Use the simple iterative method to find the root of the polynomial $f(x) = x^3 - 30x^2 + 2552$ in the vicinity of $x = 11.8620$ to four decimal places of precision. [10 marks]
- (ii) What can $\psi'(\omega)$, the slope of the iteration function at the actual root, tell us about the likelihood of convergence when the initial approximation is close to ω ? Show how the condition is derived. Can we be certain about it? [10 marks]
- (iii) How can a root-finding problem be solved as an optimization problem? Write a program fragment which uses an optimization method to find the root of $f(x)$ specified above, assuming the appropriate optimization method is available. [5 marks]

Part B

- Question 3. (i) For a system of equations $F_i(x_j) = 0$ where $i = 1 \dots m$ and $j = 1 \dots n$, which are *non-linear* in the unknowns x_j , explain an approach through which a solution can be approximated iteratively.
[13 marks]
- (ii) How might it be possible to determine that a reasonable solution for the above will be found in advance of iterative approximation?
[5 marks]
- (iii) Write a program fragment which finds an approximate solution for a non-linear system of equations. Assume an appropriate method is available.
[7 marks]
- Question 4. (i) Specify and name quadrature formulae of polynomial degrees 1, 2 and 3 that can be used in numerical integration.
[9 marks]
- (ii) Use the composite trapezoidal rule to numerically integrate $\int_0^1 e^x dx$ with intervals $h_0 = 1, h_1 = \frac{h_0}{2}, h_2 = \frac{h_1}{2}$. Note the true solution is $e - 1$.
[8 marks]
- (iii) Combine the above estimates using Richardson's deferred approach to the limit with h^2 extrapolation.
[8 marks]
- Question 5. (i) Describe in detail some parameters which should be passed to a method used to solve a system of ordinary differential equations.
[15 marks]
- (ii) Describe in detail some values which should be returned from a method used to solve a system of ordinary differential equations.
[10 marks]