
Deep Orthogonal Decomposition for surrogate modelling of time dependent PDEs

written and submitted by DAWID KOTOWSKI

A thesis presented for the degree of
Master of Science

Supervisor: PROF. DR. MARIO OHLBERGER
Faculty of Mathematics and Computer Science
University Münster
Germany, DE

submitted on September 17, 2025

Contents

0.1	Notation	1
1	Introduction	3
1.1	Time Dependent PDEs	3
1.2	Neural networks	4
1.3	Motivation	4
1.4	Literature review	4
1.5	Outline	4
2	Theoretical Background	5
2.1	Linear Algebra	5
2.2	Parametric Partial Differential Equations	7
2.2.1	Sobolev Spaces	7
2.2.2	Reduced Basis Framework	8
2.3	Neural Networks	11
2.3.1	Network concatenations	13
2.3.2	Network parallelization	14
2.3.3	Standard Neural Networks	15
2.3.4	Approximation Theory	15
2.3.5	Statistical Learning	24
3	Formulation of the General Problem	27
3.1	Intrusive Reduced Order Models	30
3.1.1	Galerkin Proper Orthogonal Decomposition	31
3.1.2	Greedy Proper Orthogonal Decomposition	32
3.1.3	Non-linear Reduced Order Models	33
3.2	Data-driven Reduced Order Models	33
3.2.1	Proper Orthogonal Decomposition Deep Learning ROM	35
3.2.2	Deep Orthogonal Decomposition based ROMs	36
3.2.3	CoLoRA-DL-ROM	38
4	Approximation Theory for Reduced Order Models	39
4.1	Error Decomposition Formulas	40
4.2	Upper Complexity Bounds	51
4.3	Comparison between existing Techniques	58
4.3.1	Comparison between POD-based and DOD-based ROMs	59
5	Numerical Experiments	61
6	Conclusion	63

Todo list

Write an Introduction	3
Introduce KnW and its relation to the continous case	8
include theorem 6.1 from quarteroni2015rb	10
Motivate this with some example.	12
maybe include proof?	14
maybe include proof?	15
Why?	16
argue this	21
Definitely add traditional convergence and computational cost.	27
Add some literature	33
add a justification of the use of the reduced space and euklidian norm as done in franco2024	36
Introduce CoLoRA DL-ROM	38
include existence of s	45
do something about the delta and varepsilon definitions	58
add quick complexity bound for the dod+dfnn architecture	58
Include all numerical experiments	61
Find increase of performance depending on latent dimension	61
Give benchmark on pass forward times	61
Discuss comparison between different DL-ROMs	61
Discuss results	63
Insert outlook for future ideas and implementation	63

0.1 Notation

We set $\mathbb{N} := \{1, 2, \dots\}$, $\mathbb{N}_0 := \{0, 1, \dots\}$ and $\mathbb{N}_{\geq n} := \{n, n+1, \dots\}$ for $n \in \mathbb{N}$. Next the set $\mathbb{R}^+ := \{x \in \mathbb{R} \mid x \geq 0\}$ will define all real numbers, which are non-negative. Furthermore for any finite set A , we will write $|A|$ to denote its cardinality. However, if $A \subset \mathbb{R}^n$ measurable for some $n \in \mathbb{N}$, we will use $|A| := \lambda(A)$, i.e. as the Lebesgue measure. If $x \in \mathbb{R}$, the gaussian ceiling function is given via $\lceil x \rceil := \min\{k \in \mathbb{Z} : k \geq x\}$.

Further if $d \in \mathbb{N}$ and $\|\cdot\|$ is a norm, then for $x \in \mathbb{R}^d$ and $r > 0$ we set $B_{r, \|\cdot\|}(x)$ as the open ball around x with radius r . Additionally $|\cdot|$ denotes the regular euclidean norm on $\mathbb{R}^d$. We usually do not differ between vectors and scalars in ubiquitous notation.

The transpose of a matrix $A \in \mathbb{R}^{d_1 \times d_2}$ for some $d_1, d_2 \in \mathbb{N}$ is given by $A^T \in \mathbb{R}^{d_2 \times d_1}$. The identity matrix will be denoted with $\mathbb{I}_d := \text{diag}(1, \dots, 1) \in \mathbb{R}^{d \times d}$ for any $d \in \mathbb{N}$. Sometimes we abbreviate $\mathbb{I} = \mathbb{I}_d$, when the dimensionality is clear.

Let $(a_n)_{n \in \mathbb{N}} \in \mathbb{R}$ be some sequence, then we say $(a_n)_n$ has (*convergence of*) *order* n , if there is $c \in \mathbb{R}$ independent of n , s.t. $a_n = c \cdot n$ for all $n \in \mathbb{N}$. We then denote $a_n = \mathcal{O}(n)$.

When writing $a \ll b$ for $a, b \in \mathbb{R}$, the reader can infer, that $a < b$, but also, that the difference is "substantial". We do not provide a rigorous definition of this, even though some literature does on occasion.

Moreover, we define the *multiindex* notation as follows: For $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}_0^d$, we write $|\alpha| := \alpha_1 + \dots + \alpha_d$ and $\alpha! := \alpha_1! \dots \alpha_d!$. If $x \in \mathbb{R}^d$, then we set

$$x^\alpha := \prod_{i=1}^d x_i^{\alpha_i}.$$

Partial Differential Equations For a function $f : X \rightarrow Y$ we denote $\text{supp}(f) := \{x \in X \mid f(x) \neq 0\}$ as the *support* of f . Let $g : Y \rightarrow Z$ be another function, then $g \circ f : X \rightarrow Z$ is their *composition*. Setting the standard topology of $\mathbb{R}^d$ lets us define $\overline{A}$ for some $A \subset \mathbb{R}^d$ as the *closure* of A , and ∂A its *boundary*. If A is non-empty, we set $\text{diam}(A) := \sup_{x, y \in A} |x - y|$ as the diameter of A .

Let $\Omega \subset \mathbb{R}^d$ be open. For $n \in \mathbb{N}_0 \cup \{\infty\}$, we denote by $C^n(\Omega)$ the set of n -times continuously differentiable functions on Ω .

For $f \in C^n(\Omega)$ and $\alpha \in \mathbb{N}_0^d$ with $|\alpha| \leq n$ we write

$$D^\alpha f := \frac{\partial^{|\alpha|} f}{\partial x_1^{\alpha_1} \partial x_2^{\alpha_2} \dots \partial x_d^{\alpha_d}}.$$

A function $f : \Omega \rightarrow \mathbb{R}^m$ for $\Omega \subset \mathbb{R}^d$ is called *Lipschitz continuous*, if there is a constant $L \geq 0$ such that

$$|f(x) - f(y)| \leq L|x - y|.$$

We further denote with

$$L^p(\Omega) := \left\{ f : \Omega \rightarrow \mathbb{R} \mid f \text{ is measurable, and } \int_{\Omega} |f(x)|^p d\mu(x) < \infty \right\}$$

the Lebesgue space and L_{loc}^p as the space of $L^p(\Omega)$ functions with local support. If not specified otherwise we write $\|\cdot\|_p := \|\cdot\|_{L^p(\Omega)}$ for all $1 \leq p \leq \infty$. Note that in critical situations we tend to be explicit about the norms.

Neural Networks Consider $d_1, d_2 \in \mathbb{N}$ and a matrix $A \in \mathbb{R}^{d_1 \times d_2}$, then the number of non-zero entries of A is given by

$$\|A\|_{\ell^0} := |\{(i, j) : A_{i,j} \neq 0\}|.$$

Let $B \in \mathbb{R}^{d_1 \times d_3}$ be some other matrix for $d_3 \in \mathbb{N}$. Then we write

$$\begin{bmatrix} A & B \end{bmatrix} \in \mathbb{R}^{d_1 \times (d_2 + d_3)} \quad \text{or} \quad \begin{bmatrix} A & | & B \end{bmatrix} \in \mathbb{R}^{d_1 \times (d_2 + d_3)},$$

for the stacking of matrices, where the latter notation is meant to symbolize a strong delineation between blocks. Similar applies to vertical stacking, if $A \in \mathbb{R}^{d_1 \times d_2}$ and $B \in \mathbb{R}^{d_3 \times d_2}$. As a special case, this applies to vectors as well.

Chapter 1

Introduction

Write an Introduction

1.1 Time Dependent PDEs

The study and analysis of partial differential equations (PDEs) began with the works of Euler, d'Alembert, Lagrange and Laplace (see (Brezis & Browder, 1998) for this introduction). As so often in mathematics it started off with a set of simple problems, such as the vibrating string in one dimension as modelled by d'Alembert (1752)

$$\frac{\partial^2}{\partial t^2}u(x, t) = \frac{\partial^2}{\partial x^2}u(x, t), \quad \text{for } u \in C^2(\mathbb{R} \times \mathbb{R}^+), x \in \mathbb{R} \text{ and } t \in \mathbb{R}^+,$$

the extension to three dimensions by Euler (1759) and Bernoulli (1762), the Laplace equation for gravitational fields (1780)

$$\Delta u(x) = 0, \quad \text{for } u \in C^2(\mathbb{R}^3) \text{ and } x \in \mathbb{R}^3,$$

and finally the heat equation, as introduced by Fourier (1810 - 1822)

$$\frac{\partial}{\partial t}u(x, t) = \Delta u(x, t), \quad \text{for } u \in C^2(\mathbb{R}^3 \times \mathbb{R}^+), x \in \mathbb{R}^3 \text{ and } t \in \mathbb{R}^+.$$

As any posed problem calls for a solution to it, there have been simultaneous developments with regard to the calculation methods; the method of separation of variables was given by Euler and d'Alembert for the study of spherical harmonics and Fourier introduced similar ideas for the heat equation. For the Laplace equation, the Green functions were introduced in 1835. Green also discovered an important principle for solvability of the (later so called) Dirichlet problem

$$\begin{cases} \Delta u(x) = 0, & \text{for } x \in \Omega \subset \mathbb{R}^3, \\ u(x) = g(x), & \text{for } x \in \partial\Omega, \end{cases}$$

for some function $g \in C^0(\mathbb{R}^3)$, which revolved around minimizing an energy functional $E : C^2(\mathbb{R}^3) \rightarrow \mathbb{R}^+$, thus founding the variational principle.

Though these were first attempts at solving such problems, the idea of Green in particular led to a few analytical problems, by the lack of mathematical rigor. Together with the contribution of Cauchy on the *initial value* problem, this laid the groundwork for the field of *Existence Theory*, which tries to establish existence results for PDEs.

Notable here are the alternating method of Schwarz (1870) using the idea of domain decomposition and especially Neumanns method of integral equations developed in 1877, which would go on to be the starting point for the weak form and subsequently the Sobolev spaces. These were a necessity after B. Levi showed in 1906, that a general minimizing sequence for the Dirichlet integral was a Cauchy sequence and would therefore converge in the right space, which was then given due to the Sobolev embedding theorems.

To tackle the problem arising from "exotic" examples and the complicated nature of some PDEs, Hadamard proposed a classification of the most common linear PDEs using its characteristic polynomial, which is given by replacing all partial derivatives $\partial/\partial x_j$ by the algebraic variable ξ_j . In that case the PDE is called

1. *elliptic*, if after a change of variables the polynomial can be written as $\xi_1^2 + \xi_2^2 + \dots + \xi_n^2$,
2. *parabolic*, if after a change of variables the polynomial can be written as $\xi_1 + \xi_2^2 + \dots + \xi_n^2$,
3. *hyperbolic*, if after a change of variables the polynomial can be written as $\xi_1^2 - (\xi_2^2 + \dots + \xi_n^2)$.

These analytical results can be expanded a lot more, but what is most interesting in terms of this paper, are the Sobolev spaces. They, and the closely connected Bochner spaces, are the relevant solution spaces to the limits of all, in some sense, consistent numerical schemes, since the solutions are at least embedded in them. This follows quite natural, as any numerical scheme is basically bounded by the necessity of being ran on a computer and thus having a finite dimensional space it defines its approximation on, and at the very least the Sobolev and Bochner spaces admit any piecewise smooth function.

1.2 Neural networks

1.3 Motivation

1.4 Literature review

1.5 Outline

Chapter 2

Theoretical Background

2.1 Linear Algebra

For this section, we will specifically set up the euclidian (Hilbert) space $(\mathbb{R}^{N_h}, \langle \cdot, \cdot \rangle_{\mathbb{G}})$, with the inner product

$$\langle u, v \rangle_{\mathbb{G}} := u^T \mathbb{G} v,$$

for some symmetric positive definite $\mathbb{G} \in \mathbb{R}^{N_h \times N_h}$ for each $u, v \in \mathbb{R}^{N_h}$ and $\|\cdot\|$ its induced norm. As preparation for the upcoming Bauer Fike Theorem, we further want to define the matrix norm for the matrix space $\mathbb{R}^{N_h \times N}$, where $N \leq N_h$, as follows

$$\|A\|_{\text{op}} := \sqrt{\lambda_{\max}(A^T \mathbb{G} A)},$$

for $\lambda_{\max}(A^T \mathbb{G} A)$ being the maximal spectral radius of $A^T \mathbb{G} A$, which exists because $A^T \mathbb{G} A$ is symmetric and positive semi definite and hence subject to the spectral theorem. This norm satisfies the wanted property mentioned in the original paper (Bauer & Fike, 1960, p. 137-141); any operator norm such that

$$\|Ax\|^2 = x^T A^T \mathbb{G} A x \leq \lambda_{\max}(A^T \mathbb{G} A) x^T x \leq \|A\|_{\text{op}}^2 |x|^2$$

for all $x \in \mathbb{R}^N$.

Theorem 2.1 (Bauer-Fike Theorem). *Let $K \in \mathbb{R}^{N_h \times N_h}$ be some diagonalizable matrices with σ_k eigenvalues and $V = (\xi_1, \dots, \xi_N) \in \mathbb{R}^{N_h \times N_h}$ be the stacked eigenvectors matrix of K and $\delta K \in \mathbb{R}^{N_h \times N_h}$ some symmetric matrix. Then if λ_k are the eigenvalues of $K + \delta K$, we have*

$$\min_k |\lambda_k - \sigma_k| \leq \kappa(V) \|(K + \delta K) - K\|_{\text{op}},$$

where $\kappa(V) = \|V\|_{\text{op}} \|V^{-1}\|_{\text{op}}$.

Proof. See (Bauer & Fike, 1960, p. 137-141). □

Furthermore, we will often discuss the notion of orthonormality with regard to a matrix. This can be understood with the following definition:

Definition 2.2 (Matrix Orthonormality). Let $\mathbb{G} \in \mathbb{R}^{N_h \times N_h}$ be a symmetric positive definite matrix for some $N_h \in \mathbb{N}$. We then say that any matrix $\mathbb{A} \in \mathbb{R}^{N_h \times N}$ is $\mathbb{G}$ -orthonormal for some $N \leq N_h$, if

$$\mathbb{I} = \mathbb{A}^T \mathbb{G} \mathbb{A}.$$

Remark 2.3. If some matrix $\mathbb{V} \in \mathbb{R}^{N_h \times N}$ can be written as $\mathbb{V} = \mathbb{A}\tilde{\mathbb{V}}$, where $\tilde{\mathbb{V}} \in \mathbb{R}^{N_A \times N}$ and $\mathbb{A} \in \mathbb{R}^{N_h \times N_A}$ being a $\mathbb{G}$ -orthonormal matrix, then $\mathbb{V}$ is $\mathbb{G}$ -orthonormal if and only if $\tilde{\mathbb{V}}$ is orthonormal in the euklidian sense. In fact,

$$\mathbb{V}^T \mathbb{G} \mathbb{V} = \tilde{\mathbb{V}}^T \mathbb{A}^T \mathbb{G} \mathbb{A} \tilde{\mathbb{V}} = \tilde{\mathbb{V}}^T \tilde{\mathbb{V}}.$$

Definition 2.4 (Orthogonal Projector). A operator $P : V \rightarrow V$, where V is a Hilbert space with inner product $\langle \cdot, \cdot \rangle_V$, is called an *orthogonal projector* onto $W \subset V$, if

1. $P^2 = P$,
2. $\text{Im}(P) = W$ and
3. $\langle Px, y \rangle_V = \langle x, Py \rangle_V$ for all $x, y \in V$.

Lemma 2.5. Let $(\mathbb{R}^{N_h}, \langle \cdot, \cdot \rangle_{\mathbb{G}})$ be the Hilbert space from above. If now $\mathbb{A} \in \mathbb{R}^{N_h \times N_A}$ is a $\mathbb{G}$ -orthonormal matrix, then

$$P := \mathbb{A}\mathbb{A}^T \mathbb{G}$$

is a orthogonal projector on $\text{span}(\mathbb{A})$ with regard to the $\mathbb{G}$ induced inner product. We refer to this as P being the $\mathbb{G}$ -orthonormal projector on $\text{span}(\mathbb{A})$.

Proof. It suffices to simply check the items of Definition (2.4); the image statement is instantaneously clear, since $\mathbb{A}\mathbb{A}^T \mathbb{G} = \mathbb{A}(\mathbb{A}^T \mathbb{G})$. The idempotency can be checked via

$$P^2 = (\mathbb{A}\mathbb{A}^T \mathbb{G})(\mathbb{A}\mathbb{A}^T \mathbb{G}) = \mathbb{A}(\mathbb{A}^T \mathbb{G} \mathbb{A}) \mathbb{A}^T \mathbb{G} = \mathbb{A}\mathbb{A}^T \mathbb{G} = P.$$

Finally set $u, v \in \mathbb{R}^{N_h}$ arbitrarily, then

$$\begin{aligned} \langle Pu, v \rangle_{\mathbb{G}} &= (Pu)^T \mathbb{G} v \\ &= u^T P^T \mathbb{G} v \\ &= u^T (\mathbb{A}\mathbb{A}^T \mathbb{G})^T v \\ &= u^T \mathbb{G} \mathbb{A} \mathbb{A}^T \mathbb{G} v \\ &= u^T \mathbb{G} P v = \langle u, P v \rangle_{\mathbb{G}}. \end{aligned}$$

□

Lemma 2.6. Let $(\mathbb{R}^{N_h}, \langle \cdot, \cdot \rangle_{\mathbb{G}})$ be the Hilbert space from above. If now $\mathbb{A} \in \mathbb{R}^{N_h \times N_A}$ is a $\mathbb{G}$ -orthonormal matrix, then

$$\|\mathbb{A}\|_{\text{op}} = 1.$$

Proof. This is directly clear, since $\|\mathbb{A}\|_{\text{op}} = \sqrt{\lambda_{\max}(\mathbb{A}^T \mathbb{G} \mathbb{A})} = \sqrt{\lambda_{\max}(\mathbb{I})} = 1$. □

These preliminaries will play an important role, since we will use such a gramian $\mathbb{G}$ of an a priori chosen basis under a discretization of a PDE to set up a (linear) projector attaining any projection onto the solution manifold under the span of a $\mathbb{G}$ -orthonormal matrix. In fact, any projector

$$\mathbb{V}\mathbb{V}^T \mathbb{G} : \mathbb{R}^{N_h} \rightarrow \mathbb{R}^N$$

for a $\mathbb{G}$ -orthonormal matrix $\mathbb{V} \in \mathbb{R}^{N_h \times N}$ and some $N \leq N_h$ can be seen as a special case for the differential geometric framework of (nonlinear) projectors on manifolds concerning Reduced Order Modelling introduced and discussed in Buchfink et al. (2023).

2.2 Parametric Partial Differential Equations

2.2.1 Sobolev Spaces

For our work with Partial Differential Equation, and subsequent error approximations, we want to introduce Sobolev spaces. These are generalized differentiable spaces, where a weak derivative is given, as an analogon to the classic derivative in the $C^n(\Omega)$ space. They are employed in order to exploit their compact embedding in the Hölder spaces, which are generalizations of $C^n(\Omega)$. We will not go in much detail about the theory of Sobolev spaces, since we mostly concern ourselves with finite dimensional subspaces, but some ground truth must be discussed.

For this subsection we primary use (Alt, 2016), and for the first upcoming definition (Dacorogna, 2004).

Definition 2.7 (Lipschitz Domain). Let $d \in \mathbb{N}$, and $\Omega \subset \mathbb{R}^d$. Then Ω is called a *Lipschitz domain* if for every $x \in \partial\Omega$ and some $r > 0$, there is $U := B_{r,|\cdot|}(x) \subset \mathbb{R}^d$ and an isomorphic map $H : Q \rightarrow B_{r,|\cdot|}(x)$ with $Q := \{x \in \mathbb{R}^d \mid \|x\|_{\ell^1} < d\}$ such that

$$H \in C^k(\overline{Q}), \quad H^{-1} \in C^k(\overline{U}), \quad H(Q_+) = U \cap \Omega, \quad H(Q_0) = U \cap \partial\Omega$$

where $Q_+ := \{x \in Q \mid x_d > 0\}$ and similarly $Q_0 := \{x \in Q \mid x_d = 0\}$, and H is only in $C^{0,1}$.

From now on, let $d \in \mathbb{N}$ and, if not given otherwise, $\Omega \subset \mathbb{R}^d$ be a Lipschitz domain.

Definition 2.8 (Weak Derivative). Let $\alpha \in \mathbb{N}_0^d$ be a multiindex. A function $u \in L_{\text{loc}}^1(\Omega)$ is said to have an α -th weak derivative, if there is a function $v_\alpha \in L_{\text{loc}}^1(\Omega)$ such that

$$\int_{\Omega} u D^\alpha \varphi = (-1)^{|\alpha|} \int_{\Omega} v_\alpha \varphi \quad \forall \varphi \in C_0^\infty(\Omega).$$

We then write $v_\alpha = D^\alpha u$.

Definition 2.9 (Sobolev Space). Let $n \in \mathbb{N}_0$ and $1 \leq p \leq \infty$. Then we define the *Sobolev space* as

$$W^{n,p}(\Omega) := \{f \in L^p(\Omega) \mid \|f\|_{W^{n,p}(\Omega)} < \infty\},$$

where the norm on this space is given via

$$\|f\|_{W^{n,p}(\Omega)} := \begin{cases} \left(\sum_{|\alpha| \leq n} \|D^\alpha f\|_{L^p(\Omega)}^p \right)^{1/p}, & \text{if } p < \infty \\ \max_{|\alpha| \leq n} \|D^\alpha f\|_{L^\infty(\Omega)}, & \text{if } p = \infty, \end{cases}$$

for any $f \in W^{n,p}(\Omega)$.

It is possible to extend this definition for $0 < n < 1$ to that of the *Sobolev-Slobodeckij spaces*. In the analysis of partial differential equations these are used to characterize the regularity of Sobolev functions, which are restricted to the boundary $\partial\Omega$. This restriction itself is called a trace operator. One of the statements of relevance for such fractal Sobolev spaces is the following:

If $\Omega \subset \mathbb{R}^d$ is a regular open set (for example, if Ω is a Lipschitz domain), and if $u \in W^{1,p}(\Omega)$, then in fact $u \mapsto u|_{\partial\Omega}$ as a trace operator is surjective from $W^{1,p}(\Omega)$ onto $W^{1-(1/p),p}(\partial\Omega)$ and

$$\|u|_{\partial\Omega}\|_{W^{1-(1/p),p}(\partial\Omega)} \leq C \|u\|_{W^{1,p}(\Omega)},$$

for some $C > 0$ independent of p and u , see (Brezis, 2011, p. 315). This of course only makes sense in light of the next definition.

Definition 2.10 (Sobolev-Slobodeckij Space). Let $0 < s < 1$ and $1 \leq p \leq \infty$. Then we define the *Sobolev-Slobodeckij space* as

$$W^{s,p}(\Omega) := \{f \in L^p(\Omega) \mid \|f\|_{W^{s,p}(\Omega)} < \infty\},$$

where the norm on this space is given via

$$\|f\|_{W^{s,p}(\Omega)} := \begin{cases} \left(\|f\|_{L^p(\Omega)}^p + \int_{\Omega} \int_{\Omega} \left(\frac{|f(x) - f(y)|}{|x - y|^{s+d/p}} \right)^p dx dy \right)^{1/p}, & \text{if } p < \infty \\ \max \left\{ \|f\|_{L^\infty(\Omega)}, \operatorname{ess\,sup}_{x,y \in \Omega} \frac{|f(x) - f(y)|}{|x - y|^s} \right\}, & \text{if } p = \infty, \end{cases}$$

for any $f \in W^{s,p}(\Omega)$.

Furthermore, a very important theorem result will be the following theorem, translating Lipschitz continuity into Sobolev-regularity. This will allow us to employ the later introduced approximation theory for neural networks from the sole position of Lipschitz continuity.

Theorem 2.11. *Let $f : \Omega \rightarrow \mathbb{R}$ be a bounded and L -Lipschitz map, then $f \in W^{1,\infty}(\Omega)$, the weak gradient of f agrees almost everywhere with the classical gradient, and*

$$\|Df(x)\|_{L^\infty(\Omega)} \leq L.$$

Proof. See (Heinonen, 2005, Theorem 4.1) and (Heinonen, 2005, Remark 4.2). $\square$

2.2.2 Reduced Basis Framework

Introduce KnW and its relation to the continous case

For this subsection we define $u(\mu_j, t_k) \in \mathbb{R}^{N_h}$ for $(\mu_j)_{j=1}^{N_s} =: \mathcal{P} \subset \Theta$ and $(t_k)_{k=1}^{N_t} =: \mathcal{T} \subset [0, T]$ to be some vectors, later representing the solutions to a parametric PDE for a given parameter and time point in a given discrete subspace of the coefficient solution manifold. Here Θ represents an arbitrary parameter space, $T > 0$ some arbitrary time point, and $N_s, N_t \in \mathbb{N}$ be the parameter and time sample size respectively. Let the inner product on $\mathbb{R}^{N_h}$ be defined here, using a symmetric positive definite matrix $\mathbb{G} \in \mathbb{R}^{N_h \times N_h}$ via

$$\langle u, v \rangle_{\mathbb{G}} := u^T \mathbb{G} v, \quad \text{for all } u, v \in \mathbb{R}^{N_h},$$

and the corresponding norm $\|\cdot\| := \langle \cdot, \cdot \rangle_{\mathbb{G}}$. This is meant to result from a choice of an inner product stemming from the coefficient vectors $u(\mu, t)$ representing that space, in other words, given by the choice of a basis on the corresponding Hilbert space, when viewing $u(\mu, t) \in \mathbb{R}^{N_h}$ as the coefficients identifying an element in that space. It is assumed, that we sample the parameters and time points uniformly and iid from their respective spaces and that $\|u(\mu, t)\| < \infty$ for all $\mu \in \Theta$ and $t \in [0, T]$ in an essential sense. Let also

$$N_{\text{data}} := N_s \cdot N_t,$$

be the total sample size.

Note, that, with the discussion surrounding the Definition (2.2) of the $\mathbb{G}$ -orthonormality, taking the error of any such gram-sensitive projector will result in the possibility of switching to a low dimensional euklidian norm, if the matrix $\mathbb{V} \in \mathbb{R}^{N_h \times N}$ for some $N \leq N_h$ is $\mathbb{G}$ -orthonormal and can be expressed as $\mathbb{V} = \mathbb{A} \tilde{\mathbb{V}}$, where $\mathbb{A} \in \mathbb{R}^{N_A \times N}$ for $N \leq N_A \leq N_h$ is also $\mathbb{G}$ -orthonormal. This is quantified by the following result:

Proposition 2.12. Let $\mathbb{V} \in \mathbb{R}^{N_h \times N}$ for some $N \leq N_h$ be $\mathbb{G}$ -orthonormal and be written as

$$\mathbb{V} = \mathbb{A} \tilde{\mathbb{V}},$$

where $\mathbb{A} \in \mathbb{R}^{N_h \times N_A}$ is also $\mathbb{G}$ -orthonormal and $\tilde{\mathbb{V}} \in \mathbb{R}^{N_A \times N}$ for $N \leq N_A \leq N_h$. Then we have, for any $u \in \mathbb{R}^{N_h}$, that

$$\|u - \mathbb{V} \mathbb{V}^T \mathbb{G} u\|^2 = c_{\mathbb{A}} + |u_{\text{pre-red}} - \tilde{\mathbb{V}} \tilde{\mathbb{V}}^T u_{\text{pre-red}}|^2,$$

with $c_{\mathbb{A}} > 0$ being a constant only depending on $\mathbb{A}$ and $u_{\text{pre-red}} = \mathbb{A}^T \mathbb{G} u$.

Proof. By Lemma (2.5), we know that $\mathbb{A} \mathbb{A}^T \mathbb{G}$ is a $\mathbb{G}$ -orthonormal projector into the span of $\mathbb{A}$, and thus its residual $u - \mathbb{A} \mathbb{A}^T \mathbb{G} u$ must be $\mathbb{G}$ -orthogonal to all columns of $\mathbb{A}$, since we can decompose $u = u_{\perp} + u_{\parallel}$, where $u_{\parallel} \in \text{span}(\mathbb{A})$ and $u_{\perp} \in \mathbb{R}^{N_h} - \text{span}(\mathbb{A})$. This gives us the possibility to use the Pythagoras decomposition:

$$\begin{aligned} \|u - \mathbb{V} \mathbb{V}^T \mathbb{G} u\|^2 &= \|u - \mathbb{A} \tilde{\mathbb{V}} \tilde{\mathbb{V}}^T \mathbb{A}^T \mathbb{G} u\|^2 \\ &= \|u - \mathbb{A} \mathbb{A}^T \mathbb{G} u\|^2 + \|\mathbb{A} \mathbb{A}^T \mathbb{G} u - \mathbb{A} \tilde{\mathbb{V}} \tilde{\mathbb{V}}^T \mathbb{A}^T \mathbb{G} u\|^2 \\ &= \|u - \mathbb{A} \mathbb{A}^T \mathbb{G} u\|^2 + |\mathbb{A}^T \mathbb{G} u - \tilde{\mathbb{V}} \tilde{\mathbb{V}}^T \mathbb{A}^T \mathbb{G} u|^2, \end{aligned}$$

where the last step is justified by showing that for any $u \in \mathbb{R}^{N_A}$

$$\|\mathbb{A} u\| = \langle \mathbb{A} u, \mathbb{A} u \rangle_{\mathbb{G}} = (\mathbb{A} u)^T \mathbb{G} \mathbb{A} u = u^T \mathbb{A}^T \mathbb{G} \mathbb{A} u = u^T u = |u|.$$

Now setting $u_{\text{pre-red}} := \mathbb{A}^T \mathbb{G} u$ gives us the wanted statement. $\square$

This proposition might seem quite unimportant, but this will be the core idea giving us a possibility to assert closeness with regard to the representations of the solution rather than the, potentially uninformative, solution coefficient vectors in $\mathbb{R}^{N_h}$ under a computationally feasible norm (the euclidian N_A -dimensional norm).

Next, we will discuss prerequisites to the Proper Orthogonal Decomposition and related concepts, such as the Kolmogorov n -width, which lets us derive many key concepts and is an important quantity in the field of ROM methods. We base this subsection on (Quarteroni et al., 2015), but import the generalizations to arbitrary inner products from (Franco et al., 2024).

Definition 2.13 (Generalized Proper Orthogonal Decomposition). Assuming $N_{\text{data}} < \infty$ and let $\mathbb{G} \in \mathbb{R}^{N_h \times N_h}$ be a mass matrix, we define by $U_j = [u(\mu_j, t_k)]_{k=1}^{N_t} \in \mathbb{R}^{N_h \times N_t}$ the snapshot matrix for a given $\mu_j \in \mathcal{P}$ and consequently $U = [U_j]_{j=1}^{N_s} \in \mathbb{R}^{N_h \times N_{\text{data}}}$ the collective snapshot matrix. Then the (*discrete*) *correlation matrix* is given by

$$K = \frac{|\Theta \times [0, T]|}{N_{\text{data}}} U \mathbb{G} U^T \in \mathbb{R}^{N_h \times N_h}. \quad (2.1)$$

K is symmetric and semi-positive definite and thus has real and positive eigenvalues $\sigma^2 \geq \dots \geq \sigma_r^2$. We further call ψ_k^μ the eigenvectors of the correlation matrix K . With these, the corresponding POD basis of dimension $N < N_h$ is given via

$$\xi_k = \frac{1}{\sigma_k} U \psi_k, \quad 1 \leq k \leq N,$$

for the N largest eigenvalues and their corresponding eigenvectors. Then stacking these eigenvectors gives us the *POD matrix*

$$\mathbb{A} := [\xi_k]_{k=1}^N \in \mathbb{R}^{N_h \times N}.$$

This matrix is $\mathbb{G}$ -orthonormal by construction. Further, if $N_s, N_t = \infty$, we define the *continuous correlation matrix* via

$$K_\infty = \int_{\Theta \times [0, T]} u(\mu, t) \mathbb{G} u^T(\mu, t) d(\mu, t) \in \mathbb{R}^{N_h \times N_h}, \quad (2.2)$$

and its real eigenvalues by $\sigma_{k, \infty}^2$.

Algorithm 1 Generalized Proper Orthogonal Decomposition (POD)

- 1: **Input:** snapshot matrix $U = [U_j]_{j=1}^{N_s} \in \mathbb{R}^{N_h \times N_{\text{data}}}$, mass matrix $\mathbb{G} \in \mathbb{R}^{N_h \times N_h}$, reduction $N < N_h$
 - 2: **Output:** POD basis matrix $\mathbb{A}^\mu = [\xi_1^\mu, \dots, \xi_N^\mu] \in \mathbb{R}^{N_h \times N}$
 - 3: **if** $N_{\text{data}} \leq N_h$ **then**
 - 4: Form the correlation matrix $C = \frac{|\Theta \times [0, T]|}{N_{\text{data}}} U^T \mathbb{G} U \in \mathbb{R}^{N_{\text{data}} \times N_{\text{data}}}$
 - 5: Solve the eigenvalue problem $C \psi_i = \sigma_i^2 \psi_i$, $i = 1, \dots, r$
 - 6: Set $\xi_i = \frac{1}{\sigma_i} U \psi_i$, $i = 1, \dots, r$
 - 7: **else**
 - 8: Form the correlation matrix $K = \frac{|\Theta \times [0, T]|}{N_{\text{data}}} U \mathbb{G} U^T \in \mathbb{R}^{N_h \times N_h}$
 - 9: Solve the eigenvalue problem $K \xi_i = \sigma_i^2 \xi_i$, $i = 1, \dots, r$
 - 10: **end if**
 - 11: $\mathbb{A} \leftarrow \text{stack}(\xi_1, \dots, \xi_N)$
-

include theorem 6.1 from quarteroni2015rb

Proposition 2.14. Let K and K_∞ and their respective eigenvalues be defined as in Definition (2.13), then letting $N_s, N_t \rightarrow \infty$, we get

$$\sum_{k > N} \sigma_k^2 \longrightarrow \sum_{k > N} \sigma_{k, \infty}^2, \quad \text{a.s.}$$

Proof. Let $N_s, N_t < \infty$, conforming to the sampling assumption, and

$$M := \text{ess sup}_{(\mu, t) \in \Theta \times \mathcal{T}} \|u(\mu, t)\| < \infty$$

as before (see introduction in this Section (2.2.2)). Taking a variational approach on this finite dimensional space, we know, that there exists a $\mathbb{G}$ -orthonormal matrix $V_\infty \in \mathbb{R}^{N_h \times N}$ such that

$$\begin{aligned} \sum_{k > N} \sigma_{k, \infty}^2 &= \int_{\Theta \times [0, T]} \|u(\mu, t) - V_\infty V_\infty^T \mathbb{G} u(\mu, t)\|^2 d(\mu, t) \\ &= \min_{W \in \mathbb{R}^{N_h \times N} : W^T \mathbb{G} W = I} \int_{\mu \times \mathcal{T}} \|u(\mu, t) - W W^T \mathbb{G} u(\mu, t)\|^2 d(\mu, t). \end{aligned}$$

Thus, since this is the optimal choice of an approximation matrix for the continuous case and $\mathbb{A} \in \mathbb{R}^{N_h \times N}$ given by Definition (2.13) is only optimal in the discrete setting and not necessarily the continuous case, we have

$$\sum_{k > N} \sigma_{k, \infty}^2 \leq \int_{\Theta \times [0, T]} \|u(\mu, t) - \mathbb{A} \mathbb{A}^T \mathbb{G} u(\mu, t)\|^2 d(\mu, t) = \sum_{k > N} \sigma_k^2.$$

We then write entry-wise for $m, l = 1, \dots, N_h$

$$\begin{aligned} [K_\infty - K]_{ml} &:= \int_{\Theta \times [0, T]} [\mathbb{G}^{1/2} u(\mu, t)]_m [\mathbb{G}^{1/2} u(\mu, t)]_l d(\mu, t) \\ &\quad - \frac{|\Theta \times [0, T]|}{N_{data}} \sum_{j=1}^{N_s} \sum_{k=1}^{N_t} [\mathbb{G}^{1/2} u(\mu_j, t_k)]_m [\mathbb{G}^{1/2} u(\mu_j, t_k)]_l. \end{aligned}$$

By the uniform iid sampling assumed and the general bound on $\|u(\mu, t)\| \leq M$ before, we know that with the entry-wise boundedness of the PDE solutions, the resulting matrix is subject to the strong law of large numbers, and thus converges to zero, given the entry-wise matrix norm $\|\cdot\|_{\text{op}}$ as $N_t, N_s \rightarrow \infty$ almost surely.

Now using Bauer-Fike's Theorem (2.1) with the $\|\cdot\|_{\text{op}}$ -norm, we have upon ordering, that for each $\sigma_{k,\infty}^2$ there is a σ_k^2 in the spectrum of K , such that

$$|\sigma_k^2 - \sigma_{k,\infty}^2| \leq \kappa(\mathbb{A}_\infty) \|K - K_\infty\|_{\text{op}},$$

where $\kappa(\mathbb{A}_\infty)$ denotes the condition number of the stacked eigenvector matrix of K_∞ , i.e. $\mathbb{A}_\infty = (\xi_{1,\infty}, \dots, \xi_{N_h,\infty})$. This number is independent of N_s and N_t . Now by convergence of $\|K - K_\infty\|_{\text{op}} \rightarrow 0$ almost surely for $N_s, N_t \rightarrow \infty$, we conclude the proof. $\square$

Proposition 2.15. Let $\mathcal{V}_N = \{W \in \mathbb{R}^{N_h \times N} \mid W^T \mathbb{G} W = I_N\}$ be the set of all N -dimensional $\mathbb{G}$ -orthonormal matrices, $N_{data} := N_t N_s$ and $\mathbb{A} = [\xi]_{i=1}^N$ be the POD matrix according to Definition (2.13), then

$$\begin{aligned} &\frac{|\Theta \times [0, T]|}{N_{data}} \sum_{i \in I, k=1, \dots, N_t} \|u(\mu_j, t_k) - \mathbb{A} \mathbb{A}^T \mathbb{G} u(\mu_j, t_k)\|^2 \\ &= \min_{W \in \mathcal{V}_N} \frac{|\Theta \times [0, T]|}{N_{data}} \sum_{i \in I, k=1, \dots, N_t} \|u(\mu_j, t_k) - W W^T \mathbb{G} u(\mu_j, t_k)\|^2 = \sum_{j=N+1}^r \sigma_k^2. \end{aligned}$$

Definition 2.16 (Linear Kolmogorov N -width). Let $N \in \mathbb{N}$, $N \leq N_h$ and $\mathcal{S}_{N_h} = \{u(\mu, t) \in \mathbb{R}^{N_h} \mid (\mu, t) \in \Theta \times [0, T]\}$ be some parametric manifold. The linear Kolmogorov N -width of $\mathcal{S}_{N_h}$ is defined as

$$d_N(\mathcal{S}_{N_h}) = \inf_{V \in \mathbb{R}^{N_h \times N}} \sup_{u \in \mathcal{S}_{N_h}} \|u - V V^T \mathbb{G} u\|.$$

Definition 2.17 (Nonlinear Kolmogorov N -width). Let $N \in \mathbb{N}$, $N \leq N_h$ and $\mathcal{S}_{N_h} = \{u(\mu, t) \in \mathbb{R}^{N_h} \mid (\mu, t) \in \Theta \times [0, T]\}$ be some parametric manifold. The nonlinear Kolmogorov N -width of $\mathcal{S}_{N_h}$ is defined as

$$\delta_N(\mathcal{S}_{N_h}) = \inf_{\psi \in C(\mathbb{R}^{N_h}, \mathbb{R}^N), \psi' \in C(\mathbb{R}^N, \mathbb{R}^{N_h})} \sup_{u \in \mathcal{S}_{N_h}} |u - \psi(\psi'(u))|.$$

2.3 Neural Networks

In this section, we will introduce the general notion of neural networks. Specifically important for us will be concatenations of perceptrons, which are a specific subclass of networks, the so called *feedforward neural network*.

Definition 2.18 (Neural Network). Let $d, L \in \mathbb{N}$, then a *neural network* Φ with input dimension d and L layers is defined via

$$\Phi = ((A_1, b_1), (A_2, b_2), \dots, (A_L, b_L)),$$

such that $N_0 = d$ and $N_1, \dots, N_L \in \mathbb{N}$, where each A_l is an $N_l \times \sum_{k=0}^{l-1} N_k$ dimensional matrix, and $b_l \in \mathbb{R}^{N_l}$.

We further define $R_\rho(\Phi) : \mathbb{R}^d \rightarrow \mathbb{R}^{N_L}$ as a *realizations of Φ with activation function ρ* for some arbitrary map $\rho : \mathbb{R} \rightarrow \mathbb{R}$ as the result of the following scheme for some input $x \in \mathbb{R}^d$:

$$\begin{aligned} x_0 &:= x, \\ x_l &:= \rho \left(A_l \begin{bmatrix} x_0^T & \cdots & x_{l-1}^T \end{bmatrix}^T + b_l \right), \quad \text{for } l = 1, \dots, L-1, \\ x_L &:= A_L \begin{bmatrix} x_0^T & \cdots & x_{L-1}^T \end{bmatrix}^T + b_L, \end{aligned}$$

where ρ acts componentwise, i.e. $\rho(y) := [\rho(y_1), \dots, \rho(y_m)]^T \in \mathbb{R}^m$ for each $y = [y_1, \dots, y_m] \in \mathbb{R}^m$. Note that we can write

$$A_l = \begin{bmatrix} A_{l,x_0} & \cdots & A_{l,x_{l-1}} \end{bmatrix},$$

with A_{l,x_k} being a $N_l \times N_k$ dimensional matrix for $k = 0, \dots, l-1$ and $l = 1, \dots, L$. This comes in handy to portray the scheme via

$$\begin{aligned} x_l &:= \rho \left(A_{l,x_0}x_0 + \cdots + A_{l,x_{l-1}}x_{l-1} + b_l \right), \quad \text{for } l = 1, \dots, L-1, \\ x_L &:= A_{L,x_{L-1}}x_{L-1} + b_L. \end{aligned}$$

Moreover call $N_\Phi := d + \sum_{l=1}^L N_l$ the *number of neurons* of the network, $L_\Phi := L$ the *number of layers*, and finally $\omega_\Phi := \sum_{l=1}^L (\|A_l\|_{\ell^0} + \|b_l\|_{\ell^0})$ the *number of active weights*.

Remark 2.19. As a note, we can rewrite known neural network architectures, such as a convolutional autoencoder, as a neural network according to Definition (2.18), see also (Gühring et al., 2020, p. 22).

Motivate this with some example.

From this we can infer a simpler architecture, describing a network which omits any connection of non-neighboring layers.

Definition 2.20 (Standard Neural Network). If a neural network $\Phi = ((A_1, b_1), \dots, (A_L, b_L))$ now has the characteristic

$$A_{l,x_i} = 0, \quad \text{for } l = 1, \dots, L, \text{ and } i = 0, \dots, l-2,$$

then we call Φ a *standard neural network*.

When considering the similarities and differences between two distinct neural networks $\Phi_1 = ((A_1, b_1), \dots, (A_L, b_L))$ and $\Phi_2 = ((\hat{A}_1, \hat{b}_1), \dots, (\hat{A}_L, \hat{b}_L))$, one can easily see, that if $[A_1]_{ij} = 0$, while $[\hat{A}_1]_{ij} \neq 0$ for some i, j , then these networks are severely different, whereas if they only differ in the specific non-zero weight, they could be recovered from each other through training. This inspires the definition of an *architecture of a network*.

Definition 2.21 (Neural Network Architecture). For some $d, L \in \mathbb{N}$, we define a *neural network architecture $\mathcal{A}$ with input dimension d and L layers* to be a binary neural network, i.e. for all entries of $A_i, b_i \in \{0, 1\}$ for $i = 1, \dots, L$ of $\mathcal{A}$.

We further say that a *neural network $\Phi = ((\hat{A}_1, \hat{b}_1), \dots, (\hat{A}_L, \hat{b}_L))$ has architecture $\mathcal{A}$* , if

1. $N_l(\Phi) = N_l$ for all $l = 1, \dots, L$, and

2. $[\hat{A}_l]_{ij} \neq 0$ implies $[A_l]_{ij} \neq 0$ for all $l = 1, \dots, L$ with $i = 1, \dots, N_l$ and $j = 1, \dots, \sum_{k=0}^{l-1} N_k$.

Since the associated realization of a neural network $R_\rho(\Phi)$ requires the definition of ρ , the activation function, we will define the most commonly used one in literature. It will help a lot with some of the constructions, since it has a simple nature, but also provides the network with the necessary non-linearity.

Definition 2.22 (ReLU). The function

$$\rho : \mathbb{R} \rightarrow \mathbb{R}; \quad x \mapsto \max(0, x),$$

is called *ReLU* (Rectified Linear Unit) *activation function*.

Given such an activation function, Kurt Hornik provided in 1990 the first contribution to a field, which would emerge centuries later: Approximation theory. He proved for any $L^p(\mu)$ function, for μ some finite measure, the existence of an arbitrarily close ReLU valued neural network (Hornik, 1991).

Theorem 2.23 (Hornik's Theorem). *Define for some continuous, unbounded and non-constant activation function ρ the set of all feedforward neural networks, as*

$$\mathcal{N}_d(\rho) := \bigcup_{L=1}^{\infty} \left\{ R_\rho(\Phi) : \mathbb{R}^d \rightarrow \mathbb{R} \mid \Phi \text{ is neural network with } L \text{ layers} \right\}.$$

Then $\mathcal{N}_d(\rho)$ lies dense in $L^p(\mu)$ for every finite measure μ , $p \in (0, \infty)$.

The proof of this statement leverages on Hahn Banach, but thus does not provide the specific architecture of the required neural network, which would be used for such an approximation. We aim significantly higher, namely at a bound on the necessary non-zero weights and layers for such an approximation, and will thus not mention the proof beyond this outline.

2.3.1 Network concatenations

In this subsection we will discuss the concatenations of multiple neural networks and the effects on the characteristic numbers of it. The complexity of a concatenated network can be given nicely, when the used activation function is

Definition 2.24 (Network Concatenation). Let $L_1, L_2 \in \mathbb{N}$ and

$$\Phi^1 = ((A_1, b_1), \dots, (A_{L_1}, b_{L_1})), \quad \Phi^2 = ((B_1, c_1), \dots, (B_{L_2}, c_{L_2})),$$

be two neural networks such that the input layer of Φ^1 has the same dimension as the output layer of Φ^2 . Then, $\Phi^1 \bullet \Phi^2$ denotes the following $L_1 + L_2 - 1$ layered network:

$$\Phi^1 \bullet \Phi^2 := \left(((B_1, c_1), \dots, (B_{L_2-1}, c_{L_2-1})), ((\tilde{A}_1, \tilde{b}_1), \dots, (\tilde{A}_{L_1}, \tilde{b}_{L_1})) \right),$$

where

$$\tilde{A}_l^1 := \left[A_{l, x_0} \mid B_{L_2} \mid A_{l, x_1} \mid \dots \mid A_{l, x_{l-1}} \right], \quad \text{and} \quad \tilde{b}_l^1 := A_{l, x_0} c_{L_2} + b_l,$$

for $l = 1, \dots, L_1$. We call $\Phi^1 \bullet \Phi^2$ the *concatenation* of Φ^1 and Φ^2 .

Lemma 2.25 (Neutral Neural Network). Let $\Phi^{\text{Id}} = ((A_1, b_1), (A_2, b_2))$ be a d -dimensional neural network, with

$$A_1 := \begin{bmatrix} \text{Id}_{\mathbb{R}^d} \\ -\text{Id}_{\mathbb{R}^d} \end{bmatrix}, \quad b_1 := 0, \quad A_2 := \begin{bmatrix} 0_{\mathbb{R}^{d \times d}} & \text{Id}_{\mathbb{R}^d} & -\text{Id}_{\mathbb{R}^d} \end{bmatrix}, \quad b_2 := 0.$$

Then $R_\rho(\Phi^{\text{Id}}) = \text{Id}_{\mathbb{R}^d}$.

Lemma 2.26. For any two neural networks Φ^1 and Φ^2 according to Definition (2.24), $\Phi^1 \odot \Phi^2 := \Phi^1 \bullet \Phi^{\text{Id}} \bullet \Phi^2$ is a neural network with d -dimensional input, $L_1 + L_2$ layers with $\omega_{\Phi^1 \odot \Phi^2} \leq 2\omega_{\Phi^1} + 2\omega_{\Phi^2}$ active weights, $N_{\Phi^1 \odot \Phi^2} \leq 2N_{\Phi^1} + 2N_{\Phi^2}$ neurons. Additionally for all $x \in \mathbb{R}^d$, we have

$$R_\rho(\Phi^1 \odot \Phi^2)(x) = R_\rho(\Phi^1) \circ R_\rho(\Phi^2)(x).$$

Proof.

maybe include proof?

Intuitively speaking the latter network must account for all "missing" links, which a direct concatenation would necessarily be missing; any layer after the "switch" from the first to the latter needs to have a correction term in the weights, additionally attributing all potential connections which the first network would impose.

Let w.l.o.g. $L_2 = 2$, since the pass through the first layers of the combined network, is equivalent to the pass through the first layers of the individual network. Then we can show by induction:

1. Base case: $L_2 = 1$. We argue with $A_{L_1}x = A_{L_1, x_0}x$ for x of output dimension of Φ^1 , since the input dimension of Φ^1 is the output of Φ^2 .

$$\begin{aligned} R_\rho(\Phi^1) \circ R_\rho(\Phi^2)(x_0) &= A_{L_1} (B_{L_2, x_0}x_0 + B_{L_2, x_1}x_1 + c_{L_2}) + b_{L_1} \\ &= A_{L_1, x_0}B_{L_2, x_0}x_0 + A_{L_1, x_0}B_{L_2, x_1}x_1 + (A_{L_1, x_0}c_{L_2} + b_{L_1}) \\ &= A_{L_1, x_0}B_{L_2} \begin{bmatrix} x_0^T \\ x_1^T \end{bmatrix}^T + (A_{L_1, x_0}c_{L_2} + b_{L_1}) \\ &= R_\rho(\Phi^1 \bullet \Phi^2)(x_0) \end{aligned}$$

2. Induction assumption: Let the statement be true for $L_2 \geq 2$ fixed.
3. Induction step: $L_2 \mapsto L_2 + 1$.

□

2.3.2 Network parallelization

We have shown, that there is a possibility to simply compose networks, as is possible with functions. To further build upon that idea, we will now introduce stacking in the context of defining a network $P(\Phi^1, \Phi^2)$ from two networks Φ^1 and Φ^2 such that $R_\rho(P(\Phi^1, \Phi^2))(x) = (R_\rho(\Phi^1)(x), R_\rho(\Phi^2)(x))$ for all $x \in \mathbb{R}^d$.

Definition 2.27 (Network Parallelization). Let $L_1, L_2 \in \mathbb{N}$ s.t. $L_1 \leq L_2$ and let

$$\Phi^1 = ((A_1, b_1), \dots, (A_{L_1}, b_{L_1})), \quad \Phi^2 = ((B_1, c_1), \dots, (B_{L_2}, c_{L_2})),$$

be two neural networks with d -dimensional input. We call

$$P(\Phi^1, \Phi^2) := ((C_1, d_1), \dots, (C_{L_2}, d_{L_2})),$$

the *parallelization of Φ^1 and Φ^2* with

$$C_l := \left[\begin{array}{c|c|c|c|c} A_{l,x_0} & A_{l,x_1} & 0 & \cdots & A_{l,x_{l-1}} & 0 \\ B_{l,x_0} & 0 & B_{l,x_1} & \cdots & 0 & B_{l,x_{l-1}} \end{array} \right], \quad d_l := \begin{bmatrix} b_l \\ c_l \end{bmatrix}, \quad \text{for } 1 \leq l < L_1,$$

$$C_{L_1} := \left[\begin{array}{c|c|c|c|c} B_{L_1,x_0} & 0 & B_{L_1,x_1} & \cdots & 0 & B_{L_1,x_{L_1-1}} & B_{L_1,x_{L_1}} & \cdots & B_{L_1,x_{L_1-1}} \end{array} \right], \quad d_{L_1} := c_{L_1},$$

for $L_1 \leq l < L_2$, and

$$C_{L_2} := \left[\begin{array}{c|c|c|c|c|c|c|c|c} A_{L_2,x_0} & A_{L_2,x_1} & 0 & \cdots & A_{L_2,x_{L_2-1}} & 0 & 0 & \cdots & 0 \\ B_{L_2,x_0} & 0 & B_{L_2,x_1} & \cdots & 0 & B_{L_2,x_{L_2-1}} & B_{L_2,x_{L_2}} & \cdots & B_{L_2,x_{L_2-1}} \end{array} \right],$$

$$d_{L_2} := \begin{bmatrix} b_{L_2} \\ c_{L_2} \end{bmatrix}.$$

A similar construction is used for the case $L_1 > L_2$.

Lemma 2.28. In the setting of Definition (2.27), we have that $P(\Phi^1, \Phi^2)$ is a neural network with d -dimensional input, $\max\{L_1, L_2\}$ layers, $\omega_{P(\Phi^1, \Phi^2)} = \omega_{\Phi^1} + \omega_{\Phi^2}$ active weights and $N_{P(\Phi^1, \Phi^2)} = N_{\Phi^1} + N_{\Phi^2} - d$ neurons. Additionally, for all $x \in \mathbb{R}^d$, we have that

$$R_\rho(P(\Phi^1, \Phi^2))(x) = (R_\rho(\Phi^1)(x), R_\rho(\Phi^2)(x)).$$

maybe include proof?

2.3.3 Standard Neural Networks

Corollary 2.29. Let ρ be the ReLU activation function. Then there are constants $C_1, C_2 > 0$ such that for any neural network Φ with d -dimensional input and $L = L(\Phi)$ layers, $N = N(\Phi)$ neurons and $M = M(\Phi)$ nonzero weights, there is a standard neural network Φ_{st} with d -dimensional input, L layers, at most $C_1 LN$ neurons and $C_2 \cdot (LN + M)$ nonzero weights such that

$$R_\rho(\Phi)(x) = R_\rho(\Phi_{\text{st}})(x)$$

for all $x \in \mathbb{R}^d$.

2.3.4 Approximation Theory

For all further theoretical results we will assume that ρ , the activation function, is the ReLU as given in Definition (2.22) and furthermore define the set of functions bounded in Sobolev norm as

$$\mathcal{F}_{n,d,p,B} := \{f \in W^{n,p}((0,1)^d) \mid \|f\|_{W^{n,p}((0,1)^d)} \leq B\}.$$

To create a decent outline for this subsection, we will shortly discuss the goal and the steps to be taken; in order to show a bound on the weights and layers needed for an approximating neural network, we need to take into account the interpolation error, we may overcome with sufficient regularity. Say we would have a function f in the Sobolev space $W^{2,2}((0,1)^d)$, then it is apparent, that at least 2 derivatives of f must exist, in a piecewise, embedded sense, which can be leveraged to get a Taylor expansion around selected points.

This procedure will deduce a quantifiable error for the network approximating the initial arbitrary Sobolev function f in Sobolev norm, as long as we can bound the error, which the polynomial approximation produces with regard to the network itself.

This leaves us with two main objectives:

1. Bound the Sobolev function at critical equidistant points, using its Taylor expansion.
2. Relate a fixed error rate of the approximation polynomial and some neural network, to the networks complexity.

These two objectives are pursued by both Yarotsky (Yarotsky, 2017) and Gühring et al (Gühring et al., 2019), even though it is worth noting, that Yarotsky only provides the special case $s = 0, B = 1$, since the proof only shows an estimation with a Taylor expansion regarding the L^∞ norm, missing the crucial step of the interpolation bound to change the norm to the Sobolev expanded counterpart and additionally uses the L^∞ norm to approximate multiplication by a neural network.

Tho we will use the statement of Gühring later on, which is the extension of Yarotsky to any choice of s, p and B , we will only give the idea of the proof, since it follows the same pattern as Yarotsky's proof, but does include a lot of further theoretical setting, which will not be of much use in the actual results presented in this paper.

Proposition 2.30. There are constants $c_1, c_2, c_3 > 0$, such that for all $\varepsilon \in (0, 1)$ there is a neural network $\Phi_\varepsilon^{\text{sq}}$ with at most $c_1 \cdot \ln(1/\varepsilon)$ nonzero weights, at most $c_2 \cdot \ln(1/\varepsilon)$ layers, at most $c_3 \cdot \ln(1/\varepsilon)$ neurons, and with a 1-dimensional input and output such that for $f(x) := x^2$

$$\|R_\rho(\Phi_\varepsilon^{\text{sq}}) - f\|_{W^{0,\infty}((0,1)^1)} \leq \varepsilon,$$

and $R_\rho(\Phi_\varepsilon^{\text{sq}}(0)) = 0$.

Proof. Define the "tooth" function $g : [0, 1] \rightarrow [0, 1]$ with

$$g(x) = \begin{cases} 2x, & \text{for } x < 1/2, \\ 2(1-x), & \text{for } x \geq 1/2, \end{cases}$$

and its composition

$$g_s(x) := \underbrace{g \circ g \circ \dots \circ g}_{s\text{-times}}(x).$$

Furthermore, from the definition of the ReLU function (refer to (2.22)), we can also set $g(x) = 2\rho(x) - 4\rho(x - 1/2) + 2\rho(x - 1)$, which is intern, the same as defining a neural network with 1-dimensional input and output via $\Phi = ((A_1, b_1), (A_2, b_2))$ with

$$A_1 = \begin{bmatrix} \text{Id}_{\mathbb{R}} \\ \text{Id}_{\mathbb{R}} \\ \text{Id}_{\mathbb{R}} \end{bmatrix}, \quad b_1 = \begin{pmatrix} 0 \\ -1/2 \\ -1 \end{pmatrix}, \quad A_2 = \begin{bmatrix} 2 & -4 & 2 \end{bmatrix}, \quad b_2 = 0.$$

Then it is evident, using Lemma (2.26), that $g_s(x) = R_\rho(\underbrace{\Phi \odot \dots \odot \Phi}_{s\text{-times}})(x)$ for all $x \in \mathbb{R}$ represents a network with $\mathcal{O}(s)$ layers, active weights and neurons. Now, with an explicit formulation of g_m for a fixed level $m \geq 1$, one can identify the function $f(x) := x^2$ using linear interpolation on equidistant points $k/(2^m)$, where $k = 0, \dots, 2^m - 1$, i.e. let $\Phi_m = \Phi \odot \dots \odot \Phi$ (m -times) be a neural network, because then for all $x \in [\frac{k}{2^m}, \frac{k+1}{2^m}]$ it is true, that

$$R_\rho(\Phi_m)(x) = \left(\frac{(k+1)^2}{2^m} - \frac{k^2}{2^m} \right) \left(x - \frac{k}{2^m} \right) + \left(\frac{k}{2^m} \right)^2.$$

Why?

Then we also know that, using the Bramble-Hilbert lemma in one-dimension (Bramble & Hilbert, 1970), the fact that $f''(x) = 2$ for all $x \in (0, 1)$, and $h = 1/2^m$, that

$$\|R_\rho(\Phi_m) - f\|_{L^\infty((0,1))} \leq \max_{k=0, \dots, 2^m-1} Ch^2 \|f''\|_{L^\infty([\frac{k}{2^m}, \frac{k+1}{2^m}])} = C' 2^{-2m}.$$

Hence setting $m = \lceil \ln(1/\varepsilon) \rceil$, with $C, C' > 0$ constants, only dependent on the domain, we achieve the wanted bound from the statement for the neural network $\Phi_\varepsilon^{\text{sq}} := \Phi_m$ with $\mathcal{O}(m) = \mathcal{O}(\ln(1/\varepsilon) + 1) \leq \mathcal{O}(2\ln(1/\varepsilon))$ layers, neurons and active weights. $\square$

This statement enables us to express multiplication using a neural network. This is straight forward, when viewing the polarization identity:

$$xy = \frac{1}{2}((x+y)^2 - x^2 - y^2), \quad \text{for } x, y \in \mathbb{R}, \quad (2.1)$$

as given by the first binomial formula.

Proposition 2.31. Let $M > 0$ and $\varepsilon \in (0, 1)$, then there is a ReLU neural network $\tilde{\times}$ with 2-dimensional input and 1-dimensional output, and constants $c_1 > 0$ and $c_2 = c_2(M) > 0$, such that for $\times : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}; (x, y) \mapsto xy$

1. $\|R_\rho(\tilde{\times}) - \times\|_{W^{0,\infty}((-M,M)^2)} \leq \varepsilon$,
2. if $x = 0$ or $y = 0$, then $R_\rho(\tilde{\times})(x, y) = 0$,
3. the number of layers, active weights and neurons of $\tilde{\times}$ is at most $c_1 \ln(1/\varepsilon) + c_2$.

Proof. Let $\delta := \varepsilon/(6M^2)$. Then we construct an approximation of the square function $f(x) = x^2$ for $x \in (0, 1)$ using Proposition (2.30) for the chosen δ , i.e. we define Φ_δ^{sq} , such that

$$\|R_\rho(\Phi_\delta^{\text{sq}}) - f\|_{W^{0,\infty}((0,1)^1)} \leq \frac{\varepsilon}{6M^2},$$

which now of course has $\mathcal{O}(\ln(1/\delta))$ many layers, active weights and neurons. Since the statement necessarily needs to implement negative numbers as well, we show with $|x| = \rho(x) + \rho(-x)$ that it is indeed possible to expand any neural network with a positive domain to a negative domain using a neural network (by simply concatenating). Thus we construct $\tilde{\times}$ via adding an addition layers on top of parallel square networks with scaled input, achieving

$$R_\rho(\tilde{\times})(x, y) = 2M^2 \left(R_\rho(\Phi_\delta^{\text{sq}}) \left(\frac{|x+y|}{2M} \right) - R_\rho(\Phi_\delta^{\text{sq}}) \left(\frac{|x|}{2M} \right) - R_\rho(\Phi_\delta^{\text{sq}}) \left(\frac{|y|}{2M} \right) \right).$$

To appreciate the third statement of our Proposition, we may now put together the necessary complexity of this network using the constants in Proposition (2.30);

1. We have 2 layers for the implementation of the absolute value, 1 for the addition in the end, and $c_0 \ln(1/\delta)$ for the square function, thus $L_{\tilde{\times}} \leq c_0 \ln(1/\varepsilon) + c_0 \ln(6M^2) + 3$.
2. We need $4 + 2 + 2$ active weights for the first matrix to distribute each (x, y) onto the parallel networks together with 6 active weights for the superposing of the input into the $|\cdot|$ -function. Another 3 are required for the last layer adding all outputs. Of course each of the different computations for the artificial scaled square function must be in parallel, thus $\omega_{\tilde{\times}} \leq 3c_0 \ln(1/\varepsilon) + 3c_0 \ln(6M^2) + 17$.

3. A similar consideration gives us $9 = (2 + 1) \cdot 3$ neurons for the first two and the last layer, thus giving us $N_{\times} \leq 3c_0 \ln(1/\varepsilon) + 3c_0 \ln(6M^2) + 9$.

Finally taking the adequate maximum, and remarking, that c_0 does not depend on M , we get the wanted bound in (2).

To make the true multiplication match the form of our implemented network, using (2.1), we may rewrite for $x, y \in (-M, M)$

$$\begin{aligned} xy &= 4M^2 \left(\frac{x}{2M} \cdot \frac{y}{2M} \right) \\ &= 4M^2 \frac{1}{2} \left(\left(\frac{x}{2M} + \frac{y}{2M} \right)^2 - \left(\frac{x^2}{2M} \right)^2 - \left(\frac{y^2}{2M} \right)^2 \right) \\ &= 2M^2 \left(\left(\frac{|x+y|}{2M} \right)^2 - \left(\frac{|x|}{2M} \right)^2 - \left(\frac{|y|}{2M} \right)^2 \right). \end{aligned}$$

Now with the deviation from each of the square approximation we get,

$$\begin{aligned} \|R_\rho(\tilde{\times}) - \times\|_{L^\infty((0,1)^2)} &\leq 2M^2 \left(3 \cdot \sup_{\frac{|x|}{2M} : x \in (-M, M)} |R_\rho(\Phi_\delta^{\text{sq}})(\tilde{x}) - \tilde{x}^2| \right) \\ &\leq 6M^2\delta = \varepsilon. \end{aligned}$$

We note a few things happening here for clarity; it is possible to simply use the reverse triangle equality to get all three term errors added, and then put them together. This can be done, since the $L^\infty((0,1)^2)$ norm appreciates only the maximal value of the error across all possible elements of the $(0,1)^2$ domain, but because the inner function is bijective, these are just the same. Additionally we use the, trivial by definition, statement

$$\|g \circ f\|_{L^\infty((0,1))} \leq \|g\|_{L^\infty((0,1))}.$$

This has shown the statement (3).

Statement (1) on the other hand, is a direct consequence of the subliminally result in Proposition (2.30). $\square$

As a last step, before stating and proving the Yarotsky Theorem, we want to discuss the construction and complexity of a network approximating any collection of equidistantly spaced function with local support in a higher dimensional domain $(0,1)^d$ for $1 \leq d < \infty$. If we take into account, that we may also want these functions to form a partition of unity on the high dimensional domain, this will be ground to tackle the complexity of the approximation of polynomials with neural networks.

Lemma 2.32. For any $d, N \in \mathbb{N}$ there is a collection of functions

$$\Psi = \left\{ \phi_m \mid m \in \{0, \dots, N\}^d \right\},$$

with $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$ for all $m = (m_1, \dots, m_d) \in \{0, \dots, N\}^d$ with the following properties:

1. $0 \leq \phi_m(x) \leq 1$ for all $\phi_m \in \Psi$ and every $x \in \mathbb{R}^d$,
2. $\sum_{\phi_m \in \Psi} \phi_m(x) = 1$ for every $x \in [0, 1]^d$,

3. $\text{supp}(\phi_m) \subset B_{1/N, \|\cdot\|_{\ell^\infty}}(m/N)$ for all $\phi_m \in \Psi$,
4. there are constants $C, c \geq 1$ such that for each $\phi_m \in \Psi$ there is a neural network Φ_m with d -dimensional input and output, at most 3 layers, $C \cdot d$ active weights and neurons, such that

$$\prod_{l=1}^d [R_\rho(\Phi_m)]_l = \phi_m$$

and $\|[R_\rho(\Phi_m)]_l\|_{L^\infty((0,1)^d)} \leq 1$ for all $l = 1, \dots, d$.

Proof. The idea of the proof, will be one of construction, where we will find adequate functions, which are easily transferable into ReLU neural networks. For this, define the functions

$$\psi : \mathbb{R} \rightarrow \mathbb{R}, \quad \psi(x) = \begin{cases} 1, & |x| < 1, \\ 0, & 2 < |x|, \\ 2 - |x| & 1 \leq |x| \leq 2, \end{cases}$$

and $\phi_m : \mathbb{R}^d \rightarrow \mathbb{R}$ as the product of shifted and scaled versions of ψ , which inherit its local support, i.e.

$$\phi_m(x) := \prod_{l=1}^d \psi\left(3N\left(x_l - \frac{m_l}{N}\right)\right),$$

for $m = (m_1, \dots, m_d) \in \{0, \dots, N\}^d$. This function satisfies our first three parts of the Lemma directly: Take one x_i for some $i = 1, \dots, d$. Now we get that $\psi(3N(x_i - (m_i/N))) \in [-2, 2]$, since ψ has support of $[-2, 2]$. This gives us

$$x \in \left[\frac{m}{N} - \frac{2}{3N}, \frac{m}{N} + \frac{2}{3N}\right],$$

yielding the third property, as we took i arbitrarily. Simultaneously we use the zero product property, and the fact, that the range of ψ is $[0, 1]$ to get the first property. Defining $z_i := 3Nx_i$, we can rewrite for any N the one-dimensional function

$$S(x_i) := \sum_{m_i=0}^N \psi\left(3N\left(x_i - \frac{m_i}{N}\right)\right) \quad \text{into the function} \quad \tilde{S}(z_i) := \sum_{m_i=0}^N \psi(z_i - 3m_i),$$

that intern gives us the wanted result of $\tilde{S}(z_i) = 1$ for any $z_i \in [0, 3N]$, because the least distance between points is 3, which means going forwards or backwards either lands one outside the support of the function, if $z_i \in [-1, 1]$, or it adds both third case values, i.e. setting $z_i \in [1, 2]$ w.l.o.g

$$\psi(z_i) + \psi(z_i - 3) = (2 - z_i) + (2 + (z_i - 3)) = 1.$$

Now we can extrapolate this result to S by a simple scaling argument and then use Fubini and the factorization of the sum over products to get

$$\sum_{m \in \{0, \dots, N\}^d} \phi_m(x) = \sum_{m_1=0}^N \cdots \sum_{m_d=0}^N \prod_{l=1}^d \psi\left(3N\left(x_l - \frac{m_l}{N}\right)\right) = \prod_{l=1}^d \left(\sum_{m_l=0}^N \psi\left(3N\left(x_l - \frac{m_l}{N}\right)\right) \right) = 1.$$

Finally we are only left with proving the forth property. For this, we need to construct a neural network Φ_ψ , that realizes ψ . Let

$$A_1 := \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad b_1 := \begin{pmatrix} 2 \\ 1 \\ -1 \\ -2 \end{pmatrix}, \quad A_2 := [0 \mid 1 \quad -1 \quad -1 \quad 1], \quad b_2 := 0,$$

define the neural network $\Phi_\psi = ((A_1, b_1), (A_2, b_2))$. With distinction by cases one can easily see, that for all $x \in \mathbb{R}$

$$\begin{aligned} R_\rho(\Phi_\psi)(x) &= A_2(\rho(A_1x + b_1)) + b_2 \\ &= A_2 \left(\begin{bmatrix} \rho(x+2) & \rho(x+1) & \rho(x-1) & \rho(x-2) \end{bmatrix}^T \right) \\ &= \rho(x+2) + \rho(x-2) - (\rho(x+1) + \rho(x-1)) \\ &= \psi(x). \end{aligned}$$

Furthermore the complexity of this network is 2 layers, 12 active weights and 6 neurons. Let $\Phi_{m,l}$ be a neural network with d -dimensional input and one-dimensional output, such that

$$R_\rho(\Phi_{m,l})(x) = 3N \left(x_l - \frac{m_l}{N} \right), \quad \text{for all } x \in \mathbb{R}^d \text{ and } m \in \{0, \dots, N\}^d.$$

Then the complexity of this network is 1 layer, 2 active weights and $d+1$ neurons. Finally realizing that the composition of both these functions, i.e. the realization of the concatenation of networks, would yield each factor for the wanted function. If we now parallelize this concatenation, we get the forth property by definition, i.e. let

$$\Phi_m := P(\Phi_\psi \odot \Phi_{m,l} \mid l = 1, \dots, d),$$

since then

$$\prod_{l=1}^d [R_\rho(\Phi_m)]_l(x) = \phi_m(x), \quad \text{for all } x \in \mathbb{R}^d.$$

This network now has d -dimensional input and output and the complexity, using both Lemma (2.26) and Lemma (2.28), of 3 layers, at most $(12+2) \cdot 2 \cdot d = 28d$ active weights, and at most $9d$ neurons. $\square$

With these auxiliary results, we may finally start with the statement and proof of the Yarotsky Theorem.

Theorem 2.33 (Yarotsky Theorem). *Let $d \in \mathbb{N}, n \in \mathbb{N}, B = 1$ and $f \in \mathcal{F}_{n,d,\infty,B}$ some arbitrary function. Then there is a constant $c = c(d, n) > 0$ with the following properties:*

For any $\varepsilon \in (0, 1)$, there is a neural network architecture $\mathcal{A}_\varepsilon = \mathcal{A}_\varepsilon(d, n, \varepsilon)$ with d -dimensional input and one-dimensional output such that, there is a neural network Φ_ε^f that has architecture $\mathcal{A}_\varepsilon$ and satisfies

$$\|f - R_\rho(\Phi_\varepsilon^f)\|_{W^{0,\infty}((0,1)^d)} \leq \varepsilon,$$

and

1. $L_{\mathcal{A}_\varepsilon} \leq c \cdot \ln(1/\varepsilon),$
2. $\omega_{\mathcal{A}_\varepsilon} \leq c \cdot \varepsilon^{-d/n} \ln(1/\varepsilon),$
3. $N_{\mathcal{A}_\varepsilon} \leq c \cdot \varepsilon^{-d/n} \ln(1/\varepsilon).$

Proof. Given the definition of the functions $\phi_m \in \Psi$ as constructed in the proof of Lemma (2.32), we want to structure the proof of this Theorem in two sub-goals:

The formulation of an approximate function f_1 for f using the partition of unity and an adequate Taylor expansion, and the subsequent approximation of that function f_1 with a neural network Φ_{f_1} using the $\tilde{\times}$ -multiplication of Proposition (2.31).

Step 1: Taylor Expansion

For any $m \in \{0, \dots, N\}^d$, consider the $(n-1)$ -Taylor polynomial for the function f at $x_0 = m/N$:

$$P_m(x) = \sum_{n: |n| < n} \frac{D^n f(x)}{n!} \Big|_{x=\frac{m}{N}} \left(x - \frac{m}{N}\right)^n, \quad \text{for } x \in \mathbb{R}^d,$$

with the conventions $n! = \prod_{l=1}^d n_l!$ and $(x - m/N)^n = \prod_{l=1}^d (x_l - \frac{m_l}{N})^{n_l}$. With the partition of unity of Lemma (2.32), we can then set any function, as a linear combination of its point-values and the partition function. Especially this allows us to define

$$f = \sum_{m \in \{0, \dots, N\}^d} \phi_m f, \quad \text{and} \quad f_1 = \sum_{m \in \{0, \dots, N\}^d} \phi_m P_m.$$

Now we can exploit the properties given by Lemma (2.32), as follows; firstly use the $\|\phi_m\|_{L_\infty((0,1)^d)} = 1$ property in Line (2.3), then one can estimate the maximal number of partitions with 2^d in Line (2.4). After this we take the simple Lagrange remainder

argue this

in Line (2.5). Lastly one can simply argue with $f \in \mathcal{F}_{n,d,\infty,B=1}$ in Line (2.6).

$$\|f - f_1\|_{L_\infty((0,1)^d)} \leq \sum_{m \in \{0, \dots, N\}^d} \|\phi_m (f - P_m)\|_{L_\infty((0,1)^d)} \quad (2.2)$$

$$\leq \sum_{m: |x_l - \frac{m_l}{N}| < \frac{1}{N}, \forall l=1, \dots, d} \|f - P_m\|_{L_\infty((0,1)^d)} \quad (2.3)$$

$$\leq 2^d \max_{m: |x_l - \frac{m_l}{N}| < \frac{1}{N}, \forall l=1, \dots, d} \|f - P_m\|_{L_\infty((0,1)^d)} \quad (2.4)$$

$$\leq \frac{2^d d^n}{n!} \left(\frac{1}{N}\right)^n \max_{n: |n|=n} \|D^n f\|_{L_\infty((0,1)^d)} \quad (2.5)$$

$$\leq \frac{2^d d^n}{n!} \left(\frac{1}{N}\right)^n \quad (2.6)$$

Now taking

$$N = \left\lceil \left(\frac{n!}{2^d d^n} \cdot \frac{\varepsilon}{2} \right)^{-\frac{1}{n}} \right\rceil$$

gives the approximation bound of

$$\|f - f_1\|_{L_\infty((0,1)^d)} \leq \frac{\varepsilon}{2}. \quad (2.7)$$

Step 2: Approximation by a Neural Network

The polynomial function f_1 may be further expressed in a different form, which allows us to separately compare it to a potential network approximation using the definition of ϕ_m , i.e. for $x \in \mathbb{R}^d$

$$f_1(x) = \sum_{m \in \{0, \dots, N\}^d} \sum_{n: |n| < n} a_{m,n} \underbrace{\left(\phi_m(x) \left(x - \frac{m}{N}\right)^n \right)}_{=: f_{m,n}(x)}. \quad (2.8)$$

This expansion now has at most $d^n(N+1)^d$ terms containing $f_{m,n}$, since

$$\left| \left\{ n \in \{0, \dots, n-1\}^d : |n| < n \right\} \right| \leq \left| \left\{ n \in \{0, \dots, n-1\}^d \right\} \right| = d^n,$$

and obviously $|\{m = \{0, \dots, N\}^d\}| = (N+1)^d$. We note, that we take $M = d+n$ for the definition of the network multiplication, i.e. $\tilde{\times} : (-M, M)^2 \rightarrow \mathbb{R}$. This is done, because we have as an obvious consequence from the fact, that if $x, y \in (-1, 1)$, that

$$|R_\rho(\tilde{\times})(x, y) - xy| \leq \delta \implies |R_\rho(\tilde{\times})(x, y)| \leq |y| + \delta,$$

for some fixed but not yet specified $\delta \in (0, 1)$. Recursively concatenating $\tilde{\times}$ for $(d+n-1)$ -times can then propagate the error to at most $M = (d+n)$ upon the last realization of $\tilde{\times}$.

As aforementioned, we can now compare each function $f_{m,n}$ pointwise to a realization of a network approximating at most $n-1$ of the multiplications needed for $\prod_{l=1}^d (x_l - (m_l/N))^{n_l}$, as we know that $\sum_{l=1}^d n_l < n$, d additional multiplications for the realization of the function ϕ_m and one multiplication for putting it together. Hence we define a neural network $\Phi_{f_1}^{m,n}$ with d -dimensional input and one dimensional output via

$$\begin{aligned} R_\rho(\Phi_{f_1}^{m,n})(x) = R_\rho(\tilde{\times}) & \left([R_\rho(\Phi_m)]_1(x_1), R_\rho(\tilde{\times}) \left([R_\rho(\Phi_m)]_2(x_2), \dots \right. \right. \\ & R_\rho(\tilde{\times}) \left([R_\rho(\Phi_m)]_d(x_d), R_\rho(\tilde{\times}) \left(x_1 - \frac{m_1}{N}, \dots \right. \right. \\ & R_\rho(\tilde{\times}) \left(x_{d-1} - \frac{m_{d-1}}{N}, R_\rho(\tilde{\times}) \left(x_d - \frac{m_d}{N}, \dots \right. \right. \\ & \left. \left. \left. \left. \left. \left. R_\rho(\tilde{\times}) \left(x_d - \frac{m_d}{N}, x_d - \frac{m_d}{N} \right) \right) \right) \right) \right) \right) \right). \end{aligned}$$

for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ where we used Φ_m as in the proof Lemma (2.32). This network contains at most $(d+n)$ concatenations of $\tilde{\times}$, and thus by Lemma (2.32) and its constant $C > 0$, Lemma (2.26), and Proposition (2.31) and its constants $c_0, c_1(d, n) > 0$,

$$L_{\Phi_{f_1}^{m,n}} \leq 3c_0(d+n) \left(\ln \left(\frac{1}{\delta} \right) + (d+n)c_1 \right), \quad \omega_{\Phi_{f_1}^{m,n}} \leq C2(d+n)d \left(\ln \left(\frac{1}{\delta} \right) + 2(d+n)C \right). \quad (2.9)$$

for a given precision level $\delta \in (0, 1)$, where the number of neurons are of the same order as the number of active weights.

The error comparison of $R_\rho(\Phi_{f_1}^{m,n})$ and $f_{m,n}$ can then be done by counting the maximal depth of usages of the network $\tilde{\times}$, which is $(d+n)$ as shown above, and the estimate of

$$\left| \left(x - \frac{m}{N} \right) \right| \leq 1, \quad |\psi(x)| \leq 1,$$

for all $x \in (0, 1)^d$. Then directly

$$\|R_\rho(\Phi_{f_1}^{m,n}) - f_{m,n}\|_{L_\infty((0,1)^d)} \leq (d+n)\delta.$$

Now since, as mentioned below Equation (2.8), f_1 has at most $d^n(N+1)^d$ terms, but since any $x \in \mathbb{R}^d$ can be only in the support of at most 2^d functions ϕ_m by the second statement in Proposition (2.31), the sum going over all m can be reduced to only those, which are non-zero, which we employ in Line (2.11). This brings $(N+1)^d \rightarrow 2^d$, supporting the equation in Line (2.12). Setting

$$\delta = \frac{\varepsilon}{2 \cdot 2^d d^n (d+n)}$$

allows us to finally assert, using the bound of Equation (2.9) for Line (2.13),

$$\|R_\rho(\Phi_{f_1}) - f_1\|_{L_\infty((0,1)^d)} \leq \sum_{m \in \{0, \dots, N\}^d} \sum_{n: |n| < n} |a_{m,n}| \|R_\rho(\Phi_{f_1}^{m,n}) - f_{m,n}\|_{L_\infty((0,1)^d)} \quad (2.10)$$

$$= \sum_{m: x \in \text{supp}(\phi_m)} \sum_{n: |n| < n} |a_{m,n}| \|R_\rho(\Phi_{f_1}^{m,n}) - f_{m,n}\|_{L_\infty((0,1)^d)} \quad (2.11)$$

$$\leq 2^d d^n \max_{m: x \in \text{supp}(\phi_m)} \max_{n: |n| < n} \|R_\rho(\Phi_{f_1}^{m,n}) - f_{m,n}\|_{L_\infty((0,1)^d)} \quad (2.12)$$

$$\leq 2^d d^n (d + n) \delta \quad (2.13)$$

$$= \frac{\varepsilon}{2}. \quad (2.14)$$

Our knowledge of δ together with at most $d^n(N+1)^d = \mathcal{O}(\varepsilon^{-d/n})$ parallel networks with one additive layer and the fact that generally we have $c' \ln(1/\delta + c'') \leq c''' \ln(1/\delta)$ for $c', c'' > 0$ and $c''' \geq c' \ln(1/c'')$, gets us the final complexity of Φ_{f_1} from the starting point of Equation (2.9), i.e.

$$L_{\Phi_{f_1}} \leq c \ln(1/\varepsilon), \quad \omega_{\Phi_{f_1}} \leq c \varepsilon^{-d/n} \ln(1/\varepsilon), \quad N_{\Phi_{f_1}} \leq c \varepsilon^{-d/n} \ln(1/\varepsilon).$$

with some constant $c = c(d, n) > 0$.

Using the triangle inequality, we can show that this network indeed satisfies the wanted precision bound with both errors stemming from Equation () and Equation () respectively:

$$\|f - R_\rho(\Phi_{f_1})\|_{L_\infty((0,1)^d)} \leq \|f - f_1\|_{L_\infty((0,1)^d)} + \|f_1 - R_\rho(\Phi_{f_1})\|_{L_\infty((0,1)^d)} \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

□

We may even expand Yarotsky to cover a more general case, where $f \in \mathcal{F}_{n,d,p,B}$ for arbitrarily $1 \leq p \leq \infty$ and $B > 0$ and the norm, to which we assert the precision, an arbitrary Sobolev norm. This extension is presented by Gühring et al. (Gühring et al., 2019).

Theorem 2.34 (Gühring Theorem). *Let $d \in \mathbb{N}, n \in \mathbb{N}_{\geq 2}, B > 0$ and $0 \leq s \leq 1$ and $f \in \mathcal{F}_{n,d,p,B}$ some arbitrary function. Then there is a constant $c = c(d, n, p, B, s) > 0$ with the following properties:*

For any $\varepsilon \in (0, 1)$, there is a neural network architecture $\mathcal{A}_\varepsilon = \mathcal{A}_\varepsilon(d, n, p, B, s, \varepsilon)$ with d -dimensional input and one-dimensional output such that, there is a neural network Φ_ε^f that has architecture $\mathcal{A}_\varepsilon$ and satisfies

$$\|f - R_\rho(\Phi_\varepsilon^f)\|_{W^{s,p}((0,1)^d)} \leq \varepsilon,$$

and

1. $L_{\mathcal{A}_\varepsilon} \leq c \cdot \ln(1/\varepsilon),$
2. $\omega_{\mathcal{A}_\varepsilon} \leq c \cdot \varepsilon^{-d/(n-s)} \ln(1/\varepsilon),$
3. $N_{\mathcal{A}_\varepsilon} \leq c \cdot \varepsilon^{-d/(n-s)} \ln(1/\varepsilon).$

Proof Sketch. Considering that all of the tools used for this proof will neither be of any use to the main results in this paper nor to the general structure, we will only sketch the general steps necessary to achieve it for the interested reader:

1. The first step is to expand the Proposition (2.30) to the $W^{1,\infty}(0,1)$ -norm. This is done by simple calculation, since an explicit construction is being made.
2. This can be applied to Proposition (2.31) in the same way. Additionally we need to bound $|R_\rho(\tilde{\times})|_{W^{1,\infty}((0,1)^2)}$, since this is required in for the bound on the collective error under the Taylor expanded function in the end.
3. The most import additional step is to bound the switch of norms from $W^{1,p}((0,1)^d)$ to $W^{s,p}((0,1)^d)$ of a function being approximated by the partition of unity, that we discussed in Lemma (2.32). This gives

$$\|f - f_N\|_{W^{s,p}((0,1)^d)} \leq C \left(\frac{1}{N}\right)^{n-s} \|f\|_{W^{n,p}((0,1)^d)},$$

for a partition of unity of size N . It is worth noting, that this factor in front is responsible for the different approximation in regard to weights and neurons, since it influences the necessary choice of δ .

4. The last step, which differs greatly from the proof of Yarotsky Theorem (2.33), is the approximation of a sum of localized polynomials with a neural network. Here the challenge lies again in the control of the derivatives.

□

2.3.5 Statistical Learning

We have now introduced the framework for analytical discussion regarding complexity and potential of neural networks. What is clearly missing however, is the practical "achievement" of a true neural network Φ fitting the function in question from some architecture $\mathcal{A}_\Phi$, that a machine learning engineer thought of. This crucial step in the process of development is referred to as *learning*. This subsection is my own work, if not cited otherwise.

In order to quantify the "falseness" of a potential attribution, we need a function, that compares a true solution to an output of the network operating with the true parameters connected to the true solution. We remark that the following two definitions are based upon the paper (Ciampiconi et al., 2024), where the interested reader can find extensive terminology and currently used loss functions and their optimization techniques.

Definition 2.35 (Feature and Label Space). Let $\mathcal{X}$ be the so-called *feature space* and $\mathcal{Y}$ the *label space* of the underlying unknown function

$$f : \mathcal{X} \rightarrow \mathcal{Y},$$

classifying each feature with a label.

The general goal of all machine learning problems is to closely approximate that classification function f .

Definition 2.36 (Loss Function). Given $f_\theta : \mathcal{X} \rightarrow \mathcal{Y}$ arbitrarily and a feature and label space as in Definition (2.35) and a sample $\{x_i, y_i\}_{i=1}^N \subset \mathcal{X} \times \mathcal{Y}$ for some $N \in \mathbb{N}$, we define the general loss function as a mapping of $f(x_i)$ with its corresponding y_i to a non-negative real number, i.e.

$$\mathcal{L}(f_\theta | \{x_1, \dots, x_N\}, \{y_1, \dots, y_N\}) \in \mathbb{R}^+,$$

such that for f being the function from Definition (2.35) for all $N \in \mathbb{N}$

$$\mathcal{L}(f | \{x_1, \dots, x_N\}, \{y_1, \dots, y_N\}) = 0.$$

Definition 2.37 (Neural Network Weights). Suppose for some $d, L \in \mathbb{N}$ we have a neural network architecture $\mathcal{A}_\Phi$ of input dimension d , L layers and output dimension N_L according to Definition (2.18) and Φ be a neural network of that architecture (see Definition (2.21)). Further let $f : \mathbb{R}^d \rightarrow \mathbb{R}^{N_L}$ be some arbitrary function. Define the *masks of $\mathcal{A}_\Phi$* via

$$I_l := \{(i, j) : [A_l]_{ij} \neq 0\}, \quad J_l := \{i : [b_l]_i \neq 0\},$$

for each layer $1 \leq l \leq L$. We further set the *admissible weight space* as

$$\Theta(\mathcal{A}_\Phi) := \prod_{l=1}^L (\mathbb{R}^{I_l} \times \mathbb{R}^{J_l}),$$

and its elements $\theta \in \Theta(\mathcal{A}_\Phi)$ with

$$\theta = \left((\theta_{l,ij}^A)_{(i,j) \in I_l}, (\theta_{l,i}^b)_{i \in J_l} \right)_{l=1}^L.$$

Then in each layer of $\Phi(\theta) = ((\hat{A}_1(\theta), \hat{b}_1(\theta)), \dots, (\hat{A}_L(\theta), \hat{b}_L(\theta)))$ the affine transformation can be uniquely retrieved via the lifting

$$\hat{A}_l(\theta)_{ij} = \begin{cases} \theta_{l,ij}^A, & (i, j) \in I_l, \\ 0, & \text{otherwise,} \end{cases} \quad \hat{b}_l(\theta)_i = \begin{cases} \theta_{l,i}^b, & i \in J_l, \\ 0, & \text{otherwise.} \end{cases}$$

Then the network Φ will be called a *fitting network with regard to f* , if $\Phi := \Phi(\theta^*)$ is given such that

$$\theta^* = \arg \min_{\theta \in \Theta(\mathcal{A}_\Phi)} \mathcal{L}(R_\rho(\Phi(\theta)) | \{x_1, \dots, x_N\}, \{f(x_1), \dots, f(x_N)\})$$

for some sample $\{x_1, \dots, x_N\}_{i=1}^N \subset \mathbb{R}^d$ with $N \in \mathbb{N}$ and a suitable choice of a loss function $\mathcal{L}$. Note, that this minimizing condition and the architecture are what, not necessarily uniquely, determines the fitting network.

Chapter 3

Formulation of the General Problem

Definitely add traditional convergence and computational cost.

Let $\Theta \subset \mathbb{R}^p$ and $\Theta' \subset \mathbb{R}^q$ be compact and denote the geometric and physical parameter spaces respectively. Further we let $\Omega \subset \mathbb{R}^d$ be some Lipschitz domain and $[0, T]$ be a time interval for some $T > 0$ of a parametric time-dependent PDE, i.e. for any $\mu \in \Theta$ and $\nu \in \Theta'$

$$u^{\mu, \nu} := u(\mu, \nu), \quad \text{in a sufficiently smooth space,}$$

is the (classical) solution of the PDE, for the choice of the space depending on the given specified problem, if

$$\begin{cases} \frac{\partial}{\partial t} u^{\mu, \nu}(x, t) + \mathcal{L}(\mu, \nu) u^{\mu, \nu}(x, t) + \mathcal{N}(u^{\mu, \nu}(x, t), \mu, \nu) = f(x, t; \mu, \nu), & (x, t) \in \Omega \times [0, T], \\ \mathcal{B}(\mu, \nu) u^{\mu, \nu}(x, t) = g_N(x, t; \mu, \nu) & (x, t) \in \partial\Omega \times [0, T], \\ u^{\mu, \nu}(x, t) = u_0^{\mu, \nu}(x) & (x, t) \in \Omega \times \{0\}, \end{cases}$$

where $\mathcal{L}$ represents a linear, and $\mathcal{N}$ represents a non-linear integro-differential operator. Using calculus of variation or distribution theory, one can then formulate a weak form of the problem and find a general space, most commonly Sobolev (for elliptic problems with coercive properties), Bochner (for parabolic problems with coercive properties on the stationary part) or discontinuous Galerkin spaces (for hyperbolic problems with enough regularity), on which a weak solution can be defined. Such a space $\mathcal{H}$ is generally some Banach space, most often a subspace of L^2 .

This weak solution will have less regularity imposed than the strong formulation, but this sometimes allows us to retrieve existence and uniqueness results. Quite famously, there are counter examples of course; the Navier-Stokes Equation in three dimensions poses one such result, where a weak form exists and numerical methods have been developed, which can provide a solution in special cases, but existence remains unproven in general. As mentioned beforehand, we will completely omit the exact transformation of the strong PDE formulation into a weak one, and simply assume that this can be done:

Say we (can) discretize the problem in the spacial domain Ω with N_h degrees of freedom, i.e. let $\mathcal{H}_h \subset \mathcal{H}$ be a Hilbert space with $\mathcal{H}_h \cong \mathbb{R}^{N_h}$, a scalar product $\langle \cdot, \cdot \rangle_{\mathcal{H}_h}$ and a corresponding norm $\|\cdot\|_{\mathcal{H}_h}$, such that the resulting full order model solution can be identified using the formulation

$$t \mapsto u_h(\cdot, t; \mu, \nu) = \sum_{j=1}^{N_h} [u_h^{\mu, \nu}(t)]_j \varphi_j(\mu) \in C^1([0, T]; \mathcal{H}_h) \quad (3.1)$$

with $\{\varphi_j(\mu)\}_{j=1}^{N_h}$ a basis of $\mathcal{H}_h$ for each $\mu \in \Theta$ and $u_h^{\mu,\nu}(t) \in \mathbb{R}^{N_h}$ for each $t \in [0, T]$, and the underlying problem for the coefficients

$$\begin{cases} M(\mu) \frac{\partial}{\partial t} u_h^{\mu,\nu}(t) + A(\mu, \nu) u_h^{\mu,\nu}(t) + N(u_h^{\mu,\nu}(t), \mu, \nu) = f(t; \mu, \nu) & t \in [0, T], \\ u_h^{\mu,\nu}(0) = u_0(\mu, \nu) & t = 0, \end{cases} \quad (3.2)$$

where we assume

- $M : \Theta \rightarrow \mathbb{R}^{N_h \times N_h}$ to send the geometrical parameter to a mass matrix,
- $A : \Theta \times \Theta' \rightarrow \mathbb{R}^{N_h \times N_h}$ to send the set to a symmetric positive definite stiffness matrix,
- $N : \mathbb{R}^{N_h} \times \Theta \times \Theta' \rightarrow \mathbb{R}^{N_h}$ to be some non-linear term,
- $f : [0, T] \times \mathbb{R}^{N_h} \times \Theta \times \Theta' \rightarrow \mathbb{R}^{N_h}$ to be the corresponding source term of the equation.

Additionally let $u_0 : \Theta \times \Theta' \rightarrow \mathbb{R}^{N_h}$ be the initial data. The mass matrix is defined for each $\mu \in \Theta$ as follows

$$M(\mu) = \begin{bmatrix} \langle \varphi_1(\mu), \varphi_1(\mu) \rangle_{\mathcal{H}_h} & \cdots & \langle \varphi_1(\mu), \varphi_{N_h}(\mu) \rangle_{\mathcal{H}_h} \\ \vdots & \ddots & \vdots \\ \langle \varphi_{N_h}(\mu), \varphi_1(\mu) \rangle_{\mathcal{H}_h} & \cdots & \langle \varphi_{N_h}(\mu), \varphi_{N_h}(\mu) \rangle_{\mathcal{H}_h} \end{bmatrix}.$$

This allows us to equip $\mathcal{S} \subset \mathbb{R}^{N_h}$, the space of the coefficients of any solution in $\mathcal{H}_h$, with the following inner product

$$\langle u_h^{\mu,\nu,t}, u_h^{\mu',\nu',t'} \rangle_M := (u_h^{\mu,\nu,t})^T M(\mu) u_h^{\mu',\nu',t'}, \quad (3.3)$$

for each $\mu, \mu' \in \Theta, \nu, \nu' \in \Theta'$ and $t, t' \in [0, T]$ and an induced norm $\|\cdot\|$ which is, in fact, the equivalent of the norm above, i.e. $\|u_h^{\mu,\nu,t}\| = \|u_h(\cdot, t; \mu, \nu)\|_{\mathcal{H}_h}$.

Note that it is indeed possible to arrive at that discretization without skipping to a weak formulation, e.g. using finite differences or finite volume schemes, but since any existence statement needs a more general space than C^n to guarantee completeness and hence imposes some identity using a basis, we need only to infer such a procedure. For the interested reader, an explicit discretization using finite volume schemes can be seen in (Haasdonk & Ohlberger, 2008).

This formulation establishes smoothness with regard to time, however we will implicitly further discretize in time using $\{t_k\}_{k=1}^{N_t} := \mathcal{T}$ time steps by gathering data only at these points. This is called *method of lines approach* (Hesthaven et al., 2022, p. 278).

Definition 3.1 (Solution Data). Following this, we may denote with

$$u_h^{\mu,\nu,t} := u_h^{\mu,\nu}(t) \in \mathbb{R}^{N_h}, \quad \text{for } \mu \in \Theta, \nu \in \Theta' \text{ and all } t \in \mathcal{T},$$

a finite dimensional *solution snapshot* and

$$u_h^{\mu,\nu} := [u_h^{\mu,\nu}(t)]_{t \in \mathcal{T}} \in \mathbb{R}^{N_h \times N_t}, \quad \text{for } \mu \in \Theta \text{ and } \nu \in \Theta',$$

with $N_t := |\mathcal{T}|$ a finite dimensional *solution trajectory* of the discretized parametric PDE. Furthermore, let $\mathcal{P}_1 \subset \Theta$, $\mathcal{P}_2 \subset \Theta'$ and $\mathcal{P} = \mathcal{P}_1 \times \mathcal{P}_2$ be some finite parameter data with respective sizes $N_{s_1}, N_{s_2} \in \mathbb{N} \cup \{\infty\}$ and $N_s = N_{s_1} N_{s_2}$.

Mainly, we stated the problem framework, to know with which class of PDEs we will be working with. Hence any regularity assumption is only given here in order to assert an outlook on the use of L^p norms for the error between the weak solution $u \in \mathcal{H}$, the finite dimensional solution $u_h \in \mathcal{H}_h$ and the approximation of that solution $\hat{u}_h \in \mathcal{H}_h$ (of reduced order) using ROM methods.

It is the latter ones, we will be concerned with in this thesis, which are however clearly embedded in L^2 , since they are finite dimensional and bounded by an upcoming assumption.

From this point forward, we want to consistently rely on the following few assumptions, which regulate the discrete problem above. The first Assumption (1) justifies a nuanced approach regarding the geometric parameter μ and the physical parameter ν . The second Assumption (2) is necessary for the derivation of any convergence result to a true parameter-to-solution map, when sampling an ever growing number of examples, while the third Assumption (3) imposes regularity on the mentioned parameter-to-solution map. Interestingly enough, it is that Assumption, which ensures the existence of a true solution of the discrete problem for any specified parameter choice. Observing of course, that this does not translate into the solvability for the upstream general problem. The last Assumption (4) gives us the ability to later use an Autoencoder for a reduction.

Assumption 1. Let $\mathcal{S} := \{u_h^{\mu,\nu,t} \in \mathbb{R}^{N_h} \mid (\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]\}$ denote the *total solution manifold* and $\mathcal{S}_{\mu,t} := \{u_h^{\mu,\nu,t} \in \mathbb{R}^{N_h} \mid \nu \in \Theta'\}$ denote the *solution submanifold* for a fixed set $(\mu, t) \in \Theta \times [0, T]$, then we assume that

1. the linear KnW of $\mathcal{S}$ decays slowly, i.e.

$$d_N(\mathcal{S}) \leq CN^{-\alpha},$$

where $0 < \alpha \leq 1$ and $C > 0$ some constant.

2. the linear KnW of $\mathcal{S}_{\mu,t}$ decays quickly for all $\mu \in \Theta$ and $t \in [0, T]$, i.e.

$$\sup_{(\mu,t) \in \Theta \times [0,T]} d_N(\mathcal{S}_{\mu,t}) \leq C'N^{-\beta},$$

where $\beta > 1$ and $C' > 0$ some constant.

This is mostly satisfied for problems in which the transport dominated or non-linear part is chained to the parameter μ , and any coercive or elliptic operator is most heavily influence by ν .

Assumption 2. Let Θ, Θ' and $[0, T]$ for $T > 0$ as above. We assume that all $N_s \in \mathbb{N}_{\geq 2}$ parameter data points according to Definition (3.1) are sampled uniformly and iid in the joint parameter space $\Theta \times \Theta'$, while a uniform grid is employed for the time variable, $\mathcal{T} = \{\Delta t, 2\Delta t, \dots, N_t \Delta t\}$, where $N_t \in \mathbb{N}_{\geq 2}$, N_{s_1} and N_{s_2} are the respective number of random samples in the parameter spaces Θ and Θ' , $N_s = N_{s_1} N_{s_2} \in \mathbb{N}$ is the total sample size, and $\Delta t = T/N_t$.

We remark, that the equidistant spacing of the time samples is a matter of convenience, and could be loosened under some constraints.

Assumption 3. Let $\mathcal{G} : \Theta \times \Theta' \times [0, T] \rightarrow \mathbb{R}^{N_h}$ be the parameter-to-solution map, mapping $(\mu, \nu, t) \mapsto u_h^{\mu,\nu,t}$. Here we assume that

1. $m = \operatorname{ess\,inf}_{(\mu,\nu,t) \in \Theta \times \Theta' \times \mathcal{T}} \|u_h(\mu, \nu, t)\| > 0$, $M = \operatorname{ess\,sup}_{(\mu,\nu,t) \in \Theta \times \Theta' \times \mathcal{T}} \|u_h(\mu, \nu, t)\| < \infty$,
2. $\mathcal{G}$ is Lipschitz-continuous with the constant $L > 0$.

A very important fact follows already directly from the presumably allowed well-posed definition of such a map; the solution $u_h(\cdot, \cdot; \mu, \nu)$ exists and is unique, in at least an L^2 -embedded sense, since the time-derivate can be relaxed using a variational argument.

Assumption 4. Let $N \in \mathbb{N}$ be fixed. We assume that for s, s' sufficiently large, there are two maps $\psi_* : \mathbb{R}^n \rightarrow \mathbb{R}^N$ and $\psi'_* : \mathbb{R}^N \rightarrow \mathbb{R}^n$ which attain the perfect nonlinear embedding of the reduced solution manifold, as given in Definition (2.17) for any $n \leq 2(p + q) + 3$, i.e. for $\mathcal{S}_N := \{q(\mu, \nu, t) = \mathbb{A}^T u(\mu, \nu, t) \mid (\mu, \nu, t) \in \Theta \times \Theta' \times \mathcal{T}\}$, where $\mathbb{A} \in \mathbb{R}^{N_h \times N}$ is the POD matrix,

$$\delta_n(\mathcal{S}_N) = 0.$$

Every following Reduced Order Model will pursue the universal (and vague) following goal.

Goal 1. Find

$$\hat{\mathcal{G}} : \Theta \times \Theta' \times [0, T] \rightarrow \mathbb{R}^{N_h}; \quad (\mu, \nu, t) \mapsto \hat{u}_h^{\mu, \nu, t},$$

such that for all $\mu \in \Theta$ and $\nu \in \Theta'$

$$\sup_{t \in [0, 1]} \|u_h^{\mu, \nu, t} - \hat{u}_h^{\mu, \nu, t}\| = \sup_{t \in [0, 1]} \|u_h(\cdot, t; \mu, \nu) - \hat{u}_h(\cdot, t; \mu, \nu)\|_{\mathcal{H}_h}$$

is small.

In general, we remind ourselves of the starting situation: $\mathcal{H}_h \subset \mathcal{H}$ is a finite dimensional approximation space of our weak problem. As mentioned before, we also have employed discretized time-steps. We therefore have something along the lines of

$$\begin{cases} M(\mu) \hat{\partial}_t u_h^{\mu, \nu}(t) + A(\mu, \nu) u_h^{\mu, \nu}(t) + N(u_h^{\mu, \nu}(t), \mu, \nu) = f(t; \mu, \nu) & t \in \mathcal{T}, \\ u_h^{\mu, \nu}(0) = u_0(\mu, \nu) & t = 0, \end{cases} \quad (3.4)$$

for $\hat{\partial}_t$ being a discrete approximation of the time-derivative and $u_h^{\mu, \nu}(t) \in \mathbb{R}^{N_h}$ for $t \in [t_k, t_{k+1})$ and some $k = 0, \dots, N_t - 1$ the piecewise constant coefficient vector for the true solution

$$u_h(\cdot, t; \mu, \nu) = \sum_{j=1}^{N_h} [u_h^{\mu, \nu}(t)]_j \varphi_j(\mu) \in \mathcal{H}_h, \text{ for each } t \in [t_k, t_{k+1}) \text{ and } (\mu, \nu) \in \Theta \times \Theta'.$$

3.1 Intrusive Reduced Order Models

In this section we want to introduce some such *intrusive* methods for the dimensionality reduction of the finite dimensional parametric PDE. The name "intrusive" means to symbolize the "intrusion" of the method into the formulation of the PDE and the specifications of its operators. Coinciding with that, there are methods for linear and non-linear problems respectively, as the more simple and light weight reduction techniques can leverage an abstract linear variational problem to achieve a (Petrov-)Galerkin projection, whereas any non-linear problem, so a problem with $N \neq 0$ according to Equation (3.2), utilize the specifications of the non-linear operator to achieve reduction.

Remark 3.2 (Historic Literature). The development of intrusive reduced order models (ROMs) dates back to the late 1970s and 1980s, when methods for projection onto low-dimensional trial spaces were systematically introduced in the context of parametrized PDEs. Early formulations of

the *reduced basis (RB) method* can be found in the control and approximation theory literature, e.g. in the pioneering works of Courant and Hilbert on variational methods (Courant & Hilbert, 1953), and in the first computational implementations by Noor and coworkers in structural mechanics (Noor, 1980). These ideas were later rigorously developed into the modern RB methodology by Patera in the mid-1980s (Patera, 1984), and subsequently advanced by Maday, Rovas, Prud'homme and others in the 1990s and 2000s (Maday et al., 2002; Prud'homme et al., 2002).

In parallel, the POD emerged as a computational tool for obtaining optimal low-rank approximations of dynamical systems, building on statistical formulations dating back to Karhunen and Loève. The introduction of POD-Galerkin for intrusive model reduction was consolidated in applied mechanics and fluid dynamics in the 1990s and early 2000s (see (Holmes et al., 1996; Kunisch & Volkwein, 2001)).

As nonlinear problems became central, the cost of evaluating nonlinear terms prompted the development of *hyper-reduction techniques*, such as the Empirical Interpolation Method (EIM) (Barraut et al., 2004) and its discrete variants (DEIM) (Chaturantabut & Sorensen, 2010), which remain core components of modern intrusive ROM frameworks.

We will start with the framework of projection-based methods, that necessitate a problem with $N = 0$.

All of the following methods will be discussed broadly, as the thesis concerns itself mostly with the later data-driven ROMs.

Definition 3.3 (Reduced Basis). Let $\mathcal{H}_n \subset \text{span}(u_h(\cdot, t; \mu, \nu) \in \mathcal{H}_h \mid (\mu, \nu, t) \in \mathcal{P}_1 \times \mathcal{P}_2 \times \mathcal{T})$ be an n -dimensional subspace of $\mathcal{H}_h$, then if $\{\hat{\varphi}_j\}_{j=1}^n$ is an orthonormalized basis of $\mathcal{H}_n$, we call this the *reduced basis*.

From this starting point, the goal of all *reduced basis methods* is to find a reduced basis $\{\hat{\varphi}_j(\mu)\}_{j=1}^n$ and $\hat{u}_h^{\mu, \nu}(t) \in \mathbb{R}^n$ for $n \ll N_h$, such that

$$\hat{u}_h(\cdot, t; \mu, \nu) = \sum_{j=1}^n [\hat{u}_h^{\mu, \nu}(t)]_j \hat{\varphi}_j(\mu) \in \mathcal{H}_n \subset \mathcal{H}_h,$$

where, in a yet vague formulation,

$$\hat{u}_h(\cdot, t; \mu, \nu) \approx u_h(\cdot, t; \mu, \nu) \quad \text{for all } \mu \in \Theta, \nu \in \Theta' \text{ and } t \in [0, T].$$

In the terminology the RB methods are a setup for many different ROMs, but most often appear for so-called projection-based ROMs, which itself are intrusive ROMs. They help with the identification of a suitable basis to project upon. Two prominent examples of such reduced basis methods in the context of projection-based ROMs can be found in the following. Both will exhibit intrinsically an *offline* and an *online* part in the algorithm, whose separation we will mostly neglect. However, when remarking something about the offline procedure, we usually talk about the part of the algorithm that can be collected a priori to its purposed deployment. Here this is the collection and reduction of dimension using a reduced basis via samples. On the other hand, online refers to its counterpart; the many-query element of the procedure, such as the evaluation of the best-guess parameter-to-solution map. This terminology is used often throughout the literature as it instinctively makes sense in a software development environment.

3.1.1 Galerkin Proper Orthogonal Decomposition

As a prerequisite for Galerkin orthogonality, we assume here that the weak form giving rise to the equation at hand derives from an abstract bilinear form in the stationary part. This is justified if the non-linear summand is zero. The source for this subsection is (Quarteroni et al., 2015, p. 137).

The POD as introduced in Definition (2.13) is one of the most common methods for a reduction in a classical sense. We can use the system above to solve for the solution using a Galerkin projection for the following Ansatz:

$$\hat{u}_h(x, t; \mu, \nu) = \sum_{j=1}^n [\hat{u}_h^{\mu, \nu}(t)]_j \xi_j(x), \quad (3.1)$$

where $x \in \Omega$, $t \in [0, 1]$, $(\mu, \nu) \in \Theta \times \Theta'$ and ξ_j for $1 \leq j \leq n$ are the POD modes, as given by the construction of the POD in Definition (2.13) with $\mathbb{G} := \mathbb{I}_{N_h}$, and $\hat{u}_h^{\mu, \nu}(t) \in \mathbb{R}^n$ is the coefficient vector, that solves the following reduced equation

$$\begin{cases} \mathbb{A}^T M(\mu) \mathbb{A} \hat{\partial}_t \hat{u}_h^{\mu, \nu}(t) + \mathbb{A}^T A(\mu, \nu) \mathbb{A} \hat{u}_h^{\mu, \nu}(t) = \mathbb{A}^T f(t; \mu, \nu) & t \in \mathcal{T}, \\ \hat{u}_h^{\mu, \nu}(0) = u_0(\mu, \nu) & t = 0. \end{cases} \quad (3.2)$$

Note, that in this construction $\mathbb{A}$ is *not* $M(\mu)$ -orthogonal, thus losing certain projection qualities with regard to the inherent L^2 -norm of the underlying Hilbert space $\mathcal{H}_h$. The parameter-to-solution map can be recovered by the following Algorithm (2).

Algorithm 2 POD-Galerkin

- 1: **Input:** $(\mu_j, \nu_j)_{j=1}^{N_s}$ data sample and $n < N_h$ reduced dimension
 - 2: **Output:** approximated parameter-to-solution-map $(\mu, \nu) \mapsto \hat{u}_h(\cdot, \cdot; \mu, \nu)$
 - 3: Compute $\{u_h^{\mu_j, \nu_j, t_{km}}\}_{m=1}^{N_t}$ from Equation (3.4) (with $N = 0$)
 - 4: $(\xi_j, \sigma_j)_{j=1}^n \leftarrow \text{POD}(u_h^{\mu_j, \nu_j, t_{km}}, n)$
 - 5: $\mathbb{A} \leftarrow \text{stack}(\xi_j, j = 1, \dots, n)$
 - 6: Build reduced Equation (3.2)
 - 7: For each (μ, ν) solve Equation (3.2)
 - 8: Recover $\hat{u}_h(\cdot, \cdot; \mu, \nu)$ via Equation (3.1)
-

3.1.2 Greedy Proper Orthogonal Decomposition

An introduction to this method in the stationary and well-posed case can be found in (Siena et al., 2022). As done previously in the subsection above, we assume $N = 0$ in Equation (3.4). This gives us a coercive abstract bilinear form for the stationary part. Additionally let $\eta : \Theta \times \Theta' \rightarrow \mathbb{R}$ be a function estimating the error of projection under a specific parameter set, i.e. such that

$$\sup_{t \in \mathcal{T}} \|u_h(\mu, \nu, t) - \hat{u}_h(\mu, \nu, t)\| \leq \eta(\mu, \nu),$$

for $(\mu, \nu) \in \mathcal{P}$. Purely using POD-Galerkin can encounter major pitfalls, namely if the range of $\mathcal{P}$ is extensive, since then calculating a singular value decomposition for the correlation matrix might get too cumbersome to provide all areas of interest in the set. Moreover, to obtain the snapshots for fixed parameters, a complete time-trajectory must be computed, which entails a lot of overhead of otherwise useless information. To overcome this (Grepl & Patera, 2005) proposed, and later (Haasdonk & Ohlberger, 2008) improved upon, the mixed POD-Greedy approach, which takes a POD over the time-trajectories (or, in the improved version, the error trajectories) while using the Greedy approach to handle the parameter samples. It follows a similar workflow, as the POD-Galerkin, since the only thing, that changes is the choice of basis, and the fact that the size of the basis is chosen according to the tolerance in the offline part of the algorithm, i.e.

$$\hat{u}_h(x, t; \mu, \nu) = \sum_{j=1}^N [\hat{u}_h^{\mu, \nu}(t)]_j \xi_j(x), \quad (3.3)$$

with $x \in \Omega$, $t \in [0, 1]$, $(\mu, \nu) \in \Theta \times \Theta'$ identifies the reduced solution of dimension N , where ξ_j are the basis functions obtained from the offline procedure in Algorithm (3) for $1 \leq j \leq N$ and the coefficient vector being the solution to

$$\begin{cases} \mathbb{V}_{\text{rb}}^T M(\mu) \mathbb{V}_{\text{rb}} \hat{\partial}_t \hat{u}_h^{\mu, \nu}(t) + \mathbb{V}_{\text{rb}}^T A(\mu, \nu) \mathbb{V}_{\text{rb}} \hat{u}_h^{\mu, \nu}(t) = \mathbb{V}_{\text{rb}}^T f(t; \mu, \nu) & t \in \mathcal{T}, \\ \hat{u}_h^{\mu, \nu}(0) = u_0(\mu, \nu) & t = 0. \end{cases} \quad (3.4)$$

Note again, that $\mathbb{V}_{\text{rb}}$ is *not* $M(\mu)$ -orthonormal. The complexity of the offline computation for this Algorithm (3) is additive, and not multiplicative, in the number of training data $\mathcal{P}$, which is a huge advantage when compared to the direct POD-Galerkin approach.

Algorithm 3 POD-Greedy-Galerkin

```

1: Input:  $\text{tol}$ ,  $\mathcal{Z} = 0$ ,  $N = 0$ ,  $(\mu_j, \nu_j)_{j=1}^{N_s}$ ,  $N_1$  and  $N_2$ 
2: Output: approximated parameter-to-solution-map  $(\mu, \nu) \mapsto \hat{u}_h(\cdot, \cdot; \mu, \nu)$ 
3: for  $j = 1, \dots, N_s$  do
4:   Compute  $\{u_h^{\mu_j, \nu_j, t_{k_m}}\}_{m=1}^{N_t}$  from Equation (3.4) (with  $N = 0$ )
5:    $(\zeta_j, \lambda_j)_{j=1}^{N_1} \leftarrow \text{POD}(\{u_h^{\mu_j, \nu_j, t_{k_m}}\}_{m=1}^{N_t}, N_1)$ 
6:    $\mathcal{Z} \leftarrow \{\mathcal{Z}, \{\zeta_1, \dots, \zeta_{N_1}\}\}$ 
7:    $N \leftarrow N + N_2$ 
8:   Compress:  $(\xi_j, \sigma_j)_{j=1}^n \leftarrow \text{POD}(\mathcal{Z}, N)$ 
9:    $(\mu_{j+1}, \nu_{j+1}) \leftarrow \arg \max_{(\mu, \nu) \in \mathcal{P}} \eta(\mu, \nu)$ 
10:  if  $\eta(\mu_{j+1}, \nu_{j+1}) > \text{tol}$  then
11:     $j \leftarrow j + 1$ 
12:  else
13:    break
14:  end if
15: end for
16:  $\mathbb{V}_{\text{rb}} \leftarrow \text{stack}(\xi_1, \dots, \xi_N)$ 
17: Build reduced Equation (3.4)
18: For each  $(\mu, \nu)$  solve Equation (3.4)
19: Recover  $\hat{u}_h(\cdot, \cdot; \mu, \nu)$  via Equation (3.3)
    
```

Remark 3.4. It would be possible to include the physical parameter into the first POD scheme in each iteration, which could give a high yield for this specific problem as the Kolmogorov n -width decay as imposed by Assumption (1) implies. However since we lose generality by setting $N = 0$ in both projection-based algorithm, this is only an argument for divergent discussions.

3.1.3 Non-linear Reduced Order Models

Add some literature

3.2 Data-driven Reduced Order Models

In contrast to the introduced intrusive ROMs above, the *data-driven* alternatives aim to circumvent the direct use of the governing equations, and, as such, allow for more leniency toward unknown

operator qualities.

Remark 3.5 (Historic Literature). The origins of data-driven ROMs can be traced back to techniques in fluid dynamics and control, where the objective was to extract coherent structures or dominant modes from experimental measurements without explicit reliance on the full Navier–Stokes equations. A first influential step was the development of the POD for experimental data, introduced in the turbulence community by Lumley (Lumley, 1967), and further popularized in the coherent structure analysis of Sirovich (Sirovich, 1987). While POD itself does not provide a dynamical law for the online phase, it laid the foundation for representing complex systems with a small number of empirical modes.

A decisive breakthrough came with the advent of the *Dynamic Mode Decomposition* (DMD) (Schmid, 2010), which may be interpreted as a data-driven approximation of the infinite-dimensional Koopman operator (Rowley et al., 2009). DMD represents temporal evolution as a best-fit linear operator acting on snapshot data, thereby yielding both modal decompositions and a surrogate model for prediction. This was one of the first systematic attempts to construct a fully non-intrusive ROM.

In parallel, regression-based approaches have been developed to learn reduced operators directly from data, often under the name of *operator inference* (Peherstorfer et al., 2016), or in the form of interpolation-based reduced models, e.g. POD with interpolation (PODI) as in Barrault et al. (Barrault et al., 2004). These methods demonstrate that the reduced dynamics can be inferred without Galerkin projection, either by statistical regression or by interpolation in parameter space.

For this section we intentionally do not give an explicit training algorithm for any of the methods presented, since it can be inferred with an optimizer of the readers choice from the discussion in Subsection (2.3.5). With readability in mind, we thus only state the architecture of the network at hand and its loss function. Most importantly, we assume that some choice of basis for $\mathcal{H}_h$ is already done a priori to the training of the network and independent of $\mu \in \Theta$ thus setting $\mathbb{G} := M(\mu)$ for all $\mu \in \Theta$ as our mass matrix. Under Equation (3.3) this also implies a μ -independent norm $\|\cdot\|$, which in fact allow us to use the POD matrix, if constructed with $\mathbb{G}$, as a true projector onto the span of $\mathbb{A}$ in an L^2 sense.

For clarity, we formulate the following as an assumption, since from experience, the reduction system across comparable architectures can get confusing, and the reader is therefore encouraged to revisit this assumption, whenever the general setting appears to be unclear.

Assumption 5 (Dimension Hierarchy). Let for all following data-driven Reduced Order Models, the following natural number hierarchy be fixed

$$n < N' < N \ll N_A \leq N_h.$$

Note, that " $\ll$ " is an unspecified relation, meaning to symbolize a "large" difference.

A general Deep Learning based ROM (DL-ROM) first appeared in architecture and training in the paper (Fresca et al., 2020); to properly approximate the potentially non-linear solution manifold of the high fidelity FOM, we employ an Autoencoder (AE), and on the reduced manifold we employ some other Deep Learning architecture, like the proposed Deep Feed-Forward Neural Network (DFNN), to learn the latent dynamics of the parametric PDE. This concludes in the following approximation of the FOM coefficient solution $u_h^{\mu,\nu,t} \in \mathbb{R}^{N_h}$ for each $\mu \in \Theta$, $\nu \in \Theta'$ and $t \in \mathcal{T}$

$$\hat{u}_h^{\mu,\nu,t} = (R_\rho(\Psi_h(\theta_D^*)) \circ R_\rho(\Phi_n(\theta_{DF}^*))) (t; \mu, \nu) \in \mathbb{R}^{N_h}, \quad (3.1)$$

where we set $\Psi_h = f_h^D$ for the decoder architecture with input dimension n and output dimension N_h (and $\Psi'_n = f_n^E$ the corresponding encoder architecture) of a convolutional AE. Then Ψ'_n denotes

the projection onto the reduced trial solution manifold and we let $\Phi_n = \Phi_n^{\text{DF}}$ to be the DFNN architecture of input dimension $p + q + 1$ and output dimension n , mapping the dynamics of the parametric PDE onto the reduced trial solution manifold.

The subsequent training is then performed via a loss function of a convex combination of both, the projection error using the encoder and the dynamics error considering both the decoder and the feed forward network, i.e. for $\theta = (\theta_D, \theta_E, \theta_{\text{DF}})$ define the optimal weights via the problem in Definition (2.37) with the loss function

$$\begin{aligned} \mathcal{L}_{\text{DL-ROM}} \left(R_\rho(\Psi_h \odot \Phi_n(\theta)) | \{ \mu_i, \nu_j, k\Delta t \}_{i,j,k}, \{ u_h^{\mu_i, \nu_j, k\Delta t} \}_{i,j,k} \right) = \\ \sum_{i=1}^{N_{s1}} \sum_{j=1}^{N_{s2}} \sum_{k=1}^{N_t} \frac{\omega_h}{2} | u_h^{\mu_i, \nu_j, k\Delta t} - (R_\rho(\Psi_h(\theta_D)) \circ R_\rho(\Phi_n(\theta_{\text{DF}}))) (\mu_i, \nu_j, k\Delta t) |^2 \\ + \frac{1 - \omega_h}{2} | R_\rho(\Psi'_n(\theta_E))(u_h^{\mu_i, \nu_j, k\Delta t}) - R_\rho(\Phi_n^{\text{DF}}(\theta_{\text{DF}}))(\mu_i, \nu_j, k\Delta t) |^2. \end{aligned} \quad (3.2)$$

3.2.1 Proper Orthogonal Decomposition Deep Learning ROM

The main idea of the POD-DL-ROM is the additional employment of an outer a priori reduction of DL-ROM problem using a POD. It can be found in (Fresca & Manzoni, 2022). Note, that we sway a bit from the source, since we are trying to achieve an estimate, which is close to the high-fidelity solution in an L^2 -sense and not a euclidian one by applying the mass matrix from the start of the section. Such a reduction yields great benefits, such as the scalability in N_h and reduction in computational cost for the offline training. It is done by employing a POD (see Algorithm (1)) on the stacked solution trajectories

$$U := \left[u_h^{\mu_1, \nu_1} | \dots | u_h^{\mu_{N_{s1}}, \nu_1} | \dots | u_h^{\mu_{N_{s1}}, \nu_{N_{s2}}} \right]^T \in \mathbb{R}^{N_h \times N_{\text{data}}},$$

with the solution trajectories of Definition (3.1), resulting in the POD matrix $\mathbb{A}_P \in \mathbb{R}^{N_h \times N}$ for the $N < N_h$ from Assumption (5). Then we can define the pre-reduced solution snapshots by computing

$$u_{\text{red}}^{\mu, \nu, t} := \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t} \in \mathbb{R}^N \quad \text{for all } (\mu, \nu, t) \in \mathcal{P}_1 \times \mathcal{P}_2 \times \mathcal{T}.$$

By $\mathbb{G}$ -orthonormality of $\mathbb{A}_P$, we can thus define the FOM approximate coefficient solution $\hat{u}_h^{\mu, \nu, t} \in \mathbb{R}^{N_h}$ for each $\mu \in \Theta$, $\nu \in \Theta'$ and $t \in \mathcal{T}$

$$\hat{u}_h^{\mu, \nu, t} = \mathbb{A}_P \cdot (R_\rho(\Psi_N(\theta_D^*)) \circ R_\rho(\Phi_n(\theta_{\text{DF}}^*))) (t; \mu, \nu) \in \mathbb{R}^{N_h}, \quad (3.3)$$

where we set $\Psi_N = f_N^D$ for the decoder architecture with input dimension n and output dimension N (and $\Psi'_n = f_n^E$ the corresponding encoder architecture) of a convolutional AE. Then f_n^E denotes the projection onto the reduced trial solution manifold and we let $\Phi_n = \Phi_n^{\text{DF}}$ to be the DFNN architecture of input dimension $p + q + 1$ and output dimension n same as before.

The loss function has now the attribute of being fed only at most N dimensional training data, and only evaluating at most N dimensional euclidian norms, hence greatly increasing performance, i.e. with P being the network parallelization operator

$$\begin{aligned} \mathcal{L}_{\text{POD-DL-ROM}} \left(R_\rho(P(\Psi_N \odot \Phi_n, \Psi'_n)(\theta)) | \{ \mu_i, \nu_j, k\Delta t \}_{i,j,k}, \{ u_{\text{red}}^{\mu_i, \nu_j, k\Delta t} \}_{i,j,k} \right) = \\ \frac{1}{N_{\text{data}}} \sum_{i=1}^{N_{s1}} \sum_{j=1}^{N_{s2}} \sum_{k=1}^{N_t} \left(\frac{\omega_h}{2} | u_{\text{red}}^{\mu_i, \nu_j, k\Delta t} - (R_\rho(\Psi_N(\theta_D)) \circ R_\rho(\Phi_n(\theta_{\text{DF}}))) (\mu_i, \nu_j, k\Delta t) |^2 \right. \\ \left. + \frac{1 - \omega_h}{2} | R_\rho(\Psi'_n(\theta_E))(u_{\text{red}}^{\mu_i, \nu_j, k\Delta t}) - R_\rho(\Phi_n(\theta_{\text{DF}}))(\mu_i, \nu_j, k\Delta t) |^2 \right). \end{aligned} \quad (3.4)$$

add a justification of the use of the reduced space and euklidian norm as done in franco2024

3.2.2 Deep Orthogonal Decomposition based ROMs

A big advantage of the POD-DL-ROM is the fact, that under the assumption of a static basis $\{\xi_j\}_{j=1}^{N_h} \subset \mathcal{H}_h$ of $\mathcal{H}_h$ and its associated mass matrix $\mathbb{G}$ and $\mathcal{H}_h$ equivalent norm $\|\cdot\|$, we may reduce the high dimensional loss function in its essence to a manageable computational complexity. However, this so far ignored the particular problem in sight; it ignores Assumption (1), which guarantees a significant reduction cost, when employing a (μ, t) -independent projector. To circumvent this Kolmogorov N -width barrier for $N \ll N_h$, one can instead of using a POD for the choice of reduced basis, invoke a Neural Network based linear projector, that fits a basis superposition dependend on μ and t , and can thus exhibit a further reduction to $N' < N$ under a similar value for the worst-case projection, i.e. the Kolmogorov N' -width.

The following technically depicts a RB method.

Definition 3.6 (Deep Orthogonal Decomposition). Recollect the dimensional hierarchy of Assumption (5). Employ a POD (according to Algorithm (1)) with $\mathbb{G}$ given as above using the stacked solution trajectories

$$U := \left[u_h^{\mu_1, \nu_1} \mid \dots \mid u_h^{\mu_{N_{s1}}, \nu_1} \mid \dots \mid u_h^{\mu_{N_{s1}}, \nu_{N_{s2}}} \right]^T \in \mathbb{R}^{N_h \times N_{\text{data}}},$$

with the solution trajectories of Definition (3.1), resulting in the POD matrix $\mathbb{A} \in \mathbb{R}^{N_h \times N_A}$. Now set up for the inner module of the Deep Orthogonal Decomposition a Neural Network architecture $\Phi_{\tilde{\mathbb{V}}}$ with input dimension $p+1$ and output dimension $N_A \times N'$ such that

$$\Phi_{\tilde{\mathbb{V}}} = \text{ORTH} \circ \text{STACK} \circ (P((R_1 \odot s), \dots, (R_N \odot s)))(\mu, t),$$

where P is denoting the network parallelization operator as given in Definition (2.27)

1. s is a Deep Feed Forward Neural Network architecture with input dimension $p+1$ and output dimension l ,
2. $R_1, \dots, R_N$ are a collection of Deep Feed Forward Neural Network architectures with input dimensions l and output dimensions N_A ,
3. $\text{STACK} : \mathbb{R}^{N_A \times N'} \rightarrow \mathbb{R}^{N_A \times N'}$ is a canonical stacking operator,
4. $\text{ORTH} : \mathbb{R}^{N_A \times N'} \rightarrow \mathbb{R}^{N_A \times N'}$ is an orthonormalization unit, that orthonormalizes any given matrix $W \in \mathbb{R}^{N_A \times N'}$ into $\tilde{W} \in \mathbb{R}^{N_A \times N'}$ such that $\text{span}(W) = \text{span}(\tilde{W})$. This is most often done using a QR decomposition (Golub & Van Loan, 2013).

The realization of the network $\Phi_{\tilde{\mathbb{V}}}$ for some $\mu \in \Theta$, $t \in [0, T]$ and any chosen neural network weight $\theta_{\text{DOD}} \in \Theta(\Phi_{\tilde{\mathbb{V}}})$ according to Definition (2.37) will be abbreviated with

$$\tilde{\mathbb{V}}_{\mu, t}(\theta_{\text{DOD}}) := R_\rho(\Phi_{\tilde{\mathbb{V}}}(\theta_{\text{DOD}}))(\mu, t). \quad (3.5)$$

For any choice of weights, this is called the *inner module of the Deep Orthogonal Decomposition*, or in short the inner DOD module.

The *Deep Orthogonal Decomposition* can hence be achieved by upward projection under the POD matrix $\mathbb{A}$ of the inner DOD module, i.e. a realization of the underlying DOD neural network for a choice of weights $\theta_{\text{DOD}} \in \Theta(\Phi_{\tilde{\mathbb{V}}})$ is attained for some $\mu \in \Theta$ and $t \in [0, T]$ by

$$\mathbb{V}_{\mu, t}(\theta_{\text{DOD}}) := \mathbb{A} \cdot \tilde{\mathbb{V}}_{\mu, t}(\theta_{\text{DOD}}) \in \mathbb{R}^{N_h \times N'}. \quad (3.6)$$

Nonetheless this does not warrant a "good" DOD, just a defined one. For this one needs to assert a qualitative loss function and some data, which we will collect by pre-reduction in the same manner as before, i.e.

$$u_{\text{pre-red}}^{\mu,\nu,t} := \mathbb{A}^T \mathbb{G} u_h^{\mu,\nu,t} \in \mathbb{R}^{N_A} \quad \text{for all } (\mu, \nu, t) \in \mathcal{P}_1 \times \mathcal{P}_2 \times \mathcal{T}.$$

Then the loss function with regard to that data can be defined as follows

$$\begin{aligned} \mathcal{L}_{\text{DOD}} \left(R_\rho(\Phi_{\tilde{\mathbb{V}}}(\theta_{\text{DOD}})) | \{ \mu_i, \nu_j, k\Delta t \}_{i,k}, \{ u_{\text{pre-red}}^{\mu_i, \nu_j, k\Delta t} \}_{i,j,k} \right) \\ = \frac{1}{N_{\text{data}}} \sum_{i=1}^{N_{s1}} \sum_{j=1}^{N_{s2}} \sum_{k=1}^{N_t} | u_{\text{pre-red}}^{\mu_i, \nu_j, k\Delta t} - \tilde{\mathbb{V}}_{\mu_i, k\Delta t}(\theta_{\text{DOD}}) \tilde{\mathbb{V}}_{\mu_i, k\Delta t}^T(\theta_{\text{DOD}}) u_{\text{pre-red}}^{\mu_i, \nu_j, k\Delta t} |^2. \end{aligned} \quad (3.7)$$

Note, that this is indeed again a N_A -dimensional loss function, thanks to the $\mathbb{G}$ -orthonormality of the POD matrix. In some abuse of notation, we will sometimes refer to the optimized version of the DOD network as the DOD.

Notice, that we have not yet performed any actual search for the approximate parameter-to-solution map, that maps a specific instance of parameters to a coefficient time-trajectory, that can then be used to build the solution map in the underlying high fidelity Hilber space approximating the FOM solution in Equation (3.1).

In this paper, we propose two novel data-driven ROMs, based on the DOD, to find an approximate solution-to-parameter map. These are the *Deep Orthogonal Decomposition with Deep Feed-Forward Neural Network* (DOD+DFNN) and the *Deep Orthogonal Decomposition based Deep Learning ROM* (DOD-DL-ROM). The former indeed symbolizes a general approach hence including the latter, but, with some abuse of notation, we admit the DOD+DFNN a naive direct Deep Feed Forward architecture for the determination of the coefficient, while the DOD-DL-ROM is an adaptation of the DL-ROM architecture introduced in the header of the section.

DOD+DFNN

Let $\Phi_{N'} := \Phi_{N'}^{\text{DF}}$ be some Deep Feed-Forward Neural Network architecture with input dimension $p + q + 1$ and output dimension N' , then the approximate parameter-to-solution map for a fixed weight $\theta = (\theta_{\text{DOD}}, \theta_{\text{DF}}) \in \Theta(\Phi_{\tilde{\mathbb{V}}}) \times \Theta(\Phi_n^{\text{DF}})$ according to Definition (2.37) can be given for each $(\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]$ via

$$\hat{u}_h^{\mu,\nu,t} = \mathbb{V}_{\mu,t}(\theta_{\text{DOD}}) \cdot R_\rho(\Phi_{N'}(\theta_{\text{DF}}))(\mu, \nu, t) \in \mathbb{R}^{N_h}. \quad (3.8)$$

Subsequently the training is done in series and not in parallel, as the optimal DOD parameter θ_{DOD}^* needs to be found first. Secondly, the $\mathbb{G}$ -orthonormality of the DOD can be leveraged with

$$u_{\text{red}^2}^{\mu,\nu,t} := \mathbb{V}_{\mu,t}(\theta_{\text{DOD}}^*)^T \mathbb{G} u_h^{\mu,\nu,t} \in \mathbb{R}^{N'} \quad \text{for all } (\mu, \nu, t) \in \mathcal{P}_1 \times \mathcal{P}_2 \times \mathcal{T},$$

to further reduce the training data and finally define the loss function for the DOD+DFNN architecture

$$\begin{aligned} \mathcal{L}_{\text{DOD+DNN}} \left(R_\rho(\Phi_{N'}(\theta_{\text{DF}})) | \{ \mu_i, \nu_j, k\Delta t \}_{i,k}, \{ u_{\text{red}^2}^{\mu_i, \nu_j, k\Delta t} \}_{i,j,k} \right) \\ = \frac{1}{N_{\text{data}}} \sum_{i=1}^{N_{s1}} \sum_{j=1}^{N_{s2}} \sum_{k=1}^{N_t} | u_{\text{red}^2}^{\mu_i, \nu_j, k\Delta t} - R_\rho(\Phi_{N'}(\theta_{\text{DF}}))(\mu_i, \nu_j, k\Delta t) |^2. \end{aligned} \quad (3.9)$$

This is a rather obvious extension of (Franco et al., 2024), but works surprisingly well. It will be the basic benchmark to overcome for the following algorithms.

DOD-DL-ROM

Let $\Psi_{N'} := f_{N'}^D$ be a decoder architecture with input dimension n and output dimension N' (and $\Psi'_n := f_n^E$ the corresponding encoder architecture), and $\Phi_n := \Phi_n^{\text{DF}}$ be some Deep Feed-Forward Neural Network architecture with input dimension $p + q + 1$ and output dimension n then the approximate parameter-to-solution map for a fixed weight $\theta = (\theta_{\text{DOD}}, \theta_D, \theta_{\text{DF}}) \in \Theta(\Phi_{\tilde{V}}) \times \Theta(f_N^D) \times \Theta(\Phi_n^{\text{DF}})$ according to Definition (2.37) can be given for each $(\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]$ via

$$\hat{u}_h^{\mu, \nu, t} = \mathbb{V}_{\mu, t}(\theta_{\text{DOD}}) \cdot (R_\rho(\Psi_{N'}(\theta_D)) \circ R_\rho(\Phi_n(\theta_{\text{DF}})))(t; \mu, \nu) \in \mathbb{R}^{N_h} \quad (3.10)$$

Same, as for the DOD+DNN, the training is done in series. One observes, that the training data can be again reduced with

$$u_{\text{red}^2}^{\mu, \nu, t} := \mathbb{V}_{\mu, t}(\theta_{\text{DOD}}^*)^T \mathbb{G} u_h^{\mu, \nu, t} \in \mathbb{R}^N \quad \text{for all } (\mu, \nu, t) \in \mathcal{P}_1 \times \mathcal{P}_2 \times \mathcal{T},$$

and then used to define the DOD-DL-ROM loss function utilizing the $\mathbb{G}$ -orthonormality of the DOD, hence

$$\begin{aligned} \mathcal{L}_{\text{DOD-DL-ROM}} \left(R_\rho(P(\Psi_{N'} \odot \Phi_n, \Psi'_n)(\theta)) | \{\mu_i, \nu_j, k\Delta t\}_{i,j,k}, \{u_{\text{red}^2}^{\mu_i, \nu_j, k\Delta t}\}_{i,j,k} \right) = \\ \frac{1}{N_{\text{data}}} \sum_{i=1}^{N_{s1}} \sum_{j=1}^{N_{s2}} \sum_{k=1}^{N_t} \left(\frac{\omega_h}{2} |u_{\text{red}^2}^{\mu_i, \nu_j, k\Delta t} - (R_\rho(\Psi_{N'}(\theta_D)) \circ R_\rho(\Phi_n(\theta_{\text{DF}})))(\mu_i, \nu_j, k\Delta t)|^2 \right. \\ \left. + \frac{1 - \omega_h}{2} |R_\rho(\Psi'_n(\theta_E))(u_{\text{red}^2}^{\mu_i, \nu_j, k\Delta t}) - R_\rho(\Phi_n^{\text{DF}}(\theta_{\text{DF}}))(\mu_i, \nu_j, k\Delta t)|^2 \right). \quad (3.11) \end{aligned}$$

Notice, that this reduction allows us to elect an algorithm, which uses only the n - and N' -dimensional euklidian norm, therefore enabling even less computational complexity for each training epoch, under the assumption of an already optimized DOD.

3.2.3 CoLoRA-DL-ROM

Introduce CoLoRA DL-ROM

Here we aim at minimizing the dimension through two major reductions; the first being the reduction of the stationary DOD to the latent dimension of the μ -invariant solution manifold of the stationary equivalent to the problem (regarding the Dirichlet boundary conditions as the initial conditions on some a priori defined boundary), and the second concerning the actual reduction with CoLoRA-layered architecture to the latent dynamic dimension $n \ll N_h$. The general workflow is thus given by

$$\hat{u}_h(t; \mu, \nu, \theta_{s\text{DOD}}^*, \theta_C) = \mathbb{A} \cdot \mathcal{C}_L(\dots (\mathcal{C}_2(\mathcal{C}_1(V_0(\mu, \theta_{s\text{DOD}}^*) \cdot \phi_0(\mu, \nu, \theta_{s\text{DOD}}^*)), \theta_{C_1}), \theta_{C_2}), \theta_{C_L}), \quad (3.12)$$

where $\mathcal{C}_i : \mathbb{R}^{N_A} \times \Theta_C \rightarrow \mathbb{R}^{N_A}$; $\mathcal{C}_i(x, \theta_{C_i}) = W_i x + A_i \text{diag}(\alpha_i(\nu, t)) B_i x + b_i$ for all $1 \leq i \leq L$. Remark, that we will still employ a pre-reduction using the rPOD matrix $\mathbb{A}$ stemming from the same snapshot matrix S as in both Options before.

Chapter 4

Approximation Theory for Reduced Order Models

This chapter provides the main results of this thesis based on the data-driven ROMs introduced in Chapter (3). As such we repeat, that in order to execute the reduction for the norm $\|\cdot\|$ defined in that chapter, we require a mass matrix, that is independent of $\mu \in \Theta$, since otherwise the orthonormality with regard to it cannot be computed under the generalized POD Algorithm (1). However, we note, that the paper (Brivio et al., 2023) we base most of this chapter on, and several other works dealing with a similar setup, instead neglect the representation in the underlying Hilbert space and use a simple euklidian norm for the coefficient vectors. For our situation though, we let

$$\hat{u}_h(x, t; \mu, \nu) = \sum_{j=1}^{N_h} [\hat{u}_h^{\mu, \nu}(t)]_j \varphi_j(x), \quad (4.1)$$

for all $x \in \Omega$, $t \in [0, T]$ and parameters $(\mu, \nu) \in \Theta \times \Theta'$, which simultaneously specifies $[\mathbb{G}]_{ij} = \langle \varphi_i, \varphi_j \rangle = [M(\mu)]_{ij}$ as the mass matrix. The norm for this chapter will be, if not stated otherwise, the norm from Equation (3.3), so here

$$\|u\| := \sqrt{u^T \mathbb{G} u},$$

for all $u \in \mathbb{R}^{N_h}$. Furthermore the solution vector $\hat{u}_h^{\mu, \nu}(t) = \hat{u}_h^{\mu, \nu, t} \in \mathbb{R}^{N_h}$ is set to be constant between the time instants in our model, i.e. for all $t \in [(k-1)\Delta t, k\Delta t)$ for some $k = 1, \dots, N_t$. As a goal of this chapter, we will provide upper bounds on the complexity of the networks making up the data-driven ROMs, where we especially outline a comparison between the POD-DL-ROM and the newly introduced DOD-DL-ROM.

In order to analyse any error occuring from a data-based approximation, we need to set up a probability space, in whose sense a convergence under an increased data set can be viewed. This is to be regarded as a clarification of Assumption (2).

Definition 4.1 (Sampling). Let $(\mathcal{P}, \mathcal{B}(\mathcal{P}))$ be a Borel-measurable parameter space and $(\Omega_{\mathbb{P}}, \mathcal{F}, \mathbb{P})$ some underlying probability space. Then we refer to an *independent sampling of size N* , if for each $i \in \{1, \dots, N\}$ the random variable mapping attaining a sample (μ_i, ν_i)

$$X_i : (\Omega_{\mathbb{P}}, \mathcal{F}, \mathbb{P}) \rightarrow (\mathcal{P}, \mathcal{B}(\mathcal{P}))$$

is measurable, its push forward $\mathbb{P}^{X_i}$ admits the same distribution $\mathbb{P}^X$, i.e.

$$\mathbb{P}[X_i \in A] = \mathbb{P}^{X_i}[A] = \mathbb{P}^X[A],$$

and the samples are independent. We further name it *iid sampling*, if the distribution $\mathbb{P}^X$ is the uniform distribution on $\mathcal{P}$.

With this in mind, we can view sampling as the realization of a random variable, that provides an element of the sample space. Therefore any mention of a probability space or a probability measure $\mathbb{P}$ refers to the sampling procedure in the probability space as given in Definition (4.1).

The upcoming proposition is a necessary result for the following chapters, as it provides us with a practical bound on the needed number of samples.

Proposition 4.2. Let $f \in L^2(\Theta \times \Theta' \times [0, T])$. Under Assumption (2), it is true, that

$$\mathbb{E} \left[\left| \int_{\Theta \times \Theta' \times [0, T]} f(\mu, \nu, t) d(\mu, \nu, t) - \frac{|\Theta \times \Theta' \times [0, T]|}{N_{data}} \sum_{i=1}^{N_{s1}} \sum_{j=1}^{N_{s2}} \sum_{k=1}^{N_t} f(\mu_i, \nu_j, k\Delta t) \right| \right] \leq \mathcal{O}(N_s^{-1/2} + N_t^{-1}).$$

Proof. See the Appendix of (Brivio et al., 2023). $\square$

4.1 Error Decomposition Formulas

To adequately start analysing the ROMs in question, one needs to recognize the different aspects of "error" inherent to their deployment. Namely, we can subdivide these separate instances of inaccuracy occurring in the process of both, the POD-based data-driven ROM and the DOD-based data-driven ROM, into an error caused by the imperfect recollection of the parameter spaces Θ and Θ' , an error caused by the imperfect projection given by the DOD or POD respectively and an error of imperfect attribution of latent dynamics on that projection space by a neural network. This separation can be achieved in a similar manner for both types of algorithms.

Before we start with the formulation and proof of these decompositions, we want to predefine some elements of the corresponding theorems, to make them less cumbersome to read.

Firstly let in general n, N', N, N_A and N_h be the fixed natural number hierarchy from Assumption (5). Let the approximate POD-based parameter-to-solution map be defined as follows

$$\mathcal{G}_{\text{POD+DNN}} : \Theta \times \Theta' \times [0, T] \rightarrow \mathbb{R}^{N_h}; \quad (\mu, \nu, t) \mapsto \mathbb{A}_P \cdot \hat{q}_P(\mu, \nu, t),$$

where we let $\mathbb{A}_P \in \mathbb{R}^{N_h \times N}$ be a POD computed by Algorithm (1) for the dimension N with the mass matrix $\mathbb{G}$, and

$$\hat{q}_P := R_\rho(\Phi_N^P(\theta_{\text{PDNN}})) \in \mathbb{R}^N$$

be the realization of some neural network Φ_N^P with input dimension $p + q + 1$ and output dimension N , and an already fixed weight $\theta_{\text{PDNN}} \in \Theta(\Phi_N^P)$. And let similarly the approximate DOD-based parameter-to-solution map be defined as follows

$$\mathcal{G}_{\text{DOD+DNN}} : \Theta \times \Theta' \times [0, T] \rightarrow \mathbb{R}^{N_h}; \quad (\mu, \nu, t) \mapsto \mathbb{V}_{\mu, t} \cdot \hat{q}_D(\mu, \nu, t),$$

where we let

$$\mathbb{V}_{\mu, t} := \mathbb{A} \cdot \tilde{\mathbb{V}}_{\mu, t}(\theta_{\text{DOD}}) \in \mathbb{R}^{N_h \times N'},$$

for an a priori fixed parameter $\theta_{\text{DOD}} \in \Theta(\Phi_{\tilde{\mathbb{V}}})$ according to Definition (3.6) and the used POD matrix $\mathbb{A} \in \mathbb{R}^{N_h \times N_A}$, and

$$\hat{q}_D := R_\rho(\Phi_{N'}^D(\theta_{\text{DDNN}})) \in \mathbb{R}^{N'}$$

be the realization of some neural network $\Phi_{N'}^D$ with input dimension $p + q + 1$ and output dimension N' , and an already fixed weight $\theta_{\text{DDNN}} \in \Theta(\Phi_{N'}^D)$.

Moreover we assume for this section to have after reordering and with $N_{\text{data}} = N_{s_1} N_{s_2} N_t$

$$\{(\mu_j, \nu_j, t_j), u_j\}_{j=1}^{N_{\text{data}}} := \left\{(\mu_i, \nu_j, k\Delta t), u_h^{\mu_i, \nu_j, k\Delta t}\right\}_{(i,j,k)=(1,1,1)}^{(N_{s_1}, N_{s_2}, N_t)} \subset (\mathcal{P}_1 \times \mathcal{P}_2 \times [0, T]) \times \mathbb{R}^{N_h}$$

a set of training data conforming to the label and feature notation in Definition (2.35), where the labels have been sampled in accordance to Assumption (2).

For both algorithms, the relative error can be given as

$$\mathcal{E}_R := \left(\int_{\Theta \times \Theta' \times [0, T]} \frac{\|u_h^{\mu, \nu, t} - \hat{u}_h^{\mu, \nu, t}\|^2}{\|u_h^{\mu, \nu, t}\|^2} d(\mu, \nu, t) \right)^{1/2}, \quad (4.1)$$

where $\hat{u}_h^{\mu, \nu, t} \in \mathbb{R}^{N_h}$ can be given either with regard to the approximate parameter-to-solution map of the DOD+DNN or the POD+DNN.

Theorem 4.3 (POD+DNN Decomposition). *Define the preliminaries such as in the introduction to this Section (4.1). Then under the Assumptions (2) and (3), we have for the relative error of the POD+DNN*

$$\mathcal{E}_R^{\text{POD+DNN}} \leq \mathcal{E}_S + \mathcal{E}_{\text{POD}} + \mathcal{E}_{\text{NN}},$$

where

1. *The sampling error*

$$\mathcal{E}_S = \mathcal{E}_S(\mathcal{G}, (\mu_i, \nu_i, t_i)_{i=1}^{N_{\text{data}}}, N)$$

satisfies $\mathcal{E}_S \rightarrow 0$ as $N_s, N_t \rightarrow \infty$, with expectation

$$\mathbb{E}[\mathcal{E}_S] = \mathcal{O}(N_s^{-1/4} + N_t^{-1}).$$

2. *The POD projection error*

$$\mathcal{E}_{\text{POD}} = \mathcal{E}_{\text{POD}}(\mathcal{G}, (\mu_i, \nu_i, t_i)_{i=1}^{N_{\text{data}}}, N)$$

satisfies $\mathcal{E}_{\text{POD}} \rightarrow \mathcal{E}_{\text{POD}, \infty}$ almost surely as $N_s, N_t \rightarrow \infty$, where

$$\mathcal{E}_{\text{POD}, \infty} = \mathcal{E}_{\text{POD}, \infty}(\mathcal{G}, N)$$

is independent of the data snapshots and proportional to the linear KnW of the manifold $\mathcal{S} = \{u_h^{\mu, \nu, t} \mid (\mu, \nu, t) \in \Theta' \times [0, T]\}$.

3. *The neural network approximation error*

$$\mathcal{E}_{\text{NN}} = \mathcal{E}_{\text{NN}}(\mathcal{G}, N, \theta_{\text{PDNN}}, \Phi_N^P)$$

is arbitrarily small depending on the size and choice of weights of the neural network Φ_N^P .

Proof. Let

$$\|\cdot\|_{L_\omega^2} := \left(\int_{\Theta \times \Theta' \times [0, T]} \frac{\|\cdot\|^2}{\|u_h^{\mu, \nu, t}\|^2} d(\mu, \nu, t) \right)^{1/2},$$

define a map $\|\cdot\|_{L_\omega^2} : L^2(\Theta \times \Theta' \times [0, T]; \mathbb{R}^{N_h}) \rightarrow \mathbb{R}^+$. Then this is a norm. We coarsely proof this, by remarking the inheritance of the triangle inequality from the $\|\cdot\|$ under the usage of the Minkowski inequality, i.e. for arbitrary $v, w \in L^2(\Theta \times \Theta' \times [0, T]; \mathbb{R}^{N_h})$

$$\begin{aligned} \|v + w\|_{L_\omega^2}^2 &= \int_{\Theta \times \Theta' \times [0, T]} \frac{\|v(\mu, \nu, t) + w(\mu, \nu, t)\|^2}{\|u_h^{\mu, \nu, t}\|^2} d(\mu, \nu, t) \\ &\leq \int_{\Theta \times \Theta' \times [0, T]} \frac{\|v(\mu, \nu, t)\|^2 + \|w(\mu, \nu, t)\|^2}{\|u_h^{\mu, \nu, t}\|^2} d(\mu, \nu, t) = \|v\|_{L_\omega^2}^2 + \|w\|_{L_\omega^2}^2, \end{aligned}$$

and the absolut homogeneity from simple qualities of the integral. Finally let $\|v\|_{L_\omega^2} = 0$ for some $v \in L^2(\Theta \times \Theta' \times [0, T])$, then assuming $v \neq 0$ yields with Assumption (3)

$$\|v\|_{L_\omega^2}^2 \geq m^{-1} \int_{\Theta \times \Theta' \times [0, T]} \|v(\mu, \nu, t)\| d(\mu, \nu, t) > 0,$$

hence reaching a contradiction. With the knowledge of $\|\cdot\|_{L_\omega^2}$ being a norm, we know it must be subject to the triangle inequality, thus

$$\mathcal{E}_R = \left(\int_{\Theta \times \Theta' \times [0, T]} \frac{\|u_h^{\mu, \nu, t} - \mathbb{A}_P \hat{q}_P(\mu, \nu, t)\|^2}{\|u_h^{\mu, \nu, t}\|^2} d(\mu, \nu, t) \right)^{1/2} \quad (4.2)$$

$$\leq \|u_h^{\mu, \nu, t} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t}\|_{L_\omega^2} + \|\mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t} - \mathbb{A}_P \hat{q}_P(\mu, \nu, t)\|_{L_\omega^2}. \quad (4.3)$$

Let $q_N(\mu, \nu, t) := \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t} \in \mathbb{R}^N$ be a reduced solution according to the POD basis. Then the natural definition of the neural network approximation error is

$$\mathcal{E}_{NN} = \left(\int_{\Theta \times \Theta' \times [0, T]} \frac{\|\mathbb{A}_P q_N(\mu, \nu, t) - \mathbb{A}_P \hat{q}_P(\mu, \nu, t)\|^2}{\|u_h^{\mu, \nu, t}\|^2} d(\mu, \nu, t) \right)^{1/2}. \quad (4.4)$$

This marks the second part in (4.3). This is only dependent on the ability of the neural network to identifying the best choice on a given linear sub-manifold of the solution manifold. The first part is concerned with the error generated from the best *possible* approximation of such a linear sub-manifold.

For this, we can bound the norm $\|\cdot\|_{L_\omega^2} \leq m^{-1} \|\cdot\|$, where $m > 0$ is the constant stemming from Assumption (3). Further we remind ourselves of the definition of the discrete correlation matrix $K = |\Theta \times \Theta \times \mathcal{T}| N_{data}^{-1} U \mathbb{G} U^T$ and its eigenvalues σ_k^2 in (2.13). Using $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$ for all $a, b \geq 0$, we get

$$\begin{aligned} &\|u_h^{\mu, \nu, t} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t}\|_{L_\omega^2} \\ &\leq m^{-1} \left(\int_{\Theta \times \Theta' \times [0, T]} \|u_h^{\mu, \nu, t} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 d(\mu, \nu, t) \right)^{1/2} \\ &\leq m^{-1} \left(\int_{\Theta \times \Theta' \times [0, T]} \|u_h^{\mu, \nu, t} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 d(\mu, \nu, t) - \sum_{k>N} \sigma_k^2 + \sum_{k>N} \sigma_k^2 \right)^{1/2} \\ &\leq m^{-1} \left(\left| \int_{\Theta \times \Theta' \times [0, T]} \|u_h^{\mu, \nu, t} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 d(\mu, \nu, t) - \sum_{k>N} \sigma_k^2 \right| + \sum_{k>N} \sigma_k^2 \right)^{1/2} \\ &\leq m^{-1} \underbrace{\left| \int_{\Theta \times \Theta' \times [0, T]} \|u_h^{\mu, \nu, t} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 d(\mu, \nu, t) - \sum_{k>N} \sigma_k^2 \right|^{1/2}}_{=: \mathcal{E}_S} + m^{-1} \underbrace{\sqrt{\sum_{k>N} \sigma_k^2}}_{=: \mathcal{E}_{POD}}. \end{aligned}$$

This concludes the estimate up to the facts about the convergence of the errors. Recalling Proposition (2.15), we can rewrite the sampling error as follows

$$\mathcal{E}_S = m^{-1} \left| \int_{\Theta \times \Theta' \times [0, T]} \|u_h^{\mu, \nu, t} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 d(\mu, \nu, t) - \frac{|\Theta \times \Theta' \times [0, T]|}{N_{data}} \sum_{j=1}^{N_{data}} \|u_j - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_j\|^2 \right|^{1/2}$$

Furthermore, we define, using the compactness postulated in Assumption (2) and the boundedness in Assumption (3), the L^2 -error map via

$$f(\mu, \nu, t) := \|u_h^{\mu, \nu, t} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 \leq M^2 \|I_{N_h} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G}\|_{\text{op}}^2 = M^2 \|I_{N_A} - \mathbb{A}_P\|_{\text{op}} < \infty.$$

Thus with $f \in L^2(\Theta \times \Theta' \times [0, T])$, we know that by Proposition (4.2), $\mathbb{E}[\mathcal{E}_S] \leq \mathcal{O}(N_s^{-1/4} + N_t^{-1})$.

By the strong law of large numbers, we can thus conclude, that $\mathcal{E}_S \rightarrow 0$ almost surely, as $N_t, N_s \rightarrow \infty$, and further by Proposition (2.14), we have that $\mathcal{E}_{POD} \rightarrow \mathcal{E}_{POD, \infty}$ as $N_t, N_s \rightarrow \infty$, where

$$\mathcal{E}_{POD, \infty} = m^{-1} \sqrt{\sum_{k>N} \sigma_{k, \infty}^2}. \quad (4.5)$$

□

Even optimistically, the upper error bound decomposition of Theorem (4.3) suggest for an ever-increasing number of data points $N_s, N_t \rightarrow \infty$, and an ever-increasing complexity for the Deep Neural Network underlying the realization of $\hat{q}_P$, an error of order of the Kolmogorov N -width of the solution manifold, since the DNN cannot overcome the unavoidable loss to even the perfect POD choice. Intuitively, the specific proportionality constant will be larger if the quotient between "small" regions of the solution space to the "large" regions of the solution space is big.

This observation can be reformulated into the following theorem.

Theorem 4.4. *Under the assumptions of Theorem (4.3), we have that*

$$\mathcal{E}_R^{POD+DNN} \geq \frac{m}{M} \mathcal{E}_{POD, \infty},$$

$$\text{where } \mathcal{E}_{POD, \infty} := m^{-1} \sqrt{\sum_{k>N} \sigma_{k, \infty}^2}.$$

Proof. We can simply start from the definition and argue, that under Proposition (2.15) the projection coefficients are chosen at best sub-optimal, i.e.

$$\mathcal{E}_R = \int_{\Theta \times \Theta' \times [0, T]} \frac{\|u_h^{\mu, \nu, t} - \mathbb{A}_P \hat{q}_N(\mu, \nu, t)\|^2}{\|u_h^{\mu, \nu, t}\|^2} d(\mu, \nu, t) \geq \int_{\Theta \times \Theta' \times [0, T]} \frac{\|u_h^{\mu, \nu, t} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t}\|^2}{\|u_h^{\mu, \nu, t}\|^2} d(\mu, \nu, t),$$

where $\mathbb{A}_P \in \mathbb{R}^{N_h \times N}$ is the POD matrix as given by Definition (2.13). Then starting from the other direction for a POD matrix $\mathbb{A}_\infty$ under the continuous correlation matrix, we get under the definition

of the proof of Theorem (4.3)

$$(\mathcal{E}_{\text{POD},\infty})^2 = m^{-2} \sum_{k>N} \sigma_{k,\infty}^2 \quad (4.6)$$

$$= m^{-2} \int_{\Theta \times \Theta' \times [0,T]} \|u_h^{\mu,\nu,t} - \mathbb{A}_\infty \mathbb{A}_\infty^T \mathbb{G} u_h^{\mu,\nu,t}\|^2 d(\mu, \nu, t) \quad (4.7)$$

$$\leq m^{-2} \int_{\Theta \times \Theta' \times [0,T]} \|u_h^{\mu,\nu,t} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu,\nu,t}\|^2 d(\mu, \nu, t) \quad (4.8)$$

$$= m^{-2} \int_{\Theta \times \Theta' \times [0,T]} \frac{\|u_h^{\mu,\nu,t} - \mathbb{A}_P \mathbb{A}_P^T \mathbb{G} u_h^{\mu,\nu,t}\|^2}{\|u_h^{\mu,\nu,t}\|^2} \|u_h^{\mu,\nu,t}\|^2 d(\mu, \nu, t) \quad (4.9)$$

$$\leq \frac{M^2}{m^2} (\mathcal{E}_R)^2, \quad (4.10)$$

where we use the Proposition (2.15) for the continuous POD matrix in Step (4.7), the optimality of the same proposition for Step (4.8) and the first equation of this proof in Step (4.10) with the rather obvious use of the bound on $\|u_h^{\mu,\nu,t}\|$ from Assumption (3). $\square$

We have now discussed some general relative error results for any POD+DNN ROM for some given neural network weights. As a next step, one can do the same for the DOD+DNN ROM. First, we need an auxiliary statement, that precedents the similarity of the definition of the DOD projection error to the POD projection error, since the former depends on the architecture of the inner DOD module and thus must be at least sub-optimal to some degree.

The key idea here being, that one can employ Horniks Theorem to ensure existence of a neural network architecture and corresponding neural network of such architecture, such that it finds $\mathbb{V}^*$ minimizing the objective functional

$$(\mu, t, \mathbb{V}) \mapsto \int_{\Theta'} \|u_h^{\mu,\nu,t} - \mathbb{V}_{\mu,t} \mathbb{V}_{\mu,t}^T \mathbb{G} u_h^{\mu,\nu,t}\|^2 d(\nu),$$

for any $(\mu, t) \in \Theta \times [0, T]$. Under triangle equality for the distance to the optimal minimizer of the objective functional, this can be separated into the mistake of fitting the network to the minimizer, which is arbitrarily small, and the mistake of the minimizer, that is nothing but the Kolmogorov N' -width under fixed (μ, t) .

Remark 4.5. Generally we can *not* employ Yarotsky Theorem (2.33) here, since even though the objective functional, as described above, can be shown to be Lipschitz continuous under our current assumptions, as given above, its minimizer with regard to the matrix entry does not necessarily inherit any regularity and Yarotsky does require Sobolev-type smoothness. However, we will set up the proof for the following lemma in such a way, that Yarotsky can be directly interchanged, if a higher regularity for the optimal objective functional is imposed.

Before we start, we would like to define the aforementioned objective functional and its minimizer, since further assumptions on the PDE can expose the needed further regularity.

The source for the following proposition and its framework is generally (Franco et al., 2024).

Definition 4.6 (Optimal DOD). Let $N', N_A, N_h \in \mathbb{N}$ be fixed by Assumption (5). Assuming we have $\mu \in \Theta, t \in [0, T]$ and $\mathbb{V} = \mathbb{A} \tilde{\mathbb{V}} \in [-1, 1]^{N_h \times N'}$, with $\mathbb{A} \in \mathbb{R}^{N_h \times N_A}$ fixed being $\mathbb{G}$ -orthonormal and hence $\|\mathbb{A}\|_{\text{op}} = 1$ by Lemma (2.6), and $\tilde{\mathbb{V}} \in [-1, 1]^{N_A \times N'}$, then we define $J : \Theta \times [0, T] \times [-1, 1]^{N_h \times N'} \rightarrow \mathbb{R}$ the *objective DOD functional* via

$$J(\mu, t, \mathbb{V}) := \int_{\mathcal{X}} \|u_h^{\mu,\nu,t} - \mathbb{V} \mathbb{V}^T \mathbb{G} u_h^{\mu,\nu,t}\|^2 d(\nu),$$

for any compact set $\mathcal{X} \subset \Theta'$ with $\dim(\mathcal{X}) = q$. Then we call a Borel measurable map $s : \Theta \times [0, T] \rightarrow [-1, 1]^{N_A \times N'}$ producing w.l.o.g. orthonormal matrices given by

$$s : (\mu, t) \mapsto \underset{\mathbb{V} \in [-1, 1]^{N_A \times N'}}{\operatorname{argmin}} J(\mu, t, \mathbb{A}\mathbb{V}),$$

the *optimal DOD* for $\mathcal{X}$, if it is uniquely defined.

Under the already given assumptions, we can pose its first existence result.

Lemma 4.7. Let $J : \Theta \times [0, T] \times [-1, 1]^{N_h \times N'} \rightarrow \mathbb{R}$ be an objective DOD functional for some $\mathcal{X} \subset \Theta'$ with $\dim(\mathcal{X}) = q$ as in Definition (4.6), then under Assumption (3), we have that J is Lipschitz, and the optimal DOD s exists.

Proof. Let $\mu, \mu' \in \Theta$, $t, t' \in [0, T]$ and $\mathbb{V}, \mathbb{V}' \in \mathbb{R}^{N_h \times N'}$ arbitrary. For the following calculation, we compose the preliminary identity for $u := u_h^{\mu, \nu, t}$ and $u' := u_h^{\mu', \nu, t'}$ for each $\mu \in \Theta, \nu \in \mathcal{X}$ and $t \in [0, T]$, and the projectors $P := \mathbb{V}\mathbb{V}^T\mathbb{G}$, $P' := \mathbb{V}'(\mathbb{V}')^T\mathbb{G}$,

$$\begin{aligned} (\mathbb{I} - P)u - (\mathbb{I} - P')u' &= u - Pu - u' + P'u' \\ &= u - Pu - u' + P'u' + Pu' - Pu' \\ &= u - u' - Pu + Pu' + P'u' - Pu' \\ &= (\mathbb{I} - P)(u - u') + (P' - P)u'. \end{aligned} \tag{4.11}$$

Then we can calculate with, the reverse triangle inequality, and our auxiliary Equation (4.11),

$$\begin{aligned} |J(\mu, t, \mathbb{V}) - J(\mu', t', \mathbb{V}')| &= \int_{\mathcal{X}} \left| \|u_h^{\mu, \nu, t} - \mathbb{V}\mathbb{V}^T\mathbb{G}u_h^{\mu, \nu, t}\|^2 - \|u_h^{\mu', \nu, t'} - \mathbb{V}'(\mathbb{V}')^T\mathbb{G}u_h^{\mu', \nu, t'}\|^2 \right| d(\nu) \\ &= \int_{\mathcal{X}} \left| \|u_h^{\mu, \nu, t} - \mathbb{V}\mathbb{V}^T\mathbb{G}u_h^{\mu, \nu, t}\| - \|u_h^{\mu', \nu, t'} - \mathbb{V}'(\mathbb{V}')^T\mathbb{G}u_h^{\mu', \nu, t'}\| \right| d(\nu) \\ &= \int_{\mathcal{X}} \left\| \left(u_h^{\mu, \nu, t} - \mathbb{V}\mathbb{V}^T\mathbb{G}u_h^{\mu, \nu, t} \right) - \left(u_h^{\mu', \nu, t'} - \mathbb{V}'(\mathbb{V}')^T\mathbb{G}u_h^{\mu', \nu, t'} \right) \right\|^2 d(\nu) \\ &\leq \int_{\mathcal{X}} \|(\mathbb{I} - \mathbb{V}\mathbb{V}^T\mathbb{G})(u_h^{\mu, \nu, t} - u_h^{\mu', \nu, t'})\|^2 d(\nu) + M^2 |\mathcal{X}| \|\mathbb{V}\mathbb{V}^T\mathbb{G} - \mathbb{V}'(\mathbb{V}')^T\mathbb{G}\|_{\text{op}} \\ &\leq |\mathcal{X}| (L^2 + 2M^2) (|\mu - \mu'| + |t - t'| + |\mathbb{V} - \mathbb{V}'|), \end{aligned}$$

where our last step is justified with the projector quality of $\mathbb{V}\mathbb{V}^T\mathbb{G}$ from Lemma (2.5), thus explicitly having eigenvalues 0 and 1, and therefore making J Lipschitz continous.

include existence of s

□

Proposition 4.8 (Optimal Deep Orthogonal Decomposition). For each geometric parameter $\mu \in \Theta$, let $\mathcal{S}_{\mu, t}^{\text{pre-red}} := \{\mathbb{A}^T \mathbb{G} u_h^{\mu, \nu, t}\}_{\nu \in \Theta'}$ be the (μ, t) -invariant pre-reduced solution submanifold, where $\mathbb{A} \in \mathbb{R}^{N_h \times N_A}$ is a $\mathbb{G}$ -orthonormal matrix. Then, under the Assumptions (2) and (3), for every $\varepsilon > 0$ there are weights $\theta_{\text{DOD}}^* \in \Theta(\Phi_{\mathbb{V}})$ for a DOD of output dimension $N_h \times N'$ according to Definition (3.6), denoting $\mathbb{V}_{\mu, t} := \mathbb{V}_{\mu, t}(\theta_{\text{DOD}}^*)$ for each $(\mu, t) \in \Theta \times [0, T]$ and a constant $c'_{\mathbb{A}} > 0$ only depending on $\mathbb{A}$ and the parameter space, such that

$$\int_{\Theta \times \Theta' \times [0, T]} \|u_h^{\mu, \nu, t} - \mathbb{V}_{\mu, t}^2 \mathbb{V}_{\mu, t}^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 d(\mu, \nu, t) \leq \varepsilon + c'_{\mathbb{A}} + \int_{\Theta \times [0, T]} d_{N'}(\mathcal{S}_{\mu, t}^{\text{pre-red}})^2 d(\mu, t).$$

Proof. Let $N', N_A, N_h \in \mathbb{N}$ be fixed by Assumption (5). We set J , the objective DOD functional for Θ' , and s , the optimal DOD for Θ' , according to Definiton (4.6), whereas the map

$$s : (\mu, t) \mapsto \underset{\tilde{\mathbb{V}} \in [-1, 1]^{N_A \times N'}}{\operatorname{argmin}} J(\mu, t, \mathbb{A}\tilde{\mathbb{V}}).$$

is bounded (given by continuity of J and compactness of its image) and therefore $s \in L^2(\Theta \times [0, T]; \mathbb{R}^{N_A \times N'})$, i.e.

$$\|s\|_{L^2(\Theta \times [0, T]; \mathbb{R}^{N_A \times N'})}^2 := \int_{\Theta \times [0, T]} |s(\mu, t)|^2 d(\mu, t) < \infty.$$

Define $[s]_{ij}$ as the canoncially lifted entry in position $(i, j) \in \{1, \dots, N_A\} \times \{1, \dots, N'\}$.

This means that by Hornik Theorem (2.23), there is a neural network architecture $\mathcal{A}^{[\tilde{\mathbb{V}}_0]_{ij}}$ with $(p+1)$ -dimensional input and one-dimensional ouput, and a neural network of that architecture $[\tilde{\mathbb{V}}_0]_{ij}$, satisfying

$$\|[s]_{ij} - R_\rho([\tilde{\mathbb{V}}_0]_{ij})\|_{L^2(\Theta \times [0, T]; \mathbb{R})}^2 \leq \varepsilon,$$

for each $(i, j) \in \{1, \dots, N_A\} \times \{1, \dots, N'\}$, and then using network parallelization as in Definition (2.27), we can set the optimal DOD neural network as

$$\tilde{\mathbb{V}}_0 := P \left(P([\tilde{\mathbb{V}}_0]_{11}, \dots, [\tilde{\mathbb{V}}_0]_{1N'}), \dots, P([\tilde{\mathbb{V}}_0]_{N_A 1}, \dots, [\tilde{\mathbb{V}}_0]_{N_A N'}) \right),$$

since it attains the wanted accuracy

$$\|s - R_\rho(\tilde{\mathbb{V}}_0)\|_{L^2(\Theta \times [0, T]; \mathbb{R}^{N_A \times N'})}^2 \leq \varepsilon.$$

To now ensure, that the provided realization of the almost DOD neural network are indeed in the pre-image of J , we need to ensure, that there is a network, which indeed attains the accuracy, but is also entrywise $[-1, 1]$. For this, let $\zeta = ((A_1, b_1), (A_2, b_2), (A_3, b_3))$ be given via

$$\begin{aligned} A_1 &:= [\mathbb{I}_{\mathbb{R}^{N_A \times N'}}], & b_1 &:= (\mathbb{I}_{\mathbb{R}^{N_A \times N'}}), & A_2 &:= [-\mathbb{I}_{\mathbb{R}^{N_A \times N'}}], & b_2 &:= (\mathbb{I}_{\mathbb{R}^{N_A \times N'}}), \\ A_3 &:= [-\mathbb{I}_{\mathbb{R}^{N_A \times N'}}], & b_3 &:= (\mathbb{I}_{\mathbb{R}^{N_A \times N'}}) \end{aligned}$$

The realization of this network is equivalent to the entry-wise operator of $x \mapsto 1 - \rho(2 - \rho(\rho(x+1))) \in [-1, 1]$ for $x \in \mathbb{R}$ on all entries of a $(N_A \times N')$ -dimensional matrix. Note, that to make this neural network truly adhere to Definition (2.18), which needs the dimensions to be natural numbers, we would have to reinterpret any matrix as a canoncially stacked vector, but we omit this for the sake of readability, and since it does not affect the number of layers or weights.

Further because the Lipschitz constant of this map is 1, we can infer with the definition $\mathbb{V}_{\mu, t} := \mathbb{A} \cdot (R_\rho(\zeta(\theta_{\text{DOD}}^*)) \circ R_\rho(\tilde{\mathbb{V}}_0(\theta_{\text{DOD}}^*))) (\mu, t)$ a new network, called the inner DOD module, with a final architecture $\Phi_{\tilde{\mathbb{V}}}$ and complexity only differing by constants, and a choice of neural network weights $\theta_{\text{DOD}}^* \in \Theta(\Phi_{\tilde{\mathbb{V}}})$, such that

$$\int_{\Theta \times [0, T]} \|\mathbb{V}_{\mu, t} - \mathbb{A}s(\mu, t)\|^2 d(\mu, t) \leq \int_{\Theta \times [0, T]} |R_\rho(\tilde{\mathbb{V}}_0)(\mu, t) - s(\mu, t)|^2 d(\mu, t) \leq \varepsilon,$$

by the 1-Lipschitz proptert of $R_\rho(\zeta(\theta_{\text{DOD}}^*))$ and the same argument as was used in the proof of Proposition (2.12). Finally, with this assertion, the fact that this, as a matrix, is indeed

in the pre-image of J and the Proposition (2.12) for the switch to the euklidian norm with $c'_\mathbb{A} := |\Theta \times \Theta' \times [0, T]| c_\mathbb{A}$, we can write with $u_{\text{pre-red}}^{\mu, \nu, t} := \mathbb{A}^T \mathbb{G} u_h^{\mu, \nu, t}$,

$$\begin{aligned} & \int_{\Theta \times \Theta' \times [0, T]} \|u_h^{\mu, \nu, t} - \mathbb{V}_{\mu, t} \mathbb{V}_{\mu, t}^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 d(\mu, \nu, t) \\ & \leq \int_{\Theta \times [0, T]} |J(\mu, t, \mathbb{V}_{\mu, t}) - J(\mu, t, \mathbb{A} s(\mu, t))| d(\mu, t) + \int_{\Theta \times [0, T]} J(\mu, t, \mathbb{A} s(\mu, t)) d(\mu, t) \\ & \leq \varepsilon + \int_{\Theta \times [0, T]} \left(\min_{\mathbb{V} \in \mathbb{R}^{N_A \times N'}} J(\mu, t, \mathbb{A} \mathbb{V}) \right) d(\mu, t) \\ & \leq \varepsilon + c'_\mathbb{A} + \int_{\Theta \times [0, T]} \left(\min_{\tilde{\mathbb{V}} \in \mathbb{R}^{N_A \times N'}} \sup_{\nu \in \Theta'} |u_{\text{pre-red}}^{\mu, \nu, t} - \tilde{\mathbb{V}} \tilde{\mathbb{V}}^T u_{\text{pre-red}}^{\mu, \nu, t}|^2 \right) d(\mu, t). \end{aligned} \quad (4.12)$$

□

Remark 4.9. We have structured the Proposition (4.8) with the L^1 -integral in mind, but we may redefine it to L^∞ and set the objective DOD functional for $\mathcal{P}_2$ for the statement and proof as

$$J(\mu, t, \mathbb{V}) := \frac{|\Theta'|}{N_{s_2}} \sum_{\nu \in \mathcal{P}_2} \|u_h^{\mu, \nu, t} - \mathbb{V} \mathbb{V}^T \mathbb{G} u_h^{\mu, \nu, t}\|^2.$$

Then the entire proof can be done in the same way, since the Lipschitz property of J in Lemma (4.7) remains true, and culminates in the following version of Proposition (4.8) for $\mathbb{V}_{\mu, t} := \mathbb{V}_{\mu, t}(\hat{\theta}_{\text{DOD}}^*)$, when discarding the very last step in Equation (4.12)

$$\begin{aligned} & \sup_{(\mu, t) \in \Theta \times [0, T]} \frac{|\Theta'|}{N_{s_2}} \sum_{\nu \in \mathcal{P}_2} \|u_h^{\mu, \nu, t} - \mathbb{V}_{\mu, t} \mathbb{V}_{\mu, t}^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 \\ & \leq \varepsilon + c'_\mathbb{A} + \sup_{(\mu, t) \in \Theta \times [0, T]} \frac{|\Theta'|}{N_{s_2}} \min_{\tilde{\mathbb{V}} \in \mathbb{R}^{N_A \times N'}} \sum_{\nu \in \mathcal{P}_2} |u_{\text{pre-red}}^{\mu, \nu, t} - \tilde{\mathbb{V}} \tilde{\mathbb{V}}^T u_{\text{pre-red}}^{\mu, \nu, t}|^2. \end{aligned}$$

Notice, that in general $\hat{\theta}_{\text{DOD}}^* \neq \theta_{\text{DOD}}^*$, since the change of the objective DOD functional induces a change to the optimal DOD and hence a potential change to the choice of neural network weights.

As another note, it is clear, that if $N_A = N_h$, we have $c'_\mathbb{A} = c_\mathbb{A} = 0$ by the definition of the POD as a canonical $\mathbb{G}$ -orthonormalization and the right-hand-side sum would be over the non-reduced quantity. This would simply constitute a case, where we completely circumvent a pre-reduction.

With that in mind, one can now build an analogon to the relative error on the POD+DNN ROM (i.e. Theorem (4.3)), since we now have a similar statement as the one of Proposition (2.15), in the form of Remark (4.9). In fact, we can even leverage Proposition (2.15) to define

$$\sigma(\mu, t)_1^2 \geq \dots \geq \sigma(\mu, t)_n^2 \quad \text{for any } n \leq r, \quad (4.13)$$

as the eigenvalues of a virtual POD (see Definition (2.13)) with the snapshot matrix

$$U := \left[u_{\text{pre-red}}^{\mu, \nu_1, t} \mid \dots \mid u_{\text{pre-red}}^{\mu, \nu_{N_{s_2}}, t} \right]^T \in \mathbb{R}^{N_A \times N_{s_2}},$$

for each $(\mu, t) \in \Theta \times [0, T]$, and a subsequently defined discrete correlation matrix

$$K(\mu, t) := \frac{|\Theta'|}{N_{s_2}} U U^T,$$

that fulfill

$$\frac{|\Theta'|}{N_{s_2}} \min_{\tilde{\mathbb{V}} \in \mathbb{R}^{N_A \times n}} \sum_{\nu \in \mathcal{P}_2} |u_{\text{pre-red}}^{\mu, \nu, t} - \tilde{\mathbb{V}} \tilde{\mathbb{V}}^T u_{\text{pre-red}}^{\mu, \nu, t}|^2 = \sum_{k > n} \sigma(\mu, t)_k^2,$$

for the $n \leq r$ from above. In the same manner, we can define a virtual infinite limit POD with the respective infinite correlation matrix and their eigenvalues

$$\sigma(\mu, t)_{1, \infty}^2 \geq \cdots \geq \sigma(\mu, t)_{n, \infty}^2 \quad \text{for any } n \leq r, \quad (4.14)$$

that fulfill

$$\min_{\tilde{\mathbb{V}} \in \mathbb{R}^{N_A \times n}} \int_{\nu \in \Theta'} |u_{\text{pre-red}}^{\mu, \nu, t} - \tilde{\mathbb{V}} \tilde{\mathbb{V}}^T u_{\text{pre-red}}^{\mu, \nu, t}|^2 d(\nu) = \sum_{k > n} \sigma(\mu, t)_{k, \infty}^2.$$

Theorem 4.10. *Define the preliminaries such as in the introduction to this Section (4.1). Further set $\mathbb{V}_{\mu, t} := \mathbb{V}_{\mu, t}(\theta_{DOD}^*)$ for the choice of weights $\theta_{DOD}^* \in \Theta(\Phi_{\tilde{\mathbb{V}}})$ as in Proposition (4.8). Then, under the Assumptions (2) and (3), we have for the relative error of the DOD+DNN*

$$\mathcal{E}_R^{DOD+DNN} \leq \mathcal{E}_S + \mathcal{E}_{\mathbb{A}} + \mathcal{E}_{DOD} + \mathcal{E}_{NN}, \quad (4.15)$$

where

1. The sampling error

$$\mathcal{E}_S = \mathcal{E}_S(\mathcal{G}, (\mu_i, \nu_i, t_i)_{i=1}^{N_{data}}, N')$$

satisfies $\mathcal{E}_S \rightarrow 0$ as $N_{s_2} \rightarrow \infty$, with expectation

$$\mathbb{E}[\mathcal{E}_S] \leq \mathcal{O}(N_{s_2}^{-1/4}).$$

2. The ambient error

$$\mathcal{E}_{\mathbb{A}} = \mathcal{E}_{\mathbb{A}}(\mathcal{G}, (\mu_i, \nu_i, t_i)_{i=1}^{N_{data}}, N_A)$$

being equal to zero, if $N_A = N_h$.

3. The DOD projection error

$$\mathcal{E}_{DOD} = \mathcal{E}_{DOD}(\mathcal{G}, (\mu_i, \nu_i, t_i)_{i=1}^{N_{data}}, N')$$

satisfies $\mathcal{E}_{DOD} \rightarrow \mathcal{E}_{DOD, \infty} + \varepsilon$ almost surely as $N_{s_2} \rightarrow \infty$ for any $\varepsilon = \varepsilon(\Phi_{\tilde{\mathbb{V}}}) > 0$ arbitrarily small depending on the size and choice of weights of the inner DOD module, where

$$\mathcal{E}_{DOD, \infty} = \mathcal{E}_{DOD, \infty}(\mathcal{G}, N')$$

is independent of the data snapshots and of order of the maximal linear KnW of the solution submanifold $\mathcal{S}_{\mu, t} = \{u_h^{\mu, \nu, t} \mid \nu \in \Theta'\}$ for any $(\mu, t) \in \Theta \times [0, T]$.

4. The neural network approximation error

$$\mathcal{E}_{NN} = \mathcal{E}_{NN}(\mathcal{G}, N', \Phi_{N'}^D)$$

is arbitrarily small depending on the size and choice of weights of the neural network $\Phi_{N'}^D$.

Proof. In its core, this proof inherits its structure and arguments from the one of Theorem (4.3), we will therefore only shorten the parts in which they are interchangeable. Let $\|\cdot\|_{L_\omega^2} : L^2(\Theta \times \Theta' \times [0, T]; \mathbb{R}^{N_h}) \rightarrow \mathbb{R}^+$ be the norm as given before, and define

$$q_{N'}(\mu, \nu, t) := \mathbb{V}_{\mu, t}^T \mathbb{G} u_h^{\mu, \nu, t} \in \mathbb{R}^{N'},$$

as the reduced solution regarding the DOD projection. Then we can apply the triangle inequality like before, to achieve

$$\mathcal{E}_R^{\text{DOD+DNN}} \leq \|u_h^{\mu, \nu, t} - \mathbb{V}_{\mu, t} \mathbb{V}_{\mu, t}^T \mathbb{G} u_h^{\mu, \nu, t}\|_{L_\omega^2} + \|\mathbb{V}_{\mu, t} q_{N'}(\mu, \nu, t) - \mathbb{V}_{\mu, t} \hat{q}_D(\mu, \nu, t)\|_{L_\omega^2}.$$

In turn, from here we can already infer the error stemming from the latent dynamics identification under the neural network realization $\hat{q}_D$, i.e.

$$\mathcal{E}_{NN} = \left(\int_{\Theta \times \Theta' \times [0, T]} \frac{\|\mathbb{V}_{\mu, t} q_{N'}(\mu, \nu, t) - \mathbb{V}_{\mu, t} \hat{q}_D(\mu, \nu, t)\|^2}{\|u_h^{\mu, \nu, t}\|^2} d(\mu, \nu, t) \right)^{1/2}. \quad (4.16)$$

Moreover, we can extrapolate a similar procedure of splitting the remaining term into the respective errors as in the proof of Theorem (4.3), where we already assume for the fixed $N' \in \mathbb{N}$, the quasi-optimal choice of weights $\theta_{\text{DOD}}^* \in \Theta(\Phi_{\tilde{\mathbb{V}}})$ for the underlying DOD neural network in accordance to Remark (4.9) for some fixed $\delta > 0$, fixed $c'_\mathbb{A}$ and its subsequently defined virtual discrete correlation eigenvalues $\sigma(\mu, t)_k$ in Equation (4.13)

$$\begin{aligned} & \|u_h^{\mu, \nu, t} - \mathbb{V}_{\mu, t} \mathbb{V}_{\mu, t}^T \mathbb{G} u_h^{\mu, \nu, t}\|_{L_\omega^2} \\ & \leq m^{-1} \left(\int_{\Theta \times [0, T]} \int_{\Theta'} \|u_h^{\mu, \nu, t} - \mathbb{V}_{\mu, t} \mathbb{V}_{\mu, t}^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 d(\nu) d(\mu, t) \right)^{1/2} \\ & = m^{-1} \left(\int_{\Theta \times [0, T]} \int_{\Theta'} \|u_h^{\mu, \nu, t} - \mathbb{V}_{\mu, t} \mathbb{V}_{\mu, t}^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 d(\nu) - \sum_{k > N'} \sigma(\mu, t)_k^2 + \sum_{k > N'} \sigma(\mu, t)_k^2 d(\mu, t) \right)^{1/2} \\ & \leq m^{-1} \left| \underbrace{\int_{\Theta \times [0, T]} \int_{\Theta'} \|u_h^{\mu, \nu, t} - \mathbb{V}_{\mu, t} \mathbb{V}_{\mu, t}^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 d(\nu) - \frac{|\Theta'|}{N_{s_2}} \sum_{\nu \in \mathcal{P}_2} \|u_h^{\mu, \nu, t} - \mathbb{V}_{\mu, t} \mathbb{V}_{\mu, t}^T \mathbb{G} u_h^{\mu, \nu, t}\|^2 d(\mu, t)}_{=:\mathcal{E}_S} \right|^{1/2} \\ & + \underbrace{m^{-1} |\Theta \times [0, T]|^{1/2} c'_\mathbb{A}}_{=:\mathcal{E}_\mathbb{A}} + \underbrace{m^{-1} |\Theta \times [0, T]|^{1/2} \delta + m^{-1} |\Theta \times [0, T]|^{1/2} \sup_{(\mu, t) \in \Theta \times [0, T]} \sqrt{\sum_{k > N'} \sigma(\mu, t)_k^2}}_{=:\mathcal{E}_{\text{DOD}}}. \end{aligned}$$

Deploying the same strategy as in the proof of Theorem (4.3), we can assert, that indeed $\mathcal{E}_S \rightarrow 0$ almost surely by the strong law of large numbers, and that $\mathbb{E}[\mathcal{E}_S] \leq \mathcal{O}(N_{s_2}^{-1/4})$ by Proposition (4.2) with $N_t = 1, N_{s_1} = 1$.

Moreover, we have from Proposition (2.14), with $N_{s_1}, N_t = 1$, that

$$\begin{aligned} \mathcal{E}_{\text{DOD}} &= \frac{|\Theta \times [0, T]|^{1/2}}{m} \left(\delta + \sup_{(\mu, t) \in \Theta \times [0, T]} \sqrt{\sum_{k > N'} \sigma(\mu, t)_k^2} \right) \\ &\xrightarrow{N_{s_2} \rightarrow \infty} \frac{|\Theta \times [0, T]|^{1/2}}{m} \sup_{(\mu, t) \in \Theta \times [0, T]} \sqrt{\sum_{k > N'} \sigma(\mu, t)_k^2} + \frac{\delta |\Theta \times [0, T]|^{1/2}}{m} =: \mathcal{E}_{\text{DOD}, \infty} + \varepsilon, \end{aligned}$$

which can be arbitrarily improved for a more complex choice of the inner DOD module architecture. This dependency on the complexity is suggested by Yarotsky Theorem (2.33), since the L^∞ error correlates with the $\mathcal{H}_h$ representation error by the compactness of the time-parameter space, that is given by Assumption (2), but is only proveable under additional assumptions as argued in Remark (4.5).

Finally, we know that by Remark (4.9) $c'_\mathbb{A} = 0$ implying $\mathcal{E}_\mathbb{A} = 0$, if $N_A = N_h$, which thus concludes the proof. $\square$

To connect this line of thought to the prior discussion for the POD+DNN, we can now pose a similar lower bound on the error for the DOD+DNN ROM technique. For practical reasoning, the reader should keep in mind, that the result is closely connected to the complexity of the neural network of the inner DOD module, even if the following theorem omits such an appreciation, since the connection given through Yarotsky Theorem (2.33) is not immediate under the preliminaries alone.

Theorem 4.11. *Under the assumptions of Theorem (4.10), we have that*

$$\mathcal{E}_R^{DOD+DNN} \geq \frac{m}{M} \mathcal{E}'_{DOD,\infty},$$

where we set

$$\mathcal{E}'_{DOD,\infty} := m^{-1} \int_{(\mu,t) \in \Theta \times [0,T]} \sqrt{\sum_{k>N'} \sigma(\mu,t)_{k,\infty}^2} d(\mu,t),$$

and $\sigma(\mu,t)_{k,\infty}$ the eigenvalues for a non-reduced continuous correlation matrix.

Proof. The proof will again follow a similar structure as the one of Theorem (4.4). In fact, we can again infer the sub-optimal choice of the projection coefficients under any neural network realization

$$\mathcal{E}_R = \int_{\Theta \times \Theta' \times [0,T]} \frac{\|u_h^{\mu,\nu,t} - \mathbb{V}_{\mu,t} \hat{q}_{N'}(\mu,\nu,t)\|^2}{\|u_h^{\mu,\nu,t}\|^2} d(\mu,\nu,t) \geq \int_{\Theta \times \Theta' \times [0,T]} \frac{\|u_h^{\mu,\nu,t} - \mathbb{V}_{\mu,t} \mathbb{V}_{\mu,t}^T \mathbb{G} u_h^{\mu,\nu,t}\|^2}{\|u_h^{\mu,\nu,t}\|^2} d(\mu,\nu,t),$$

where $\mathbb{V}_{\mu,t} \in \mathbb{R}^{N_h \times N'}$ is the DOD matrix realization for any $(\mu,t) \in \Theta \times [0,T]$ as given by Definition (3.6). Then again starting from the other direction for a virtual POD matrix $\mathbb{A}_{\mu,t}^\infty \in \mathbb{R}^{N_h \times N'}$ for each $(\mu,t) \in \Theta \times [0,T]$ under the continuous correlation matrix, obtained as a limit of

$$U := \left[u_h^{\mu,\nu_1,t} | \dots | u_h^{\mu,\nu_{N_{s_2}},t} \right]^T \in \mathbb{R}^{N_h \times N_{s_2}}$$

for $N_{s_2} \rightarrow \infty$, which exists and is well defined via the Proposition (2.15) and has eigenvalues

$\sigma(\mu, t)_{1,\infty}^2 \geq \dots \geq \sigma(\mu, t)_{N',\infty}^2$, we get using the definition of the proof of Theorem (4.10)

$$(\mathcal{E}'_{\text{DOD},\infty})^2 = m^{-2} \int_{(\mu,t) \in \Theta \times [0,T]} \sum_{k > N'} \sigma(\mu, t)_{k,\infty}^2 d(\mu, t) \quad (4.17)$$

$$= m^{-2} \int_{(\mu,t) \in \Theta \times [0,T]} \int_{\Theta'} \|u_h^{\mu,\nu,t} - \mathbb{A}_{\mu,t}^\infty (\mathbb{A}_{\mu,t}^\infty)^T \mathbb{G} u_h^{\mu,\nu,t}\|^2 d(\nu) d(\mu, t) \quad (4.18)$$

$$\leq m^{-2} \int_{(\mu,t) \in \Theta \times [0,T]} \int_{\Theta'} \|u_h^{\mu,\nu,t} - \mathbb{V}_{\mu,t} \mathbb{V}_{\mu,t}^T \mathbb{G} u_h^{\mu,\nu,t}\|^2 d(\nu) d(\mu, t) \quad (4.19)$$

$$= m^{-2} \int_{\Theta \times \Theta' \times [0,T]} \frac{\|u_h^{\mu,\nu,t} - \mathbb{V}_{\mu,t} \mathbb{V}_{\mu,t}^T \mathbb{G} u_h^{\mu,\nu,t}\|^2}{\|u_h^{\mu,\nu,t}\|^2} \|u_h^{\mu,\nu,t}\|^2 d(\mu, \nu, t) \quad (4.20)$$

$$\leq \frac{M^2}{m^2} (\mathcal{E}_R)^2, \quad (4.21)$$

where we use the Proposition (2.15) for the continuous POD matrix in Step (4.18), the optimality of the same proposition for Step (4.19) and the first equation of this proof in Step (4.21) with the rather obvious use of the bound on $\|u_h^{\mu,\nu,t}\|$ from Assumption (3). Notice, that this statement works independent of the pre-reduction problem, since $\mathbb{A}_{\mu,t}^\infty$ is just the optimal of any $(N_h \times N')$ -dimensional matrix-based projector, and subsequently does not suffer from the pre-reduction. $\square$

Note, that this result barely differs from Theorem (4.4) for the POD+DNN ROM lower bound, and that we use a different version for the DOD error, than we have for the proof of Theorem (4.10), because the average over the Kolmogorov N' -widths in $\Theta \times [0, T]$ has more comparability to the Kolmogorov N -width decay of the whole solution manifold as a lower bound.

Of course, the optimality step in the proof suggests the use of a "good" DOD matrix and is hence closely related to the complexity of the inner DOD module, but this explains the lack of any mention of a choice for the neural network weights; this simply works for any matrix.

4.2 Upper Complexity Bounds

In this section, we will specify under the Assumption (2), (3), (4) and (5) the upper complexity bounds for our three main algorithms: the POD-DL-ROM, the DOD+DFNN and the DOD-DL-ROM, as introduced in Section (3.2).

However, to achieve actual complexity bounds for the latter two ROMs, we will necessarily need to impose one further assumption for the regularity of the optimal objective functional as discussed in Remark (4.5). This will be inserted in an upcoming part, before the discussion of the DOD-DL-ROM upper complexity bound as it is optional for the assertion of the complexity bound of the main Theorem (4.13). Indeed, if we omit any new assumptions, we may achieve the same Probably-Almost-Correct bound for the relative error of the DOD-DL-ROM, however without any information about the maximal complexity of the inner DOD module.

For the following discussion, we want to again set some preliminaries to shorten the theorems layout. Firstly, we assume the setup of the previous Section (4.1), where we recall defining a sample to be used for all subsequent algorithms both POD-based and DOD-based neural networks, respectively marked POD+DNN and DOD+DNN. In fact, set up N and N' as follows

$$N = \arg \min_{n \in \mathbb{N}} \left(\sum_{k > n} \sigma_k^2 \leq \frac{m^2}{9} \varepsilon^2 \right), \quad N' = \arg \min_{n \in \mathbb{N}} \left(\sup_{(\mu,t) \in \Theta \times [0,T]} \sum_{k > n} \sigma(\mu, t)_k^2 \leq \frac{m^2}{16|\Theta \times [0, T]|} \varepsilon^2 \right) \quad (4.1)$$

with $\varepsilon > 0$ yet unspecified, and σ_k^2 according to Definition (2.13) and $\sigma(\mu, t)_k^2$ to Equation (4.13). Note, that under Assumption (1) the hierarchy of Assumption (5) is still preserved, if the choice of ε is small enough and N_{data} is large enough.

From this starting point, refine the definition of the approximate parameter-to-solution map for the POD+DNN into

$$\mathcal{G}_{\text{POD-DL-ROM}} : \Theta \times \Theta' \times [0, T] \rightarrow \mathbb{R}^{N_h}; \quad (\mu, \nu, t) \mapsto \mathbb{A}_P \cdot (\psi_N \circ \phi_n)(\mu, \nu, t),$$

the POD-DL-ROM under weights $\theta_{\text{POD-DL-ROM}} = (\theta_D, \theta_{\text{DF}})$ with

$$\phi_n := R_\rho(\Phi_n(\theta_{\text{DF}})) : \mathbb{R}^{p+q+1} \rightarrow \mathbb{R}^n, \psi_N := R_\rho(\Psi_N(\theta_D)) : \mathbb{R}^n \rightarrow \mathbb{R}^N,$$

from the Subsection (3.2.1). Note, that $\mathbb{A}_P \neq \mathbb{A}$ in general, since like before $\mathbb{A}_P \in \mathbb{R}^{N_h \times N}$ and $\mathbb{A} \in \mathbb{R}^{N_h \times N_A}$. Similarly, refine the approximate parameter-to-solution map for the DOD+DNN into

$$\mathcal{G}_{\text{DOD+DFNN}} : \Theta \times \Theta' \times [0, T] \rightarrow \mathbb{R}^{N_h}; \quad (\mu, \nu, t) \mapsto \mathbb{V}_{\mu, t} \cdot \phi_{N'}(\mu, \nu, t),$$

the DOD+DFNN under weights $\theta_{\text{DOD+DFNN}} = (\theta_{\text{DOD}}, \theta_{\text{DF}_2})$ with

$$\begin{aligned} \mathbb{V}_{\mu, t} &:= \mathbb{A} \cdot R_\rho(\Phi_{\tilde{\mathbb{V}}}(\theta_{\text{DOD}}))(\mu, t) \in \mathbb{R}^{N_h \times N'}, \quad \text{for each } (\mu, t) \in \Theta \times [0, T], \text{ and} \\ \phi_{N'} &:= R_\rho(\Phi_{N'}(\theta_{\text{DF}_2})) : \mathbb{R}^{p+q+1} \rightarrow \mathbb{R}^{N'} \end{aligned}$$

from the Subsection (3.2.2) and the DOD Definition (3.6), and

$$\mathcal{G}_{\text{DOD-DL-ROM}} : \Theta \times \Theta' \times [0, T] \rightarrow \mathbb{R}^{N_h}; \quad (\mu, \nu, t) \mapsto \mathbb{V}_{\mu, t} \cdot (\psi_{N'} \circ \phi_n)(\mu, \nu, t),$$

the DOD-DL-ROM under weights $\theta_{\text{DOD-DL-ROM}} = (\theta_{\text{DOD}}, \theta_{\text{D}_2}, \theta_{\text{DF}_3})$ with

$$\begin{aligned} \mathbb{V}_{\mu, t} &:= \mathbb{A} \cdot R_\rho(\Phi_{\tilde{\mathbb{V}}}(\theta_{\text{DOD}}))(\mu, t) \in \mathbb{R}^{N_h \times N'}, \quad \text{for each } (\mu, t) \in \Theta \times [0, T], \text{ and} \\ \psi_{N'} &:= R_\rho(\Psi_{N'}(\theta_{\text{D}_2})) : \mathbb{R}^n \rightarrow \mathbb{R}^{N'}, \quad \phi_n := R_\rho(\Phi_n(\theta_{\text{DF}_3})) : \mathbb{R}^{p+q+1} \rightarrow \mathbb{R}^n, \end{aligned}$$

from the Subsection (3.2.2) and the DOD Definition (3.6).

Furthermore, let $\chi_* \in C^s(\mathbb{R}^n; \mathbb{R}^{N'})$ and $\chi'_* \in C^{s'}(\mathbb{R}^{N'}; \mathbb{R}^n)$ respectively be the maps attaining the perfect embedding Assumption (4), with $s \gg s' \geq 2$. Since $N' < N$ in general, the perfect embedding is implied for N aswell; let $\psi_* \in C^s(\mathbb{R}^n; \mathbb{R}^N)$ and $\psi'_* \in C^{s'}(\mathbb{R}^N; \mathbb{R}^n)$ be the maps attaining the equivalent for N .

Finally set constants

$$C'_1 := \sup_{|\alpha| \leq s'} \sup_{v \in \mathbb{R}^{N'}} |D^\alpha \chi_*(v)|, \quad C'_2 := \sup_{|\alpha| \leq s} \sup_{w \in \mathbb{R}^n} |D^\alpha \chi'_*(w)|,$$

and C_1, C_2 analogously for ψ_* and ψ'_* .

Theorem 4.12. *Let the preliminaries of the POD-DL-ROM be given as in the introduction of this Section (4.2). Then there is $N_{\text{data}} = N_{\text{data}}(\delta, \varepsilon)$, a constant $c = c(\Theta, \Theta', T, L, C_1, C_2, p, q, n, s, s')$ and a choice of weights for the POD-DL-ROM approximate parameter-to solution map $\mathbb{A}_P \cdot \hat{q}_N := \mathbb{A}_P \cdot (\psi_N \circ \phi_n) : \mathbb{R}^{p+q+1} \rightarrow \mathbb{R}^{N_h}$ such that*

$$\mathbb{P} \left[\mathcal{E}_R^{\text{POD-DL-ROM}} < \varepsilon \right] > 1 - \delta,$$

and the complexity is given by

- $L_{\Psi_N} = c \log(\varepsilon^{-1})$ layers,
- $\omega_{\Psi_N} = cN\varepsilon^{-n/(s-1)} \log(\varepsilon^{-1})$ active weights,

for the Decoder, and

- $L_{\Phi_n} = c \log(\varepsilon^{-1})$ layers,
- $\omega_{\Phi_n} = cn\varepsilon^{-(p+q+1)} \log(\varepsilon^{-1})$ active weights,

for the Deep Feed-Forward Neural Network.

Proof. We can directly bound $\mathcal{E}_S = \mathcal{E}_S(N_{s_1}, N_{s_2}, N_t)$ defined in the proof of Theorem (4.3) independently of N , under the Assumption (2) by the Weak Law of Large Numbers, i.e. for all $1 > \delta > 0$ and $\varepsilon > 0$ there are N_{s_1}, N_{s_2}, N_t , such that

$$\mathbb{P}[\mathcal{E}_S((N_{s_1}, N_{s_2}, N_t)) < \varepsilon/3] > 1 - \delta. \quad (4.2)$$

This result provides the "probability" part of the PAC (probably almost correct) bound, which we want to show. In other words, provided this, the other two error types can be bounded without regard to the probability space underlying the sampling procedure. Bounding $\mathcal{E}_{\text{NN}}$ and $\mathcal{E}_{\text{POD}}$ by $\varepsilon/3$ respectively will then provide the wanted result. For this, by choosing N as in Equation (4.1), we simply rewrite

$$\mathcal{E}_{\text{POD}} := m^{-1} \sqrt{\sum_{k>N} \sigma_k^2} \leq \frac{\varepsilon}{3}. \quad (4.3)$$

To now bound the neural network approximation error, we will need to simply rewrite the error in terms of a form accommodating a use of Gühring or Yarotsky Theorem;

$$\begin{aligned} \mathcal{E}_{\text{NN}} &= \left(\int_{\Theta \times \Theta' \times [0, T]} \frac{\|\mathbb{A}_P q_N(\mu, \nu, t) - \mathbb{A}_P \hat{q}_N(\mu, \nu, t)\|^2}{\|u_h^{\mu, \nu, t}\|^2} d(\mu, \nu, t) \right)^{1/2} \\ &\leq m^{-1} \left(\int_{\Theta \times \Theta' \times [0, T]} \|\mathbb{A}_P q_N(\mu, \nu, t) - \mathbb{A}_P \hat{q}_N(\mu, \nu, t)\|^2 d(\mu, \nu, t) \right)^{1/2} \\ &\leq m^{-1} \left(|\Theta \times \Theta' \times [0, T]| \sup_{(\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]} \|\mathbb{A}_P q_N(\mu, \nu, t) - \mathbb{A}_P \hat{q}_N(\mu, \nu, t)\|^2 \right)^{1/2} \\ &\leq m^{-1} \left(|\Theta \times \Theta' \times [0, T]| \cdot \|\mathbb{A}_P\|_{\text{op}}^2 \sup_{(\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]} \|q_N(\mu, \nu, t) - \hat{q}_N(\mu, \nu, t)\|^2 \right)^{1/2} \\ &= m^{-1} |\Theta \times \Theta' \times [0, T]|^{1/2} \sup_{(\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]} \|q_N(\mu, \nu, t) - \hat{q}_N(\mu, \nu, t)\|. \end{aligned} \quad (4.4)$$

whereas we split the latter error into both parts, since $\hat{q}_N = \psi_N \circ \phi_n$ per definition. This means employing two types of error estimations in the matter of finding the respective optimal maps for the feed forward estimation on the latent dynamics space via $\phi_* : \mathbb{R}^{(p+q+1)} \rightarrow \mathbb{R}^n$ and the decoder lift via $\psi_* : \mathbb{R}^n \rightarrow \mathbb{R}^N$.

- Let $\mathcal{S}_N := \{q(\mu, \nu, t) = \mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t} \in \mathbb{R}^N \mid (\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]\}$, then the image of the optimal embedding $\mathcal{V}_n = \psi'_*(\mathcal{S}_N) \in \mathbb{R}^n$ is such that $\text{diam}(\mathcal{V}_n) \leq C_1 L \cdot \text{diam}(\Theta \times \Theta' \times [0, T])$,

given the Assumption (3) and the preliminaries. In fact,

$$\sup_{w, w' \in \mathcal{V}_n} |w - w'| = \sup_{v, v' \in \mathcal{S}_N} |\psi'_*(v) - \psi'_*(v')| \quad (4.5)$$

$$\leq C_1 \sup_{v, v' \in \mathcal{S}_N} |v - v'| \quad (4.6)$$

$$\leq C_1 \sup_{(\mu, \nu, t), (\mu', \nu', t') \in \Theta \times \Theta' \times [0, T]} |\mathbb{A}_P^T \mathbb{G}(u_h^{\mu, \nu, t} - u_h^{\mu', \nu', t'})| \quad (4.7)$$

$$\leq C_1 \sup_{(\mu, \nu, t), (\mu', \nu', t') \in \Theta \times \Theta' \times [0, T]} |\mathbb{A}_P^T \mathbb{G}|_{\text{op}} |u_h^{\mu, \nu, t} - u_h^{\mu', \nu', t'}| \quad (4.8)$$

$$\leq C_1 L \cdot \text{diam}(\Theta \times \Theta' \times [0, T]). \quad (4.9)$$

Here, we have used the definition of $\mathcal{V}_n$ as the image of ψ'_* in Step (4.5), the fact, that

$$C_1 \geq \sup_{v \in \mathcal{S}_N} |D^\alpha \psi'_*|$$

for $|\alpha| = 1$ and its implication under the mean value theorem (Rudin, 1976, p. 107) in Step (4.6) and the fact, that $|\mathbb{A}_P^T \mathbb{G}|_{\text{op}} = \|\mathbb{A}_P\|_{\text{op}} = 1$ by $\mathbb{G}$ -orthonormality of $\mathbb{A}_P$, together with the Lipschitz-quality of $\mathcal{G}$ from Assumption (3) in Step (4.8). This argument allows us to find that $\psi_* \in \mathcal{F}_{s, n, \infty, C_1 L \cdot \text{diam}(\Theta \times \Theta' \times [0, T])}$, after unilateral prior transformation $x \mapsto x/|x| \in (0, 1)^n$ of any input, for the target space of the optimal function required by Theorem Gühring (2.34). Thus there are neural network weights θ_D for $\psi_N = R_\rho(\Psi_N(\theta_D)) : \mathbb{R}^n \rightarrow \mathbb{R}^N$ fitting the network realization to that optimal function, such that

$$\|\psi_* - \psi_N\|_{W^{1, \infty}((0, 1)^d)} \leq \frac{m}{6} |\Theta \times \Theta' \times [0, T]|^{-1/2} \varepsilon,$$

with $L_{\Psi_N} = c \log(\varepsilon^{-1})$ layers and $\omega_{\Psi_N} = cN\varepsilon^{-n/(s-1)} \log(\varepsilon^{-1})$ active weights. For the sake of completeness, we mention that one would build N -many entry-wise neural networks in parallel and hence attain for each $\omega_{[\psi_N]_1} = c\varepsilon^{-n/(s-1)} \log(\varepsilon^{-1})$ active weights under Gühring, then employ the Lemma (2.28) for the parallelization of these. Note, that this does not influence the order of the layer number. A similar approach was already argued in the proof of Proposition (4.8). This statement is equivalent to

$$\sup_{v \in \mathcal{V}_N} \|\psi_*(v) - \psi_N(v)\| < \frac{m}{6} |\Theta \times \Theta' \times [0, T]|^{-1/2} \varepsilon, \quad (4.10)$$

$$\text{ess sup}_{v, v' \in \mathcal{V}_N} \frac{|(\psi_* - \psi_N)(v) - (\psi_* - \psi_N)(v')|}{|v - v'|} < \frac{m}{6} |\Theta \times \Theta' \times [0, T]|^{-1/2} \varepsilon, \quad (4.11)$$

by definition of the derivative and the fact, that Sobolev norms are the sum of both the equations above. Additionally this implies that, ψ_N is Lipschitz with constant $C_3 := C_2 + m/6 \cdot |\Theta \times \Theta' \times [0, T]|^{-1/2}$, since

$$\frac{|(\psi_* - \psi_N)(v) - (\psi_* - \psi_N)(v')|}{|v - v'|} \geq \frac{|\psi_N(v) - \psi_N(v')|}{|v - v'|} - \frac{|\psi_*(v) - \psi_*(v')|}{|v - v'|}$$

by inverse triangle inequality for any $v, v' \in \mathcal{V}_N$ and subsequently employing the C_2 -bound for ψ_* from the preliminaries of this Section (4.2).

- Now let $\phi_*(\mu, \nu, t) = \psi'_*(q(\mu, \nu, t))$ for $(\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]$, which is Lipschitz continuous, with constant LC_1 . In fact, we have for any pair $(\mu, \nu, t), (\mu', \nu', t') \in \Theta \times \Theta' \times [0, T]$,

$$\begin{aligned} |\phi_*(\mu, \nu, t) - \phi_*(\mu', \nu', t')| &= |\psi'_*(\mathbb{A}_P^T \mathbb{G} u_h^{\mu, \nu, t}) - \psi'_*(\mathbb{A}_P^T \mathbb{G} u_h^{\mu', \nu', t'})| \\ &\leq C_1 |\mathbb{A}_P^T \mathbb{G}(u_h^{\mu, \nu, t} - u_h^{\mu', \nu', t'})| \\ &\leq C_1 L \cdot (|\mu - \mu'| + |\nu - \nu'| + |t - t'|), \end{aligned}$$

by the very same arguments as in the calculation for the diameter of $\mathcal{V}_n$ above. The Lipschitz continuity of this map then allows us to employ Theorem (2.11), which guarantees that $\phi_* \in W^{1,\infty}(\mathcal{S}_N)$, since $\text{diam}(\mathcal{S}_N) \leq L \cdot \text{diam}(\Theta \times \Theta' \times [0, T])$ by the same arguments as in the $\mathcal{V}_n$ -case. Hence, after entry-wise normalization, $\phi_* \in \mathcal{F}_{1,p+q+1,\infty,\text{diam}(\mathcal{S}_N)}$.

Now we employ Theorem Yarotski (2.33) to assert the existence of neural network weights θ_{DF} such that its realization $\phi_n := R_\rho(\Phi_n(\theta_{\text{DF}})) : \mathbb{R}^{p+q+1} \rightarrow \mathbb{R}^n$ attains

$$\|\phi_* - \phi_n\|_{W^{0,\infty}(\Theta \times \Theta' \times [0, T])} < \frac{m}{6C_3} |\Theta \times \Theta' \times [0, T]|^{-1/2} \varepsilon, \quad (4.12)$$

with $L_{\Phi_n} = c \log(\varepsilon^{-1})$ layers and $\omega_{\Phi_n} = cn\varepsilon^{-(p+q+1)} \log(\varepsilon^{-1})$ active weights, again with the argument of neural network parallelization.

Starting from the last inequality in (4.4) we can perform the split in a clean fashion using the triangle inequality and reminding ourselves of the Assumption (4), which guarantees $q = \psi_*(\psi'_*)$, and thus gives with both neural network estimates in Equations (4.10) and (4.12)

$$\begin{aligned} & \sup_{(\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]} \|q(\mu, \nu, t) - \hat{q}_N(\mu, \nu, t)\| \\ & \leq \sup_{(\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]} (\|\psi_*(\psi'_*(\mu, \nu, t)) - \psi(\phi_*(\mu, \nu, t))\| + \|\psi(\phi_*(\mu, \nu, t)) - \psi(\phi(\mu, \nu, t))\|) \\ & \leq \sup_{v \in \mathcal{V}_N} \|\psi_*(v) - \psi(v)\| + C_3 \sup_{(\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]} (\|\psi(\phi_*(\mu, \nu, t)) - \psi(\phi(\mu, \nu, t))\|) \\ & < \frac{m}{6} |\Theta \times \Theta' \times [0, T]|^{-1/2} \varepsilon + C_3 \frac{m}{6C_3} |\Theta \times \Theta' \times [0, T]|^{-1/2} \varepsilon = \frac{\varepsilon}{3} m |\Theta \times \Theta' \times [0, T]|^{-1/2}, \end{aligned}$$

which cancels with $m^{-1} |\Theta \times \Theta' \times [0, T]|^{1/2}$ to get

$$\mathcal{E}_{\text{NN}} < \frac{\varepsilon}{3}. \quad (4.13)$$

Now putting together all three estimates (4.2), (4.3) and (4.13) with the reasoning of Theorem (4.3) finally gives

$$\mathcal{E}_R \leq \mathcal{E}_S + \mathcal{E}_{\text{POD}} + \mathcal{E}_{\text{NN}} < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon,$$

with higher probability than $1 - \delta$. $\square$

Building upon the discussion in the beginning of this Section (4.2), we may explore the implication of a more regular optimal DOD for $\mathcal{X} = \mathcal{P}_2$ from Definition (4.6). Indeed, if such an assumption is given, then Yarotsky Theorem can be invoked for the implied construction of the inner DOD module in Remark (4.9).

Assumption 6. Let the objective DOD functional $J : \Theta \times [0, T] \times [-1, 1]^{N_h \times N'}$ be given for $\mathcal{X} = \mathcal{P}_2$ as in Definition (4.6), and the corresponding optimal DOD

$$s : \Theta \times [0, T] \mapsto [0, 1]^{N_h \times N'}; \quad (\mu, t) \mapsto \underset{\tilde{\mathbf{V}} \in [-1, 1]^{N_A \times N'}}{\text{argmin}} \quad J(\mu, t, \mathbb{A}\tilde{\mathbf{V}})$$

be $\hat{L}$ -Lipschitz.

This assumption will now provide us with the ability to use Yarotsky Theorem retroactively in Proposition (4.8) to achieve bounds on the complexity for the inner DOD module. However, the relative error bound, we discussed in Theorem (4.10) needs an assertion for N_A to be close to N_h to achieve the actual benefit of a (μ, t) -dependent projector.

As a matter of fact, in the following we set $N_A = N_h$ to avoid cumbersome notation and make use of the Assumption (1) later on. This is within the framework of Assumption (5), but note, that this can be relaxed in a similar way to the POD error $\mathcal{E}_{\text{POD}}$, as has been done in Theorem (4.12). In fact, this constitutes that N_A must be chosen in the same order as N has been for the POD-DL-ROM, since the ambient error $c_{\mathbb{A}}$ from Proposition (2.12) has the same inhibition of scaling with the total POD eigenvalues.

Theorem 4.13. *Let the preliminaries of the DOD-DL-ROM be given as in the introduction of this Section (4.2), $N_A = N_h$ explicitly and the Assumption (6) be true. Then there is $N_{\text{data}} = N_{\text{data}}(\delta, \varepsilon)$, a constant $c = c(\Theta, \Theta', T, L, \hat{L}, C_1, C_2, p, q, n, s, s')$ and a choice of weights for the DOD-DL-ROM approximate parameter-to solution map $(\mu, \nu, t) \mapsto \mathbb{V}_{\mu, t} \cdot \hat{q}_{N'} := \mathbb{V}_{\mu, t} \cdot (\psi_{N'} \circ \phi_n) : \mathbb{R}^{p+q+1} \rightarrow \mathbb{R}^{N_h}$ such that*

$$\mathbb{P} \left[\mathcal{E}_R^{\text{DOD-DL-ROM}} < \varepsilon \right] > 1 - \delta,$$

and the complexity is given by

- $L_{\Phi_{\tilde{V}}} = c \log(\varepsilon^{-1})$ layers,
- $\omega_{\Phi_{\tilde{V}}} = c N_A N' \varepsilon^{-(p+1)} \log(\varepsilon^{-1})$ active weights,

for the inner DOD module,

- $L_{\Psi_{N'}} = c \log(\varepsilon^{-1})$ layers,
- $\omega_{\Psi_{N'}} = c N' \varepsilon^{-n/(s-1)} \log(\varepsilon^{-1})$ active weights,

for the Decoder, and

- $L_{\Phi_n} = c \log(\varepsilon^{-1})$ layers,
- $\omega_{\Phi_n} = c n \varepsilon^{-(p+q+1)} \log(\varepsilon^{-1})$ active weights,

for the Deep Feed-Forward Neural Network.

Proof. Similar as to the proof of Theorem (4.12), we will try to bound each quantity of the statement in Theorem (4.10), but with the catch, that $\mathcal{E}_{\text{DOD}}$ is stochastic in nature, and thus we will employ Remark (4.9) to find an auxiliary term, which is independent of the data, to which the approximation of the neural net of the DOD is sufficiently "close". Here, we instantaneously remark, that taking $N_A = N_h$ implies $\mathcal{E}_{\mathbb{A}} = 0$, and we will not mention this contribution again.

The sampling error can be bounded in the same way as before, i.e. we can directly bound $\mathcal{E}_S = \mathcal{E}_S(N_{s_2})$ independently of N' , under the Assumption (2) by the Weak Law of Large Numbers, i.e. for all $1 > \delta > 0$ and $\varepsilon > 0$ there is N_{s_2} , such that

$$\mathbb{P}[\mathcal{E}_S < \varepsilon/4] > 1 - \delta. \quad (4.14)$$

Now recall the definition of $\mathcal{E}_{\text{DOD}}$ in the proof of Theorem (4.10) and further setting the $\delta > 0$ from the upstream Remark (4.9) realizing an "optimal" discrete approximation for a fixed data size N_{s_2} as

$$\delta = \frac{\varepsilon m}{4|\Theta \times [0, T]|^{1/2}}$$

concluding, under the additional use of definition of N' in Equation (4.1), in the following approximation for the DOD error:

$$\begin{aligned}\mathcal{E}_{\text{DOD}} &= \frac{|\Theta \times [0, T]|^{1/2}}{m} \left(\delta + \sup_{(\mu, t) \in \Theta \times [0, T]} \sqrt{\sum_{k > N'} \sigma(\mu, t)_k^2} \right) \\ &< \frac{\varepsilon}{4} + \left(\frac{|\Theta \times [0, T]|^{1/2}}{m} \sup_{(\mu, t) \in \Theta \times [0, T]} \sqrt{\sum_{k > N'} \sigma(\mu, t)_k^2} \right) \\ &\leq \frac{\varepsilon}{4} + \left(\frac{|\Theta \times [0, T]|^{1/2}}{m} \cdot \frac{\varepsilon m}{4|\Theta \times [0, T]|^{1/2}} \right) = \frac{\varepsilon}{2}\end{aligned}\tag{4.15}$$

Under the Assumption (6), we get that a neural network can be indeed found, such that it realizes

$$\| [s]_{ij} - R_\rho([\tilde{V}_0]_{ij}) \|_{L^\infty(\Theta \times [0, T]; \mathbb{R})} \leq \delta,$$

for each $(i, j) \in \{1, \dots, N_A\} \times \{1, \dots, N'\}$ inside the proof of Proposition (4.8) under the specifications of Remark (4.9), with at most

- $L_{\Phi_{[\tilde{V}_0]_{ij}}} = c'' \log(\varepsilon^{-1})$ layers and
- $\omega_{\Phi_{[\tilde{V}_0]_{ij}}} = c'' \varepsilon^{-(p+1)} \log(\varepsilon^{-1})$ active weights,

for some $c'' = c''(\Theta, T, p, \hat{L}) > 0$ by Yarotsky Theorem (2.33), since the $\hat{L}$ -Lipschitz property of the optimal DOD justifies $W^{1, \infty}$ -regularity with a $\hat{L}$ bound by Theorem (2.11). Then with the parallelization of the next argument of the proof, we get no change in the layer bound, but we multiply the weights approximation with N' and with N_A by Lemma (2.28). Lastly, the concatenation of the neural network of $\tilde{V}_0$ with ζ for normalization only changes the layers by a constant according to Lemma (2.26), thus achieving for the inner DOD module at most

- $L_{\Phi_{\tilde{V}}} = c' \log(\varepsilon^{-1})$ layers and
- $\omega_{\Phi_{\tilde{V}}} = c' N_A N' \varepsilon^{-(p+1)} \log(\varepsilon^{-1})$ active weights,

with $c' = 3 + c''$.

Moreover we can use the exact same arguments as in the bound for $\mathcal{E}_{\text{NN}}$ for the POD-DL-ROM as in the proof of Theorem (4.12), which yields

$$\mathcal{E}_{\text{NN}} < \frac{\varepsilon}{4}.\tag{4.16}$$

For sake of uniformity, we then take $c > 0$ to be the maximal of each individual constant of all three separate neural networks. Note, that here we imply, that the reduction is achieved with the replacement of N with N' in each of the arguments (and also the use of potentially different weights and architectures), hence also getting slightly different values for the number of layers and weights. Finally putting each Equation, i.e. (4.14), (4.15) and (4.16), together, we get

$$\mathcal{E}_R \leq \mathcal{E}_S + \mathcal{E}_{\text{DOD}} + \mathcal{E}_{\text{NN}} < \frac{\varepsilon}{4} + \frac{\varepsilon}{2} + \frac{\varepsilon}{4} = \varepsilon,$$

with probability greater than $1 - \delta$ concluding our proof. $\square$

In the next step, we can follow up this complexity PAC bound with the same statement for the slightly simpler architecture of the DOD+DFNN. It is clear, that the omission of the reduction by an autoencoder simplifies the reduction procedure by one step. This does not change basically anything regarding the proof, as the choice of N' ensures a bound here too.

Theorem 4.14. *Let the preliminaries of the DOD+DFNN be given as in the introduction of this Section (4.2), $N_A = N_h$ explicitly and the Assumption (6) be true. Then there is $N_{data} = N_{data}(\delta, \varepsilon)$, a constant $c = c(\Theta, \Theta', T, L, \hat{L}, p, q, N')$ and a choice of weights for the DOD+DFNN approximate parameter-to solution map $(\mu, \nu, t) \mapsto \mathbb{V}_{\mu, t} \cdot \hat{q}_{N'} := \mathbb{V}_{\mu, t} \cdot \phi_{N'} : \mathbb{R}^{p+q+1} \rightarrow \mathbb{R}^{N_h}$ such that*

$$\mathbb{P} \left[\mathcal{E}_R^{DOD+DFNN} < \varepsilon \right] > 1 - \delta,$$

and the complexity is given by

- $L_{\Phi_{\tilde{\mathbb{V}}}} = c \log(\varepsilon^{-1})$ layers,
- $\omega_{\Phi_{\tilde{\mathbb{V}}}} = c N_A N' \varepsilon^{-(p+1)} \log(\varepsilon^{-1})$ active weights,

for the inner DOD module, and

- $L_{\Phi_{N'}} = c \log(\varepsilon^{-1})$ layers,
- $\omega_{\Phi_{N'}} = c N' \varepsilon^{-(p+q+1)} \log(\varepsilon^{-1})$ active weights,

for the Deep Feed-Forward Neural Network.

do something about the delta and varepsilon definitions

add quick complexity bound for the dod+dfnn architecture

4.3 Comparison between existing Techniques

In this section, we will compare the presented data-driven ROM techniques with each other in the context of the Theorems (4.12), (4.13) and (4.14), or in other words; we will examine the differences of the upper complexity bounds for the POD-based neural network approach and both DOD-based neural network approaches. As another step, these complexities will be coarsely compared to other techniques, not yet presented in the paper, which have been proposed in the context of data-driven Reduced Order Modelling. For this, we will recite some results from relevant literature and argue the advantages and disadvantages of the DOD-based ROMs in comparison.

Within this framework, we will play around with potentially sharper assumptions, especially hypothetically improving regularity of the parameter-to-solution map $\mathcal{G}$ in Assumption (3) and the regularity of the optimal DOD s in Assumption (6). These two regularities namely determine the effectivity of the neural net under Yartosky (and/or Gühring) Theorem (2.33). To have a substantial difference in the use of the DOD, as opposed to the POD, we want to focus the readers attention on Assumption (1), which ensures

1. For $\{\sigma_{k,\infty}^2\}_k$ the eigenvalues of the POD matrix given under the continuous correlation matrix of the entire solution manifold $\mathcal{S} = \{u_h^{\mu,\nu,t} \mid (\mu, \nu, t) \in \Theta \times \Theta' \times [0, T]\}$ a slow decay, i.e.

$$\sqrt{\sum_{k>n} \sigma_{k,\infty}^2} \leq C n^{-\alpha},$$

for $C > 0$ a constant and $0 < \alpha < 1$, and

2. For $\{\sigma(\mu, t)_{k,\infty}^2\}_k$ the eigenvalues of the (μ, t) -fixed POD matrix given under the continuous non-reduced correlation matrix of the (μ, t) -dependent solution manifold $\mathcal{S}_{\mu,t} = \{u_h^{\mu,\nu,t} \mid \nu \in \Theta'\}$ a quick decay, i.e.

$$\sup_{(\mu,t) \in \Theta \times [0,T]} \sqrt{\sum_{k>n} \sigma(\mu, t)_{k,\infty}^2} \leq C' n^{-\beta},$$

for $C' > 0$ a constant and $\beta \geq 1$.

This fact derives as a direct consequence from Proposition (2.14) and the definitions of the continous correlation matrices in Definition (2.13). With these two bounds, we can presumably take for very large data $N_s, N_t \rightarrow \infty$ and $N_A = N_h$

$$N = \mathcal{O}\left(\varepsilon^{-\frac{1}{\alpha}}\right), \quad N' = \mathcal{O}\left(\varepsilon^{-\frac{1}{\beta}}\right), \quad (4.1)$$

for the guaranteed PAC bounds of the POD-DL-ROM, the DOD+DFNN and the DOD-DL-ROM of fixed precision $\varepsilon > 0$ as is dictated in the beginning of Section (4.2). In fact, we know that by Proposition (2.14) $\sum_{k>n} \sigma_k^2 \rightarrow \sum_{k>n} \sigma_{k,\infty}^2$ for $N_s, N_t \rightarrow \infty$, and therefore

$$N = \arg \min_{n \in \mathbb{N}} \left\{ \sum_{k>n} \sigma_k^2 \geq \frac{m^2}{9} \varepsilon^2 \right\} \xrightarrow{N_s, N_t \rightarrow \infty} \arg \min_{n \in \mathbb{N}} \left\{ C n^{-2\alpha} \geq \frac{m^2}{9} \varepsilon^2 \right\} = \mathcal{O}(\varepsilon^{-\frac{1}{\alpha}}),$$

and a similar argument for N' . These rough estimates for the order of N and N' then let us get an idea for the upcoming comparisons, which mainly will involve the weighing of the higher complexity cost of the DOD-based ROMs against the lighter, but potentially less reductive, POD-based methods. Note, that the assumption of $N_A = N_h$ could be relaxed, if the pre-reduced (μ, t) -dependent solution manifold also has fast KnW decay.

4.3.1 Comparison between POD-based and DOD-based ROMs

Within the subsection, our purpose will be to list the differences between the POD-based techniques for model order reduction and the DOD-based ones depending on the regularity imposed on the optimal DOD s as given by Definition (4.6), for $N_A = N_h$, and the regularity imposed on the parameter-to-solution map $\mathcal{G}$. As such, assume additionally to the framework of the Section (4.2),

1. $\mathcal{G} \in W^{r,\infty}(\Theta \times \Theta' \times [0, T]; \mathbb{R}^{N_h})$ for some $r \geq 1$,
2. $s \in W^{m,\infty}(\Theta \times [0, T]; \mathbb{R}^{N_h \times N'})$ for some $m \geq 1$.

Let also $0 < \varepsilon < 1$ in general for the sake of monotonicity later on. To cluster the total complexity of a ROM, we can just add the respective layers and weights of each neural network contained in each method. Such a procedure leaves us with

$$\begin{aligned} - \omega_{\text{POD-DL-ROM}} &= \mathcal{O}\left(N \varepsilon^{-n/(s-1)} \log(\varepsilon^{-1}) + n \varepsilon^{-(p+q+1)/r} \log(\varepsilon^{-1})\right), \\ - \omega_{\text{DOD-DL-ROM}} &= \mathcal{O}\left(N_h N' \varepsilon^{-(p+1)/m} \log(\varepsilon^{-1}) + N' \varepsilon^{-n/(s-1)} \log(\varepsilon^{-1}) + n \varepsilon^{-(p+q+1)/r} \log(\varepsilon^{-1})\right), \\ - \omega_{\text{DOD+DFNN}} &= \mathcal{O}\left(N_h N' \varepsilon^{-(p+1)/m} \log(\varepsilon^{-1}) + N' \varepsilon^{-(p+q+1)/r} \log(\varepsilon^{-1})\right), \end{aligned}$$

and the same layer bound for all of them by insertion of the regularity in the respective proofs of the upper complexity bounds, where Yarosky Theorem (2.33) has been used. We remark, that in Assumption (4), we set $n \leq 2(p+q) + 3$ in general, and $s \gg s' \geq 2$ in the preliminaries of

Section (4.2). This quantifies all relevant complexity numbers. Namely, we want to analyse three distinct cases, starting with a general agnostic one, where r, m are close to 1, in a sense that $n/(s-1) < (p+1)/m$ and $n/(s-1) = (p+q+1)/r$. This directly results in

$$\begin{aligned}\omega_{\text{POD-DL-ROM}} &= \mathcal{O}\left(N\varepsilon^{-n/(s-1)}\log(\varepsilon^{-1}) + n\varepsilon^{-(p+q+1)/r}\log(\varepsilon^{-1})\right) \\ &\lesssim \mathcal{O}\left(N\varepsilon^{-n/(s-1)}\log(\varepsilon^{-1})\right) \\ &\ll \mathcal{O}\left(N_h N' \varepsilon^{-(p+1)/m}\log(\varepsilon^{-1})\right) \\ &\lesssim \omega_{\text{DOD+DFNN}},\end{aligned}$$

where we used, that $N \ll N_h$ and the fact that $n \ll N'(\beta, \varepsilon), N(\alpha, \varepsilon)$ by Assumption (5). The same heuristic can be done for DOD-DL-ROM active weights bound, since essentially the number of weights are dominated by the DOD neural network contribution anyways. The situation changes, however, if we can assert a high regularity for the optimal DOD s , i.e. if $m \rightarrow \infty$ such that $(p+1)/m \approx 0$, since then the contribution of the DOD to the complexity is minimal and we get

$$\begin{aligned}\omega_{\text{DOD-DL-ROM}} &\approx \mathcal{O}\left(N' \varepsilon^{-n/(s-1)}\log(\varepsilon^{-1}) + n\varepsilon^{-(p+q+1)/r}\log(\varepsilon^{-1})\right) \\ &\lesssim \mathcal{O}\left(N\varepsilon^{-n/(s-1)}\log(\varepsilon^{-1})\right) \\ &\approx \omega_{\text{POD-DL-ROM}}.\end{aligned}$$

This can be easily attributed to the fact, that $N' + n \lesssim N$, because of Equation (4.1). Here, we have also still prescribed the dominance of the decoder complexity over the forward pass, therefore awarding the DOD-DL-ROM with a huge bonus due to the inherently better approximation under the smooth optimal decoder map $\zeta_* \in C^s(\mathbb{R}^n; \mathbb{R}^{N'})$. In fact,

$$\begin{aligned}\omega_{\text{DOD-DL-ROM}} &\approx \mathcal{O}\left(N' \varepsilon^{-n/(s-1)}\log(\varepsilon^{-1}) + n\varepsilon^{-(p+q+1)/r}\log(\varepsilon^{-1})\right) \\ &\lesssim \mathcal{O}\left(N\varepsilon^{-n/(s-1)}\log(\varepsilon^{-1})\right) \\ &\approx \omega_{\text{POD-DL-ROM}}.\end{aligned}$$

Chapter 5

Numerical Experiments

Include all numerical experiments

Find increase of performance depending on latent dimension

Give benchmark on pass forward times

Discuss comparison between different DL-ROMs

Chapter 6

Conclusion

Discuss results

Insert outlook for future ideas and implementation

Bibliography

- Alt, H. W. (2016, July). *Linear functional analysis*. Springer London. <https://doi.org/10.1007/978-1-4471-7280-2>
- Barraut, M., Maday, Y., Nguyen, N. C., & Patera, A. T. (2004). An 'empirical interpolation' method: Application to efficient reduced-basis discretization of partial differential equations. *Comptes Rendus Mathematique*, 339(9), 667–672. <https://doi.org/10.1016/j.crma.2004.08.006>
- Bauer, F. L., & Fike, C. T. (1960). Norms and exclusion theorems. *Numerische Mathematik*, 2, 137–141. <https://api.semanticscholar.org/CorpusID:121278235>
- Bramble, J. H., & Hilbert, S. R. (1970). Bounds for a class of linear functionals with applications to Hermite interpolation. *Numerische Mathematik*, 16(4), 362–369. <https://doi.org/10.1007/BF02165086>
- Brezis, H. (2011). *Functional analysis, sobolev spaces and partial differential equations* (1st ed.). Springer New York. <https://doi.org/10.1007/978-0-387-70914-7>
- Brezis, H., & Browder, F. (1998). Partial differential equations in the 20th century. *Advances in Mathematics*, 135(1), 76–144. <https://doi.org/https://doi.org/10.1006/aima.1997.1713>
- Brivio, S., Fresca, S., Franco, N. R., & Manzoni, A. (2023). Error estimates for pod-dl-roms: A deep learning framework for reduced order modeling of nonlinear parametrized pdes enhanced by proper orthogonal decomposition. <https://arxiv.org/abs/2305.04680>
- Buchfink, P., Glas, S., & Haasdonk, B. (2023). Approximation bounds for model reduction on polynomially mapped manifolds. *arXiv preprint arXiv:2312.00724*.
- Chaturantabut, S., & Sorensen, D. C. (2010). Nonlinear model reduction via discrete empirical interpolation. *SIAM Journal on Scientific Computing*, 32(5), 2737–2764. <https://doi.org/10.1137/090766498>
- Ciampiconi, L., Elwood, A., Leonardi, M., Mohamed, A., & Rozza, A. (2024). A survey and taxonomy of loss functions in machine learning. <https://arxiv.org/abs/2301.05579>
- Courant, R., & Hilbert, D. (1953). *Methods of mathematical physics, vol. i*. Interscience Publishers.
- Dacorogna, B. (2004, November). *Introduction to the calculus of variations*. <https://doi.org/10.1142/p361>
- Franco, N. R., Manzoni, A., Zunino, P., & Hesthaven, J. S. (2024). Deep orthogonal decomposition: A continuously adaptive data-driven approach to model order reduction [arXiv:2404.18841]. <https://arxiv.org/abs/2404.18841>
- Fresca, S., Dede, L., & Manzoni, A. (2020). A comprehensive deep learning-based approach to reduced order modeling of nonlinear time-dependent parametrized pdes. <https://arxiv.org/abs/2001.04001>
- Fresca, S., & Manzoni, A. (2022). Pod-dl-rom: Enhancing deep learning-based reduced order models for nonlinear parametrized pdes by proper orthogonal decomposition. *Computer Methods in Applied Mechanics and Engineering*, 388, 114181. <https://doi.org/10.1016/j.cma.2021.114181>
- Golub, G. H., & Van Loan, C. F. (2013). *Matrix computations*. Johns Hopkins University Press. <https://doi.org/10.56021/9781421407944>

- Grepl, M. A., & Patera, A. T. (2005). A posteriori error bounds for reduced-basis approximations of parametrized parabolic partial differential equations. *ESAIM: Mathematical Modelling and Numerical Analysis*, 39, 157–181. <https://doi.org/10.1051/m2an:2005006>
- Gühring, I., Kutyniok, G., & Petersen, P. (2019). Error bounds for approximations with deep relu neural networks in $W^{s,p}$ norms. <https://arxiv.org/abs/1902.07896>
- Gühring, I., Raslan, M., & Kutyniok, G. (2020). Expressivity of deep neural networks. <https://arxiv.org/abs/2007.04759>
- Haasdonk, B., & Ohlberger, M. (2008). Reduced basis method for finite volume approximations of parametrized linear evolution equations. *ESAIM: Mathematical Modelling and Numerical Analysis*, 42, 277–302. <https://doi.org/10.1051/m2an:2008001>
- Heinonen, J. (2005). Lectures on lipschitz analysis [[Online; accessed 20-September-2018]].
- Hesthaven, J., Pagliantini, C., & Rozza, G. (2022). Reduced basis methods for time-dependent problems. *Acta Numerica*, 31, 265–345. <https://doi.org/10.1017/S0962492922000058>
- Holmes, P., Lumley, J. L., & Berkooz, G. (1996). *Turbulence, coherent structures, dynamical systems and symmetry*. Cambridge University Press.
- Hornik, K. (1991). Approximation capabilities of multilayer feedforward networks. *Neural Networks*, 4(2), 251–257. [https://doi.org/https://doi.org/10.1016/0893-6080\(91\)90009-T](https://doi.org/https://doi.org/10.1016/0893-6080(91)90009-T)
- Kunisch, K., & Volkwein, S. (2001). Galerkin proper orthogonal decomposition methods for a general equation in fluid dynamics. *SIAM Journal on Numerical Analysis*, 40(2), 492–515. <https://doi.org/10.1137/S0036142900382612>
- Lumley, J. L. (1967). The structure of inhomogeneous turbulent flows. In A. M. Yaglom & V. I. Tatarsky (Eds.), *Atmospheric turbulence and radio wave propagation* (pp. 166–178). Nauka.
- Maday, Y., Patera, A. T., & Rovas, D. V. (2002). A blackbox reduced-basis output bound method for noncoercive linear problems. In *Proceedings of the 12th international conference on domain decomposition methods*. Domain Decomposition Press.
- Noor, A. K. (1980). Recent advances in reduction methods for nonlinear problems. *Computers & Structures*, 12(1), 31–44. [https://doi.org/10.1016/0045-7949\(80\)90101-6](https://doi.org/10.1016/0045-7949(80)90101-6)
- Patera, A. T. (1984). A spectral element method for fluid dynamics: Laminar flow in a channel expansion. *Journal of Computational Physics*, 54(3), 468–488. [https://doi.org/10.1016/0021-9991\(84\)90128-1](https://doi.org/10.1016/0021-9991(84)90128-1)
- Peherstorfer, B., Willcox, K., & Gunzburger, M. (2016). Optimal model management for multifidelity monte carlo estimation. *SIAM Journal on Scientific Computing*, 38(5), A3163–A3194. <https://doi.org/10.1137/15M1046472>
- Prud’homme, C., Rovas, D. V., Veroy, K., Machiels, L., Maday, Y., Patera, A. T., & Turinici, G. (2002). Reliable real-time solution of parametrized partial differential equations: Reduced-basis output bound methods. *Journal of Fluids Engineering*, 124(1), 70–80. <https://doi.org/10.1115/1.1448332>
- Quarteroni, A., Manzoni, A., & Negri, F. (2015). *Reduced basis methods for partial differential equations*. Springer. <https://doi.org/10.1007/978-3-319-15431-2>
- Rowley, C. W., Mezić, I., Bagheri, S., Schlatter, P., & Henningson, D. S. (2009). Spectral analysis of nonlinear flows. *Journal of Fluid Mechanics*, 641, 115–127. <https://doi.org/10.1017/S0022112009992059>
- Rudin, W. (1976). *Principles of mathematical analysis* (3rd ed.). McGraw-Hill.
- Schmid, P. J. (2010). Dynamic mode decomposition of numerical and experimental data. *Proceedings of the 61st Annual Meeting of the APS Division of Fluid Dynamics*.
- Siena, P., Girfoglio, M., & Rozza, G. (2022). An introduction to pod-greedy-galerkin reduced basis method. <https://arxiv.org/abs/2203.08532>

- Sirovich, L. (1987). Turbulence and the dynamics of coherent structures. part i: Coherent structures. *Journal of Fluid Mechanics*, 175, 419–427. <https://doi.org/10.1017/S0022112087000199>
- Yarotsky, D. (2017). Error bounds for approximations with deep relu networks. *Neural Networks*, 94, 103–114. <https://doi.org/https://doi.org/10.1016/j.neunet.2017.07.002>