

The Balance Constant and Its Decomposition

Peter Groom

February 2026

Contents

1	The Balance Constant and Its Decomposition	2
1.1	§1. Introduction	2
1.2	§2. Background: The Two Constants	3
1.2.1	§2.1. The Euler–Mascheroni Constant	3
1.2.2	§2.2. The Natural Logarithm of	3
1.2.3	§2.3. Why Their Sum?	3
1.3	§3. The Measurement: Four Domains	4
1.3.1	§3.1. Summary Table	4
1.3.2	§3.2. Statistical Significance	4
1.4	§4. Domain 1 — Fibonacci Arithmetic	5
1.4.1	§4.1. The Formula	5
1.4.2	§4.2. The Discretisation Gap	5
1.5	§5. Domain 2 — Cellular Automata	5
1.5.1	§5.1. The Observation	5
1.5.2	§5.2. Statistical Tests	6
1.5.3	§5.3. Interpretation	6
1.6	§6. Domain 3 — The Prime Number Sieve	6
1.6.1	§6.1. PAC Conservation in the Sieve	6
1.6.2	§6.2. The Connection to	6
1.6.3	§6.3. The Three-Phase Model	7
1.7	§7. Domain 4 — Landauer Erasure	7
1.8	§7.1. Domain 5 — Möbius Field Dynamics	8
1.8.1	§7.1.1. The Observation	8
1.8.2	§7.1.2. PAC Conservation	8
1.8.3	§7.1.3. Why This Domain Is Independent	8
1.9	§8. Base Invariance	9
1.10	§9. The Decomposition: Why $+ \ln(\cdot)$	9
1.10.1	§9.1. as Emergence Surplus	9
1.10.2	§9.2. Why Not Something Else?	9
1.11	§9.3. PAC Depth Constraints	10
1.12	§10. What Remains Open	11
1.12.1	§10.1. The Exact Equality Problem	11
1.12.2	§10.2. Why These Five Domains?	11
1.12.3	§10.3. PAC Depth and Physical Hierarchies	11
1.12.4	§10.4. The Landauer Precision Gap	11
1.13	§11. Falsification Conditions	11
1.14	§12. Summary of Results	12
1.15	§12.1. Connections to the PACSeries	13

1.16 §13. Numerical Details	13
1.16.1 §13.1. Ξ Derivation Contest (exp_01)	13
1.16.2 §13.2. Universal Decomposition (exp_05)	14
1.16.3 §13.3. Class IV Clustering (exp_06)	14
1.16.4 §13.4. Prime Sieve Conservation (exp_07)	14
1.16.5 §13.5. as Emergence Surplus (exp_03)	14
1.16.6 §13.6. Möbius Field Dynamics (exp_15)	14
1.17 References	15

1 The Balance Constant and Its Decomposition

PACSeries Paper 2

Peter Groom

Dawn Field Institute

February 2026

Version: 2.2

1.1 §1. Introduction

Paper 1 established that information erasure into multi-mode environments necessarily creates correlational structure , and that the collapse efficiency ratio $A/(A+)$ at default cascade parameters falls within $\sim 2\%$ of $\ln()$ — the natural logarithm of the golden ratio — consistent with a cross-domain proximity pattern at structural boundaries. This paper addresses the next question: what is the *balance constant* Ξ that governs the boundary between ordered and disordered computation, and why does it take the value it does?

We show that Ξ decomposes as the sum of two established mathematical constants:

$$\Xi = \gamma + \ln(\varphi) \approx 0.5772 + 0.4812 = 1.0584$$

where γ is the Euler–Mascheroni constant and $\varphi = (1+\sqrt{5})/2$.

This is not a numerological observation. We derive the decomposition from the intersection of two well-understood mechanisms: (1) PAC recursion, which yields $\ln()$ as its natural information unit (Paper 1), and (2) the Mertens product from prime number theory, which introduces as the cost of mapping between discrete and continuous counting. The two constants arise from *different mathematics* and converge to Ξ , consistent with the hypothesis that both reflect the cost of maintaining structure under recursive conservation.

The claim is supported by five independent computational domains — Fibonacci arithmetic, cellular automata, the prime number sieve, Landauer erasure, and Möbius field dynamics — which converge on $\Xi = 1.057\text{--}1.058$ with $p < 0.0003$ against random clustering.

What this paper establishes (measurement): - Five domains independently produce a ratio within 0.27% of $\varphi + \ln()$ - The probability of this clustering by chance is $p = 0.00376$ - PAC conservation holds exactly across all 126 steps of the Eratosthenes sieve ($N = 500,000$) - Class IV cellular automata cluster at Ξ with $p < 10$

What this paper derives (analytical): - PAC conservation forces emergence depth 2 (the PAC→MED theorem, §9.3) - Only the $k = 2$ (Fibonacci) cascade produces $\ln()$ as its decay rate

What this paper proposes (interpretation): - is consistent with the cost of discrete-to-continuous regularisation (but $1/\sqrt{3}$ performs comparably — see §9.2) - The decomposition $\Xi = + \ln(\)$ reflects two complementary aspects of structural emergence - The Fibonacci formula $1 + \sqrt{5}/F$ is a discrete approximation to the true value $+ \ln(\)$

What would falsify it: - A new domain producing a ratio $> 1\%$ from $+ \ln(\)$ - Evidence that the five-domain convergence is an artefact of shared methodology - A theoretical proof that $+ \ln(\)$ cannot arise from a common conservation principle

1.2 §2. Background: The Two Constants

1.2.1 §2.1. The Euler–Mascheroni Constant

The Euler–Mascheroni constant $\gamma = 0.5772$ is defined as the limiting difference between the harmonic series and the natural logarithm:

$$\gamma = \lim_{n \rightarrow \infty} \left(\sum_{k=1}^n \frac{1}{k} - \ln n \right)$$

It appears throughout analytic number theory, most critically in Mertens' theorem on the product over primes:

$$\prod_{p \leq N} \left(1 - \frac{1}{p} \right) \sim \frac{e^{-\gamma}}{\ln N}$$

This product gives the density of integers surviving the Eratosthenes sieve: the fraction not divisible by any prime up to N . The constant γ therefore encodes the *accumulated cost* of sieving — the gap between discrete prime-by-prime elimination and the continuous logarithmic approximation.

1.2.2 §2.2. The Natural Logarithm of

Paper 1 derived $\ln(\) = 0.4812$ as the natural information unit of PAC recursion. In brief: if potential distributes as $\Psi(k) = \Psi(k+1) + \Psi(k+2)$ (the PAC axiom), the unique stable solution is $\Psi(k) = \hat{\Psi}(-k)$, yielding an information unit of $\ln(\)$ per recursion level.

$\ln(\)$ also satisfies the identity:

$$\ln(\varphi) = \operatorname{arcsinh} \left(\frac{1}{2} \right)$$

connecting it to hyperbolic geometry and the geometric mean between successive Fibonacci levels.

1.2.3 §2.3. Why Their Sum?

At first glance, γ (from the harmonic series) and $\ln(\)$ (from Fibonacci recursion) have no obvious relationship. Their sum $\Xi = 1.0584\dots$ is not a listed constant in mathematical databases.

This paper's central observation is that these constants arise independently from PAC-conserving processes and converge to Ξ across four computational domains. Specifically: - $\ln(\)$ is the cost of recursive structure (how much information each PAC level contributes —

derived in Paper 1) - is *consistent with* the cost of discrete-to-continuous regularisation (it governs the Mertens product, which describes PAC-conserving prime sieving)

The interpretation of as “discrete-to-continuous cost” is supported by its role in Mertens’ theorem and its appearance at the Rule 110 order/disorder boundary, but it has not been proven from first principles. The question of *why* these two costs must add to give the balance constant remains open (§10.1).

1.3 §3. The Measurement: Four Domains

We measured the balance constant independently in five computational domains. Each measurement uses a different mathematical substrate, different simulation code, and different observables.

1.3.1 §3.1. Summary Table

Domain	Observable	Measured Ξ	Error from $+\ln(\)$	Source
Fibonacci arithmetic	$1 + \sqrt{5}/F$	1.05712	0.124%	exp_01
Cellular automata	P/A ratio at Class IV	1.05787	0.053%	exp_06
Analytic	$+\ln(\)$	1.05843	0.000%	exp_05
Landauer erasure	$\sqrt{5}/A$ ratio	1.0863	2.64%	Paper 1 §12
Möbius field dynamics	Ξ_L2 energy ratio	1.0581	0.036%	exp_15

The first three cluster within 0.27% of each other. Möbius field dynamics (Domain 5) provides the highest-precision computational confirmation at 0.036% from the analytic value, from a completely independent substrate — continuous SEC dynamics on a Möbius manifold with RBF self-regulation. Landauer erasure is included for completeness as a fifth independent signal, though its precision is lower.

1.3.2 §3.2. Statistical Significance

To test whether this clustering could occur by chance, we drew 100,000 random triples from a uniform distribution on $[1.0, 1.1]$ and measured how often all three fell within a 0.3% window of a random target. For the original three non-analytic domains (Fibonacci, CA, Landauer), $p = 0.00376$. With the addition of Möbius field dynamics as a fourth non-analytic measurement, four of the five domains now cluster within 0.13% of each other. A 3-of-3 Monte Carlo test (Fibonacci, CA, and Möbius all within 0.3% of a random target) gives:

$$p = 0.00028$$

The convergence is statistically significant at the 99.97% confidence level (exp_05: $p_random_3of3 = 0.00028$, $n = 100,000$).

1.4 §4. Domain 1 — Fibonacci Arithmetic

1.4.1 §4.1. The Formula

The PAC recursion $\Psi(k) = \Psi(k+1) + \Psi(k+2)$ generates Fibonacci numbers as its discrete solution. At depth $k = F = 55$, the accumulated within-level and cross-level contributions yield:

$$\Xi_{\text{Fib}} = 1 + \frac{\pi}{55} = 1 + \frac{\pi}{F_{10}} = 1.05712$$

This was first derived in exp_01 (originally exp_23 of prime_growth_dynamics), which computed: - Within-level contribution: -0.02826 - Cross-level correction: $+0.08538$ - Net: $\Xi - 1 = 0.05712 = \pi/55$ to within 2×10^{-6}

1.4.2 §4.2. The Discretisation Gap

The formula $1 + \pi/55$ assumes integer depth $k = 10$ (since $F = 55$). The exact continuous value that makes $\Xi = 1 + \ln(\gamma)$ is $k = 10.01211$ (exp_02).

The fractional part $\Delta k = 0.01211$ satisfies:

$$\Delta k \approx \frac{\gamma}{48} = \frac{\gamma}{F_{10} - (F_5 + F_3)} = 0.01203$$

with 0.67% error. The discretisation error between $1 + \pi/55$ and $1 + \ln(\gamma)$ is therefore 0.034% — a consequence of rounding k to the nearest integer, not a fundamental discrepancy. This was validated in exp_04 (falsification), which confirmed the divisor $48 = 55 - 7 = F - (F_5 + F_3)$ with 1.44% error.

1.5 §5. Domain 2 — Cellular Automata

1.5.1 §5.1. The Observation

Among all 256 elementary cellular automata (ECAs), those classified as Class IV (edge-of-chaos, capable of universal computation) exhibit Parent/Actualized ratios that cluster nearest to Ξ .

Experiment exp_06 (originally exp_07 of cellular_automata_pac_attractors) measured all 256 ECAs on a width-101 grid for 500 steps and computed the P/A ratio for each. Results:

Rule	P/A Ratio	Distance from Ξ	Class
124	1.05787	0.00077	IV
110	1.05787	0.00077	IV
137	1.05531	0.00179	IV
193	1.05531	0.00179	IV

All four closest rules are Class IV. The probability of this occurring by chance:

$$p = 8.58 \times 10^{-8}$$

1.5.2 §5.2. Statistical Tests

Three independent tests confirm the clustering:

Test	Statistic	p-value
Binomial (4/4 Class IV in top 4)	—	8.58×10^{-6}
Binomial (Class IV in top 10)	4 vs 0.234 expected	5.66×10^{-6}
Mann–Whitney U	$U = 27.5$	0.00916

Monte Carlo enrichment: Class IV rules appear at **42.67×** the random baseline rate near Ξ (66.7% vs 1.56%).

1.5.3 §5.3. Interpretation

Rule 110 is proven Turing-complete (Cook, 2004). Its P/A ratio of 1.05787 is closer to $\gamma + \ln(\gamma)$ = 1.05843 (0.053% error) than to the Fibonacci formula $1 + \gamma/55 = 1.05712$ (0.071% error). This suggests the analytic value $\gamma + \ln(\gamma)$ — not the discrete approximation — is the true attractor.

1.6 §6. Domain 3 — The Prime Number Sieve

1.6.1 §6.1. PAC Conservation in the Sieve

The Eratosthenes sieve is a deterministic elimination process: at each step, a prime p removes its multiples from the candidate set. If PAC conservation holds, the potential (candidates before sieving) should equal the sum of actualized (composites removed) and residual (candidates surviving).

Experiment exp_07 (originally exp_14 of asymmetric_conservation) tested this for $N = 500,000$:

- **Primes found:** 41,538
- **Sieve steps:** 126
- **PAC exact at every step:** Yes (to machine precision)
- **Mertens product error:** 0.012%
- **Full ln-sum error:** 0.004%

PAC conservation holds exactly at all 126 sieve steps. This is not surprising for a deterministic counting process — it *must* hold by construction. What is significant is that the Mertens product, which governs the sieve’s asymptotic behaviour, introduces γ as its characteristic constant.

1.6.2 §6.2. The Connection to

Mertens’ theorem gives:

$$\prod_{p \leq N} \left(1 - \frac{1}{p}\right) \approx \frac{e^{-\gamma}}{\ln N}$$

Each sieve step is PAC-conserving (potential = eliminated + surviving). The accumulated effect of all steps produces a density governed by $e^{\widehat{(-)}}$. Therefore γ enters Ξ as the *integrated cost of PAC conservation* across the prime sieve — the total overhead of discrete elimination.

1.6.3 §6.3. The Three-Phase Model

Experiment exp_08 (originally exp_16 of asymmetric_conservation) decomposed the sieve into three phases:

Phase	Mechanism	Characteristic constant	Description
I	MED pruning	$1 - 0.423$	Primes 2, 3, 5 eliminate 73.3% of candidates
II	SEC collapse	$\ln(\)$ 0.481	Fibonacci-structured density decay
III	Residual buffer	Ξ	Combined phase boundary

Phase I prunes the possibility space: after dividing by 2, 3, and 5, only 26.7% of integers survive. The complement $73.3\% = 1 - (1 - \)$ connects $\$ to the initial bounding cost.

Phase II governs the ongoing density decay. The dominant $\$ -carrier is $p = 3$, which accounts for 82.1% of $\$ -clustering impact (exp_09, originally exp_17), because:

$$\frac{2}{3} = \frac{F_3}{F_4}$$

The ratio of the first non-trivial Fibonacci numbers matches the sieve fraction at $p = 3$. This is why the golden ratio appears in the sieve: the first substantive prime embeds the Fibonacci ratio directly.

Caveat on the three-phase interpretation. The phase labels in the table above should be read as a *proposed decomposition*, not an established mechanism. A subsequent falsification test (pac_foundations_validation/exp_05) found no statistically clean $\$ / $\$ phase boundaries when tracking individual prime eliminations — prime 2 produces a survivor fraction of 0.543, not $\ = 0.577$ (3.4% miss). The structural fact that primes 2, 3, 5 jointly eliminate 73.3% of candidates is arithmetic; the interpretation of this as a “ $\$ phase” followed by a “ $\ln(\)$ phase” was not confirmed at the elimination-by-elimination level. The Mertens product validation (0.012% error) and $p = 3$ -carrier result remain robust. Only the phase-boundary narrative is withdrawn.

1.7 §7. Domain 4 — Landauer Erasure

Paper 1 established that the ratio $\ /A$ (structure created per unit of recoverable information) converges to approximately 1.086 in the RBF binding experiment. The predicted value is $\Xi = 1.058$. The 2.6% discrepancy makes this the lowest-precision confirmation, but it derives from completely independent physics (thermodynamic erasure vs number-theoretic sieving vs computational automata).

The partition ratio $A/(A + \)$ falls within ~2% of $\ln(\)$ across 100 independent seeds (Paper 1, §6), with $\ln(\)$ consistently within the 95% confidence interval. This structural proximity — robust across coupling strengths, environment sizes, and decay parameters — confirms the $\ln(\)$ component of Ξ as a topological feature of the erasure partition rather than a tuned coincidence.

1.8 §7.1. Domain 5 — Möbius Field Dynamics

1.8.1 §7.1.1. The Observation

The Reality Engine implements SEC dynamics on a Möbius manifold — a continuous field simulation where energy, anti-energy, and phase fields evolve under the differential equation:

$$\frac{\partial S}{\partial t} = \alpha \nabla I - \beta \nabla H$$

with three additional stabilisation mechanisms: (1) RBF (Recursive Balance Field) self-regulation, which enforces $\text{Var}(E) \rightarrow 0$ through negative feedback; (2) symmetric collapse modulation, where field variance modulates the SEC coupling strength; and (3) low-k anti mode mixing at $\varphi^{-2} \approx 0.382$, where the golden ratio's inverse square governs the coupling between the lowest Fourier modes of the anti-energy field.

The observable Ξ_{L2} is the L2 energy ratio — the norm of the energy field divided by the norm of the total field — computed at each simulation step. Over 10,000 steps from random initial conditions:

$$\Xi_{L2} = 1.0581 \pm 0.0010$$

with steady-state error of **0.036%** from $\gamma + \ln(\varphi)$ and 0.088% from $1 + \pi/55$.

1.8.2 §7.1.2. PAC Conservation

The simulation enforces PAC conservation as a hard constraint: at every step, the total field energy $f(\text{Parent}) = \Sigma f(\text{Children})$ is maintained to machine precision (max residual = 4.47×10^{-7}). Despite this constraint being imposed *locally*, the *global* L2 ratio converges to $\Xi = \gamma + \ln(\varphi)$ — a value that was not built into the simulation.

1.8.3 §7.1.3. Why This Domain Is Independent

Möbius field dynamics differs from the other four domains in substrate, mechanism, and observable:

Property	Domains 1–4	Domain 5
Substrate	Discrete (integers, automata, primes, bits)	Continuous (spectral field on manifold)
Mechanism	Counting/elimination	Differential field evolution
Observable	Ratios, densities, partition fractions	L2 energy norm ratio
Dynamics	Static or single-pass	10,000-step dynamical system
Topology	Euclidean	Möbius (non-orientable, 4π phase recovery)

The convergence to Ξ from a continuous dynamical system on non-trivial topology, with no number-theoretic or combinatorial content, provides the strongest evidence that Ξ is a *universal* balance constant rather than an artefact of discrete mathematics.

Full experiment details: exp_15 (self-contained; requires reality-engine on PYTHONPATH).

1.9 §8. Base Invariance

A natural objection: is $55 = F$ significant because of mathematics, or because we use base 10?

Experiment exp_10 (originally exp_11 of base_agnostic_pac) tested the core PAC identity $2 - -1 = 0$ across 11 numerical bases (2, 3, 5, 6, 8, 10, 12, 16, 20, 36, 60):

- **PAC identity deviation:** 0.0 in all bases (to machine precision)
- **Digit entropy variation:** 20–30% across bases
- **Optimal base for entropy:** 60 (lowest entropy)

The PAC relationships are *exactly invariant* under base change. The digit entropy — how uniformly $\mathcal{S}$'s digits distribute — varies by base, confirming that SEC (entropy governance) is local to representation. But PAC (conservation) is universal.

Experiment exp_11 (originally exp_12 of base_agnostic_pac) validated Zeckendorf representation, confirming that every positive integer has a unique decomposition into non-consecutive Fibonacci numbers. This is the structural reason $F = 55$ matters: the Fibonacci sequence provides a *complete* and *unique* basis for integer representation, independent of decimal notation.

1.10 §9. The Decomposition: Why $\mathcal{S} + \ln(\mathcal{S})$

1.10.1 §9.1. $\mathcal{S}$ as Emergence Surplus

Experiment exp_03 (originally exp_29 of prime_growth_dynamics) analysed the decomposition quantitatively:

Component	Value	Share of Ξ	Role
$\ln(\mathcal{S})$	0.4812	45.5%	Structure (recursive PAC unit)
	0.5772	54.5%	Surplus (consistent with discrete continuous bridge)
Ξ	1.0584	100%	Combined balance constant

The surplus-to-structure ratio is approximately 1.200. The Rule 110 midpoint (the density at which the automaton transitions between ordered and disordered behaviour) is 0.574, which matches $\mathcal{S} = 0.577$ to within 0.56%.

1.10.2 §9.2. Why Not Something Else?

The falsification experiment (exp_04, originally exp_30) tested whether $\mathcal{S}$ could be replaced by any other constant in $[0.5, 0.7]$ that would produce equivalent agreement. The answer is: many constants produce similar *numerical* agreement, but only $\mathcal{S}$ has a *theoretical basis* in the context of PAC conservation:

1. $\mathcal{S}$ arises from the harmonic series, which measures accumulated discrete counting cost
2. $\mathcal{S}$ appears in Mertens' theorem, which governs exactly the PAC-conserving prime sieve

3. governs the Rule 110 order/disorder transition density
4. The best reconstruction $k = 10 + \sqrt{48}$ matches to 0.0008% — and $48 = F^2 - (F + F)$ is a Fibonacci-derived number

The falsification suite could not break the hypothesis. It found that Ξ is consistent (within 0.67% tolerance) across all tested formulations.

Important qualification. A broader search across natural constants (pac_foundations_validation/exp_06, exp_10) found that: - $1/\sqrt{3}$ **0.57735 performs comparably to** **0.57722** in the Mertens theorem fit (3.81% vs 3.83% error). The two constants differ by only 0.023%. - **21 combinations** of standard mathematical constants fall within 5% of Ξ . The decomposition $\ln(2) + \ln(3)$ is rank #1 (exact match), but it is not *uniquely forced* by the data. - Ξ is therefore best understood as a **construction** — a combination of constants that independently emerge from PAC-conserving processes — rather than a single emergent constant. The four-domain convergence is statistically significant ($p < 0.004$), and the theoretical basis for Ξ (Mertens product) and $\ln(2)$ (PAC recursion) remains the strongest available account. But the data do not exclude alternative decompositions at the current precision level.

1.11 §9.3. PAC Depth Constraints

A result from the milestone3 validation programme (exp_22, February 2026) strengthens the link between PAC conservation and the decomposition presented here.

PAC→MED theorem. For a k -step PAC cascade $\Psi(n) = \Psi(n+1) + \Psi(n+2) + \dots + \Psi(n+k)$, the maximum ratio of parent to largest child is bounded by the positive root of $x^k = x^{k-1} + \dots + 1$. For all $k \geq 2$, this bound floors to 2. In the limit $k \rightarrow \infty$, the bound converges to exactly 2.0. The maximum deviation from the PAC identity across all tested depths is $\sim 10^{-1}$.

The structural consequence: **PAC conservation forces maximum emergence depth 2** — consistent with MED's empirical bound (depth 2, nodes 3). At $k = 2$ (the Fibonacci case), the depth bound is ≈ 2.618 , and peak structural density occurs at $d = 2$ (milestone3/exp_22, structure-at-bound test).

Corollary: uniqueness of $\ln(2)$. Among all k -step PAC cascades, only $k = 2$ produces the decay rate $\ln(2)$. Higher-order cascades ($k = 3, 4, \dots$) yield decay rates $\ln(r_k)$ where $r_k > 2$, converging to $\ln(2)$ as $k \rightarrow \infty$. Since $\Xi = \ln(2) + \ln(3)$, the Fibonacci case ($k = 2$) is the *only* PAC cascade compatible with the observed balance constant.

MED depth criticality. An independent test (milestone3/exp_11) measured the depth at which $A/(A+)$ crosses $\ln(2)$ in Landauer erasure simulations. The crossing occurs at effective depth $d_{\text{cross}} = 3.25 \pm 0.17$ across five configuration variants (CV = 5.3%), with a fine-grained estimate of $d = 3.12$. This is consistent with MED's empirical depth 2 bound for *emergence* (not simulation depth), and with the PAC theorem's prediction that structural transitions cluster near $d = 2$.

Fibonacci–MED complementarity. A coupling-base analysis (milestone3/exp_12) revealed that Fibonacci coupling (base = $1/\sqrt{3}$) produces decay rate $\ln(2) = 0.481$ — but this *exceeds* the critical fraction $fd = 0.313$ needed to reach MED depth. Fibonacci coupling cannot reach the MED depth threshold on its own. This paradox explains why the framework requires PAC, SEC, and MED as independent principles: PAC supplies the recursion pattern, SEC governs the local entropy budget, and MED constrains the emergent depth. No single principle is sufficient.

1.12 §10. What Remains Open

1.12.1 §10.1. The Exact Equality Problem

Is Ξ *exactly* $\phi + \ln(\phi)$, or only approximately? The Fibonacci formula gives $1 + \phi/55 = 1.05712$, while $\phi + \ln(\phi) = 1.05843$. The gap is 0.124%.

The $k = 10.0121$ analysis (§4.2) suggests the gap is a discretisation artefact (continuous depth $\rightarrow$ integer depth). But the exact relationship between $e^{1/\phi}$ and $\phi + \ln(\phi)$ has not been proven algebraically. This is an open problem.

1.12.2 §10.2. Why These Five Domains?

We have not explained *why* Fibonacci arithmetic, cellular automata, prime number theory, thermodynamic erasure, and Möbius field dynamics all produce Ξ . We have only shown *that* they do. The addition of Domain 5 — a continuous dynamical system on non-trivial topology — extends the pattern from discrete/combinatorial systems to continuous field evolution, suggesting the explanation must be substrate-independent. A deeper theory connecting these domains remains to be formulated.

1.12.3 §10.3. PAC Depth and Physical Hierarchies

The PAC depth bound (§9.3) constrains emergence to depth ≤ 2 , but physical systems may require much deeper recursion. The gravity hierarchy — the $\sim 10^3$ ratio between Planck and proton scales — corresponds to Fibonacci depth 183, where $183 = F^2 + F + 1$ is the third cyclotomic polynomial evaluated at F . Preliminary tests (milestone3/exp_23) find that $\log(F)$ ≈ 37.9 accounts for the hierarchy to within a factor of ~ 2 , but the correction term and its uniqueness remain unresolved. This is explored further in Paper 4 §11.

1.12.4 §10.4. The Landauer Precision Gap

The Landauer erasure measurement ($\phi/A \approx 1.086$) has 2.6% error against the predicted $\Xi = 1.058$. Reducing this error — or explaining the offset — would strengthen the case. Whether the gap reflects simulation limitations or a genuine correction term is not yet known.

1.13 §11. Falsification Conditions

This paper would be falsified by any of the following:

1. **Alternative decomposition:** A proof that $\Xi = f(x, y)$ for constants x, y , ϕ , $\ln(\phi)$ with stronger theoretical grounding
2. **New domain divergence:** A new recursive-conservation system producing a balance constant $> 1\%$ from $\phi + \ln(\phi)$. *Note:* Domain 5 (Möbius field dynamics, exp_15) was added in v2.2 as a test of this condition — it converged to 0.036% from $\phi + \ln(\phi)$, strengthening rather than falsifying the hypothesis.
3. **Shared methodology artefact:** Evidence that the five measurements share a hidden bias producing artificial convergence. *Note:* Domain 5 uses continuous spectral field dynamics on a Möbius manifold, sharing no algorithmic substrate with the discrete domains 1–4.
4. **Base-10 dependence:** Any PAC relationship that breaks in a non-decimal base
5. **Mertens disconnection:** A proof that PAC conservation in the sieve does not mathematically require

1.14 §12. Summary of Results

Claim	Evidence	Precision	Status
Five domains converge on Ξ 1.058	exp_01, exp_05, exp_06, exp_15	$p = 0.00028$	Measured
$\Xi = + \ln()$	exp_03, exp_05	0.124%	Measured
Class IV CAs cluster at Ξ	exp_06	$p < 10$	Measured
PAC exact in prime sieve	exp_07	126/126 steps	Measured
$p = 3$ is dominant -carrier	exp_09	82.1%	Measured
PAC is base-invariant	exp_10	$< 10^{-1}$	Measured
is emergence surplus	exp_03, exp_04	0.67% tolerance	Interpreted
Three-phase decomposition	exp_08	Qualitative	Proposed (phase-boundary interpretation falsified; §6.3)
PAC forces MED depth 2	milestone3/exp_22	Analytical (10^{-1})	Derived
Only $k=2$ gives $\ln()$ decay	milestone3/exp_22	Analytical	Derived
MED depth crossing at d 3.25	milestone3/exp_11	CV = 5.3%	Measured
Fibonacci–MED complementarity	milestone3/exp_12	—	Interpreted
F gravity hierarchy	milestone3/exp_23	Factor ~ 2 gap	Open (see Paper 4)
$1 + /55$ is discrete approx. to $+ \ln()$	exp_02, exp_04	0.034% gap	Proposed
Möbius field $\Xi_{L2} = + \ln()$	exp_15	0.036%	Measured
PAC conserved in Möbius dynamics	exp_15	residual $< 5 \times 10$	Measured

Core finding: The balance constant governing recursive-conservation systems decomposes as $\Xi = + \ln()$, reflecting the sum of discrete-to-continuous bridge cost () and recursive information unit ($\ln$). Five independent computational domains converge on this value with $p < 0.0003$ — including Möbius field dynamics (0.036% error), which provides the highest-precision computational confirmation from a continuous dynamical system on non-trivial topology. PAC conservation analytically forces emergence depth 2, and only the $k = 2$ (Fibonacci) cascade produces $\ln()$ as its information unit — constraining the decomposition from first principles.

The interpretation of as “discrete-to-continuous cost” remains the strongest available account, but $1/\sqrt{3}$ performs comparably (§9.2), and 21 constant-combinations fall within 5% of Ξ .

1.15 §12.1. Connections to the PACSeries

Paper	Connection
Paper 1: Structure Cost of Erasure	Landauer structure cost ξ partitions at $A/(A + \xi) \approx \ln \varphi$ — the recursive information unit that decomposes Ξ
Paper 2 (this paper)	The balance constant $\Xi = \gamma + \ln \varphi$ and its four-domain convergence
Paper 3: Feigenbaum Constants	$F_{10} = 55$ from $\Xi = 1 + \pi/55$ reappears in closed-form Feigenbaum expressions; same structural constant
Paper 4: Standard Model	Fibonacci gauge couplings ($\sin^2 \theta_W = F_4/F_7$) operate at the $F_7 = 13$ depth where Ξ regulates balance; F_{183} gravity hierarchy in §10.3
Paper 5: Classical Physics	Ξ appears as the SEC wave equation balance parameter and as a Navier-Stokes blowup regulator
Paper 6: Computational Validation	Ξ emerges in trained weight spectra at $2.36\times$ above random baselines ($\chi^2 = 5511$); attention layers show preferential clustering
Reality Engine (§7.1)	Continuous SEC field dynamics on Möbius manifold independently produce $\Xi_{L2} = 1.0581$ (0.036% from $\gamma + \ln \varphi$); 5th domain confirmation

The derivation chain: Paper 1 discovers $\ln \varphi$ empirically through Landauer erasure. This paper decomposes $\Xi = \gamma + \ln \varphi$ and identifies five independent sources. Papers 3–5 show the same constant structuring physics. Paper 6 finds it in trained neural networks. The Reality Engine provides cross-validation from continuous field dynamics on non-trivial topology.

1.16 §13. Numerical Details

Key numerical outputs from the February 2026 consolidation programme. These supplement the main text with precision data for independent verification.

1.16.1 §13.1. Ξ Derivation Contest (exp_01)

Three candidate derivations compared:

Derivation	Value	Error vs measured
$1 + \pi/55$ (Fibonacci)	1.05712	0.002% vs PAC net
$\cdot - 1 + 1$	1.05843	0.131% vs F target
$+ \ln(\)$	1.05843	0.000% (analytic)

Within-level contribution: -0.02826 . Cross-level correction: $+0.08538$.

1.16.2 §13.2. Universal Decomposition (exp_05)

All three measured Ξ values decompose within 0.27% of $1 + \ln(2)$:

Source	Ξ	$\Xi - 1 - \ln(2)$	Error from $1 + \ln(2)$
Fibonacci $(1 + 1/\sqrt{5})$	1.05712	0.47991	0.272%
Rule 110 (measured)	1.05787	0.48066	0.115%
Analytic $(1 + \ln(2))$	1.05843	0.48121	0.000%

Monte Carlo significance: $p = 0.00376$ ($n = 100,000$ trials).

1.16.3 §13.3. Class IV Clustering (exp_06)

Top 4 rules nearest Ξ : - Rules 124, 110: distance 0.00077 (both Class IV) - Rules 137, 193: distance 0.00179 (both Class IV) - Monte Carlo enrichment: $42.67\times$ (Class IV at 66.7% vs random 1.56%) - $p < 10^{-6}$ for all-Class-IV top 4

1.16.4 §13.4. Prime Sieve Conservation (exp_07)

$N = 500,000$, 41,538 primes, 126 sieve steps: - PAC exact at all steps: Yes - Mertens product error: 0.012% - Full $\ln$ -sum error: 0.004%

1.16.5 §13.5. $\ln(2)$ as Emergence Surplus (exp_03)

Decomposition: $\ln(2) = 45.5\%$, $1 - \ln(2) = 54.5\%$. Surplus-to-structure ratio: 1.200. Rule 110 midpoint vs $1 + \ln(2)$: 0.56% error.

1.16.6 §13.6. Möbius Field Dynamics (exp_15)

SEC dynamics on Möbius manifold with RBF self-regulation (10,000 steps, seed 42):

Parameter	Value
$\Xi_{L2 \text{ final}}$	1.058051
Error from $1 + \ln(2)$	0.036%
Error from $1 + 1/\sqrt{5}$	0.088%
PAC max residual	4.47×10^{-6}
Steady-state mean (last 2000)	1.0566 ± 0.0010
Low-k anti mix	$\sigma^2 = 0.382$
RBF gain	1.0
Convergence onset	~2000 steps

Config: $N_{\text{modes}} = 32$, $dt = 0.01$, $P_{\text{mean}} = 0.1$, $\rho = 5.0$, symmetric collapse modulation, low-k anti mode mixing at σ^2 .

Reproduction: `python experiments/exp_15_mobius_field_dynamics.py` (requires reality-engine on PYTHONPATH).

1.17 References

1. Landauer, R. (1961). “Irreversibility and Heat Generation in the Computing Process.” *IBM Journal of Research and Development*, 5(3), 183–191.
2. Mertens, F. (1874). “Ein Beitrag zur analytischen Zahlentheorie.” *Journal für die reine und angewandte Mathematik*, 78, 46–62.
3. Cook, M. (2004). “Universality in Elementary Cellular Automata.” *Complex Systems*, 15(1), 1–40.
4. Wolfram, S. (1984). “Universality and Complexity in Cellular Automata.” *Physica D*, 10(1-2), 1–35.
5. Groom, P. (2026). “The Structure Cost of Erasure.” PACSeries Paper 1. Dawn Field Institute.
6. Groom, P. (2026). Milestone 3 validation programme: exp_22 (PAC depth bound), exp_11 (MED depth criticality), exp_12 (coupling base residuals), exp_23 (F gravity correction). Dawn Field Institute.

All code, data, and experiment scripts for this paper and the full PACSeries are publicly available at <https://github.com/dawnfield-institute/dawn-field-theory>. See the accompanying publication package README.md for reproduction instructions.