

Math & Theoretical Physics Notes

Dawoud Roger

—

E-mail: dawoudroger@gmail.com

ABSTRACT: In an attempt to better remember and learn physics and scientific english, this collection of personal notes aimed at my future self was created, covering theoretical physics and the math I will be using to learn it.

It will be a set of notes divided in two parts, in the first part I will take notes of subjects studied this past academic year on math methods and in the second I'll take notes of self-studied more fundamental subjects.

Contents

I	Part I : Mathematical methods	3
1.	Differential equations	3
α .	Bernoulli's differential equation	3
β .	Characteristic's equation	3
γ .	Wronskian	4
δ .	Laplace Problem	5
ϵ .	Green's functions	5
2.	Complex analysis	7
α .	Generalities	8
β .	Integration	9
II	Group Theory and representations	11
1.	On Groups	11
2.	On Representations	13
3.	SU(2)	13
4.	SU(3)	13
III	Differential geometry	14
1.	Topology	14

I Part I : Mathematical methods

1. Differential equations

Here I display my favourite equations and resolution methods, more can be found at [1].

α . Bernoulli's differential equation

Let us consider a Bernoulli differential equation defined as

$$\frac{d}{dx}y(x) = a(x)y(x) + b(x)y^n(x), \text{ where } n \in \mathbb{R}$$

rewritten as :

$$y^{-n}(x) \frac{d}{dx}y(x) = a(x)y^{1-n}(x) + b(x)$$

The resolution of such an equation begins by the substitution of the function $y(x)$ by a quite trivial function

$$u(x) = y^{1-n}(x)$$

such a substitution leads to a simpler equation which we know how to solve

$$\frac{1}{1-n} \frac{d}{dx}u(x) = a(x)u(x) + b(x)$$

$$\frac{d}{dx}u(x) = p(x)u(x) + q(x)$$

β . Characteristic's equation

Let us consider the following equation :

$$a(x, y, z) \frac{\partial(x, y, z)}{\partial x} f + b(x, y, z) \frac{\partial(x, y, z)}{\partial y} f + c(x, y, z) \frac{\partial(x, y, z)}{\partial z} f = 0$$

We can define the equation above as the derivative of $f(x, y, z)$ by a variable along which it is constant

$$\frac{df}{ds} = a(x, y, z) \frac{\partial(x, y, z)}{\partial x} f + b(x, y, z) \frac{\partial(x, y, z)}{\partial y} f + c(x, y, z) \frac{\partial(x, y, z)}{\partial z} f = 0$$

Leading to the so-simple equations

$$a(x, y, z) = \frac{dx}{ds} \text{ and } b(x, y) = \frac{dy}{ds} \text{ and } c(x, y) = \frac{dz}{ds}$$

Naturally the next to be taken to determine $s(x, y, z)$ is to isolate $ds(x, y, z)$

$$ds(x, y, z) = \frac{dx}{a(x, y, z)} = \frac{dy}{b(x, y, z)} = \frac{dz}{c(x, y, z)}$$

By isolating the three RHS¹ terms 2 by 2 (one of the combinations may be omitted as it carries redundant information), we find

$$\frac{dy}{dx} = \frac{b(x, y, z)}{a(x, y, z)} \text{ and } \frac{dy}{dz} = \frac{b(x, y, z)}{c(x, y, z)}$$

and the solutions to these differential equations are none other than two families of characteristic lines along which $f(x, y)$ is conserved. These sets of solutions are sets that depend on the integration constants that shall be determined using boundary conditions².

Finally³ :

$$f(x, y, z) = g(\Psi(x, y, z), \Phi(x, y, z))$$

γ. Wronskian

When studying a physical problem, it is interesting to find the family of functions that are solution to a category of Differential equation. The use of the Wronskian allows to determine (starting from $n-1$ sets of solutions) the last set of solutions to an n -th order linear homogeneous differential equation, not any set of solutions but a linearly independent family that would complete the $n-1$ collection of families leading to us having the "basis" of solutions to n order linear homogeneous differential equations. Let us consider this equation

$$f^{(n)}(x) + p_{n-1}(x)f^{(n-1)}(x) + \dots + p_1(x)f'(x) + p_0(x)f(x) = 0$$

The Wronskian is the following determinant

$$W(f_1, f_2, \dots, f_n)(x) = \begin{vmatrix} f_1(x) & f_2(x) & \dots & f_n(x) \\ f_1'(x) & f_2'(x) & \dots & f_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ f_1^{(n-1)}(x) & f_2^{(n-1)}(x) & \dots & f_n^{(n-1)}(x) \end{vmatrix}$$

According to Abel's⁴ identity

$$\frac{dW}{dx} = -p_{n-1}(x)W(x) \rightarrow W(x) = C \exp\left(-\int p_{n-1}(x) \frac{d}{dx}\right)$$

¹Right Hand Side.

²Dirichlet boundary conditions specify conditions on the solution itself, whereas Neumann boundary conditions specify conditions on the derivatives of the solution, the combinations of these conditions are diverse and may be found on https://en.wikipedia.org/wiki/Mixed_boundary_condition.

³this method can be applied to an N -variable function but I only explained the three variable one as the difference between each N -dimensional implementation of it is that we only generate $N-1$ equations thus $N-1$ characteristic sets of lines.

⁴what a Man !

By developing the determinant on the last column we obtain :

$$\sum_{k=0}^{n-1} M_k(x) f_n^{(k)}(x) = C \exp \left(- \int p_{n-1}(x) \frac{d}{dx} \right)$$

Which should in theory be solvable.

δ. Laplace Problem

Laplace's problem⁵ is famous, elegant, useful and omnipresent in the physics I've encountered up until now. Be it in Electromagnetism, elementary Gravity problems, and of course in the use of Green's functions.

$$\Delta f = 0$$

the 1D problem is trivial, so are the 2D and 3D ones to be fair, yet ... here we go :

1-dimensional problem

$$\frac{\partial^2 f}{\partial x^2}(x) = 0 \implies f(x) = Ax + B, \text{ where } A, B \in \mathbb{R}$$

2-dimensional problem

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) f(x, y) = 0 \iff \frac{\partial^2(x, y)}{\partial x^2} f = - \frac{\partial^2(x, y)}{\partial y^2} f$$

and as we can, we will decompose $f(x, y) = X(x)Y(y)$

$$\frac{\frac{\partial^2 X}{\partial x^2}(x)}{X(x)} = - \frac{\frac{\partial^2 Y}{\partial y^2}(y)}{Y(y)}$$

and as both the LHS and the RHS are linearly independent because of their different variables, we can therefore guess that they are equal to a constant K , whose real or imaginary nature will influence the results we get.

3-dimensional problem

The 3-dimensional version of the above problems are almost exactly the same, only the number of solutions obtained varies. The idea of the laplace problem resolution is centered around variable separation and if needed more than one arbitrary constant may be used to dully seperate the variable.

ε. Green's functions

Green's method is a way to solve a problem through the definition of what is called a Green function $G(\vec{r} - \vec{\tilde{r}})$ which holds a significant physical relevance. The first step consists of the rewritting of our n -dimensional physical equation as integrals

$$\Delta \phi(\vec{r} - \vec{\tilde{r}}) = k \rho(\vec{r} - \vec{\tilde{r}}) \longrightarrow \int G(\vec{r} - \vec{\tilde{r}}) \rho(\vec{\tilde{r}}) d\vec{\tilde{r}} = k \int \delta^n(\vec{r} - \vec{\tilde{r}}) \rho(\vec{\tilde{r}}) d\vec{\tilde{r}}$$

⁵For further information and/or details check out the *GOAT*'s lecture notes : Wietze HERREMAN.

This leads to the much simpler equation

$$\Delta G(\vec{r} - \tilde{\vec{r}}) = k\delta^n(\vec{r} - \tilde{\vec{r}})$$

It is interesting to notice that if for example we take ϕ a gravitational potential for a punctual star of mass M located at the origin of our frame, we therefore have⁶ $k = 4\pi GM$ ⁷ and $\rho(\vec{r}) = \delta^3(\vec{r})$.

1-dimensional problem

A one dimensionnal Green function problem gets simplified to an equation of the form :

$$\frac{\partial^2 G}{\partial r^2}(r - \tilde{r}) = \delta(r - \tilde{r})$$

We first solve the previous equation for $x \neq 0$

$$G_{\pm}|_{r \neq \tilde{r}} = C_{\pm}(r - \tilde{r}) + B_{\pm}, \text{ with : } C_{\pm}, B_{\pm} \in \mathbb{R}$$

Boundary conditions allow us to determine either (or both) integration constant(s), and another way of determining the hypothetical undertermined integration constant is through the integration over the parameter of the dirac delta.

$$\int_{\gamma} \frac{\partial^2 G}{\partial r^2}(r - \tilde{r})dr = \int_{\gamma} \delta(r - \tilde{r})dr, \text{ where } \gamma : r \in [-\varepsilon; \varepsilon]$$

$$\frac{\partial G}{\partial r}(\varepsilon) - \frac{\partial G}{\partial r}(-\varepsilon) = C_+ - C_- = 1$$

and using further conditions and information on our physical system it is possible to determine the constants $C_{\pm}$ and $B_{\pm}$. This cas is very specific as the absence of a solution due to an "infinite density" at $r = \tilde{r}$ makes the r axis disconnected, and so there is no reason to make the terms at $\tilde{r}^-$ and $\tilde{r}^+$ be the same.

2-dimensional problem

A two dimensionnal Green function problem⁸ gets simplified to an equation of the form :

$$\Delta G(\vec{r}) = r^{-1} \frac{\partial}{\partial r} \left(r \frac{\partial G}{\partial r} \right) = k\delta^2(\vec{r})$$

Solving this problem as a Laplace problem first

$$G = C \ln(|\vec{r} - \tilde{\vec{r}}|) + B$$

then integrating over a constant r surface (a circle)

$$\int \Delta G d^2 S = \int k\delta^2(\vec{r} - \tilde{\vec{r}}) d^2 S = k$$

⁶the G used in k stands for the universal gravitational constant.

⁷in the case of other physical theories such as Electricity or Magnetism, the constants k differ (q/ε_0 in case of the scalar electric potential).

⁸in cylindrical coordinates.

$$k = \oint R d\theta \left[\vec{\nabla} G|_{r=R} \cdot \vec{e}_r \right] = 2\pi C$$

$$C = \frac{k}{2\pi}$$

Finally using the boundary conditions as $r \rightarrow \infty$ we determine the value of B

$$G(\vec{r} - \vec{\tilde{r}}) = B + k \frac{\ln |\vec{r} - \vec{\tilde{r}}|}{2\pi}$$

3-dimensional problem

A three dimensional Green problem simplifies to

$$\Delta G(\vec{r} - \vec{\tilde{r}}) = k \delta^3(\vec{r} - \vec{\tilde{r}})$$

once again, let us solve the homogeneous equation where $\vec{r} \neq \vec{\tilde{r}}$:

$$r^{-2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial G}{\partial r} \right) = 0 \implies G(\vec{r} - \vec{\tilde{r}}) = -\frac{C}{\|\vec{r} - \vec{\tilde{r}}\|} + B$$

We input this in the volume integral

$$\int_V \Delta G dV = \oint_{\partial V} \vec{\nabla} G \cdot d\vec{S}$$

$$C = \frac{k}{4\pi}$$

whereas B is determined by our G value at $\|\vec{r} - \vec{\tilde{r}}\| \rightarrow \infty$

$$G(\vec{r} - \vec{\tilde{r}}) = -\frac{k}{4\pi \|\vec{r} - \vec{\tilde{r}}\|} + B$$

Applying these results to our previous example regarding the star of mass M (in [subsection 6](#)), we get

$$\phi(\vec{r}) = - \int \left[\frac{GM}{\|\vec{r} - \vec{\tilde{r}}\|} - B \right] \delta^3(\vec{\tilde{r}}) d\vec{\tilde{r}}$$

leading to the well-known result⁹:

$$\phi(\vec{r}) = -\frac{GM}{\|\vec{r}\|} + B$$

2. Complex analysis

Complex analysis¹⁰ is the study of functions of complex variables. To put it simply we will focus on functions such as $f(z) \in \mathbb{C}$ with $z \in \mathbb{C}$.

⁹still for a punctual mass M positionned at the center of the frame.

¹⁰[\[2\]](#) is a good ressource to study Complex Analysis.

α . Generalities

Some general notations commonly used

$$f(z) = f(x, y) = u(x, y) + \mathbf{i}v(x, y), \text{ where } : u, v \in \mathbb{R}$$

$f(z)$ is a **holomorphic** function in Ω an open subset of $\mathbb{C}$ if the equation above holds $\forall z \in \Omega$.

It is important when doing analysis to have good definitions of derivation. It just so happens that for a matter of consistency we define it the same way we did in real analysis (we simply replace x by z), only we add the condition that the usual (following) equality must hold no matter the path taken

$$f'(z_0) = \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} = \frac{df(z)}{dz}$$

Therefore if we take a path along either x or iy , we must have (for functions we will call **analytical**) equality of the derivatives along both paths :

$$\begin{aligned} f'(z_0) &= \lim_{x \rightarrow x_0} \frac{u(x, y_0) - u(x_0, y_0)}{x - x_0} + \mathbf{i} \lim_{x \rightarrow x_0} \frac{v(x, y_0) - v(x_0, y_0)}{x - x_0} = \frac{\partial u(x, y)}{\partial x} + \mathbf{i} \frac{\partial v(x, y)}{\partial x} \\ f'(z_0) &= \lim_{y \rightarrow y_0} \frac{u(x_0, y) - u(x_0, y_0)}{\mathbf{i}y - \mathbf{i}y_0} + \mathbf{i} \lim_{y \rightarrow y_0} \frac{v(x_0, y) - v(x_0, y_0)}{\mathbf{i}y - \mathbf{i}y_0} = -\mathbf{i} \frac{\partial u(x, y)}{\partial y} + \frac{\partial v(x, y)}{\partial y} \end{aligned}$$

Leading to the well-known Cauchy-Riemann (CR) conditions of analyticity for a complex function

$$\boxed{\frac{\partial u(x, y)}{\partial x} = \frac{\partial v(x, y)}{\partial y} \quad \& \quad \frac{\partial u(x, y)}{\partial y} = -\frac{\partial v(x, y)}{\partial x}} \quad (2..1)$$

We can define the derivatives

$$\partial_z = \frac{1}{2} \left(\frac{\partial}{\partial x} - \mathbf{i} \frac{\partial}{\partial y} \right) \quad \text{and} \quad \partial_{\bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x} + \mathbf{i} \frac{\partial}{\partial y} \right)$$

leading to the re-writing of the CR conditions as

$$\frac{\partial f}{\partial \bar{z}} = 0$$

We call **analytical** function any function that can be decomposed into a Laurent series

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n = \sum_{n=-\infty}^{-1} a_n (z - z_0)^n + \sum_{n=0}^{\infty} a_n (z - z_0)^n$$

we will refer to the LH term the singular part of $f(z)$ and the RH term the analytical part of $f(z)$.

A **meromorphic** function is one such that the derivative is defined on $(\Omega \setminus \{z_i\}_{i \in \mathbb{Z}}) \subset \mathbb{C}$, implying it isn't derivable on $\{z_i\}_{i \in \mathbb{Z}}$ and we call this set of points **singular points** (basically singularities). There are three kinds of singularities :

- **n -th order pole**, which is characterized by $f(z) \propto (z - z_0)^{-n}$
- **Essential** singularities, characterized by infinite terms proportional to $(z - z_0)^{-1}$, and therefore a non vanishing singular part of the Laurent series decomposition of the function.
- **Apparent** singularities that aren't actually singularities.

β . Integration

A few results that are very practical to physicists. I omitted a few to speed my writing up and get to more interesting and relevant subjects faster but I will come back to this section.

Cauchy's Theorem : When integrating a holomorphic function over a closed contour, it is expected because of the Cauchy theorem for its value to be null.

$$\oint_{\Gamma} dz f(z) = 0$$

The holomorphy condition means the absence of singularities inside the contour as no singularities of $f(z)$ exist on Ω anyways. The contour containing no singularities is therefore called homotopic to 0.

Cauchy's generalized Theorem : When integrating a meromorphic function over a closed contour homotopic to zero, it is expected because of Cauchy's generalized theorem for its value to be :

$$\oint_{\Gamma} dz f(z) = \sum_i \oint_{\Gamma_i} dz f(z)$$

where each Γ_i is a contour around one of the singularities contained in Γ .

Jordan's lemma : Let us consider a circular contour $\mathcal{C}_{\mathcal{R}}$ (not necessarily closed) then the following equation holds

$$\begin{aligned} \lim_{R \rightarrow 0} \sup_{z \in \mathcal{C}_{\mathcal{R}}} |zf(z)| = 0 &\implies \lim_{R \rightarrow 0} \int_{\mathcal{C}_{\mathcal{R}}} dz f(z) = 0 \\ \lim_{R \rightarrow \infty} \sup_{z \in \mathcal{C}_{\mathcal{R}}} |zf(z)| = 0 &\implies \lim_{R \rightarrow \infty} \int_{\mathcal{C}_{\mathcal{R}}} dz f(z) = 0 \end{aligned}$$

This lemma is very practical for the determination of integrals using the upcoming *Residue Theorem*.

Residue Theorem : Let $f(z)$ be a meromorphic function on $\Omega \subset \mathbb{C}$, let Γ be a closed contour surrounding N singularities of $f(z)$, and Γ doesn't cut through any singularities of $f(z)$. The integral

$$\oint_{\Gamma > 0} dz f(z) = 2i\pi \sum_{i=1}^N \text{Res}(f(z), z_i)$$

where $\text{Res}(f(z), z_i) = a_{-1}$, the term proportional to $(z - z_i)^{-1}$. For n -th order poles, there is another residue formula :

$$\text{Res}(f(z), z_i) = \lim_{z \rightarrow z_i} \frac{1}{(n-1)!} \frac{d^{n-1}}{dz^{n-1}} [(z - z_i)^n f(z)]$$

Lebesgue's dominated convergence theorem : Let $f_n(z)$ be a sequence of measurable functions, assuming it converges pointwise (simplement) almost everywhere to a function $f(z)$ on a measurable set A . If there exists a function $g(z)$ that is integrable (sommable) over A such that $|f_n(z)| \leq g(z) \quad \forall n \in \mathbb{N}$ almost everywhere, then :

$$\lim_{n \rightarrow \infty} \int_A dz f_n(z) = \int_A dz f(z), \text{ where } f(z) = \lim_{n \rightarrow \infty} f_n(z)$$

II Group Theory and representations

This section is heavily inspired from ([3],[4], [5] and [6]) as they are my self-study material¹.

1. On Groups

Group : A group $(\mathcal{G}, \circ)$ is a set $\mathcal{G}$ equipped with an internal composition law " $\circ$ ", that respects axioms of :

i) Associativity : $ab \circ c = a \circ bc$

ii) Existence of a neutral element : $a \circ e = e \circ a = a$

iii) Existence of an inverse element : $a \circ a^{-1} = e$

where obviously $a, b, c, a^{-1}, e \in \mathcal{G}$.

Notice that commutativity isn't a requirement and that if a group has a commutative operation it is said to be **abelian**.

Topological Group : A topological group is a topological space that is also a group, such that the group operation $\circ$ is continuous, same goes for the inversion mapping

$$\begin{aligned} \mathcal{G} \times \mathcal{G} &\rightarrow \mathcal{G} \\ (a, b) &\mapsto a \circ b \end{aligned}$$

$$\begin{aligned} \mathcal{G} &\rightarrow \mathcal{G} \\ a &\mapsto a^{-1} \end{aligned}$$

Homomorphism : A homomorphism φ is a mapping from a group $(\mathcal{G}, \circ)$ to a group $(\mathcal{H}, \star)$, such that

$$\varphi(a \circ b) = \varphi(a) \star \varphi(b), \quad \forall a, b \in \mathcal{G}$$

If said homomorphism is bijective, then it is also an **isomorphism**.

If $\mathcal{H} = \mathcal{G}$ and φ is bijective, it is then said to be an **automorphism**.

An **inner automorphism** is a mapping of the form $g' \rightarrow gg'g^{-1}$.

Center : The center of a group $(\mathcal{G}, \circ)$ is the set of elements that commute with all other elements of $(\mathcal{G}, \circ)$:

$$Z_{\mathcal{G}} = \{g \in \mathcal{G} | g \circ a = a \circ g, \forall a \in \mathcal{G}\}$$

¹[5, 6] are especially good respectively for problems to solve and physics examples.

Starting now I will often be simplifying $(\mathcal{G}, \circ)$ to "the group $\mathcal{G}$ ".

Subgroup : A subgroup $\mathcal{H}$ of $\mathcal{G}$, is a subset of $\mathcal{G}$ stable under the internal composition law defined on $\mathcal{G}$. Simply put, it is a subset of $\mathcal{G}$ that is also a group under the same law.

Cosets : Let us consider $\mathcal{H}$ a subgroup of $\mathcal{G}$. let us consider the effect "on the left" of $a \in \mathcal{G}$ on $\mathcal{H}$.

$$a\mathcal{H} = \{a \circ h \mid \forall h \in \mathcal{H}\}$$

$a\mathcal{H}$ is called a left coset of $\mathcal{G}$.

We can also define $\mathcal{G}/\mathcal{H}$ the group of cosets of $\mathcal{H} \subset \mathcal{G}$, also called factor group of $\mathcal{G}$ by $\mathcal{H}$.

Invariant/normal subgroup : A subgroup $\mathcal{H}$ of $\mathcal{G}$ is said to be invariant if $a\mathcal{H}a^{-1} = \mathcal{H}$, $\forall a \in \mathcal{G}$.

By simply multiplying both sides of the previous equation with a to the right we show that right and left cosets are equal if $\mathcal{H}$ is invariant (normal).

$$a\mathcal{H}a^{-1}a = a\mathcal{H} = \mathcal{H}a$$

In conclusion, for each invariant subgroup of $\mathcal{G}$, there exists $\text{card}(\mathcal{G})$ cosets on each side that are equal to their opposite side coset.

In the context of the definition of $\mathcal{G}/\mathcal{H}$, it is interesting to note that the $\mathcal{H}$ s in question are invariant subgroups for consistency let us consider two elements of $\mathcal{G}/\mathcal{H}$

$$(a\mathcal{H})(b\mathcal{H}) = (ab)\mathcal{H}, \quad \text{where } a, b \in \mathcal{G}$$

We must have in order for this equality to be true $\mathcal{H}b = b\mathcal{H}$ so that

$$(a\mathcal{H})(b\mathcal{H}) = a(\mathcal{H}b)\mathcal{H} = (ab)\mathcal{H}\mathcal{H} = (ab)\mathcal{H}$$

Also, $\text{card}(\mathcal{G})$ is the number of elements in $\mathcal{G}$, in case it is (in)finite $\mathcal{G}$ would then be called a(n) (in)finite group.

The **kernel**² of a homomorphism φ from $\mathcal{G}$ to $\mathcal{G}'$ is the set of elements of $\mathcal{G}$ that get mapped to the identity of $\mathcal{G}'$

$$\ker(\varphi) = \{g \in \mathcal{G} \mid \varphi(g) = e_{\mathcal{G}'}\}$$

It is quite obvious from simple calculations, very similar to the ones used to prove that the elements of the factor group must be cosets of invariant subgroups, that the kernel of a homomorphism is an invariant subgroup of the initial group.

²Some of the definitions and notions I introduce seem disconnected from Physics, they are not. Take conjugacy classes for example, across one traces/characters are the same, and so are some other invariants that may be observables (from what I've picked up until one at least).

Let $n \in \ker(\varphi)$, and $g, g^{-1} \in \mathcal{G}$

$$\varphi(gng^{-1}) = \varphi(g)\varphi(n)\varphi(g^{-1}) = \varphi(g)\varphi(g^{-1})e_{\mathcal{G}'} = e_{\mathcal{G}'}$$

Therefore $gng^{-1} \in \ker(\varphi)$, put differently :

$$\ker(\varphi) = g \ker(\varphi) g^{-1}.$$

The **conjugacy class** of $x \in \mathcal{G}$ is the set of $gxg^{-1} \in \mathcal{G}$, $\forall g \in \mathcal{G}$.

2. On Representations

Representation : A representation $(\mathcal{E}, U)$ of the group $\mathcal{G}$ on the vector space $\mathcal{E}$ is a homomorphism :

$$\begin{aligned} U : \mathcal{G} &\rightarrow \text{GL}(V) \\ g &\mapsto U(g) \end{aligned}$$

If said homomorphism is an isomorphism then the representation is said to be faithful.

The mapping $g \mapsto U(g)$ must be continuous if $\mathcal{G}$ is a topological group and $\mathcal{E}$ is a topological vector space.

We shall call the vector space $\mathcal{E}$ (as do some of the authors in the referenced books) the **representation space** of $\mathcal{G}$, often called representation for simplicity. The dimension of the representation is that of the vector space it acts on, naturally. In the case of unitarity of the representation of each element of $\mathcal{G}$, i.e. $U^\dagger(g) = U^{-1}(g)$, $\forall g \in \mathcal{G}$, it is said that the representation is **unitary**.

Choosing a specific basis of $\mathcal{E}$, the operators $U(g)$ are represented by matrix elements denoted $u_j^i(g)$ for which the following equation holds³ :

$$u_j^i(g) = u_k^i(g)u_j^k(g)$$

Character : The characters χ_U of a representation of $\mathcal{G}$ are the traces of the matrix representations

$$\text{Tr } U(g) \equiv u_i^i(g)$$

Let $(\mathcal{E}_1, U_1)$ and $(\mathcal{E}_2, U_2)$ of the same group $(\mathcal{G}, \circ)$ ⁴ are said to be equivalent if there exists an isomorphism T of $\mathcal{E}_1$ on $\mathcal{E}_2$ such that :

$$TU_1(g) = U_2(g)T \quad \forall g \in \mathcal{G}$$

3. SU(2)

4. SU(3)

³Einstein's summation convention is used throughout these notes !

⁴Yeah I just wrote it rigorously, surprised ?

III Differential geometry

1. Topology

In this section the notes will mostly contain definitions (just like the first subsection of the group theory section $\dots$), I will almost exclusively use [7] which was lent to me by K.NOUI, I'm very grateful to him, it was unavailable at the university library.

This section is an introduction to the following sections.

Bibliography

- [1] T. Nakayama and H. Shima, *Higher Mathematics for Physics and Engineering: Mathematical Methods for Contemporary Physics*, Springer Berlin Heidelberg, Berlin, Heidelberg (2010), [10.1007/b138494](https://doi.org/10.1007/b138494).
- [2] W. Appel, *Mathématiques pour la physique et les physiciens !*, Éditions H&K, Paris, 5e éd. revue, corrigée et (encore) augmentée ed. (2017).
- [3] A. Rougé, *Introduction à la physique subatomique*, Les cours de l'École polytechnique, Ellipses, Paris (1997).
- [4] H. Georgi, *Lie algebras in particle physics: from isospin to unified theories*, no. 54 in Frontiers in physics, Westview press, Boulder, Colo, 2nd ed ed. (1999).
- [5] V.A. Rubakov and S.S. Wilson, *Classical Theory of Gauge Fields*, Princeton University Press, Princeton (2009).
- [6] J. Fuchs and C. Schweigert, *Symmetries, Lie algebras and representations: a graduate course for physicists*, Cambridge monographs on mathematical physics, Cambridge university press, Cambridge (1997).
- [7] M. Nakahara, *Geometry, topology and physics*, Graduate student series in physics, A. Hilger, Bristol New York (1990).