

Elementary Differential Equations and Boundary Value Problems

Twelfth Edition

Boyce

Chapter 2

First-Order Differential Equations

Section 2.4

Differences Between Linear and Nonlinear Differential Equations

Linear vs. Nonlinear Ordinary Differential Equations

- Recall that a first-order ODE has the form $y' = f(t, y)$, and is linear if f is linear in y , and nonlinear if f is nonlinear in y .
- Examples: $y' = ty - e^t$ (linear), $y' = ty^2$ (non-linear)
- In this section, we will see that first-order linear and nonlinear equations differ in a number of ways, including:
 - The theory describing existence and uniqueness of solutions, and corresponding domains, are different.
 - Solutions to linear equations can be expressed in terms of a general solution, which is not usually the case for nonlinear equations.
 - Linear equations have explicitly defined solutions while nonlinear equations typically do not, and nonlinear equations may or may not have implicitly defined solutions.
- For both types of equations, numerical and graphical construction of solutions are important.

Existence and Uniqueness Theorem for First-Order Linear Equations

- Consider the linear first order initial value problem:

$$y' + p(t)y = g(t), \quad y(0) = y_0$$

If the functions p and g are continuous on an open interval containing the point $t = t_0$, then there exists a unique function that satisfies the initial value problem for each t in I .

- **Proof outline:** Use Ch 2.1 discussion and results:

$$y = \frac{1}{\mu(t)} \left(\int_{t_0}^t \mu(s) g(s) ds + y_0 \right), \quad \mu(t) = \exp \int_{t_0}^t p(s) ds$$

Existence and Uniqueness Theorem for First-Order Nonlinear Equations

- Consider the nonlinear first order initial value problem:

$$y' = f(t, y), \quad y(t_0) = y_0$$

- Let the functions f and $\partial f / \partial y$ be continuous in some rectangle $\alpha < t < \beta, \gamma < y < \delta$ containing the point (t_0, y_0) . Then in some interval $t_0 - h < t < t_0 + h$ in the rectangle there is a unique solution $y = \phi(t)$ of the initial value problem.
- Proof discussion:** Since there is no general formula for the solution of arbitrary nonlinear first order initial value problems, this proof is difficult, and beyond the scope of this course.
- The conditions stated in the theorem are sufficient but not necessary to guarantee existence of a solution, and continuity of f ensures existence but not uniqueness of $y = \phi(t)$

Example 2.4.1: Linear Initial Value Problems

Use Theorem 2.4.1 to find an interval in which the initial value problem

$$ty' + 2y = 4t^2, \quad y(1) = 2$$

has a unique solution. Next, do the same when the initial condition is changed to $y(-1) = 2$.

- Rewrite the original equation in standard form:

$$y' + \left(\frac{2}{t}\right)y = 4t \text{ where } p(t) = \frac{2}{t} \text{ and } g(t) = 4t$$

- The solution to this initial value problem is defined for $t > 0$, the interval on which $p(t) = 2/t$ is continuous.
- If the initial condition is $y(-1) = 2$, then the solution is given by same expression as above, but is defined on $t < 0$.
- In either case, the Existence and Uniqueness Theorem guarantees that solution is unique on corresponding interval.

Example 2.4.2: Nonlinear Initial Value Problem

Apply Theorem 2.4.2 to the initial value problem

$$\frac{dy}{dx} = \frac{3x^2 + 4x + 2}{2(y-1)}, \quad y(0) = -1$$

- The functions f and $\partial f / \partial y$ are given by

$$f(x, y) = \frac{3x^2 + 4x + 2}{2(y-1)}, \quad \frac{\partial f}{\partial y}(x, y) = -\frac{3x^2 + 4x + 2}{2(y-1)^2},$$

and are continuous except on line $y = 1$.

- Thus we can draw an open rectangle about $(0, -1)$ in which f and $\partial f / \partial y$ are continuous, as long as it doesn't cover $y = 1$.
- How wide is the rectangle? Recall the solution defined for $x > -2$, with

$$y = 1 - \sqrt{x^3 + 2x^2 + 2x + 4}$$

Example 2.4.2: Change Initial Condition

- Our nonlinear initial value problem is

$$\frac{dy}{dx} = \frac{3x^2 + 4x + 2}{2(y-1)}, \quad y(0) = -1$$

with

$$f(x, y) = \frac{3x^2 + 4x + 2}{2(y-1)}, \quad \frac{\partial f}{\partial y}(x, y) = -\frac{3x^2 + 4x + 2}{2(y-1)^2},$$

which are continuous except on line $y = 1$.

- If we change the initial condition to $y(0) = 1$, then the Existence and Uniqueness Theorem is not satisfied. Solving this new initial value problem, we obtain:

$$y = 1 \pm \sqrt{x^3 + 2x^2 + 2x}, \quad x > 0$$

- Thus a solution exists but is not unique.

Example 2.4.3: Nonlinear Initial Value Problem

Apply Theorem 2.4.2 to the following initial value problem and then solve the problem.

$$y' = y^{1/3}, y(0) = 0 \quad (t \geq 0)$$

- The functions f and $\partial f / \partial y$ are given by


$$f(t, y) = y^{1/3}, \frac{\partial f}{\partial y}(t, y) = \frac{1}{3} y^{-2/3}$$

- Thus f is continuous everywhere, but $\frac{\partial f}{\partial y}$ doesn't exist at $y = 0$, and hence the Existence and Uniqueness Theorem does not apply.
- Solutions exist, but are not unique.

Example 2.4.3: Multiple Solutions

Separating variables, solving, and including the initial conditions, we obtain not one, but two solutions:

$$y^{-1/3} dy = dt \Rightarrow \frac{3}{2} y^{2/3} = t + c \Rightarrow y = \pm \left(\frac{2}{3} t \right)^{3/2}, \quad t \geq 0$$



The solution family for different values of t_0

Example 2.4.4: Nonlinear Initial Value Problem

Solve the initial value problem:

$$y' = y^2, y(0) = 1$$

- The functions f and $\partial f / \partial y$ are given by

$$f(t, y) = y^2, \frac{\partial f}{\partial y}(t, y) = 2y$$

- Thus f and $\partial f / \partial y$ are continuous at $t = 0$, so the Existence and Uniqueness Theorem guarantees that solutions exist and are unique.
- Separating variables and solving, we obtain

$$y^{-2} dy = dt \Rightarrow -y^{-1} = t + c \Rightarrow y = -\frac{1}{t + c} \Rightarrow y = \frac{1}{1 - t}$$

- The solution $y(t)$ is defined on $(-\infty, 1)$. Note that the singularity at $t = 1$ is not obvious from original IVP statement.

Interval of Existence: Linear Equations

- Per the Existence and Uniqueness Theorem, the solution of a linear initial value problem $y' + p(t)y = g(t)$, $y(t_0) = y_0$ exists throughout any interval about $t = t_0$ on which p and g are continuous.
- Vertical asymptotes or other discontinuities of a solution can only occur at points of discontinuity of p or g .
- However, a solution may be differentiable at points of discontinuity of p or g . See Chapter 2.1: Example 3 of text.
- Compare these comments with Example 1 and with previous linear equations in Chapter 1 and Chapter 2.

Interval of Existence: Nonlinear Equations

- In the nonlinear case, the interval on which a solution exists may be difficult to determine.
- The solution $y = \phi(t)$ exists as long as $[t, \phi(t)]$ remains within a rectangular region indicated in Theorem 2.4.2. This is what determines the value of h in that theorem. Since $\phi(t)$ is usually not known, it may be impossible to determine this region.
- In any case, the interval on which a solution exists may have no simple relationship to the function f in the differential equation $y' = f(t, y)$, in contrast with linear equations.
- Furthermore, any singularities in the solution may depend on the initial condition as well as the equation.
- Compare these comments to the preceding examples.

General Solutions

- For a first order linear equation, it is possible to obtain a solution containing one arbitrary constant, from which all solutions follow by specifying values for this constant.
- For nonlinear equations, such general solutions may not exist. That is, even though a solution containing an arbitrary constant may be found, there may be other solutions that cannot be obtained by specifying values for this constant.
- Consider $y' = y^2$ in example 4 where the solution $y = -\frac{1}{t+c}$ contains an arbitrary constant, but does not include all solutions to the differential equation since the function $y = 0$ is a solution of the differential equation which cannot be obtained by specifying a value for c in solution above.

Explicit Solutions: Linear Equations

- Per the Existence and Uniqueness Theorem, a solution of a linear initial value problem

$$y' + p(t)y = g(t), y(t_0) = y_0$$

exists throughout any interval about $t = t_0$ on which p and g are continuous, and this solution is unique.

- The solution has an explicit representation,

$$y = \frac{\int_{t_0}^t \mu(s) g(s) ds + y_0}{\mu(t)}, \quad \text{where } \mu(t) = e^{\int_{t_0}^t p(s) ds},$$

and can be evaluated at any appropriate value of t , as long as the necessary integrals can be computed.

Explicit Solution Approximation

- For linear first order equations, an explicit representation for the solution can be found, as long as necessary integrals can be solved.
- If integrals can't be solved, then numerical methods are often used to approximate the integrals.

Implicit Solutions: Nonlinear Equations

- For nonlinear equations, explicit representations of solutions may not exist.
- As we have seen, it may be possible to obtain an equation which implicitly defines the solution. If equation is simple enough, an explicit representation can sometimes be found.
- Otherwise, numerical calculations are necessary in order to determine values of y for given values of t . These values can then be plotted in a sketch of the integral curve.
- Recall the examples from earlier in the chapter and consider the following example

$$y' = \frac{y \cos x}{1 + 3y^3}, y(0) = 1 \quad \Rightarrow \quad \ln y + y^3 = \sin x + 1$$

Copyright

Copyright © 2021 John Wiley & Sons, Inc.

All rights reserved. Reproduction or translation of this work beyond that permitted in Section 117 of the 1976 United States Act without the express written permission of the copyright owner is unlawful. Request for further information should be addressed to the Permissions Department, John Wiley & Sons, Inc. The purchaser may make back-up copies for his/her/their own use only and not for distribution or resale. The Publisher assumes no responsibility for errors, omissions, or damages, caused by the use of these programs or from the use of the information contained herein.