

# Elementary Differential Equations and Boundary Value Problems

**Twelfth Edition**

**Boyce**

## Chapter 3

### Second-Order Linear Differential Equations

# Section 3.1 Homogeneous Differential Equations with Constant Coefficients

# 2<sup>nd</sup> Order Linear Homogeneous Equations-Constant Coefficients

- A **second order ordinary differential equation** has the general form

$$\frac{d^2y}{dt^2} = f\left(t, y, \frac{dy}{dt}\right)$$

where  $f$  is some given function.

- This equation is said to be **linear** if  $f$  is linear in  $y$  and  $y'$ :

$$y'' + p(t)y' + q(t)y = g(t)$$

Otherwise the equation is said to be **nonlinear**.

- A second order linear equation often appears as

$$P(t)y'' + Q(t)y' + R(t)y = G(t)$$

- If  $g(t)$  or  $G(t) = 0$  for all  $t$ , then the equation is called **homogeneous**. Otherwise the equation is **nonhomogeneous**.

# Homogeneous Equations, Initial Values

- In Sections 3.5 and 3.6, we will see that once a solution to a homogeneous equation is found, then it is possible to solve the corresponding nonhomogeneous equation, or at least express the solution in terms of an integral.
- The focus of this chapter is thus on homogeneous equations; and in particular, those with constant coefficients:

$$ay'' + by' + cy = 0$$

We will examine the variable coefficient case in Chapter 5.

- Initial conditions typically take the form

$$y(t_0) = y_0, \quad y'(t_0) = y'_0,$$

- Thus solution passes through  $(t_0, y_0)$ , and the slope of solution at  $(t_0, y_0)$  is equal to  $y'_0$ .

# Example 3.1.1: General Solution

- Find the general solution for the second order linear differential equation:

$$y'' - y = 0$$

- Two solutions of this equation are

$$y_1(t) = e^t, \quad y_2(t) = e^{-t}$$

- Other solutions include

$$y_3(t) = 3e^t, \quad y_4(t) = 5e^{-t}, \quad y_5(t) = 3e^t + 5e^{-t}$$

- Based on these observations, we see that there are infinitely many solutions of the form

$$y(t) = c_1 e^t + c_2 e^{-t}$$

- It will be shown in Section 3.2 that all solutions of the differential equation above can be expressed in this form.

# Example 3.1.1: Initial Conditions

- Now consider the following initial value problem for our equation:

$$y'' - y = 0, \quad y(0) = 2, \quad y'(0) = -1$$

- We have found a general solution of the form

$$y(t) = c_1 e^t + c_2 e^{-t}$$

- Using the initial equations,

$$\left. \begin{array}{l} y(0) = c_1 + c_2 = 2 \\ y'(0) = c_1 - c_2 = -1 \end{array} \right\} \Rightarrow c_1 = \frac{1}{2}, c_2 = \frac{3}{2}$$

- Thus

$$y(t) = \frac{1}{2} e^t + \frac{3}{2} e^{-t}$$

# The Characteristic Equation

- To solve the 2<sup>nd</sup> order equation with constant coefficients,

$$ay'' + by' + cy = 0$$

we begin by assuming a solution of the form  $y = e^{rt}$ .

- Substituting this into the differential equation, we obtain

$$ar^2e^{rt} + bre^{rt} + ce^{rt} = 0$$

- Simplifying,

$$e^{rt}(ar^2 + br + c) = 0$$

and hence

$$ar^2 + br + c = 0$$

- This last equation is called the **characteristic equation** of the differential equation.
- We then solve for  $r$  by factoring or using quadratic formula.

# The General Solution

- Using the quadratic formula on the characteristic equation

$$ar^2 + br + c = 0,$$

we obtain two solutions,  $r_1$  and  $r_2$ .

- There are three possible results:

- The roots  $r_1, r_2$  are real and  $r_1 \neq r_2$ .
- The roots  $r_1, r_2$  are real and  $r_1 = r_2$ .
- The roots  $r_1, r_2$  are complex.

$$r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

- In this section, we will assume  $r_1, r_2$  are real and  $r_1 \neq r_2$ .
- In this case, the general solution has the form

$$y(t) = c_1 e^{r_1 t} + c_2 e^{r_2 t}$$



# Initial Conditions

- For the initial value problem

$$ay'' + by' + cy = 0 \text{ where } y(t_0) = y_0 \text{ and } y'(t_0) = y'_0$$

we use the general solution

$$y(t) = c_1 e^{r_1 t} + c_2 e^{r_2 t}$$

together with the initial conditions to find  $c_1$  and  $c_2$ . That is,

$$\left. \begin{aligned} c_1 e^{r_1 t_0} + c_2 e^{r_2 t_0} &= y_0 \\ c_1 r_1 e^{r_1 t_0} + c_2 r_2 e^{r_2 t_0} &= y'_0 \end{aligned} \right\} \Rightarrow c_1 = \frac{y'_0 - y_0 r_2}{r_1 - r_2} e^{-r_1 t_0}, c_2 = \frac{y_0 r_1 - y'_0}{r_1 - r_2} e^{-r_2 t_0}$$

- Since we are assuming  $r_1 \neq r_2$ , it follows that a solution of the form  $y = e^{rt}$  to the above initial value problem will always exist, for any set of initial conditions.

## Example 3.1.2 General Solution

- Consider the linear differential equation

$$y'' + 5y' + 6y = 0$$

- Assuming an exponential solution leads to the characteristic equation:

$$y(t) = e^{rt} \Rightarrow r^2 + 5r + 6 = 0 \Leftrightarrow (r + 2)(r + 3) = 0$$

- Factoring the characteristic equation yields two solutions:  $r_1 = -2$  and  $r_2 = -3$
- Therefore, the general solution to this differential equation has the form

$$y(t) = c_1 e^{-2t} + c_2 e^{-3t}$$

# Example 3.1.3 Particular Solution

- Consider the initial value problem

$$y'' + 5y' + 6y = 0, \quad y(0) = 2, \quad y'(0) = 3$$

- From the preceding example, we know the general solution has the form:


$$y(t) = c_1 e^{-2t} + c_2 e^{-3t}$$

- With derivative:  $y'(t) = -2c_1 e^{-2t} - 3c_2 e^{-3t}$

- Using the initial conditions:

$$\left. \begin{array}{l} c_1 + c_2 = 2 \\ -2c_1 - 3c_2 = 3 \end{array} \right\} \Rightarrow c_1 = 9, \quad c_2 = -7$$

- Thus  $y(t) = 9e^{-2t} - 7e^{-3t}$



# Example 3.1.4: Initial Value Problem

- Consider the initial value problem

$$4y'' - 8y' + 3y = 0, \quad y(0) = 2, \quad y'(0) = \frac{1}{2}$$

- Then  $y(t) = e^{rt} \Rightarrow 4r^2 - 8r + 3 = 0 \Leftrightarrow (2r - 3)(2r - 1) = 0$


- Factoring yields two solutions,  $r_1 = \frac{3}{2}$  and  $r_2 = \frac{1}{2}$   $y(t) = -\frac{1}{2}e^{3t/2} + \frac{5}{2}e^{t/2}$

- The general solution has the form

$$y(t) = c_1 e^{3t/2} + c_2 e^{t/2}$$

- Using initial conditions:

$$\left. \begin{aligned} c_1 + c_2 &= 2 \\ \frac{3}{2}c_1 + \frac{1}{2}c_2 &= \frac{1}{2} \end{aligned} \right\} \Rightarrow c_1 = -\frac{1}{2}, c_2 = \frac{5}{2}$$



## Example 3.1.5: Find Maximum Value

For the initial value problem in Example 3.3, to find the maximum value attained by the solution, we set  $y'(t) = 0$  and solve for  $t$ :

Set  $y' = 0$  and multiply by  $e^{3t}$  to find  $t_{max}$  which satisfies

$e^t = \frac{7}{6}$ , hence:

$$t_{max} = \ln\left(\frac{7}{6}\right) \cong 0.15415$$

The corresponding maximum value  $y_{max}$  is given by:

$$y_{max} = 9e^{-2t_m} - 7e^{-3t_m} = \frac{108}{49} \cong 2.20408$$

# Copyright

## **Copyright © 2021 John Wiley & Sons, Inc.**

All rights reserved. Reproduction or translation of this work beyond that permitted in Section 117 of the 1976 United States Act without the express written permission of the copyright owner is unlawful. Request for further information should be addressed to the Permissions Department, John Wiley & Sons, Inc. The purchaser may make back-up copies for his/her/their own use only and not for distribution or resale. The Publisher assumes no responsibility for errors, omissions, or damages, caused by the use of these programs or from the use of the information contained herein.