

# Elementary Differential Equations and Boundary Value Problems

**Twelfth Edition**

**Boyce**

## Chapter 3

### Second-Order Linear Differential Equations

# Section 3.4 Repeated Roots; Reduction of Order

# Repeated Roots; Reduction of Order

- Recall our 2<sup>nd</sup> order linear homogeneous ODE

$$ay'' + by' + cy = 0$$

where  $a$ ,  $b$  and  $c$  are constants

- Assuming an exponential solution leads to characteristic equation:

$$y(t) = e^{rt} \Rightarrow ar^2 + br + c = 0$$

- Quadratic formula (or factoring) yields two solutions,  $r_1$  and  $r_2$ :

$$r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

- When  $b^2 - 4ac = 0$ , then  $r_1 = r_2 = \frac{-b}{2a}$ , with both roots yielding the same solution:

$$y_1(t) = ce^{-bt/(2a)}$$

# Finding the Second Solution Using Multiplying Factor $v(t)$

- We know that if  $y_1(t)$  is a solution, then  $y_2(t) = cy_1(t)$  is a solution.
- Since  $y_1$  and  $y_2$  are linearly dependent, we can generalize this approach and multiply by a function  $v(t)$ , and determine conditions for which  $y_2$  is a solution:

$$y_1(t) = e^{-bt/(2a)} \text{ a solution } \square \quad \text{try } y_2(t) = v(t)e^{-bt/(2a)}$$

$$y_2(t) = v(t)e^{-bt/(2a)}$$

$$y_2'(t) = v'(t)e^{-bt/(2a)} - \frac{b}{2a}v(t)e^{-bt/(2a)}$$

$$y_2''(t) = v''(t)e^{-bt/(2a)} - \frac{b}{2a}v'(t)e^{-bt/(2a)} - \frac{b}{2a}v'(t)e^{-bt/(2a)} + \frac{b^2}{4a^2}v(t)e^{-bt/(2a)}$$

$$y_2'' = v''(t)e^{-bt/(2a)} - \frac{b}{a}v'(t)e^{-bt/(2a)} + \frac{b^2}{4a^2}v(t)e^{-bt/(2a)}$$

# Finding the Multiplying Factor $v(t)$

- Substituting derivatives into the original ODE, we can find an expression for  $v$ :

$$e^{-bt/(2a)} \left\{ a \left[ v''(t) - \frac{b}{a} v'(t) + \frac{b^2}{4a^2} v(t) \right] + b \left[ v'(t) - \frac{b}{2a} v(t) \right] + cv(t) \right\} = 0$$

$$av''(t) - bv'(t) + \frac{b^2}{4a} v(t) + bv'(t) - \frac{b^2}{2a} v(t) + cv(t) = 0$$

$$av''(t) + \left( \frac{b^2}{4a} - \frac{b^2}{2a} + c \right) v(t) = 0$$

$$av''(t) + \left( \frac{b^2}{4a} - \frac{2b^2}{4a} + \frac{4ac}{4a} \right) v(t) = 0 \Leftrightarrow av''(t) + \left( \frac{-b^2}{4a} + \frac{4ac}{4a} \right) v(t) = 0$$

$$av''(t) - \left( \frac{b^2 - 4ac}{4a} \right) v(t) = 0$$

$$v''(t) = 0 \Rightarrow v(t) = k_3 t + k_4$$

# General Solution

- To find our general solution, we have:

$$\begin{aligned}y(t) &= k_1 e^{-bt/(2a)} + k_2 v(t) e^{-bt/(2a)} \\&= k_1 e^{-bt/(2a)} + (k_3 t + k_4) e^{-bt/(2a)} \\&= c_1 e^{-bt/(2a)} + c_2 t e^{-bt/(2a)}\end{aligned}$$

- Thus the general solution for repeated roots is

$$y(t) = c_1 e^{-bt/(2a)} + c_2 t e^{-bt/(2a)}$$

# Wronskian

- The general solution is

$$y(t) = c_1 e^{-bt/(2a)} + c_2 t e^{-bt/(2a)}$$

- Thus every solution is a linear combination of

$$y_1(t) = e^{-bt/(2a)}, y_2(t) = t e^{-bt/(2a)}$$

- The Wronskian of the two solutions is

$$\begin{aligned} W(y_1, y_2)(t) &= \begin{vmatrix} e^{-bt/(2a)} & t e^{-bt/(2a)} \\ -\frac{b}{2a} e^{-bt/(2a)} & \left(1 - \frac{bt}{2a}\right) e^{-bt/(2a)} \end{vmatrix} \\ &= e^{-bt/a} \left(1 - \frac{bt}{2a}\right) + e^{-bt/a} \left(\frac{bt}{2a}\right) \\ &= e^{-bt/a} \neq 0 \quad \text{for all } t \end{aligned}$$

- Thus  $y_1$  and  $y_2$  form a fundamental solution set for equation.

## Example 3.4.1 (part one)

- Solve the differential equation:

$$y'' + 4y' + 4y = 0$$

- Assuming an exponential solution leads to characteristic equation:

$$y(t) = e^{rt} \Rightarrow r^2 + 4r + 4 = 0 \Leftrightarrow (r + 2)^2 = 0 \Leftrightarrow r = -2$$

- So one solution is  $y_1(t) = e^{-2t}$  and a second solution is found:

$$y_2(t) = v(t)e^{-2t}$$

$$y_2'(t) = v'(t)e^{-2t} - 2v(t)e^{-2t}$$

$$y_2''(t) = v''(t)e^{-2t} - 4v'(t)e^{-2t} + 4v(t)e^{-2t}$$

- Substituting these into the differential equation and simplifying yields  $v''(t) = 0$ ,  $v'(t) = k_1$ ,  $v(t) = k_1t + k_2$  where  $k_1$  and  $k_2$  are arbitrary constants.




## Example 3.4.1 (part two)

- Letting  $k_1 = 1$  and  $k_2 = 0$ ,  $v(t) = t$  and  $y_2(t) = te^{-2t}$
- So the general solution is

$$y(t) = c_1 e^{-2t} + c_2 t e^{-2t}$$

- Note that both  $y_1$  and  $y_2$  tend to 0 as  $t \rightarrow \infty$  regardless of the values of  $c_1$  and  $c_2$
- The figure shows three solutions of this equation with different sets of initial conditions.
  - $y(0) = 2, y'(0) = 1$  (top)
  - $y(0) = 1, y'(0) = 1$  (middle)
  - $y(0) = 1/2, y'(0) = 1$  (bottom)



## Example 3.4.2 (part one)

- Find the solution of the initial value problem

$$y'' - y' + \frac{1}{4}y = 0, \quad y(0) = 2, \quad y'(0) = \frac{1}{3}$$

- Assuming exponential solution leads to characteristic equation:

$$y(t) = e^{rt} \Rightarrow r^2 - r + \frac{1}{4} = 0 \Leftrightarrow (r - \frac{1}{2})^2 = 0 \Leftrightarrow r = \frac{1}{2}$$


- Thus the general solution is

$$y(t) = c_1 e^{t/2} + c_2 t e^{t/2}$$

- Using the initial conditions:

$$\left. \begin{aligned} c_1 &= 2 \\ \frac{1}{2}c_1 + c_2 &= \frac{1}{3} \end{aligned} \right\} \Rightarrow c_1 = 2, \quad c_2 = -\frac{2}{3}$$

- Thus  $y(t) = 2e^{t/2} - \frac{2}{3}te^{t/2}$



## Example 3.4.2 (part two)


- Suppose that the initial slope in the previous problem was increased

$$y(0) = 2, \quad y'(0) = 2$$

- The solution of this modified problem is

$$y(t) = 2e^{t/2} + te^{t/2}$$

- Notice that the coefficient of the second term is now positive. This makes a difference in the graph, since the exponential function is raised to a positive power.



# Reduction of Order

- The method used so far in this section also works for equations with nonconstant coefficients:

$$y'' + p(t)y' + q(t)y = 0$$

- That is, given that  $y_1$  is solution, try  $y_2 = v(t)y_1$ :

$$y_2(t) = v(t)y_1(t)$$

$$y_2'(t) = v'(t)y_1(t) + v(t)y_1'(t)$$

$$y_2''(t) = v''(t)y_1(t) + 2v'(t)y_1'(t) + v(t)y_1''(t)$$

- Substituting these into the original ODE and collecting terms,

$$y_1v'' + (2y_1' + py_1)v' + (y_1'' + py_1' + qy_1)v = 0$$

- Since  $y_1$  is a solution to the differential equation, this last equation reduces to a first order equation in  $v$ :

$$y_1v'' + (2y_1' + py_1)v' = 0$$

## Example 3.4.3: Reduction of Order

- Given the variable coefficient equation and solution  $y_1$ ,

$$2t^2 y'' + 3ty' - y = 0, \quad t > 0; \quad y_1(t) = t^{-1},$$

use reduction of order method to find a second solution:

$$y_2(t) = v(t) t^{-1}$$

$$y_2'(t) = v'(t) t^{-1} - v(t) t^{-2}$$

$$y_2''(t) = v''(t) t^{-1} - 2v'(t) t^{-2} + 2v(t) t^{-3}$$

- Substituting these into the original ODE and collecting terms,

$$2t^2(v''t^{-1} - 2v't^{-2} + 2vt^{-3}) + 3t(v't^{-1} - vt^{-2}) - vt^{-1} = 0$$

$$\Leftrightarrow 2v''t - 4v' + 4vt^{-1} + 3v' - 3vt^{-1} - vt^{-1} = 0$$

$$\Leftrightarrow 2tv'' - v' = 0$$

$$\Leftrightarrow 2tu' - u = 0, \quad \text{where } u(t) = v'(t)$$

## Example 3.4.3: Finding $v(t)$

- To solve

$$2tu' - u = 0, \quad u(t) = v'(t)$$

for  $u$ , we can use the separation of variables method:

$$2t \frac{du}{dt} - u = 0 \quad \Leftrightarrow \quad \int \frac{du}{u} = \int \frac{1}{2t} dt \quad \Leftrightarrow \quad \ln|u| = \frac{1}{2} \ln|t| + C$$

$$\Leftrightarrow |u| = |t|^{1/2} e^C \quad \Leftrightarrow u = ct^{1/2}, \text{ since } t > 0.$$

- Thus  $v' = ct^{1/2}$

and hence

$$v(t) = \frac{2}{3} ct^{3/2} + k$$

## Example 3.4.3: General Solution

- Since  $v(t) = \frac{2}{3}ct^{3/2} + k$

$$y_2(t) = \left( \frac{2}{3}ct^{3/2} + k \right) t^{-1} = \frac{2}{3}ct^{1/2} + k t^{-1}$$

- Recall  $y_1(t) = t^{-1}$
- So we can neglect the second term of  $y_2$  to obtain

$$y_2(t) = t^{1/2}$$

- The Wronskian of  $y_1(t)$  and  $y_2(t)$  can be computed

$$W[y_1, y_2](t) = \frac{3}{2}t^{-3/2} \neq 0 \text{ for } t > 0$$

- Hence the general solution to the differential equation is

$$y(t) = c_1 t^{-1} + c_2 t^{1/2}$$

# Copyright

## **Copyright © 2021 John Wiley & Sons, Inc.**

All rights reserved. Reproduction or translation of this work beyond that permitted in Section 117 of the 1976 United States Act without the express written permission of the copyright owner is unlawful. Request for further information should be addressed to the Permissions Department, John Wiley & Sons, Inc. The purchaser may make back-up copies for his/her/their own use only and not for distribution or resale. The Publisher assumes no responsibility for errors, omissions, or damages, caused by the use of these programs or from the use of the information contained herein.