

Elementary Differential Equations and Boundary Value Problems

Twelfth Edition

Boyce

Chapter 3

Second Order Linear Differential Equations

Section 3.6 Variation of Parameters

Variation of Parameters

- Recall the nonhomogeneous equation

$$y'' + p(t)y' + q(t)y = g(t)$$

where p , q , g are continuous functions on an open interval I .

- The associated homogeneous equation is

$$y'' + p(t)y' + q(t)y = 0$$

- In this section we will learn the **variation of parameters** method to solve the nonhomogeneous equation. As with the method of undetermined coefficients, this procedure relies on knowing solutions to the homogeneous equation.
- Variation of parameters is a general method, and requires no detailed assumptions about solution form. However, certain integrals must be evaluated, and this can present difficulties.

Example 3.6.1: Variation of Parameters

- Find the general solution to:

$$y'' + 4y = 8 \tan t, \quad -\pi/2 < t < \pi/2$$

- We cannot use the undetermined coefficients method since $g(t)$ is a quotient of $\sin t$ or $\cos t$, instead of a sum or product.
- Recall that the solution to the homogeneous equation is

$$y_C(t) = c_1 \cos(2t) + c_2 \sin(2t)$$

- To find a particular solution to the nonhomogeneous equation, we begin with the form

$$y(t) = u_1(t) \cos(2t) + u_2(t) \sin(2t)$$

- Then

$$y'(t) = u_1'(t) \cos(2t) - 2u_1(t) \sin(2t) + u_2'(t) \sin(2t) + 2u_2(t) \cos(2t)$$

- or $y'(t) = -2u_1(t) \sin(2t) + 2u_2(t) \cos(2t) + u_1'(t) \cos(2t) + u_2'(t) \sin(2t)$

Example 3.6.1: Derivatives and 2nd Equation

- From the previous slide,

$$y'(t) = -2u_1(t)\sin(2t) + 2u_2(t)\cos(2t) + u_1'(t)\cos(2t) + u_2'(t)\sin(2t)$$

- We need two equations to solve for u_1 and u_2 . The first equation is the differential equation. To get a second equation, we will require:

$$u_1'(t)\cos(2t) + u_2'(t)\sin(2t) = 0$$

- Then

$$y'(t) = -2u_1(t)\sin(2t) + 2u_2(t)\cos(2t)$$

- Next,

$$y''(t) = -2u_1'(t)\sin(2t) - 4u_1(t)\cos(2t) + 2u_2'(t)\cos(2t) - 4u_2(t)\sin(2t)$$

Example 3.6.1: Two Equations for u_1 and u_2

- Recall that our differential equation is

$$y'' + 4y = 8 \tan t$$

- Substituting y'' and y into this equation, we obtain

$$\begin{aligned} & -2u_1'(t)\sin(2t) - 4u_1(t)\cos(2t) + 2u_2'(t)\cos(2t) - 4u_2(t)\sin(2t) \\ & + 4(u_1(t)\cos(2t) + u_2(t)\sin(2t)) = 8 \tan t \end{aligned}$$

- This equation simplifies to

$$-2u_1'(t)\sin(2t) + 2u_2'(t)\cos(2t) = 8 \tan t$$

- Thus, to solve for u_1 and u_2 , we have the two equations:

$$-2u_1'(t)\sin(2t) + 2u_2'(t)\cos(2t) = 8 \tan t$$

$$u_1'(t)\cos(2t) + u_2'(t)\sin(2t) = 0$$

Example 3.6.1: Solve for u_1'

- To find u_1 and u_2 , we first need to solve for u_1' and u_2'

$$-2u_1'(t)\sin(2t) + 2u_2'(t)\cos(2t) = 8\tan t$$

$$u_1'(t)\cos(2t) + u_2'(t)\sin(2t) = 0$$

- From second equation, $u_2'(t) = -u_1'(t)\frac{\cos 2t}{\sin 2t}$

- Substituting this into the first equation,

$$-2u_1'(t)\sin(2t) + 2\left[-u_1'(t)\frac{\cos(2t)}{\sin(2t)}\right]\cos(2t) = 8\tan t$$

$$-2u_1'(t)\sin^2(2t) - 2u_1'(t)\cos^2(2t) = 8\tan t \sin(2t)$$

$$-2u_1'(t)\left[\sin^2(2t) + \cos^2(2t)\right] = 8\left[\frac{2\sin^2 t \cos t}{\cos t}\right]$$

$$u_1'(t) = -8\sin^2 t$$

Example 3.6.1: Solve for u_1 and u_2

- From the previous slide,

$$u_1'(t) = -8 \sin^2 t, \quad u_2'(t) = -u_1'(t) \frac{\cos 2t}{\sin 2t}$$

- Then

$$u_2'(t) = 8 \sin^2 t \frac{\cos(2t)}{\sin(2t)} = 4 \frac{\sin t (2 \cos^2 t - 1)}{\cos t} = 4 \sin t \left(2 \cos t - \frac{1}{\cos t} \right)$$

- Thus

$$u_1(t) = \int u_1'(t) dt = 4 \sin t \cos t - 4t + c_1$$

$$u_2(t) = \int u_2'(t) dt = 4 \ln(\cos t) - 4 \cos^2 t + c_2$$

Example 3.6.1: General Solution

- Recall our equation and homogeneous solution y_C :

$$y'' + 4y = 8 \tan t, \quad y_C(t) = c_1 \cos(2t) + c_2 \sin(2t)$$

- Using the expressions for u_1 and u_2 on the previous slide, the general solution to the differential equation is

$$\begin{aligned} y(t) &= u_1(t) \cos 2t + u_2(t) \sin 2t + y_C(t) \\ &= (4 \sin t \cos t) \cos(2t) + (4 \ln(\cos t) - 4 \cos^2 t) \sin(2t) + c_1 \cos(2t) + c_2 \sin(2t) \\ &= -2 \sin(2t) - 4t \cos(2t) + 4 \ln(\cos t) \sin(2t) + c_1 \cos(2t) + c_2 \sin(2t) \end{aligned}$$

Variation of Parameters Summary

- Suppose y_1, y_2 are fundamental solutions to the homogeneous equation associated with the nonhomogeneous equation $y'' + p(t)y' + q(t)y = g(t)$, where we note that the coefficient on y'' is 1.
- To find u_1 and u_2 , we need to solve the equations

$$u_1'(t)y_1(t) + u_2'(t)y_2(t) = 0$$

$$u_1'(t)y_1'(t) + u_2'(t)y_2'(t) = g(t)$$

- Doing so, and using the Wronskian, we obtain

$$u_1'(t) = -\frac{y_2(t)g(t)}{W[y_1, y_2](t)}, \quad u_2'(t) = \frac{y_1(t)g(t)}{W[y_1, y_2](t)}$$

- Thus

$$u_1(t) = -\int \frac{y_2(t)g(t)}{W[y_1, y_2](t)} dt + c_1, \quad u_2(t) = \int \frac{y_1(t)g(t)}{W[y_1, y_2](t)} dt + c_2$$

Theorem 3.6.1

- Consider the equations

$$y'' + p(t)y' + q(t)y = g(t) \quad (1)$$

$$y'' + p(t)y' + q(t)y = 0 \quad (2)$$

- If the functions p , q and g are continuous on an open interval I , and if y_1 and y_2 are fundamental solutions to Eq. (2), then a particular solution of Eq. (1) is

$$Y(t) = -y_1(t) \int_{t_0}^t \frac{y_2(s)g(s)}{W[y_1, y_2](s)} ds + y_2(t) \int_{t_0}^t \frac{y_1(s)g(s)}{W[y_1, y_2](s)} ds$$

and the general solution is

$$y = c_1 y_1(t) + c_2 y_2(t) + Y(t)$$

Copyright

Copyright © 2021 John Wiley & Sons, Inc.

All rights reserved. Reproduction or translation of this work beyond that permitted in Section 117 of the 1976 United States Act without the express written permission of the copyright owner is unlawful. Request for further information should be addressed to the Permissions Department, John Wiley & Sons, Inc. The purchaser may make back-up copies for his/her/their own use only and not for distribution or resale. The Publisher assumes no responsibility for errors, omissions, or damages, caused by the use of these programs or from the use of the information contained herein.