

Elementary Differential Equations and Boundary Value Problems

Twelfth Edition

Boyce

Chapter 3

Second-Order Linear Differential Equations

Section 3.3 Complex Roots of the Characteristic Equation

Complex Roots of Characteristic Equation

- Recall our discussion of the equation

$$ay'' + by' + cy = 0$$

where a , b and c are constants

- Assuming an exponential solution leads to characteristic equation:

$$y(t) = e^{rt} \Rightarrow ar^2 + br + c = 0$$

- Application of the quadratic formula (or factoring) yields two solutions, r_1 and r_2 :

$$r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

- If $b^2 - 4ac < 0$, then the roots are conjugate complex numbers:

$$r_1 = \lambda + i\mu \text{ and } r_2 = \lambda - i\mu$$

Thus

$$y_1(t) = e^{(\lambda + i\mu)t}, \quad y_2(t) = e^{(\lambda - i\mu)t}$$

Euler's Formula; Complex Valued Solutions

- Substituting it into the Taylor series for e^t , we obtain **Euler's formula**:

$$e^{it} = \sum_{n=0}^{\infty} \frac{(it)^n}{n!} = \sum_{k=0}^{\infty} \frac{(-1)^k t^{2k}}{(2k)!} + i \sum_{k=0}^{\infty} \frac{(-1)^k t^{2k+1}}{(2k+1)!}$$

- Generalizing Euler's formula, we obtain

$$e^{i\mu t} = \cos(\mu t) + i \sin(\mu t)$$

- Then

$$e^{(\lambda+i\mu)t} = e^{\lambda t} e^{i\mu t} = e^{\lambda t} (\cos(\mu t) + i \sin(\mu t)) = e^{\lambda t} \cos(\mu t) + i e^{\lambda t} \sin(\mu t)$$

- Therefore

$$y_1(t) = e^{(\lambda+i\mu)t} = e^{\lambda t} \cos(\mu t) + i e^{\lambda t} \sin(\mu t)$$

$$y_2(t) = e^{(\lambda-i\mu)t} = e^{\lambda t} \cos(\mu t) - i e^{\lambda t} \sin(\mu t)$$

Real-Valued Solutions

- Our two solutions thus far are complex-valued functions:

$$y_1(t) = e^{\lambda t} \cos(\mu t) + ie^{\lambda t} \sin(\mu t)$$

$$y_2(t) = e^{\lambda t} \cos(\mu t) - ie^{\lambda t} \sin(\mu t)$$

- We would prefer to have real-valued solutions, since our differential equation has real coefficients.
- To achieve this, recall that linear combinations of solutions are themselves solutions:

$$y_1(t) + y_2(t) = 2e^{\lambda t} \cos(\mu t)$$

$$y_1(t) - y_2(t) = 2ie^{\lambda t} \sin(\mu t)$$

- Ignoring constants, we obtain the two solutions

$$y_3(t) = e^{\lambda t} \cos(\mu t), \quad y_4(t) = e^{\lambda t} \sin(\mu t)$$

Real-Valued Solutions: The Wronskian

- Thus we have the following real-valued functions:

$$u(t) = e^{\lambda t} \cos(\mu t), \quad v(t) = e^{\lambda t} \sin(\mu t)$$

- Computation of the Wronskian gives:

$$W[u, v](t) = \mu e^{2\lambda t}$$

- Thus y_3 and y_4 form a fundamental solution set for our ODE, and the general solution can be expressed as

$$y = c_1 e^{\lambda t} \cos(\mu t) + c_2 e^{\lambda t} \sin(\mu t)$$

Example 3.3.1 (part one)

- Find the general solution of the differential equation:

$$y'' + y' + 9.25y = 0$$

- The characteristic equation is $r^2 + r + 9.25 = 0$ with roots given by:

$$r_1 = -\frac{1}{2} + 3i, \quad r_2 = -\frac{1}{2} - 3i$$

- Therefore, separating the real and imaginary components,

$$\lambda = -\frac{1}{2}, \mu = 3$$

and thus the general solution is

$$y = e^{-t/2} (c_1 \cos(3t) + c_2 \sin(3t))$$

Example 3.3.1 (part two)

- Using the general solution just determined

$$y = e^{-t/2} (c_1 \cos(3t) + c_2 \sin(3t))$$


- We can determine the particular solution that satisfies the initial conditions $y(0) = 2$ and $y'(0) = 8$

- So
$$\left. \begin{aligned} y(0) &= c_1 = 2 \\ y'(0) &= -\frac{1}{2}c_1 + 3c_2 = 8 \end{aligned} \right\} \Rightarrow c_1 = 2, c_2 = 3$$

- Thus the solution of this IVP is

$$y = e^{-t/2} (2 \cos(3t) + 3 \sin(3t))$$

- The solution is a decaying oscillation



Example 3.3.2

- Find the solution to the initial value problem

$$16y'' - 8y' + 145y = 0, \quad y(0) = -2, \quad y'(0) = 1$$


- Then $y(t) = e^{rt} \Rightarrow 16r^2 - 8r + 145 = 0 \Leftrightarrow r = \frac{1}{4} \pm 3i$
- Thus the general solution is $y(t) = c_1 e^{t/4} \cos(3t) + c_2 e^{t/4} \sin(3t)$

- And $y(0) = c_1 = -2$
 $y'(0) = \frac{1}{4}c_1 + 3c_2 = 1$ so, $c_1 = -2; c_2 = \frac{1}{2}$

- The solution of the IVP is

$$y(t) = -2e^{t/4} \cos(3t) + \frac{1}{2}e^{t/4} \sin(3t)$$

- The solution is a growing oscillation




Example 3.3.3

- Find the general solution of $y'' + 9y = 0$
- Then $y(t) = e^{rt} \Rightarrow r^2 + 9 = 0 \Leftrightarrow r = \pm 3i$
- Therefore $\lambda = 0, \mu = 3$

and thus the general solution is

$$y(t) = c_1 \cos(3t) + c_2 \sin(3t)$$

- Because $\lambda = 0$, there is no exponential factor in the solution, so the amplitude of each oscillation is constant. The figure shows the graph with two sets of initial conditions.



Copyright

Copyright © 2021 John Wiley & Sons, Inc.

All rights reserved. Reproduction or translation of this work beyond that permitted in Section 117 of the 1976 United States Act without the express written permission of the copyright owner is unlawful. Request for further information should be addressed to the Permissions Department, John Wiley & Sons, Inc. The purchaser may make back-up copies for his/her/their own use only and not for distribution or resale. The Publisher assumes no responsibility for errors, omissions, or damages, caused by the use of these programs or from the use of the information contained herein.