

Elementary Differential Equations and Boundary Value Problems

Twelfth Edition

Boyce

Chapter 2

First-Order Differential Equations

Section 2.2

Separable Differential Equations

Definition of Separable Differential Equations

- In this section we examine a subclass of linear and nonlinear first order equations. Consider the first order equation

$$\frac{dy}{dx} = f(x, y)$$

- We can rewrite this in the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0$$

- For example, we can let $M(x, y) = -f(x, y)$ and $N(x, y) = 1$ to do this, but there may be other ways as well. In differential form,

$$M(x, y)dx + N(x, y)dy = 0$$

- If M is a function of x only and N is a function of y only, then

$$M(x)dx + N(y)dy = 0$$

- In this case, the equation is called **separable**.

Example 2.2.1: Finding a General Solution for a Separable Equation

- Solve the following first order nonlinear equation:

$$\frac{dy}{dx} = \frac{x^2}{1-y^2}$$

- Separating variables, and using calculus, we obtain

$$(1-y^2)dy = (x^2)dx$$


$$\int (1-y^2)dy = \int (x^2)dx$$

$$y - \frac{1}{3}y^3 = \frac{1}{3}x^3 + c$$

$$3y - y^3 = x^3 + c$$

Example 2.2.1: Graphing Solutions for a Separable Equation

The general solution $3y - y^3 = x^3 + c$ defines the solution y implicitly. A graph showing the direction field and implicit plots of several solution curves for the differential equation is shown below:



Example 2.2.2: Finding an Implicit General Solution

- Solve the following first order nonlinear equation:

$$\frac{dy}{dx} = \frac{3x^2 + 4x + 2}{2(y-1)}$$

- Separating variables and using calculus, we obtain

$$2(y-1)dy = (3x^2 + 4x + 2)dx$$

$$2\int (y-1)dy = \int (3x^2 + 4x + 2)dx$$

$$y^2 - 2y = x^3 + 2x^2 + 2x + c$$

The equation above defines the solution y implicitly. An explicit expression for the solution can be found (next slide)

Example 2.2.2: Rearrange to create an explicit general solution in terms of “ y ”

$$y^2 - 2y = x^3 + 2x^2 + 2x + c$$

The general equation above defines the solution y implicitly. An explicit expression for the solution can be found in this case:

$$y^2 - 2y - (x^3 + 2x^2 + 2x + c) = 0 \Rightarrow y = \frac{2 \pm \sqrt{4 + 4(x^3 + 2x^2 + 2x + c)}}{2}$$
$$y = 1 \pm \sqrt{x^3 + 2x^2 + 2x + C}$$

Example 2.2.2: Initial Value Condition (0, -1)

- Suppose we seek a solution satisfying $y(0) = -1$. Using the implicit expression of y , we obtain

$$y^2 - 2y = x^3 + 2x^2 + 2x + C$$
$$(-1)^2 - 2(-1) = C \Rightarrow C = 3$$

- The implicit particular solution is $y^2 - 2y = x^3 + 2x^2 + 2x + 3$
- An explicit expression of y is:

$$y = 1 \pm \sqrt{x^3 + 2x^2 + 2x + 4}$$

- Of which only $y = 1 - \sqrt{x^3 + 2x^2 + 2x + 4}$ satisfies the $y(0) = -1$ initial condition.


Example 2.2.2: Solution Domain

If initial condition is $y(0) = -1$, then we choose the positive sign, instead of negative sign, on the square root term:

$$y = 1 + \sqrt{x^3 + 2x^2 + 2x + 4}$$

The solution is valid over x -values for which the quantity under the radical is positive: $x > -2$

- domain shown in green in the graph
- vertical tangent at domain boundary point $(-2, 1)$



Example 2.2.3: Implicit Solution of an Initial Value Problem

- Consider the following initial value problem:

$$\frac{dy}{dx} = \frac{4x - x^3}{4 + y^3}, \quad y(0) = 1$$

- Separating variables and using calculus, we obtain

$$(4 + y^3) dy = (4x - x^3) dx$$

$$\int (4 + y^3) dy = \int (4x - x^3) dx$$

$$4y + \frac{1}{4}y^4 = 2x^2 - \frac{1}{4}x^4 + c$$

$$y^4 + 16y + x^4 - 8x^2 = C \quad \text{where } C = 4c$$

- Using the initial condition, $y(0) = 1$, it follows that $C = 17$.

$$y^4 + 16y + x^4 - 8x^2 = 17$$

Example 2.2.3: General Solution vs. Particular Solution Through Point (0,1)

Thus the general solution is

$$y^4 + 16y + x^4 - 8x^2 = c$$

and the solution through (0,1) is

$$y^4 + 16y + x^4 - 8x^2 = 17$$


Vertical asymptotes for this particular solution will be found where:

$$y = \sqrt[3]{-4} \approx -1.5874$$

$$x \approx \pm 3.3488$$

Example 2.2.3: Graph of Solutions

- The graph of the solution through $(0, 1)$ is shown in green
- The points identified with open dots correspond to the solution domain boundaries where the tangent lines are vertical.



Parametric Equations

- The differential equation: $\frac{dy}{dx} = \frac{F(x, y)}{G(x, y)}$

is sometimes easier to solve if x and y are thought of as dependent variables of the independent variable t and rewriting the single differential equation as the system of differential equations:

$$\frac{dy}{dt} = F(x, y) \text{ and } \frac{dx}{dt} = G(x, y)$$

Chapter 9 is devoted to the solution of systems such as these.

Copyright

Copyright © 2021 John Wiley & Sons, Inc.

All rights reserved. Reproduction or translation of this work beyond that permitted in Section 117 of the 1976 United States Act without the express written permission of the copyright owner is unlawful. Request for further information should be addressed to the Permissions Department, John Wiley & Sons, Inc. The purchaser may make back-up copies for his/her own use only and not for distribution or resale. The Publisher assumes no responsibility for errors, omissions, or damages, caused by the use of these programs or from the use of the information contained herein.