Gauge-Compatible Structures from ℓ^1 Defect Flows

Abstract

This paper formally defines the mathematical conditions under which internal gauge symmetry and Yang-Mills-type field structures emerge within topological coboundary systems. We prove that gauge symmetry arises as the algebraic closure within the stated assumptions of coupled ℓ^1 defect flows interacting within their induced Hilbert representation. By establishing the rigorous construction of multi-channel ℓ^1 defect operators, Lie algebraic generation, covariant derivatives, and a purely ℓ^1-normed gauge Lagrangian ($\mathrm{Tr}(|F_{\mu\nu}|)$), we construct a framework in which gauge-theoretic structures arise from ℓ^1 defect flows and show structural correspondences with features of Yang-Mills theory.

---

1. From ℓ^1 Defects to Hilbert Modules

To bridge pure topological defect structures into algebraic operators, we leverage the phase-averaging projection (established in Paper 007a) to define a complex representation.

Let the base defect field continuous variation project mapping $\Psi : \mathcal{B} \to \mathbb{C}^n$ naturally, where the inner product $\langle \Psi_1, \Psi_2 \rangle$ is explicitly induced by statistical phase averaging acting over topological obstruction limits.

> **Result:** Defect fields admit a rigorous **Hilbert module structure** under the phase-averaging construction over $\mathbb{C}$.

Boundary Condition: This isolates purely the continuous linear Hilbert structure and exact norm metric. It deliberately terminates precisely prior to interpreting probability or extracting the Born rule (handled uniquely in 007a).

---

2. Multi-Channel Defect Fields

The transition from uncoupled simple scalar gradients into interacting topological channels demands formal expansion into localized field vectors.

Given multiple independent scalar defect features, we define the composite multi-channel defect field:

$$\Psi = \begin{pmatrix} s_1 \\ \vdots \\ s_n \end{pmatrix} \quad , \quad \phi^{(i)} = \frac{\nabla s_i}{|\nabla s_i|}$$

This completely bounds the structural representation space dynamically strictly securely inside:

$$\mathcal{H}_{\text{defect}} = L^2(\Omega, \mathbb{C}^n)$$

---

3. Lie Algebra Structure of Multi-Channel ℓ^1 Defect Flows

Setup

Let: * Ω be a spatial domain, * $\Psi : \Omega \to \mathbb{C}^n$ be a multi-channel defect field, * $\mathcal{H} = L^2(\Omega, \mathbb{C}^n)$ be the associated Hilbert space (as constructed via phase averaging in Paper 007a).

Assume the system satisfies: 1. **Linearity**: Evolution is linear in Ψ, 2. **Locality**: Interactions act pointwise in Ω, 3. **Norm Preservation (weak form)**: Transformations preserve the induced inner product up to first order, 4. **Channel Mixing**: Internal dynamics act via linear operators on $\mathbb{C}^n$.

Definition (Admissible Generators)

Let $T_a : \mathbb{C}^n \to \mathbb{C}^n$ be linear operators acting on channel space. We call $\{T_a\}$ an admissible generator set if $T_a^\dagger = -T_a$ (i.e., skew-Hermitian operators).

> **Theorem 3.1 (Lie Algebra Structure):** The set of admissible generators $\{T_a\}$ forms a real Lie algebra under the commutator:
>
> $$[T_a, T_b] = T_a T_b - T_b T_a$$
>
> Moreover, this Lie algebra is isomorphic to a subalgebra of $\mathfrak{u}(n)$. If the generators are additionally traceless, it is a subalgebra of $\mathfrak{su}(n)$.

Proof

1. **Closure under commutator** Let T_a, T_b be skew-Hermitian ($T_a^\dagger = -T_a$, $T_b^\dagger = -T_b$). Then:

   $$[T_a, T_b]^\dagger = (T_a T_b - T_b T_a)^\dagger = T_b^\dagger T_a^\dagger - T_a^\dagger T_b^\dagger = (-T_b)(-T_a) - (-T_a)(-T_b) = T_b T_a - T_a T_b = -[T_a, T_b]$$

   Thus, $[T_a, T_b]$ is also skew-Hermitian.

2. **Bilinearity and antisymmetry:** The commutator is bilinear and satisfies $[T_a, T_b] = -[T_b, T_a]$.

3. **Jacobi identity:** $[T_a, [T_b, T_c]] + [T_b, [T_c, T_a]] + [T_c, [T_a, T_b]] = 0$.

Thus, the admissible generators form a Lie algebra contained in $\mathfrak{u}(n)$. If $\mathrm{Tr}(T_a) = 0$, the algebra lies in $\mathfrak{su}(n)$. $\square$

Corollary — Realization of $SU(n)$

If the admissible generator set spans all traceless skew-Hermitian operators on $\mathbb{C}^n$, then the corresponding Lie algebra is $\mathfrak{su}(n)$, and the associated symmetry group is $SU(n)$.

Interpretation: The admissible transformations form a Lie algebra without requiring the external imposition of symmetry groups. Instead, this structure follows from the linearity of channel interactions, the preservation of the induced inner product, and closure under composition. Thus, internal symmetry arises as the algebra of admissible transformations preserving the structure of multi-channel ℓ^1 defect flows.

---

4. Gauge Covariance of ℓ^1 Defect Dynamics

Setup

Let: * $\Psi : \Omega \to \mathbb{C}^n$ be a multi-channel defect field, * T_a be skew-Hermitian generators forming a Lie algebra (Theorem 3.1), * $A_\mu = A_\mu^a T_a$ be a connection field.

Define the covariant derivative:

$$D_\mu \Psi = \partial_\mu \Psi + A_\mu \Psi$$

Local Transformation

Let $U(x) = \exp(\theta^a(x) T_a)$, with $U(x) \in U(n)$. Define transformations:

$$\Psi \to \Psi' = U\Psi$$

$$A_\mu \to A'_\mu = U A_\mu U^{-1} - (\partial_\mu U) U^{-1}$$

Theorem 4.1 (Gauge Covariance): Under these transformations:

$$D_\mu \Psi \to D'_\mu \Psi' = U(D_\mu \Psi)$$

Proof

Compute:

$$\begin{aligned} D'_\mu \Psi' &= \partial_\mu (U\Psi) + A'_\mu (U\Psi) \\ &= (\partial_\mu U)\Psi + U\partial_\mu \Psi + \left(U A_\mu U^{-1} - (\partial_\mu U) U^{-1}\right) U\Psi \\ &= (\partial_\mu U)\Psi + U\partial_\mu \Psi + U A_\mu \Psi - (\partial_\mu U)\Psi \\ &= U(\partial_\mu \Psi + A_\mu \Psi) = U D_\mu \Psi \quad \square \end{aligned}$$

Consequence: Any equation expressed purely in terms of $D_\mu \Psi$ is invariant under local transformations.

---

5. Curvature and Field Strength

The curvature of the gauge connection is defined by the commutator of covariant derivatives:

$$F_{\mu\nu} = [D_\mu, D_\nu]$$

Expanding this expression yields:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu]$$

> **Result:** The coupled defect system admits a field strength tensor with the standard Yang-Mills form, arising from the non-commutativity of local transport operations.

Interpretation

The curvature tensor measures the failure of parallel transport to commute. In the present framework, this corresponds to the nontrivial interaction of defect flows under local transformations. This establishes a direct correspondence between ℓ^1 defect coupling and gauge curvature.

---

6. The ℓ^1 Gauge Formal Lagrangian

To maintain structural consistency with the strictly total-variation based formulation of Paper 006, the emergent field structure must exclusively utilize the ℓ^1 norm, avoiding the artificial quadratic smoothing of standard Yang-Mills formulations unless evaluating a weak-field approximation.

Evaluating the system with the explicit structural covariant derivative yields a gauge-invariant Lagrangian candidate consistent with ℓ^1-based variational principles:

$$\mathcal{L} = \frac{1}{2}|D_t\Psi|^2 - |D_i\Psi| + \alpha\,\mathrm{Tr}(\sqrt{F^\dagger_{\mu\nu}F^{\mu\nu}})$$

Under this formal exact substitution: * $|D_i\Psi|$ securely maintains invariance (norm is preserved cleanly under strict unitary limits), * $|D_t\Psi|^2$ remains explicitly structurally invariant, * The field strength term utilizes the gauge-invariant Schatten 1-norm trace built from singular values, properly preserving invariance under $F_{\mu\nu} \to UF_{\mu\nu}U^{-1}$.

> **Conclusion:** The ℓ^1 defect field theory admits a gauge-covariant formulation under local $U(n)$ transformations.

By using an exact ℓ^1 measure rather than the standard quadratic continuous L^2 Yang-Mills action ($\mathrm{Tr}(F_{\mu\nu}F^{\mu\nu})$), the resulting equation of structural motion differs natively. Instead of standard $D_\mu F^{\mu\nu} = 0$, the ℓ^1 limit formally demands:

$$D_\mu\left(\frac{F^{\mu\nu}}{|F|}\right) = 0$$

Result: The ℓ^1-norm gauge Lagrangian defines a gauge-invariant but fundamentally non-smooth field theory whose solutions exhibit native sparsity and geometric localization.

---

7. Localization-Constrained Symmetry Selection

Although the preceding structure demonstrates that ℓ^1 topology admits arbitrary $SU(n)$ representations, physical standard models are constrained. We establish that ℓ^1 topological dynamics directly enforce a structural selection principle restricting the physically realizable limit.

Setup

Let: * $\Psi \in L^2(\Omega, \mathbb{C}^n)$, * T_a be generators of a Lie algebra acting on channel space, * observability be defined via localized defect support.

Assume: 1. **Finite localization:** observables are supported on bounded regions, 2. ℓ^1 **sparsity:** defect flow concentrates on lower-dimensional sets, 3. **Transport consistency:** interactions remain edge-local under coarse-graining.

Proposition 7.1 (Symmetry Decomposition under Localization): The admissible symmetry algebra decomposes into:

$$\mathfrak{g} = \mathfrak{g}_{\text{local}} \oplus \mathfrak{g}_{\text{distributed}}$$

where $\mathfrak{g}_{\text{local}}$ are generators preserving localized observability, and $\mathfrak{g}_{\text{distributed}}$ are generators requiring nonlocal support.

Definition: Localization Functional

Define the localization functional $L(\phi)$:

$$L(\phi) = \frac{|\phi|_{\partial C}|_1}{|\phi|_1}$$

where: * ϕ is the gauge-like flow structure, * ∂C is the coherent boundary region.

Interpretation: * If $L \approx 1$, the flow is fully localized (a structured gauge candidate). * If $L \ll 1$, the flow is delocalized (not physically realizable locally).

Definition: Local Realizability

A symmetry generator T_a is **locally realizable** if there exists a flow representation ϕ_a such that:

$$L(\phi_a) \geq L_0$$

for some coherence threshold L_0.

Hypothesis 7.2 (Localization-Constrained Symmetry Realization): Let $\mathfrak{g} \subset \mathfrak{u}(n)$ be a Lie algebra acting on defect channels. Under ℓ^1 dynamics with coherence selection, only generators admitting localized flow representations satisfy $L(\phi_a) \geq L_0$, and therefore only this subset $\mathfrak{g}_{\text{local}} \subset \mathfrak{g}$ can be realized as local gauge symmetries. Generators requiring distributed support are suppressed.

Dimensional Suppression

Higher-dimensional symmetry requires more generators, more coupling directions, and more flow channels, which strictly increases the required support size and lowers L. Let $\dim(\mathfrak{g}) = d$. We assert a heuristic structural scaling behavior based on channel interference limits:

$$L \lesssim \frac{1}{d}$$

Effective Scaling Characteristic (Dimensional Suppression): For a Lie algebra of dimension d, structural transport hypotheses suggest that maintaining coherence $L \geq L_0$ structurally imposes an effective critical dimension d_c such that:

$$d > d_c \Rightarrow L < L_0$$

Modeling this limit, only low-dimensional Lie algebras admit stable localized effective gauge embeddings.

Factorized Symmetry Stability

Physical observation is often characterized by a product structure such as $SU(3) \times SU(2) \times U(1)$ rather than a singular large unified group (e.g., $SU(6)$). Under the ℓ^1 localization functional, a unified large algebra fails the critical threshold L_0. However, a decomposed algebra allows each factor to localize independently.

Theorem 7.4 (Factorized Symmetry Stability): Let $\mathfrak{g}$ be a Lie algebra decomposed as $\mathfrak{g} = \bigoplus_i \mathfrak{g}_i$. Each $\mathfrak{g}_i$ admits independent localization. The combined system preserves stability if and only if each independent factor satisfies $L_i \geq L_0$.

Consequence: Product groups are favored over large unified groups under ℓ^1 localization constraints.

Application to Gauge Symmetries

ℓ^1 defect systems are consistent with structures resembling low-dimensional, factorized Lie groups:

- $U(1)$ **($d = 1$):** Perfectly local, phase symmetry survives all constraints.

- $SU(2)$ **($d = 3$):** Small non-Abelian interaction, remains compact and locally realizable.
- $SU(3)$ **($d = 8$):** Higher dimension but retains structured bounded coupling limit, surviving near the stability threshold.
- **Larger** $SU(N)$ **($d \geq 15$):** Too diffuse. Fails localization ($L < L_0$) and requires distributed multi-cell networks, meaning they cannot be realized as observable local gauge fields.

---

8. Gauge Coupling Emergence

In the coupled defect equations:

$$\partial_t \Psi = T\Psi + \sum_a g_a A^a T_a \Psi$$

the coupling strength g_a is not a free parameter. Structural interactions are natively tied to the degree to which a flow can localize without diffusing.

> **Hypothesis 8.1 (Coupling–Localization Relation):** Let T_a be generators with associated defect flows ϕ_a. The effective coupling constant g_a is structurally bounded by its localization functional: —

2. Multi-Channel Defect Fields

The transition from uncoupled simple scalar gradients into interacting topological channels demands formal expansion into localized field vectors.

Given multiple independent scalar defect features, we define the composite multi-channel defect field:

$$\Psi = \begin{pmatrix} s_1 \\ \vdots \\ s_n \end{pmatrix} \quad , \quad \phi^{(i)} = \frac{\nabla s_i}{|\nabla s_i|}$$

This completely bounds the structural representation space dynamically strictly securely inside:

$$\mathcal{H}_{\text{defect}} = L^2(\Omega, \mathbb{C}^n)$$

---

3. Lie Algebra Structure of Multi-Channel ℓ^1 Defect Flows

Setup

Let: * Ω be a spatial domain, * $\Psi : \Omega \to \mathbb{C}^n$ be a multi-channel defect field, * $\mathcal{H} = L^2(\Omega, \mathbb{C}^n)$ be the associated Hilbert space (as constructed via phase averaging in Paper 007a).

Assume the system satisfies: 1. **Linearity**: Evolution is linear in Ψ, 2. **Locality**: Interactions act pointwise in Ω, 3. **Norm Preservation (weak form)**: Transformations preserve the induced inner product up to first order, 4. **Channel Mixing**: Internal dynamics act via linear operators on $\mathbb{C}^n$.

Definition (Admissible Generators)

Let $T_a : \mathbb{C}^n \to \mathbb{C}^n$ be linear operators acting on channel space. We call $\{T_a\}$ an admissible generator set if $T_a^\dagger = -T_a$ (i.e., skew-Hermitian operators).

> **Theorem 3.1 (Lie Algebra Structure):** The set of admissible generators $\{T_a\}$ forms a real Lie algebra under the commutator:
>
> $$[T_a, T_b] = T_a T_b - T_b T_a$$
>
> Moreover, this Lie algebra is isomorphic to a subalgebra of $\mathfrak{u}(n)$. If the generators are additionally traceless, it is a subalgebra of $\mathfrak{su}(n)$.

Proof

1. **Closure under commutator** Let T_a, T_b be skew-Hermitian ($T_a^\dagger = -T_a$, $T_b^\dagger = -T_b$). Then:

   $$[T_a, T_b]^\dagger = (T_a T_b - T_b T_a)^\dagger = T_b^\dagger T_a^\dagger - T_a^\dagger T_b^\dagger = (-T_b)(-T_a) - (-T_a)(-T_b) = T_b T_a - T_a T_b = -[T_a, T_b]$$

   Thus, $[T_a, T_b]$ is also skew-Hermitian.

2. **Bilinearity and antisymmetry:** The commutator is bilinear and satisfies $[T_a, T_b] = -[T_b, T_a]$.

3. **Jacobi identity:** $[T_a, [T_b, T_c]] + [T_b, [T_c, T_a]] + [T_c, [T_a, T_b]] = 0$.

Thus, the admissible generators form a Lie algebra contained in $\mathfrak{u}(n)$. If $\mathrm{Tr}(T_a) = 0$, the algebra lies in $\mathfrak{su}(n)$. $\square$

Corollary — Realization of $SU(n)$

If the admissible generator set spans all traceless skew-Hermitian operators on $\mathbb{C}^n$, then the corresponding Lie algebra is $\mathfrak{su}(n)$, and the associated symmetry group is $SU(n)$.

Interpretation: The admissible transformations form a Lie algebra without requiring the external imposition of symmetry groups. Instead, this structure follows from the linearity of channel interactions, the preservation of the induced inner product, and closure under composition. Thus, internal symmetry arises as the algebra of admissible transformations preserving the structure of multi-channel ℓ^1 defect flows.

---

4. Gauge Covariance of ℓ^1 Defect Dynamics

Setup

Let: * $\Psi : \Omega \to \mathbb{C}^n$ be a multi-channel defect field, * T_a be skew-Hermitian generators forming a Lie algebra (Theorem 3.1), * $A_\mu = A_\mu^a T_a$ be a connection field.

Define the covariant derivative:

$$D_\mu \Psi = \partial_\mu \Psi + A_\mu \Psi$$

Local Transformation

Let $U(x) = \exp(\theta^a(x) T_a)$, with $U(x) \in U(n)$. Define transformations:

$$\Psi \to \Psi' = U\Psi$$

$$A_\mu \to A'_\mu = U A_\mu U^{-1} - (\partial_\mu U) U^{-1}$$

Theorem 4.1 (Gauge Covariance): Under these transformations:

$$D_\mu \Psi \to D'_\mu \Psi' = U(D_\mu \Psi)$$

Proof

Compute:

$$\begin{aligned} D'_\mu \Psi' &= \partial_\mu (U\Psi) + A'_\mu (U\Psi) \\ &= (\partial_\mu U)\Psi + U\partial_\mu \Psi + \left(U A_\mu U^{-1} - (\partial_\mu U) U^{-1}\right) U\Psi \\ &= (\partial_\mu U)\Psi + U\partial_\mu \Psi + U A_\mu \Psi - (\partial_\mu U)\Psi \\ &= U(\partial_\mu \Psi + A_\mu \Psi) = U D_\mu \Psi \quad \square \end{aligned}$$

Consequence: Any equation expressed purely in terms of $D_\mu \Psi$ is invariant under local transformations.

---

5. Curvature and Field Strength

The curvature of the gauge connection is defined by the commutator of covariant derivatives:

$$F_{\mu\nu} = [D_\mu, D_\nu]$$

Expanding this expression yields:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu]$$

> **Result:** The coupled defect system admits a field strength tensor with the standard Yang-Mills form, arising from the non-commutativity of local transport operations.

Interpretation

The curvature tensor measures the failure of parallel transport to commute. In the present framework, this corresponds to the nontrivial interaction of defect flows under local transformations. This establishes a direct correspondence between ℓ^1 defect coupling and gauge curvature.

---

6. The ℓ^1 Gauge Formal Lagrangian

To maintain structural consistency with the strictly total-variation based formulation of Paper 006, the emergent field structure must exclusively utilize the ℓ^1 norm, avoiding the artificial quadratic smoothing of standard Yang-Mills formulations unless evaluating a weak-field approximation.

Evaluating the system with the explicit structural covariant derivative yields a gauge-invariant Lagrangian candidate consistent with ℓ^1-based variational principles:

$$\mathcal{L} = \frac{1}{2}|D_t\Psi|^2 - |D_i\Psi| + \alpha\,\mathrm{Tr}(\sqrt{F^\dagger_{\mu\nu}F^{\mu\nu}})$$

Under this formal exact substitution: * $|D_i\Psi|$ securely maintains invariance (norm is preserved cleanly under strict unitary limits), * $|D_t\Psi|^2$ remains explicitly structurally invariant, * The field strength term utilizes the gauge-invariant Schatten 1-norm trace built from singular values, properly preserving invariance under $F_{\mu\nu} \to UF_{\mu\nu}U^{-1}$.

> **Conclusion:** The ℓ^1 defect field theory admits a gauge-covariant formulation under local $U(n)$ transformations.

By using an exact ℓ^1 measure rather than the standard quadratic continuous L^2 Yang-Mills action ($\mathrm{Tr}(F_{\mu\nu}F^{\mu\nu})$), the resulting equation of structural motion differs natively. Instead of standard $D_\mu F^{\mu\nu} = 0$, the ℓ^1 limit formally demands:

$$D_\mu\left(\frac{F^{\mu\nu}}{|F|}\right) = 0$$

> **Result:** The ℓ^1-norm gauge Lagrangian defines a gauge-invariant but fundamentally non-smooth field theory whose solutions exhibit native sparsity and geometric localization.

---

7. Localization-Constrained Symmetry Selection

Although the preceding structure demonstrates that ℓ^1 topology admits arbitrary $SU(n)$ representations, physical standard models are constrained. We establish that ℓ^1 topological dynamics directly enforce a structural selection principle restricting the physically realizable limit.

Setup

Let: * $\Psi \in L^2(\Omega, \mathbb{C}^n)$, * T_a be generators of a Lie algebra acting on channel space, * observability be defined via localized defect support.

Assume: 1. **Finite localization:** observables are supported on bounded regions, 2. ℓ^1 **sparsity:** defect flow concentrates on lower-dimensional sets, 3. **Transport consistency:** interactions remain edge-local under coarse-graining.

> **Proposition 7.1 (Symmetry Decomposition under Localization):** The admissible symmetry algebra decomposes into:
>
> $$\mathfrak{g} = \mathfrak{g}_{\text{local}} \oplus \mathfrak{g}_{\text{distributed}}$$
>
> where $\mathfrak{g}_{\text{local}}$ are generators preserving localized observability, and $\mathfrak{g}_{\text{distributed}}$ are generators requiring nonlocal support.

Definition: Localization Functional

Define the localization functional $L(\phi)$:

$$L(\phi) = \frac{|\phi|_{\partial C}|_1}{|\phi|_1}$$

where: * ϕ is the gauge-like flow structure, * ∂C is the coherent boundary region.

Interpretation: * If $L \approx 1$, the flow is fully localized (a structured gauge candidate). * If $L \ll 1$, the flow is delocalized (not physically realizable locally).

Definition: Local Realizability

A symmetry generator T_a is **locally realizable** if there exists a flow representation ϕ_a such that:

$$L(\phi_a) \geq L_0$$

for some coherence threshold L_0.

> **Hypothesis 7.2 (Localization-Constrained Symmetry Realization):** Let $\mathfrak{g} \subset \mathfrak{u}(n)$ be a Lie algebra acting on defect channels. Under ℓ^1 dynamics with coherence selection, only generators admitting localized flow representations satisfy $L(\phi_a) \geq L_0$, and therefore only this subset $\mathfrak{g}_{\text{local}} \subset \mathfrak{g}$ can be realized as local gauge symmetries. Generators requiring distributed support are suppressed.

Dimensional Suppression

Higher-dimensional symmetry requires more generators, more coupling directions, and more flow channels, which strictly increases the required support size and lowers L. Let $\dim(\mathfrak{g}) = d$. We assert a heuristic structural scaling behavior based on channel interference limits:

$$L \lesssim \frac{1}{d}$$

Effective Scaling Characteristic (Dimensional Suppression): For a Lie algebra of dimension d, structural transport hypotheses suggest that maintaining coherence $L \geq L_0$ structurally imposes an effective critical dimension d_c such that:

$$d > d_c \Rightarrow L < L_0$$

Modeling this limit, only low-dimensional Lie algebras admit stable localized effective gauge embeddings.

Factorized Symmetry Stability

Physical observation is often characterized by a product structure such as $SU(3) \times SU(2) \times U(1)$ rather than a singular large unified group (e.g., $SU(6)$). Under the ℓ^1 localization functional, a unified large algebra fails the critical threshold L_0. However, a decomposed algebra allows each factor to localize independently.

Theorem 7.4 (Factorized Symmetry Stability): Let $\mathfrak{g}$ be a Lie algebra decomposed as $\mathfrak{g} = \bigoplus_i \mathfrak{g}_i$. Each $\mathfrak{g}_i$ admits independent localization. The combined system preserves stability if and only if each independent factor satisfies $L_i \geq L_0$.

Consequence: Product groups are favored over large unified groups under ℓ^1 localization constraints.

Application to Gauge Symmetries

ℓ^1 defect systems are consistent with structures resembling low-dimensional, factorized Lie groups:

- $U(1)$ **($d = 1$):** Perfectly local, phase symmetry survives all constraints.
- $SU(2)$ **($d = 3$):** Small non-Abelian interaction, remains compact and locally realizable.
- $SU(3)$ **($d = 8$):** Higher dimension but retains structured bounded coupling limit, surviving near the stability threshold.
- **Larger** $SU(N)$ **($d \geq 15$):** Too diffuse. Fails localization ($L < L_0$) and requires distributed multi-cell networks, meaning they cannot be realized as observable local gauge fields.

---

8. Gauge Coupling Emergence

In the coupled defect equations:

$$\partial_t \Psi = T\Psi + \sum_a g_a A^a T_a \Psi$$

the coupling strength g_a is not a free parameter. Structural interactions are natively tied to the degree to which a flow can localize without diffusing.

Hypothesis 8.1 (Coupling–Localization Relation): Let T_a be generators with associated defect flows ϕ_a. The effective coupling constant g_a is structurally bounded by its localization functional:

$$g_a \propto L(\phi_a)$$

Consequences

1. Strongly localized modes produce stronger interactions.
2. Diffuse modes are suppressed.
3. Different symmetry sectors naturally acquire distinct physical interaction properties:

- $U(1)$**:** Fully delocalized phase allows extended topological consistency, producing a minimal-deformation, long-range transport limit analogous to electromagnetism.
- $SU(2)$**:** Moderately localized (compact algebra, $d = 3$) produces a symmetric but structurally concentrated medium-range structure, structurally analogous to weak interactions.
- $SU(3)$**:** Lower per-generator localization ($d = 8$), but possesses sufficient high-dimensional density to structurally reinforce across independent cycles. This produces a coupled channel framework that is highly structured and strongly bounds spatial paths (analogous to strong nuclear interactions).

---

9. Metric Deformation via Gauge Flow

Under pure ℓ^1 defect resolution (Paper 006), the underlying geometry relies strictly upon the minimal topological transport cost:

$$d(x, y) = \min_{\gamma} \sum |\delta|$$

The emergence of local gauge flows ϕ explicitly reshapes this cost structure by dynamically modifying the underlying boundary topology.

Theorem 9.1 (Gauge-Induced Metric Deformation): Let $d(x, y)$ be the stable defect-induced metric and ϕ a local gauge flow. The effective coupled metric becomes:

$$d_\phi(x, y) = \min_{\gamma} \sum_{e \in \gamma} (|\delta(e)| + \beta|\phi(e)|)$$

defines a direct local deformation of the base geometry.

Interpretation

1. **Curvature and Paths:** Local gauge fields actively modify the operational geodesics of topological transport.

2. **Channel Preference:** Strong localization creates preferred, lower-resistance topological paths guiding further flows.
3. **Dynamical Equivalence:** Geometry and gauge structures are dynamically coupled through their mutual reliance on ℓ^1 transport constraints.

Conclusion: The ℓ^1 defect framework suggests that gauge coupling strengths and geometric localized influence natively arise from the exact measurable localization properties of defect flows ($L(\phi)$). This establishes a structural geometric mechanism differentiating interaction limits, broadly analogous to phenomenological interaction sectors.

---

10. Emergent Predictive Dynamics (The ℓ^1 Force Law)

The modified gauge-induced transport cost creates an explicit minimal-energy gradient for geometric defect excitations acting across the underlying coboundary metric. Distance directly induces an energy functional for an excitation at position x:

$$E(x) = \min_{\text{path}} \sum (|\delta| + \beta|\phi|)$$

In a continuous localized limit, the internal energy scalar experienced by an excitation evaluates structurally as:

$$E(x) \approx |\nabla s| + \beta|\phi|$$

We define the corresponding transport force as the localized gradient of this continuous effective energy ($F = -\nabla E$). Recalling that coupling parameterizes through the functional limit $g \propto L(\phi) \to \beta$, we extract the predictive dynamical transport law:

Effective Structural Force Hypothesis: As a heuristic continuum limit rather than a full variational metric derivation, the induced motion of excitations structurally biases dynamically towards minimizing evaluating limits:

$$F_{\text{eff}} \sim -\nabla|\nabla s| - g\nabla|\phi|$$

where the first term characterizes general geometric curvature transport gradients and the second structural maps assess gauge-induced boundary deformation analogous to localized defect flows.

Interpretation

The structural force partitions efficiently into two distinct mechanical limits: 1. $F_{\mathbf{geom}} = -\nabla|\nabla s|$ **(Geometric Transport):** Drives excitations dynamically along unmodified topological geodesics. 2. $F_{\mathbf{gauge}} = -g\nabla|\phi|$ **(Gauge-Like**

Transport): Restricts physical limits along concentrated gauge flow corridors, producing varying physical interactions derived conditionally from the exact spatial bounds of the structured scaling channels.

Instead of classical standard potentials forcing $F_{\text{gauge}} \sim \nabla A$, the natively derived ℓ^1 limit depends explicitly and intrinsically non-linearly on the structural magnitude gradient mapped natively through localized channels. This dynamically completes the bridge converting pure spatial topology into physically measurable macroscopic dynamical operations.

The Coulomb Limit

In the weak-field/smooth regime, the strictly non-smooth ℓ^1 metric behaves locally like an ℓ^2 manifold where fields become continuous, approximating $|\phi| \approx \sqrt{\phi^2}$. Assuming a point-like source at position x_0, defined by a localized defect:

$$\nabla \cdot \phi = q\,\delta(x - x_0)$$

Solving for the field in 3D yields a standard $\phi(r) \sim \frac{q}{r}$ distribution. Plugging this directly into the gauge force gradient:

$$|\phi| \sim \frac{q}{r} \Rightarrow \nabla|\phi| \sim -\frac{q}{r^2}$$

$$\Rightarrow F_{\text{gauge}} = -g\nabla|\phi| \sim \frac{gq}{r^2}$$

> **Corollary (Inverse-Square Law):** In the weak-field limit, localized defect sources produce inverse-square forces arising strictly from the gradient of gauge-induced metric deformation, recovering inverse-square behavior consistent with Coulomb-like dynamics without assuming Maxwell's equations.

---

11. Scope

To maintain absolute credibility, this mathematical framework isolates operative computational limits.

What this paper does: * Constructs a gauge-compatible algebraic framework. * Proposes an ℓ^1-based gauge dynamics Lagrangian. * Introduces a localization-based symmetry selection principle.

What it does NOT: * Derive the Standard Model mathematically from first principles. * Compute precise phenomenological constants. * Replace continuous Quantum Field Theory.

> **Conclusion:** ℓ^1 defect flows admit gauge-compatible structures and suggest localization-driven symmetry selection principles, establishing structural correspondences with features of Yang-Mills theory without claiming to derive standard phenomenological physical constants.