Observer-relative control of cochain residuals

Jeremy H. Carroll

2026-10-23

Version 2.1.3

Abstract

We study finite candidate-selection policies constrained by several coordinate observers. Each observer specifies a projection, positive weights, a space of removable differences and a closure map. The policy bounds the observer-relative quotient residual, the closure residual and their ratio. We distinguish two coordinate estimates: a weighted norm bounds the coordinates of its input, whereas a quotient seminorm bounds coordinates only of a selected minimizing residual. Complete coordinate coverage does not remove this distinction. A one-edge example has zero quotient residual and arbitrarily large raw coordinates, even with complete observation. We give the corrected estimates, counterexamples to stronger transfer claims and a finite ranking rule. The supplied Lean development treats integer weighted norms; a numerical reference implementation additionally evaluates projected quotient problems.

1. Introduction

Several observation maps can impose simultaneous constraints on a proposed repair. Such constraints are relative to the declared maps: they do not establish stability under all possible measurements or transformations. The mathematical question is what these inequalities actually control.

Finite-dimensional quotient norms and linear-program duality are standard [1]. The cochain condition used for closure residuals is standard as well [2]. Our purpose is to specify a policy and identify the distinction between coordinate coverage, residual bounds and bounds on the original field. In particular, a bound on distance to a subspace is not a bound on the vector itself.

2. Observer data and diagnostics

Let $\omega \in \mathbb{R}^m$ and let $\mathcal{O}$ be a finite nonempty family of observers. Observer o selects a coordinate set I_o, with coordinate restriction π_o, positive weights $w_{o,k}$ on those coordinates, a linear map B_o into the observed coordinate space, and a linear closure map D_o. Assume $D_o B_o = 0$. Choose positive face weights $v_{o,j}$ when that space is nonzero.

Define

$$\Phi_o(\omega) = \min_a \sum_{k \in I_o} w_{o,k} |(\pi_o \omega - B_o a)_k|, \qquad \Gamma_o(\omega) = \sum_j v_{o,j} |(D_o \pi_o \omega)_j|.$$

The first minimum is attained because the image of B_o is a closed subspace of a finite-dimensional normed space. Write $r_o(\omega) = \pi_o \omega - B_o a_o^*$ for a selected minimizing residual. The closure residual is unchanged by subtracting $B_o a_o^*$.

The family has complete coordinate coverage if $\bigcup_o I_o = \{1, \dots, m\}$. This is a condition on the coordinate restrictions. For arbitrary linear projections, a positivity statement about coordinate weights is not a substitute for checking the common kernel of the observation maps.

3. A finite-observer policy

Fix nonnegative tolerances ϵ_o, positive ceilings M and R, and nonnegative closure limits G and δ. For a candidate ω' require, for every observer,

$$\Phi_o(\omega') \le \Phi_o(\omega) + \epsilon_o, \qquad \Phi_o(\omega') \le M,$$

$$\Gamma_o(\omega') \le \Gamma_o(\omega) + \delta, \qquad \Gamma_o(\omega') \le G.$$

We also specify a resolution-dependent ratio

$$R_o(\omega') = \begin{cases} \Gamma_o(\omega')/\Phi_o(\omega'), & \Phi_o(\omega') > \epsilon_o, \\ 0, & \Phi_o(\omega') \le \epsilon_o \text{ and } \Gamma_o(\omega') \le \epsilon_o, \\ +\infty, & \Phi_o(\omega') \le \epsilon_o < \Gamma_o(\omega'), \end{cases}$$

and require $R_o(\omega') \le R$. This is a policy definition. Its middle branch can differ from the exact positive-denominator ratio: for $\Phi = 1/4000, \Gamma = 1/2000$ and $\epsilon = 1/1000$, it is zero although $\Gamma/\Phi = 2$. Thus this test alone does not establish an exact ratio ceiling.

After T transitions satisfying the first inequality,

$$\Phi_o(\omega_T) \le \Phi_o(\omega_0) + \sum_{t=0}^{T-1} \epsilon_{o,t}.$$

This follows by summing the step inequalities. A fixed positive tolerance gives a possible increase proportional to T, rather than a uniform infinite-time bound. A uniform bound requires the explicit ceiling or a bounded cumulative tolerance. Strict non-increase is obtained by setting each tolerance to zero.

4. Coordinate estimates and their limits

Proposition 4.1: minimizing residuals. If the quotient-residual comparison in Section 3 holds, then for each observer and each of its observed coordinates,

$$w_{o,k}|r_o(\omega')_k| \le \Phi_o(\omega') \le \Phi_o(\omega) + \epsilon_o.$$

Proof. The first term is one nonnegative summand in the weighted norm of the minimizing residual. The second inequality is the assumed policy comparison.

The conclusion concerns $r_o(\omega')$, not $\pi_o\omega'$. Different observers can select different minimizing residuals. Coordinate coverage alone does not turn these observer-relative residuals into a single raw-coordinate bound.

Counterexample 4.2: complete coverage with unbounded raw coordinates. Take one directed edge with $B = (-1, 1)$, identity coordinate observation and weight one. Every scalar field $\omega_t = (t)$ is $B(0, t)^T$, so

$$\Phi(\omega_t) = 0, \qquad \Gamma(\omega_t) = 0$$

when there are no faces. The observer covers the only coordinate. A transition from zero to ω_t passes the residual comparisons and zero-ratio convention for every t, while $|\omega_t| = |t|$ is arbitrarily large. The statement $|\omega'_k| \le \Phi_o(\omega) + \epsilon_o$ is therefore false for quotient residuals, even with complete coordinate coverage.

There is a valid raw-coordinate theorem for weighted norms. Define

$$N_o(c) = \sum_{k \in I_o} w_{o,k}|c_k|.$$

Proposition 4.3: weighted input norms. Suppose coverage is complete and $N_o(c') \le N_o(c) + \epsilon_o$ for every observer. For every coordinate k choose an observer with $k \in I_o$. Then

$$w_{o,k}|c'_k| \le N_o(c) + \epsilon_o.$$

Proof. The coordinate term is at most the full weighted input norm, which is bounded by the assumed comparison. If $w_{o,k} \ge 1$, the same budget bounds $|c'_k|$. More generally divide by the specified positive weight.

This applies when no quotient minimization is performed, or when the supplied vectors are explicitly the residuals to which the weighted norms apply. It does not justify replacing N_o by a quotient seminorm on arbitrary raw fields.

5. Transfer examples and candidate ranking

Complete coverage does not prohibit transfer between observed components. On two disjoint cycle coordinates with weights $(2, 1)$, changing $(1, 0)$ to $(0, 1)$ reduces the weighted norm from two to one. Both coordinates are observed, yet the defect changes location. To forbid that transfer, a policy must include appropriate componentwise or partition constraints.

Coverage also matters for input norms. Observers that see only the first two of three coordinates assign zero observed norm to $(0, 0, K)$ for every K. No bound on the third coordinate follows from those observations.

For a finite set of candidates that passed the specified tests, one possible ranking score is

$$L(o) = \max_{a \in \mathcal{O}} \Phi_a(\omega'_o) + \frac{\lambda}{|\mathcal{O}|} \sum_{a \in \mathcal{O}} (S_{a,o} + R_a(\omega'_o)) + c_o,$$

where $\lambda \geq 0$, $c_o \geq 0$ is a supplied cost, and $S_{a,o} \geq 0$ is a supplied constraint penalty. An implementation without a clause evaluator may use $S_{a,o} = \min(\Phi_a(\omega'_o), 1)$ as a surrogate; that surrogate is not a count of violated clauses. Select a least-score candidate. An empty surviving set produces no selection. These choices specify one algorithm, not an optimal controller over an unspecified future trajectory.

6. Numerical and formal evidence

code/mod_observer_robust_control_core.py implements finite-observer comparisons and a floating-point linear program when a matrix B_o is supplied. This linear program is a numerical computation, not an exact rational solver. The numerical observer object checks matrix shapes but does not independently certify $D_o B_o = 0$; that identity is a hypothesis of the cochain interpretation and must be checked separately for supplied matrices. The numerical controller also permits its configured comparison margin on ratio and absolute ceilings. To interpret those checks mathematically, use the corresponding inflated ceilings or set the comparison margin to zero; the observer's residual tolerance remains a separate parameter.

lean4/ObserverRobustControlCore.lean and lean4/JointDetectorAdequacy.lean verify statements about integer vectors, natural-number weights and weighted input norms. Their coordinate estimate instantiates Proposition 4.3, not a raw-coordinate theorem for a quotient linear program. The zero-weight entries in their finite examples represent omitted coordinates. The supplied census records the actual declaration set and admitted kernel axioms.

The tests include the quotient/raw-coordinate counterexample and distinguish numerical LP observations from exact arithmetic identities. A successful finite test run does not establish statements about undeclared observers.

7. Discussion

The policy controls precisely the diagnostics it compares. Quotient residuals ignore removable components; raw weighted norms do not. Complete coordinate coverage helps obtain coordinate estimates for raw input norms, but it cannot bound arbitrary components removed by quotient minimization. Neither condition prevents all transfers, constraint deletion or divergence under an unbounded cumulative tolerance. Those properties require additional hypotheses and checks.

Publication information

Version DOI: 10.5281/zenodo.23604797. Concept DOI: 10.5281/zenodo.20652483. The public record identifies the October 23, 2026 edition as version 2.1.3.

References

1. S. Boyd and L. Vandenberghe, *Convex Optimization*, Cambridge University Press, 2004. Author-hosted book.
2. A. Hatcher, *Algebraic Topology*, Cambridge University Press, 2002, Section 3.1. Author-hosted book.