



CESAB
CENTRE DE SYNTHÈSE ET D'ANALYSE
SUR LA BIODIVERSITÉ

Can we trust the result?

Publication bias, robustness and sensitivity

Damien Beillouin
CIRAD, UPR HortSys

SOUTIENS INSTITUTIONNELS



Learning objectives

By the end of this session, you will be able to:

- 1 List the ways a set of studies can be a **biased sample** of the evidence.
- 2 Read a **contour-enhanced funnel plot** — and explain why asymmetry $\neq$ publication bias.
- 3 Test for small-study effects and time-lag bias **with dependent effect sizes**, and know which classic methods to avoid.
- 4 Run **sensitivity analyses** (leave-one-study-out, influence, study quality) and report robustness.

Are our 28 studies a fair sample?

A meta-analysis assumes the studies it contains are **representative** of all studies done.

But studies with **significant** results are more likely to be:

- **published** → *publication bias*
- published **quickly** → *time-lag bias*
- published **in English** → *language bias*
- published **more than once** → *multiple-publication bias*
- **cited** (and found) → *citation bias*

The consequence


The pooled effect can be **overestimated** — sometimes an effect appears where there is none.

The best cure is prevention

Search grey literature, theses, reports and non-English studies (Monday's sessions). The tests we see today can only **detect** and **probe** bias — never repair it.

Publication bias


The logic of the funnel plot



Large studies (top) cluster near the true effect; small ones (bottom) scatter **symmetrically** → a funnel.

Where would the missing studies be?
Small **non-significant** studies (between the dotted lines) stay in drawers → a corner is **missing**.

Our funnel, with significance contours



- Shaded bands: significance of each comparison (grey background: $p < 0.01$)
- Dashed line: the multilevel mean
- Top: 83 comparisons from 2 studies with $SE < 0.02$ — suspiciously precise: check the extraction

If missing studies were in the **white** zone (non-significant), publication bias is a plausible cause. If they would sit in significant zones, look for other causes (Peters et al. 2008).

Asymmetry $\neq$ publication bias

YOUR CALL

Our funnel is asymmetric. List other reasons than publication bias.

- **Heterogeneity:** small studies differ (other crops, pests, methods) — true small-study effects.
- **Artefacts:** the variance of $\ln RR$ depends on the mean; extraction errors; pseudo-replication.
- **Chance:** with few studies, funnels are often asymmetric.

Testing for small-study effects: the classic way

Egger's test (Egger et al. 1997): regress each effect size on its standard error. A slope $\neq 0$ means small studies give different results. Our pests: $p = 0.00037$.

Two problems

1. It treats our 382 comparisons as **independent** — they come from 28 studies.
2. For lnRR, the SE is computed **from the means themselves** → an artificial link between effect and SE, even without any bias.

Testing for small-study effects: the multilevel way

Add a precision measure as a **moderator** of the multilevel model (Nakagawa et al. 2022), based on **sample size only**:

$$\sqrt{1/n_T + 1/n_C}$$

```
1 rma.mv(yi, vi,  
2       mods = ~ sqrt(1/n_t + 1/n_c),  
3       random = ~ 1 | study_id / es_id,  
4       test = "t", dfs = "contain",  
5       data = dat)
```

Precision measure	Slope p
SE (artefact-prone)	0.018
$\sqrt{1/n_T + 1/n_C}$	0.10
same, cluster-robust (CR2)	0.18

With the recommended measure: **no robust evidence** of a small-study effect.

If there were bias, would the conclusion change?


Estimate	Pest abundance [95% CI]
Main model	-27% [-40; -11]
PET : effect of an infinitely large study	-43% [-61; -18]
PEESE	-36% [-50; -16]

Rule (Stanley and Doucouliagos 2014): if the PET intercept is significant (here yes), report **PEESE**.

Adjusted estimates are, if anything, **stronger**: the pest reduction is **not** an artefact of publication bias.

With strong heterogeneity, PET-PEESE tends to **over-correct**: treat it as a sensitivity check, not as “the true effect”. Alternative: **selection models** [Vevea and Hedges (1995); `selmodel()` in metafor], which model the probability of publication given the p-value.

Time-lag bias



Do early studies report stronger effects than later ones?

Year as a moderator in the multilevel model: slope = -0.0029 per year ($p = 0.87$).

No evidence of a decline effect — with only 28 studies, the test has little power.

Also possible: include year and SE together (Nakagawa et al. 2022).

Two classics to avoid (or to use with care)

Fail-safe N (Rosenthal)

“How many null studies would make the result non-significant?”

Our data: **4,098,124** studies.

Meaningless: assumes missing studies have zero effect, ignores heterogeneity and dependence, estimates no bias. **Do not report it.**

Trim-and-fill


Imputes “missing” studies to make the funnel symmetric.

Assumes asymmetry = publication bias, performs badly with heterogeneity, ignores dependence.

→ at most a **sensitivity analysis** on study-level data, never “the corrected effect”.

Robustness

Does one study drive the result?



Leave-one-study-out: refit the model 28 times, dropping one whole study each time.

- Mean ranges from **-29%** to **-23%**
- The CI never crosses 0

Cook's distance by study: largest = 0.32, no study stands out from the others.

The conclusion does not depend on any single study.

A robustness table for your paper

Analysis	Pest abundance	Conclusion changes?
Main multilevel model (28 studies)	-27% [-40; -11]	—
Only comparisons meeting all validity criteria† (13 studies; 130 comparisons not appraisable)	-27% [-39; -13]; validity as moderator $p = 0.74$	No
Leave-one-study-out (range of means)	-29% to -23%	No
Small-study effect (sample-size based)	slope $p = 0.10$; PEESE -36%	No
Time-lag (year as moderator)	slope ≈ 0 ($p = 0.87$)	No (low power)

† Critical appraisal coded by Jones et al. (2021): enough sampling units in both groups, plots in their state long enough, intervention and comparator < 1.1 km apart. This is **internal validity** of each study (risk of bias, cf. CEE critical appraisal) — a different question from publication bias.

Also worth testing: another effect-size metric, the imputation of missing SDs, the correlation ρ assumed within studies. Report **all** sensitivity analyses — including those that change the result.

Practical · 33 min + debrief

Time	Task
15:22	Contour-enhanced funnel plot (code given: interpret)
15:27	Classic vs multilevel Egger (the lnRR trap); PET-PEESE
15:37	Sensitivity : leave-one-study-out, validity
15:47	<i>Your turn</i> : natural enemies – or the optional parts (time-lag, Cook, fail-safe N)
15:55	Debrief (all together)

Open **meta_analysis_course.Rproj** (the course kit), then **td_bias.R**. Instructions and folded solutions: the practical web page.

KEY MESSAGES

- 1 | Bias is best **prevented** by the search, not corrected by statistics.
- 2 | **Asymmetry $\neq$ publication bias**: read the direction, use contour-enhanced funnels.
- 3 | With several effect sizes per study, use **multilevel** tests — for lnRR with a **sample-size-based** precision; drop fail-safe N.
- 4 | Report a **robustness table**: leave-one-study-out, influence, study quality, analytical choices.

In the online notebook

Each point of this session developed further, with the same data, more pitfalls and exercises with solutions:

- Publication bias and sensitivity
- Data extraction and critical appraisal
- Reading a meta-analysis critically

literaturesynthesis.github.io/notebook

References

- Egger, M., G. Davey Smith, M. Schneider, and C. Minder. 1997. [Bias in meta-analysis detected by a simple, graphical test](#). *BMJ* 315:629–634.
- Jones, S. K., A. C. Sánchez, S. D. Juventia, and N. Estrada-Carmona. 2021. [A global database of diversified farming effects on biodiversity and yield](#). *Scientific Data* 8:212.
- Koricheva, J., J. Gurevitch, and K. Mengersen, editors. 2013. *Handbook of meta-analysis in ecology and evolution*. Princeton University Press, Princeton.
- Nakagawa, S., M. Lagisz, M. D. Jennions, J. Koricheva, D. W. A. Noble, T. H. Parker, A. Sánchez-Tójar, Y. Yang, and R. E. O'Dea. 2022. [Methods for testing publication bias in ecological and evolutionary meta-analyses](#). *Methods in Ecology and Evolution* 13:4–21.
- Ouédraogo, D.-Y. 2024. Risks of bias in meta-analyses. Lecture, FRB-CESAB training course on meta-analyses and systematic reviews.
- Peters, J. L., A. J. Sutton, D. R. Jones, K. R. Abrams, and L. Rushton. 2008. [Contour-enhanced meta-analysis funnel plots help distinguish publication bias from other causes of asymmetry](#). *Journal of Clinical Epidemiology* 61:991–996.
- Stanley, T. D., and H. Doucouliagos. 2014. [Meta-regression approximations to reduce publication selection bias](#). *Research Synthesis Methods* 5:60–78.
- Vevea, J. L., and L. V. Hedges. 1995. [A general linear model for estimating effect size in the presence of publication bias](#). *Psychometrika* 60:419–435.
- Viechtbauer, W., and M. W.-L. Cheung. 2010. [Outlier and influence diagnostics for meta-analysis](#). *Research Synthesis Methods* 1:112–125.